



COMPOSITIO MATHEMATICA

Cluster categories for marked surfaces: punctured case

Yu Qiu and Yu Zhou

Compositio Math. **153** (2017), 1779–1819.

[doi:10.1112/S0010437X17007229](https://doi.org/10.1112/S0010437X17007229)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865



Cluster categories for marked surfaces: punctured case

Yu Qiu and Yu Zhou

Dedicated to Idun Reiten on the occasion of her seventy-fifth birthday

ABSTRACT

We study cluster categories arising from marked surfaces (with punctures and non-empty boundaries). By constructing skewed-gentle algebras, we show that there is a bijection between tagged curves and string objects. Applications include interpreting dimensions of Ext^1 as intersection numbers of tagged curves and Auslander–Reiten translation as tagged rotation. An important consequence is that the cluster(-tilting) exchange graphs of such cluster categories are connected.

1. Introduction

1.1 Overall

Cluster algebras were introduced by Fomin and Zelevinsky [FZ02] around 2000, with quiver mutation as the combinatorial aspect. Derksen *et al.* [DWZ08] further developed quiver mutation to mutation of quivers with potential. During the last decade, the cluster phenomenon was spotted in various areas in mathematics, as well as in physics, including geometric topology and representation theory. On one hand, the geometric aspect of cluster theory was explored by Fomin *et al.* [FST08] after the pioneering work of Fock and Goncharov [FG06, FG09]. They constructed a quiver $Q_{\mathbf{T}}$ (and later, Labardini-Fragoso [Lab09a, IL12] gave a corresponding potential $W_{\mathbf{T}}$) from any (tagged) triangulation \mathbf{T} of a marked surface \mathbf{S} . Moreover, they showed that mutation of quivers (with potential) is compatible with flip of triangulations. There are a lot of known results about cluster algebras in the surface case:

- Felikson *et al.* [FST12] classified cluster algebras of finite mutation type, and established that they are all from marked surfaces except for a few cases;
- Musiker *et al.* [MSW13] constructed two canonical bases by two types of collections of curves;
- Musiker *et al.* [MSW11], Musiker and Williams [MW13], and Canakci and Schiffler [CS13, CS15a, CS15b] gave combinatorial formulae for cluster variables and relations;
- Mills [Mil16] showed that there exist maximal green sequences for quivers with potential associated with triangulated marked surfaces except for once-punctured closed surfaces (cf. [ACCERV13]).

Received 24 May 2015, accepted in final form 5 January 2017, published online 15 June 2017.

2010 Mathematics Subject Classification 16G20, 18E30, 16E30, 57M50 (primary), 16G70 (secondary).

Keywords: cluster categories, intersection numbers, cluster exchange graphs, skewed-gentle algebras.

This work is supported by the Research Council of Norway, FRINAT grant number 231000.

This journal is © Foundation Compositio Mathematica 2017.

On the other hand, the categorification of cluster algebras leads to representations of quivers, due to Buan *et al.* [BMRRT06]. Later, Amiot [Ami09] introduced generalized cluster categories via Ginzburg dg algebras associated with quivers with potential. Then there is an associated cluster category $\mathcal{C}(\mathbf{T})$ for each triangulation \mathbf{T} of \mathbf{S} .

Several works have been done concerning the cluster categories associated with triangulations of surfaces. Namely, for some special cases:

- Caldero *et al.* [CCS06] realized the cluster category of type A_n by a regular polygon with $n + 3$ vertices (i.e. a disk with $n + 3$ vertices on its boundary);
- Schiffler [Sch08] realized the cluster category of type D_n by a regular polygon with n vertices and one puncture in the center (i.e. a disk with n vertices on its boundary and one puncture in its interior).

In the unpunctured case:

- Assem *et al.* [ABCP12] proved that the Jacobian algebra of such a quiver with potential is a gentle algebra and gave a bijection between arcs that are not in the triangulation and string modules of the associated gentle algebras;
- Brüstle and Zhang [BZ11] generalized the bijection of [ABCP12] to a bijection between the set of curves and valued closed curves and the set of indecomposable objects in the associated cluster category. Under this bijection, they described irreducible morphisms, the Auslander–Reiten (AR) translation and AR-triangles in the cluster category by geometric terms in the surface. They also gave a bijection between triangulations of the surface and cluster tilting objects in the cluster category such that flip of an arc is compatible with mutation;
- based on Brüstle and Zhang’s work, Zhang *et al.* [ZZZ13] proved that the intersection number of two curves is equal to the dimension of Ext^1 of the corresponding objects and gave a geometric model of torsion pairs and their mutations;
- Canakci and Schroll [CS14] described a basis for Ext^1 in the cluster category and also computed a basis for Ext^1 in the module category of the corresponding Jacobian algebra by distinguishing different types of crossings between curves;
- Marsh and Palu [MP14] showed that Calabi–Yau reduction (introduced in [IY08]) can be interpreted as cutting along curves without self-intersections in the surface;
- the present authors [Qiu16, QZ14] also investigated other categories, which are used to define cluster categories, see the formula (3.1), and obtained similar structures/formulae.

For general cases:

- for each ideal triangulation without self-folded triangles, Labardini-Fragoso [Lab09b] associated a representation of the quiver with potential $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ to each curve without self-intersections and proved that mutation of representations is compatible with flip of triangulations;
- Brüstle and Qiu [BQ15] made an effort to understand a basic functor in the cluster category, i.e. the shift (or the AR-translation in this case), in terms of an element, the tagged rotation, in the tagged mapping class group of the marked surface. Their motivation lies in the study of the Seidel–Thomas braid group.

Notice that most of the above works deal only with the unpunctured case. This is because: (i) the usual flip does not work for self-folded triangles (cf. Figure 5) and (ii) the associated

quivers with potential are much more complicated in the punctured case and their Jacobian algebras are not gentle (cf. [IL12]). Fomin *et al.* solved (i) by introducing the notion of tagging.

In this paper, we aim to study the cluster categories associated to marked surfaces with punctures and with non-empty boundaries. The main tool is skewed-gentle algebras (a special kind of clannish algebras), which were developed in [Bon92, Cra89, Den00, Gei99, GdlP99]. The essential results are summarized as follows.

THEOREM 1.1 (Theorems 4.16, 4.19, 5.2, 5.5 and 5.7). *Let \mathbf{S} be a marked surface with punctures and with non-empty boundary. Given an admissible triangulation \mathbf{T} of \mathbf{S} (see Definition 3.7), let $\mathcal{C}(\mathbf{T})$ be the associated cluster category. Then there is a bijection*

$$\begin{aligned} X^{\mathbf{T}}: \mathbf{C}^{\times}(\mathbf{S}) &\rightarrow \mathfrak{S}(\mathbf{T}) \\ (\gamma, \kappa) &\mapsto X_{(\gamma, \kappa)}^{\mathbf{T}} \end{aligned}$$

from the set $\mathbf{C}^{\times}(\mathbf{S})$ of tagged curves in the surface \mathbf{S} to the set $\mathfrak{S}(\mathbf{T})$ of string objects in the category $\mathcal{C}(\mathbf{T})$ (see Definitions 3.1 and 4.6), satisfying the following.

- (i) For every admissible triangulation \mathbf{T}' of \mathbf{S} , there is an equivalence $\Theta: \mathcal{C}(\mathbf{T}) \simeq \mathcal{C}(\mathbf{T}')$, such that $\Theta \circ X^{\mathbf{T}} = X^{\mathbf{T}'}$.
- (ii) For any tagged curve $(\gamma, \kappa) \in \mathbf{C}^{\times}(\mathbf{S})$, we have $X_{\rho(\gamma, \kappa)}^{\mathbf{T}} \cong X_{(\gamma, \kappa)}^{\mathbf{T}}[1]$, where ρ is the tagged rotation (see Definition 3.2).
- (iii) For any two tagged curves $(\gamma_1, \kappa_1), (\gamma_2, \kappa_2)$ (not necessarily distinct), we have

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}}),$$

where Int denotes the intersection number (see Definition 3.3).

- (iv) The exchange graph $\text{CEG}(\mathcal{C}(\mathbf{T}))$ of cluster tilting objects in $\mathcal{C}(\mathbf{T})$ is isomorphic to the exchange graph $\text{EG}^{\times}(\mathbf{S})$ of tagged triangulations of \mathbf{S} and hence it is connected.

1.2 Context

In § 2 we recall notions and notation about skewed-gentle algebras that we will use throughout the paper. In § 3, we recall the background of cluster categories associated with triangulated marked surfaces. In § 4, we study the skewed-gentle algebra associated with an admissible triangulation and give a correspondence between tagged curves and string objects. The relation between such correspondences from different admissible triangulations is also studied. In § 5, we give homological interpretations of geometric objects from marked surfaces, namely tagged rotation, intersection numbers, and exchange graphs of tagged triangulations. An example is presented in § 6 to demonstrate some of the notions/results in the paper. The technical proof of the main theorem, Theorem 5.5, is given in § 7. In Appendix A we discuss some properties of admissible triangulations and in Appendix B we recall Derksen *et al.*'s mutation of decorated representations.

1.3 Conventions

Throughout this article, \mathbf{k} denotes an algebraically closed field. For any \mathbf{k} -algebra A , an A -module means a finitely generated left A -module and we denote by $\text{mod } A$ the category of all A -modules. For a finite set I , we denote by $|I|$ the number of elements in I . For an object X in a triangulated category \mathcal{C} , we denote:

- by $\text{add } X$ the full subcategory of \mathcal{C} consisting of direct summands of direct sums of copies of X ;
- by X^\perp the full subcategory of \mathcal{C} consisting of objects Y with $\text{Hom}_{\mathcal{C}}(X, Y) = 0$;
- by $\mathcal{C}/(X)$ the additive quotient category of \mathcal{C} by $\text{add } X$.

2. Preliminaries on skewed-gentle algebras

We recall from [Bon92, Cra89, Den00, Gei99, GdlP99] some notions, notation and results about skewed-gentle algebras used in this paper.

2.1 Skewed-gentle algebras

A *biquiver* is a tuple (Q_0, Q_1, Q_2) , where Q_0 is the set of vertices, Q_1 is the set of solid arrows, and Q_2 is the set of dashed arrows. Let $s, t : Q_1 \cup Q_2 \rightarrow Q_0$ be the start/terminal functions of arrows. We call an arrow α in $Q_1 \cup Q_2$ a loop if $s(\alpha) = t(\alpha)$.

In this paper, we always assume that a biquiver $Q = (Q_0, Q_1, Q_2)$ satisfies:

- each arrow in Q_2 is a dashed loop;
- there is at most one loop in Q_2 at each vertex;
- there is no loop in Q_1 .

Let Q_0^{Sp} be the subset of Q_0 consisting of vertices where there is a dashed loop in Q_2 .


Skewed-gentle algebras, modeled on gentle algebras, were introduced in [GdlP99] as a certain class of clannish algebras defined in [Cra89].

DEFINITION 2.1. A pair (Q, Z) of a biquiver Q and a set Z of compositions ab of arrows a, b in Q_1 is called skewed-gentle if the following conditions hold.

- For each vertex $p \in Q_0^{\text{Sp}}$, there is at most one arrow $\alpha \in Q_1$ ending at p and at most one arrow $\beta \in Q_1$ starting at p , with $\beta\alpha \in Z$ (if both exist).
- For each vertex $p \notin Q_0^{\text{Sp}}$, there are at most two arrows $\alpha_1, \alpha_2 \in Q_1$ ending at p and at most two arrows $\beta_1, \beta_2 \in Q_1$ starting at p , and they can be labeled in such a way that $\beta_1\alpha_1 \in Z$, $\beta_2\alpha_2 \in Z$, $\beta_1\alpha_2 \notin Z$, and $\beta_2\alpha_1 \notin Z$.

An algebra Λ is called a skewed-gentle algebra if Λ is Morita equivalent to $\mathbf{k}Q/(R)$ for a skewed-gentle pair (Q, Z) , where $R = Z \cup \{\varepsilon^2 - \varepsilon \mid \varepsilon \in Q_2\}$.

Example 2.2. Let Q be the following biquiver with $Z = \{ba, cb, ac\}$.




Then (Q, Z) is a skewed-gentle pair and hence $\mathbf{k}Q/(R)$ is a skewed-gentle algebra, where $R = Z \cup \{\varepsilon_1^2 - \varepsilon_1, \varepsilon_4^2 - \varepsilon_4\}$.

2.2 Letters

Let (Q, Z) be a skewed-gentle pair. Following [Cra89, Gei99], we associate a new biquiver $\widehat{Q} = (\widehat{Q}_0, \widehat{Q}_1, \widehat{Q}_2)$ with $Q = (Q_0, Q_1, Q_2)$ by adding two new vertices i_+ and i_- and two new solid arrows $a_{i_\pm} : i \rightarrow i_\pm$ for each vertex $i \in Q_0$. That is:

- $\widehat{Q}_0 = Q_0 \cup \{i_{\pm} \mid i \in Q_0\}$;
- $\widehat{Q}_1 = Q_1 \cup \{a_{i_{\pm}} : i \rightarrow i_{\pm} \mid i \in Q_0\}$;
- $\widehat{Q}_2 = Q_2$.

For example, the biquiver \widehat{Q} associated with the biquiver Q in Example 2.2 is as shown by the following diagram.



For any arrow α in \widehat{Q} , we define a *direct letter* α and an *inverse letter* α^{-1} , which are mutually inverse. Let L be the set of all letters. The functions s, t can be extended to L by setting $s(\alpha^{-1}) = t(\alpha)$ and $t(\alpha^{-1}) = s(\alpha)$. For each $i \in Q_0 \subset \widehat{Q}_0$, let $L(i) := \{l \in L \mid s(l) = i\}$. We divide $L(i)$ into two disjoint subsets $L_+(i)$ and $L_-(i)$ with linear orders such that the subset $L_{\theta}(i)$ has one of the following forms:

- $\{a_{i_{\theta}}\}$;
- $\{a_{i_{\theta}} > \alpha\}$;
- $\{\beta^{-1} > a_{i_{\theta}}\}$;
- $\{\beta^{-1} > a_{i_{\theta}} > \alpha\}$;
- $\{\varepsilon^{-1} > a_{i_{\theta}} > \varepsilon\}$,

for $\theta \in \{\pm\}$, some solid arrows α and β in $Q_1 \subset \widehat{Q}_1$ and some dashed arrow ε in $Q_2 = \widehat{Q}_2$, satisfying that:

- for any two solid arrows γ and δ in $Q_1 \subset \widehat{Q}_1$ with $t(\delta) = s(\gamma) = k$, we have that $\gamma\delta \in Z$ if and only if γ and δ^{-1} are both in $L_+(k)$ or $L_-(k)$.

Observe that in the set $L_{\theta}(i)$, the inverse of an arrow in Q is always greater than the arrow $a_{i_{\theta}}$, and an arrow in Q is always smaller than the arrow $a_{i_{\theta}}$. Notice that if $L(i) \neq \{a_{i_+}, a_{i_-}\}$, then there are exactly two possible choices for the pair $(L_+(i), L_-(i))$.

Example 2.3. In Example 2.2, one possible choice of the subsets $L_{\theta}(i)$ is:

- $L_+(1) = \{c^{-1} > a_{1_+} > a\}$ and $L_-(1) = \{\varepsilon_1^{-1} > a_{1_-} > \varepsilon_1\}$;
- $L_+(2) = \{a^{-1} > a_{2_+} > b\}$ and $L_-(2) = \{a_{2_-}\}$;
- $L_+(3) = \{b^{-1} > a_{3_+} > c\}$ and $L_-(3) = \{a_{3_-} > d\}$;
- $L_+(4) = \{d^{-1} > a_{4_+}\}$ and $L_-(4) = \{\varepsilon_4^{-1} > a_{4_-} > \varepsilon_4\}$.

Note that the inverse letters $a_{i_{\theta}}^{-1}$ are not listed here because they are not in any set $L_{\theta}(i)$. However they can still appear as first letters in words (see below).

2.3 Words

A word \mathbf{m} is a sequence $\omega_m \cdots \omega_2 \omega_1$ of letters in L satisfying that for any $1 \leq j \leq m - 1$, $\omega_j^{-1} \in L_\theta(i)$ and $\omega_{j+1} \in L_{\theta'}(i)$ for different $\theta, \theta' \in \{\pm\}$ and some $i \in Q_0 \subset \widehat{Q}_0$. We call ω_1 the first letter of \mathbf{m} and ω_m the last letter of \mathbf{m} . Since both $L_\theta(i)$ and $L_{\theta'}(i)$ are subsets of $L(i)$, we have $t(\omega_j) = s(\omega_{j+1}) = i$. The functions s, t can be generalized to the set of words by $s(\mathbf{m}) := s(\omega_1)$ and $t(\mathbf{m}) := t(\omega_m)$. The inverse of a word $\mathbf{m} = \omega_m \cdots \omega_2 \omega_1$ is defined as $\mathbf{m}^{-1} := \omega_1^{-1} \omega_2^{-1} \cdots \omega_m^{-1}$. The product \mathbf{nm} of two words $\mathbf{m} = \omega_m \cdots \omega_2 \omega_1$ and $\mathbf{n} = \omega_{m+r} \cdots \omega_{m+2} \omega_{m+1}$ is defined to be $\omega_{m+r} \cdots \omega_{m+2} \omega_{m+1} \omega_m \cdots \omega_2 \omega_1$ if this is again a word.

A letter is called *punctured* if it is of the form a_{i_θ} or $a_{i_\theta}^{-1}$ such that $L_\theta(i) = \{\varepsilon^{-1} > a_{i_\theta} > \varepsilon\}$ for some dashed arrow ε . A word \mathbf{m} is called *left inextensible* (respectively *right inextensible*) if there is no letter l such that $l\mathbf{m}$ (respectively $\mathbf{m}l$) is again a word. A word is called *maximal* if it is both left and right inextensible. It is obvious that a word $\mathbf{m} = \omega_m \cdots \omega_1$ is right (respectively left) inextensible if and only if ω_1 (respectively ω_m) is of the form $a_{i_\theta}^{-1}$ (respectively a_{i_θ}). In a word, except for its first letter (respectively last letter), there are no letters of the form $a_{i_\theta}^{-1}$ (respectively a_{i_θ}). In particular, each word contains at most two punctured letters.

Example 2.4. In Example 2.2 with the disjoint subsets given in Example 2.3, the punctured letters are $a_{1-}, a_{1-}^{-1}, a_{4-},$ and a_{4-}^{-1} . The sequence $\mathbf{m} = a_{3+} d^{-1} \varepsilon_4^{-1} d c^{-1}$ is a word with $s(\mathbf{m}) = 1$ and $t(\mathbf{m}) = 3_+$, which is left inextensible but not right inextensible.

2.4 Orders

The linear orders in the sets $L_\pm(i), i \in Q_0 \subset \widehat{Q}_0$, induce a partial order \geq on the set of words, such that $\mathbf{m} > \mathbf{r}$ if and only if $\mathbf{m} = \omega_m \cdots \omega_2 \omega_1$ and $\mathbf{r} = \nu_r \cdots \nu_2 \nu_1$ satisfy $\omega_j \cdots \omega_1 = \nu_j \cdots \nu_1$ and $\omega_{j+1} > \nu_{j+1}$ for some $j \geq 0$.

For each vertex $i \in Q_0$, let $W_\pm(i)$ be the set of left inextensible words whose first letter is in $L_\pm(i)$. By construction, $W_\pm(i)$ are linearly ordered. For each word \mathbf{m} in $W_\theta(i)$, we use $[+]\mathbf{m}$ to denote its successor (if it exists) and use $[-]\mathbf{m}$ to denote its predecessor (if it exists). So we have $[-]\mathbf{m} > \mathbf{m} > [+]\mathbf{m}$. In case \mathbf{m} is right inextensible, we set $\mathbf{m}[+] := ([+]\mathbf{m}^{-1})^{-1}$ and $\mathbf{m}[-] := ([-]\mathbf{m}^{-1})^{-1}$.

Let \mathbf{m} be a maximal word. By [Gei99], $[+](\mathbf{m}[+]) = ([+]\mathbf{m})[+]$ provided both of them exist; denote by $[+]\mathbf{m}[+]$ one (or both) of them.

Example 2.5. In Example 2.2 with the disjoint subsets given in Example 2.3, any word in the set $W_+(2)$ starts with one of the letters $b, a^{-1},$ or a_{2+} . A word in $W_+(2)$ starting with a_{2+} has to be a_{2+} since it is already left inextensible. Moreover, any word in $W_+(2)$ that starts with a^{-1} is bigger than a_{2+} and any word that starts with b is less than a_{2+} . Furthermore, we have $[+]a_{2+} = a_{3-}b$ and $[-]a_{2+} = a_{2-}a\varepsilon_1a^{-1}$.

2.5 Admissible words

For technical reasons, we introduce a special letter ε^* for each dashed loop ε and a map F on letters which sends the elements in $\{\varepsilon^{-1} > a_{i_\theta} > \varepsilon\}$ to ε^* and preserves the other letters.

A maximal word $\mathbf{m} = \omega_m \cdots \omega_1$ is called *admissible* if the following conditions hold.

(A1) For each $\omega_i = \varepsilon$ with ε a dashed loop, we have that $\omega_1^{-1} \cdots \omega_{i-1}^{-1} > \omega_m \cdots \omega_{i+1}$, and for each $\omega_i = \varepsilon^{-1}$ with ε a dashed loop, we have that $\omega_1^{-1} \cdots \omega_{i-1}^{-1} < \omega_m \cdots \omega_{i+1}$.

(A2) If \mathbf{m} contains two punctured letters, then $F(\mathbf{m})$ is not a proper power of $F(\mathbf{m}')$ for any maximal word \mathbf{m}' containing two punctured letters, where

$$F(\mathbf{m}) := F(\omega_m) \cdots F(\omega_1)F(\omega_2^{-1}) \cdots F(\omega_{m-1}^{-1}).$$

Let \mathfrak{X} be the set of admissible words. Note that if \mathfrak{m} is in \mathfrak{X} then so is its inverse \mathfrak{m}^{-1} . Let $\overline{\mathfrak{X}}$ be the set of all equivalence classes in \mathfrak{X} with respect to $\mathfrak{m} \simeq \mathfrak{m}^{-1}$. But when we say an element \mathfrak{m} in $\overline{\mathfrak{X}}$, we always mean that \mathfrak{m} is a representative in an equivalence class.

Example 2.6. In Example 2.2 with the disjoint subsets given in Example 2.3, consider the word $\mathfrak{m} = a_{3-}c^{-1}\varepsilon_1^{-1}cd^{-1}a_{4-}^{-1}$. It is clear that \mathfrak{m} is maximal. Since $a_{4-}dc^{-1} < a_{3-}c^{-1}$, \mathfrak{m} satisfies (A1). Moreover, \mathfrak{m} contains only one punctured letter a_{4-}^{-1} , so (A2) holds automatically. Hence \mathfrak{m} is admissible, i.e. $\mathfrak{m} \in \mathfrak{X}$.

2.6 Known results

In this subsection, we collect some results on indecomposable modules of skewed-gentle algebras and homomorphism spaces between them.

Let $\mathfrak{m} \in \overline{\mathfrak{X}}$. Associate an indeterminate with each punctured letter in \mathfrak{m} and let $A_{\mathfrak{m}}$ be the \mathbf{k} -algebra generated by these indeterminates x with relations $x^2 = x$. A one-dimensional module N of $A_{\mathfrak{m}}$ is an algebra homomorphism $N : A_{\mathfrak{m}} \rightarrow \mathbf{k}$. It is determined (up to isomorphism) by the values $N(x) \in \{0, 1\}$. More precisely:

- if \mathfrak{m} contains no punctured letters, then $A_{\mathfrak{m}} = \mathbf{k}$ and there is one one-dimensional module $N = \mathbf{k}$;
- if \mathfrak{m} contains one punctured letter, then $A_{\mathfrak{m}} = \mathbf{k}[x]/(x^2 - x)$ and there are two one-dimensional modules: $N = \mathbf{k}_a$ with $\mathbf{k}_a(x) = a$, for $a \in \{0, 1\}$. Moreover, we have

$$\dim_{\mathbf{k}} \text{Hom}_{A_{\mathfrak{m}}}(\mathbf{k}_u, \mathbf{k}_v) = \delta_{u,v} \quad \forall u, v \in \{0, 1\}; \tag{2.1}$$

- if \mathfrak{m} contains two punctured letters, then $A_{\mathfrak{m}} = \mathbf{k}\langle x, y \rangle / (x^2 - x, y^2 - y)$ and there are four one-dimensional modules: $N = \mathbf{k}_{a,b}$ with $\mathbf{k}_{a,b}(x) = a$ and $\mathbf{k}_{a,b}(y) = b$, for $a, b \in \{0, 1\}$. Moreover, we have

$$\dim_{\mathbf{k}} \text{Hom}_{A_{\mathfrak{m}}}(\mathbf{k}_{u,u'}, \mathbf{k}_{v,v'}) = \delta_{u,v}\delta_{u',v'} \quad \forall u, v, u', v' \in \{0, 1\}. \tag{2.2}$$

CONSTRUCTION 2.7. For each pair (\mathfrak{m}, N) with $\mathfrak{m} = \omega_m \cdots \omega_1 \in \overline{\mathfrak{X}}$ and a one-dimensional $A_{\mathfrak{m}}$ -module N , the associated representation $M = M(\mathfrak{m}, N)$ of Q bounded by R is constructed as follows.

- For each vertex $i \in Q_0$, let $I_i = \{1 \leq j \leq m - 1 \mid t(\omega_j) = i\}$.
- Let M_i be a vector space of dimension $|I_i|$, say with base vectors $z_j, j \in I_i$.
- If $\omega_{j+1} = \alpha$ an arrow in Q_1 , define $M_{\alpha}(z_j) = z_{j+1}$, if $\omega_{j+1} = \alpha^{-1}$, with α an arrow in Q_1 , define $M_{\alpha}(z_{j+1}) = z_j$.
- If $\omega_{j+1} = \varepsilon$ an arrow in Q_2 , define $M_{\varepsilon}(z_j) = M_{\varepsilon}(z_{j+1}) = z_{j+1}$, if $\omega_{j+1} = \varepsilon^{-1}$, with ε an arrow in Q_2 , define $M_{\varepsilon}(z_{j+1}) = M_{\varepsilon}(z_j) = z_j$.
- If ω_1 (respectively ω_m) is punctured with indeterminate x , define $M_{\varepsilon_{t(\omega_1)}}(z_1) = N(x)z_1$ (respectively $M_{\varepsilon_{s(\omega_m)}}(z_{m-1}) = N(x)z_{m-1}$), where ε_i denotes the dashed loop at i .
- All other components of M_{β} are zero, for any $\beta \in Q_1 \cup Q_2$.

Example 2.8. Let $\mathfrak{m} = a_{3-}c^{-1}\varepsilon_1^{-1}cd^{-1}a_{4-}^{-1}$ be the admissible word in Example 2.6. Since \mathfrak{m} contains one punctured letter a_{4-}^{-1} , we have $A_{\mathfrak{m}} = \mathbf{k}[x]/(x^2 - x)$ and there are two different one-dimensional $A_{\mathfrak{m}}$ -modules \mathbf{k}_u for $u = 0, 1$. By Construction 2.7, the associated representations

$M(\mathfrak{m}, \mathbf{k}_u)$ are as shown by the following diagram.

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \nearrow 0 & & \searrow 0 & \\
 \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \leftarrow \\ \rightarrow \end{pmatrix} \mathbf{k}^2 & & & & \mathbf{k}^2 \begin{pmatrix} \rightarrow \\ \leftarrow \end{pmatrix} \begin{matrix} \\ u \end{matrix} \\
 & & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & & \\
 & \longleftarrow & & \longrightarrow &
 \end{array}$$

THEOREM 2.9 [Bon92, Cra89, Den00]. *Let $\Lambda = \mathbf{k}Q/(R)$ be a skewed-gentle algebra. Then $(\mathfrak{m}, N) \mapsto M(\mathfrak{m}, N)$ is an injective map from the set of pairs (\mathfrak{m}, N) , with $\mathfrak{m} \in \tilde{\mathfrak{X}}$ and N a one-dimensional $A_{\mathfrak{m}}$ -module (up to isomorphism), to the set of indecomposable representations $M(\mathfrak{m}, N)$ (up to isomorphism) of Q bounded by R .*

Remark 2.10. In fact, Bondarenko [Bon92], Crawley-Boevey [Cra89], and Deng [Den00] proved the result above for general clannish algebras, where the map can be upgraded to a bijection by enlarging the set $\tilde{\mathfrak{X}}$ and taking N to be an arbitrary indecomposable $A_{\mathfrak{m}}$ -module. Furthermore, any indecomposable module M , which is not in the image of the injective map in the above theorem, is in a homogeneous tube or in a tube of rank 2 and does not sit in the bottom of the tube. So in particular $\text{Hom}(M, \tau M) \neq 0$ for such an indecomposable module M .

The Auslander–Reiten translation τ can be interpreted by the order of words.

THEOREM 2.11 [Gei99]. *For any $\mathfrak{m} = \omega_m \cdots \omega_1 \in \tilde{\mathfrak{X}}$ and any one-dimensional $A_{\mathfrak{m}}$ -module N , if $M(\mathfrak{m}, N)$ is not projective, then*

$$\tau M(\mathfrak{m}, N) = \begin{cases} M([\] \mathfrak{m} [\], \mathbf{k}) & \text{if } \mathfrak{m} \text{ contains no punctured letters and } N = \mathbf{k}, \\ M([\] \mathfrak{m}, \mathbf{k}_{1-a}) & \text{if only } \omega_1 \text{ is punctured and } N = \mathbf{k}_a, \\ M(\mathfrak{m}, \mathbf{k}_{1-a, 1-b}) & \text{if both } \omega_1 \text{ and } \omega_m \text{ are punctured and } N = \mathbf{k}_{a,b}. \end{cases}$$

For technical reasons, we also consider a trivial word 1_i corresponding to each vertex $i \in Q_0 \subset \hat{Q}_0$. Let $\mathfrak{m} = \omega_m \cdots \omega_1$ be a word in $\tilde{\mathfrak{X}}$. For any integers i, j with $0 \leq i < j \leq m + 1$, we consider the subword $\mathfrak{m}_{(i,j)}$ of \mathfrak{m} between i and j defined as

$$\mathfrak{m}_{(i,j)} = \begin{cases} \omega_{j-1} \cdots \omega_{i+1} & \text{if } i < j - 1, \\ 1_{t(\omega_i)} & \text{if } i = j - 1, \end{cases}$$

where $1_{t(\omega_0)} := 1_{s(\omega_1)}$.

Let $\mathfrak{m} = \omega_m \cdots \omega_1$ and $\mathfrak{r} = \nu_r \cdots \nu_1$ be two words in $\tilde{\mathfrak{X}}$. A pair $((i, j), (h, l))$ of pairs of integers i, j, h, l with $0 \leq i < j \leq m + 1$ and $0 \leq h < l \leq r + 1$ is called an *int-pair* from \mathfrak{m} to \mathfrak{r} if one of the following conditions holds:

- $\mathfrak{m}_{(i,j)} = \mathfrak{r}_{(h,l)}$, $\omega_i^{-1} < \nu_h^{-1}$ and $\omega_j < \nu_l$;
- $\mathfrak{m}_{(i,j)} = (\mathfrak{r}_{(h,l)})^{-1}$, $\omega_i^{-1} < \nu_l$ and $\omega_j < \nu_h^{-1}$;

where if an inequality contains at least one of $\omega_0, \omega_{m+1}, \nu_0$ and ν_{r+1} then we assume that it holds automatically. Let $H^{\mathfrak{m}, \mathfrak{r}}$ be the set of int-pairs from \mathfrak{m} and \mathfrak{r} .

Example 2.12. Let $\mathfrak{m} = a_{4+} a_{4-}^{-1}$ and $\mathfrak{r} = a_{3-} c^{-1} \varepsilon_1^{-1} c d^{-1} a_{4-}^{-1}$ be two admissible words for the skewed-gentle pair in Example 2.2 with the disjoint subsets given in Example 2.3. Then $H^{\mathfrak{m}, \mathfrak{r}}$ contains only one element $((0, 2), (0, 2))$ for which $\mathfrak{m}_{(0,2)} = a_{4-}^{-1} = \mathfrak{r}_{(0,2)}$ with $\omega_0^{-1} < \nu_0^{-1}$ and $\omega_2 = a_{4+} < d^{-1} = \nu_2$.

Notation 2.13. Let (\mathfrak{m}, N_1) and (\mathfrak{r}, N_2) be two pairs, where $\mathfrak{m}, \mathfrak{r} \in \overline{\mathfrak{X}}$ and N_1 (respectively N_2) is a one-dimensional module of $A_{\mathfrak{m}}$ (respectively $A_{\mathfrak{r}}$). For each int-pair $J = ((i, j), (h, l))$ in $H^{\mathfrak{m}, \mathfrak{r}}$, denote by A_J the \mathbf{k} -algebra generated by the indeterminates associated with punctured letters contained in $\mathfrak{m}_{(i,j)}$ (or equivalently in $\mathfrak{r}_{(k,l)}$). Then A_J is a subalgebra of $A_{\mathfrak{m}}$ and $A_{\mathfrak{r}}$ and hence both N_1 and N_2 can be regarded as A_J -modules.

THEOREM 2.14 [Gei99]. *Under Notation 2.13, we have*

$$\dim_{\mathbf{k}} \text{Hom}_{\Lambda}(M(\mathfrak{m}, N_1), M(\mathfrak{r}, N_2)) = \sum_{J \in H^{\mathfrak{m}, \mathfrak{r}}} \dim_{\mathbf{k}} \text{Hom}_{A_J}(N_1, N_2).$$

3. Background on cluster categories for marked surfaces

3.1 Jacobian algebras and Ginzburg dg algebras

Let Q be a finite quiver and W a potential on Q , that is, a sum of cycles in Q . The *Jacobian algebra* of the quiver with potential (Q, W) is the quotient

$$\mathcal{P}(Q, W) := \widehat{\mathbf{k}Q} / \overline{\partial W},$$

where $\widehat{\mathbf{k}Q}$ is the complete path algebra of Q , $\partial W = \langle \partial_a W : a \in Q_1 \rangle$ and $\overline{\partial W}$ is the closure of ∂W in $\widehat{\mathbf{k}Q}$ (cf. [DWZ08]).

The Jacobian algebra is the 0th cohomology of its refinement, the *Ginzburg dg algebra* $\Gamma = \Gamma(Q, W)$ of (Q, W) (see the construction in [Kel12, §7.2]). There are three categories associated with Γ , namely:

- the *finite dimensional derived category* $\mathcal{D}_{fd}(\Gamma)$ of Γ , which is a 3-Calabi–Yau category;
- the *perfect derived category* $\text{per}(\Gamma)$ of Γ , which contains $\mathcal{D}_{fd}(\Gamma)$;
- the *cluster category* $\mathcal{C}(\Gamma)$ of Γ , which is the (triangulated) 2-Calabi–Yau quotient category

$$\mathcal{C}(\Gamma) := \text{per}(\Gamma) / \mathcal{D}_{fd}(\Gamma). \tag{3.1}$$

Furthermore there is a canonical cluster tilting object T_{Γ} in $\mathcal{C}(\Gamma)$ induced by the silting object Γ in $\text{per} \Gamma$ such that

$$\mathcal{C}(\Gamma) / (T_{\Gamma}) \simeq \text{mod } \text{End}_{\mathcal{C}(\Gamma)}(T_{\Gamma})^{\text{op}} \cong \text{mod } \mathcal{P}(Q, W). \tag{3.2}$$

See [Ami09, Theorem 3.5] and [KR07, §2.1, Proposition (c)].

For a vertex i of Q , let $\mu_i(Q, W)$ be the mutation of (Q, W) at i in the sense of [DWZ08], see also Appendix B. By [KY11], there exists a canonical triangulated equivalence

$$\tilde{\mu}_i : \mathcal{C}(\Gamma(Q, W)) \simeq \mathcal{C}(\Gamma(\mu_i(Q, W))). \tag{3.3}$$

3.2 Quivers with potential from marked surfaces

Throughout the article, \mathbf{S} denotes a *marked surface* with non-empty boundary in the sense of [FST08], that is, a compact connected oriented surface \mathbf{S} with a finite set \mathbf{M} of marked points on its boundary $\partial \mathbf{S}$ and a finite set \mathbf{P} of punctures in its interior $\mathbf{S} \setminus \partial \mathbf{S}$ such that the following conditions hold:

- each connected component of $\partial \mathbf{S}$ contains at least one marked point;
- \mathbf{S} is not closed, i.e. $\partial \mathbf{S} \neq \emptyset$;




FIGURE 1. The completions of curves.

– the rank

$$n = 6g + 3p + 3b + m - 6 \tag{3.4}$$

of the surface is positive, where g is the genus of \mathbf{S} , b the number of boundary components, $m = |\mathbf{M}|$ the number of marked points and $p = |\mathbf{P}|$ the number of punctures.

DEFINITION 3.1 (Curves and tagged curves). Let \mathbf{S} be a marked surface with non-empty boundary.

- An (ordinary) *curve* in \mathbf{S} is a continuous function $\gamma : [0, 1] \rightarrow \mathbf{S}$ satisfying:
 - * both $\gamma(0)$ and $\gamma(1)$ are in $\mathbf{M} \cup \mathbf{P}$;
 - * for any $0 < t < 1$, $\gamma(t)$ is in $\mathbf{S} \setminus (\partial\mathbf{S} \cup \mathbf{P})$;
 - * γ is not null-homotopic or homotopic to a boundary segment.
- The inverse of a curve γ is defined as $\gamma^{-1}(t) := \gamma(1 - t)$ for $t \in [0, 1]$.
- For two curves γ_1, γ_2 , $\gamma_1 \sim \gamma_2$ means that γ_1 is homotopic to γ_2 relative to $\{0, 1\}$ (i.e. fixing the endpoints). Define an equivalence relation \simeq on the set of curves in \mathbf{S} that $\gamma_1 \simeq \gamma_2$ if and only if either $\gamma_1 \sim \gamma_2$ or $\gamma_1^{-1} \sim \gamma_2$. Denote by $\mathbf{C}(\mathbf{S})$ the set of equivalence classes of curves in \mathbf{S} with respect to \simeq .
- Let γ be a curve in $\mathbf{C}(\mathbf{S})$ such that at least one of its endpoints is a puncture. Then define its *completion* $\bar{\gamma}$ as in Figure 1.
- A *tagged curve* is a pair (γ, κ) , where γ is a curve in \mathbf{S} and $\kappa : \{t \mid \gamma(t) \in \mathbf{P}\} \rightarrow \{0, 1\}$ is a map, satisfying the following conditions:
 - (T1) γ does not cut out a once-punctured monogon by a self-intersection (including endpoints), cf. Figure 2;
 - (T2) if $\gamma(0), \gamma(1) \in \mathbf{P}$, then the completion $\bar{\gamma}$ is not a proper power of a closed curve in the sense of the multiplication in the fundamental group of \mathbf{S} .

Note that $\kappa(t) \in \mathbf{P}$ implies $t \in \{0, 1\}$. Moreover, write $\kappa = \emptyset$ when $\{t \mid \gamma(t) \in \mathbf{P}\} = \emptyset$ by convention.




FIGURE 2. Once-punctured monogons.




FIGURE 3. The tagged rotations of three tagged curves in \mathbf{S} .

- The inverse of a tagged curve (γ, κ) is defined as $(\gamma, \kappa)^{-1} := (\gamma^{-1}, \kappa^{-1})$, where $\kappa^{-1}(t) := \kappa(1 - t)$.
- For two tagged curves $(\gamma_1, \kappa_1), (\gamma_2, \kappa_2)$, $(\gamma_1, \kappa_1) \sim (\gamma_2, \kappa_2)$ means that $\gamma_1 \sim \gamma_2$ and $\kappa_1 = \kappa_2$. Define an equivalence relation \simeq on the set of tagged curves in \mathbf{S} that $(\gamma_1, \kappa_1) \simeq (\gamma_2, \kappa_2)$ if and only if either $(\gamma_1, \kappa_1) \sim (\gamma_2, \kappa_2)$ or $(\gamma_1, \kappa_1) \sim (\gamma_2, \kappa_2)^{-1}$. Denote by $\mathbf{C}^\times(\mathbf{S})$ the set of equivalence classes of tagged curves in \mathbf{S} with respect to \simeq .

DEFINITION 3.2 (Tagged rotation [BQ15]). The rotation $\rho(\gamma)$ of a curve γ in $\mathbf{C}(\mathbf{S})$ is the curve obtained from γ by moving every endpoint of γ that is in \mathbf{M} along the boundary anticlockwise to the next marked point. The tagged rotation $(\gamma', \kappa') = \rho(\gamma, \kappa)$ of a tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})$ consists of the curve $\gamma' = \rho(\gamma)$ and the map κ' defined by $\kappa'(t) = 1 - \kappa(t)$ for t with $\gamma(t) \in \mathbf{P}$, cf. Figure 3.

DEFINITION 3.3 (Intersection numbers). For any two curves $\gamma_1, \gamma_2 \in \mathbf{C}(\mathbf{S})$:

- let $\gamma_1 \cap \gamma_2 = \{(t_1, t_2) \mid \gamma_1(t_1) = \gamma_2(t_2) \notin \mathbf{P} \cup \mathbf{M}\} \subset (0, 1)^2$ be the set of interior intersections between γ_1 and γ_2 ;
- the intersection number between them is defined to be

$$\text{Int}(\gamma_1, \gamma_2) := \min\{|\gamma'_1 \cap \gamma'_2| \mid \gamma'_1 \sim \gamma_1, \gamma'_2 \sim \gamma_2\}.$$

For any two tagged curves (γ_1, κ_1) and $(\gamma_2, \kappa_2) \in \mathbf{C}^\times(\mathbf{S})$:

- let $\mathfrak{P}(\gamma_1, \gamma_2) = \{(t_1, t_2) \mid \gamma_1(t_1) = \gamma_2(t_2) \in \mathbf{P}\} \subset \{0, 1\}^2$ be the set of intersections between γ_1 and γ_2 at \mathbf{P} ;
- A pair (t_1, t_2) in $\mathfrak{P}(\gamma_1, \gamma_2)$ is called a tagged intersection between (γ_1, κ_1) and (γ_2, κ_2) if:
 - * $\kappa_1(t_1) \neq \kappa_2(t_2)$; and
 - * when $\gamma_1|_{t_1 \rightarrow (1-t_1)} \sim \gamma_2|_{t_2 \rightarrow (1-t_2)}$, we have $\gamma_1(1 - t_1) = \gamma_2(1 - t_2)$ belongs to \mathbf{P} and $\kappa_1(1 - t_1) \neq \kappa_2(1 - t_2)$, where $\gamma|_{0 \rightarrow 1} = \gamma$ and $\gamma|_{1 \rightarrow 0} = \gamma^{-1}$;




FIGURE 4. The punctured intersections.

- let $\mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))$ be the set of tagged intersections between (γ_1, κ_1) and (γ_2, κ_2) ;
- the intersection number between them is defined to be

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) := \text{Int}(\gamma_1, \gamma_2) + |\mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))|.$$

We explain the intersection number of two tagged curves in some special cases.

Example 3.4. Let (γ_1, κ_1) and (γ_2, κ_2) be two tagged curves in $\mathbf{C}^\times(\mathbf{S})$.

- If all the endpoints of γ_1 and γ_2 are in \mathbf{M} , then

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = \text{Int}(\gamma_1, \gamma_2).$$

- If γ_1 and γ_2 are not in the same equivalence class in $\mathbf{C}(\mathbf{S})$ (i.e. $\gamma_1 \not\sim \gamma_2$ and $\gamma_1 \not\sim \gamma_2^{-1}$), then

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = \text{Int}(\gamma_1, \gamma_2) + |\{(t_1, t_2) \mid \gamma_1(t_1) = \gamma_2(t_2) \in \mathbf{P}, \kappa_1(t_1) \neq \kappa_2(t_2)\}|.$$

- If $\gamma_1 \sim \gamma_2$ whose endpoints are two different punctures, then

$$|\mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))| = \begin{cases} 2 & \text{if } \kappa_1(0) \neq \kappa_2(0) \text{ and } \kappa_1(1) \neq \kappa_2(1), \\ 0 & \text{otherwise.} \end{cases}$$

- If the two tagged curves are as in Figure 4 where $\gamma_1 \sim \gamma_2^{-1}$ and

$$\kappa_a(t) = \begin{cases} 1 & \text{if } a = 1 \text{ and } t = 0, \\ 0 & \text{otherwise,} \end{cases}$$

then $(0, 0) \in \mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))$. This is because $\kappa_1(0) = 1 \neq \kappa_2(0) = 0$ and $\gamma_1|_{0 \rightarrow 1} \sim \gamma_2|_{0 \rightarrow 1}$. For the pair $(0, 1)$, we also have $\kappa_1(0) = 1 \neq \kappa_2(1) = 0$. But since $\gamma_1|_{0 \rightarrow 1} \sim \gamma_2|_{1 \rightarrow 0}$, we need to compare the values of $\kappa_1(1)$ and $\kappa_2(0)$. Because $\kappa_1(1) = 0 = \kappa_2(0)$, the pair $(0, 1) \notin \mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))$. It is easy to see that neither $(1, 0)$ nor $(1, 1)$ is in $\mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))$. Hence in this case we have

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = 1.$$

We also mention that

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_1, \kappa_1)) = 2.$$

Since (γ_1, κ_1) has self-intersections, it is not a tagged arc in the sense of [FST08] but it is in $\mathbf{C}^\times(\mathbf{S})$.

Remark 3.5. Note that we have less restriction for curves and tagged curves than Fomin *et al.*, that is, we allow self-intersections. More precisely:




FIGURE 5. The self-folded triangle and the corresponding tagged version.

- the curves γ in $\mathbf{C}(\mathbf{S})$ without self-intersections (i.e. $\text{Int}(\gamma, \gamma) = 0$) are precisely the arcs in the sense of [FST08, Definition 2.2];
- the tagged curves (γ, κ) in $\mathbf{C}^\times(\mathbf{S})$ without self-intersections (i.e. $\text{Int}((\gamma, \kappa), (\gamma, \kappa)) = 0$) are precisely the tagged arcs in the sense of [FST08, Definition 2.4];
- for two curves γ_1 and γ_2 in $\mathbf{C}(\mathbf{S})$ without self-intersections, we have that $\text{Int}(\gamma_1, \gamma_2) = 0$ if and only if they are compatible in the sense of [FST08, Definition 7.1];
- for two tagged curves (γ_1, κ_1) and (γ_2, κ_2) in $\mathbf{C}^\times(\mathbf{S})$ without self-intersections, we have that $\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = 0$ if and only if they are compatible in the sense of [FST08, Definition 7.4].

By Remark 3.5, the following definitions of ideal triangulations and tagged triangulations are equivalent to the original ones in [FST08].

DEFINITION 3.6 (Ideal triangulations and tagged triangulations [FST08]). Let \mathbf{S} be a marked surface with non-empty boundary.

- An *ideal triangulation* is a maximal collection \mathbf{T} of curves in $\mathbf{C}(\mathbf{S})$ such that $\text{Int}(\gamma_1, \gamma_2) = 0$ for any $\gamma_1, \gamma_2 \in \mathbf{T}$.
- A *tagged triangulation* is a maximal collection \mathbf{T} of tagged curves in $\mathbf{C}^\times(\mathbf{S})$ such that $\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = 0$ for any $(\gamma_1, \kappa_1), (\gamma_2, \kappa_2) \in \mathbf{T}$.

Any ideal/tagged triangulation \mathbf{T} of \mathbf{S} consists of n ordinary/tagged curves (see [FST08, Proposition 2.10, Theorem 7.9]), where n is the rank of \mathbf{S} (cf. (3.4)). We require $n > 0$ and exclude the case of a once-punctured monogon (where $n = 1$) in the proofs. However, all the results hold in this case by direct checking and thus we will not exclude this case in the statements.

A triangle in \mathbf{T} has three distinct sides unless it is a *self-folded triangle* as in the left picture of Figure 5, where we call α the *folded side* and β the *remaining side*.

The *flip* of an ideal triangulation \mathbf{T} , with respect to a curve α in \mathbf{T} , is the unique ideal triangulation \mathbf{T}' (if it exists) that shares all curves in \mathbf{T} but α . One can always flip an ideal triangulation with respect to a curve unless it is the folded side of a self-folded triangle. To overcome this shortcoming, Fomin *et al.* [FST08] introduced tagged triangulations, with tagged flips, so that every tagged triangulation can be flipped with respect to any tagged curve in it. The exchange graph of tagged triangulations with tagged flips is denoted by $\text{EG}^\times(\mathbf{S})$, that is, the graph whose vertices are tagged triangulations and whose edges are tagged flips.

For each curve γ in $\mathbf{C}(\mathbf{S})$ with $\text{Int}(\gamma, \gamma) = 0$, we define its tagged version γ^\times to be (γ, \emptyset) unless γ is a loop enclosing a puncture, as β in the left picture of Figure 5. In that case β^\times , as in the right picture of Figure 5, is defined to be (α, κ) , where α is the unique curve without self-intersections




FIGURE 6. The quiver associated with a non-self-folded triangle.

enclosed by β and $\kappa(t) = 1$ for t with $\alpha(t) \in \mathbf{P}$. In this way, each ideal triangulation \mathbf{T} induces a tagged triangulation \mathbf{T}^\times consisting of the tagged versions of all curves in \mathbf{T} .

For each ideal triangulation \mathbf{T} , there is an associated quiver with potential $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ (cf. [FST08, IL12]). In the paper, we only study $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ in the case when \mathbf{T} is an admissible triangulation in the following sense (see Figure 14 for example).

DEFINITION 3.7. An ideal triangulation \mathbf{T} is called admissible if every puncture in \mathbf{P} is contained in a self-folded triangle in \mathbf{T} .

In particular, in such a triangulation, the folded side of each self-folded triangle connects a marked point in \mathbf{M} and a puncture in \mathbf{P} .

In an admissible triangulation \mathbf{T} , for a curve $\alpha \in \mathbf{T}$, let $\pi_{\mathbf{T}}(\alpha)$ be the curve defined as follows: if α is the folded side of a self-folded triangle in \mathbf{T} (see the left picture of Figure 5), then $\pi_{\mathbf{T}}(\alpha)$ is the corresponding remaining side (i.e. β in the left picture of Figure 5); if there is no such triangle, set $\pi_{\mathbf{T}}(\alpha) = \alpha$. The associated quiver with potential $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ is given by the following data (see Figure 6):

- the vertices of $Q_{\mathbf{T}}$ are labeled by the curves in \mathbf{T} ;
- there is an arrow from i to j whenever there is a non-self-folded triangle in \mathbf{T} having $\pi_{\mathbf{T}}(i)$ and $\pi_{\mathbf{T}}(j)$ as edges with $\pi_{\mathbf{T}}(j)$ following $\pi_{\mathbf{T}}(i)$ in the clockwise orientation (which is induced by the orientation of \mathbf{S}). For instance, the quiver for a non-self-folded triangle is shown in Figure 6;
- each subset $\{i, j, k\}$ of \mathbf{T} with $\pi_{\mathbf{T}}(i), \pi_{\mathbf{T}}(j), \pi_{\mathbf{T}}(k)$ forming an interior non-self-folded triangle in \mathbf{T} yields a unique three-cycle up to cyclic permutation. The potential $W_{\mathbf{T}}$ is the sum of all such three-cycles.

Then by §3.1, there is an associated cluster category, denoted by $\mathcal{C}(\mathbf{T})$.

3.3 Correspondence

The objects and morphisms in $\mathcal{C}(\mathbf{T})$ are expected to correspond to curves and intersection numbers, respectively. A cluster tilting object $T = \bigoplus_{j=1}^n T_j$ in a cluster category \mathcal{C} is an object satisfying $\text{Ext}_{\mathcal{C}}^1(T, X) = 0$ if and only if $X \in \text{add } T$. The mutation μ_i at the i th indecomposable direct summand acts on a cluster tilting object $T = \bigoplus_{j=1}^n T_j$, by replacing T_i with the unique indecomposable object $T'_i \not\cong T_i$ satisfying that $(T \setminus T_i) \oplus T'_i$ is a cluster tilting object.

In the unpunctured case, we have the following known results.

THEOREM 3.8 [BZ11]. *If \mathbf{S} is unpunctured, then there is a bijection between the set of curves and valued closed curves in \mathbf{S} and the set of indecomposable objects in $\mathcal{C}(\mathbf{T})$. Under such a bijection:*

- rotation of curves is compatible with shift of objects;
- triangulations of \mathbf{S} one-to-one correspond to cluster tilting objects in $\mathcal{C}(\mathbf{T})$ while flip of triangulations is compatible with mutation of cluster tilting objects.

THEOREM 3.9 [ZZZ13]. *If \mathbf{S} is unpunctured, then for any two curves γ, δ , we have*

$$\text{Int}(\gamma, \delta) = \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_\gamma, X_\delta),$$

where $X : \eta \mapsto X_\eta$ is the bijection in Theorem 3.8.

In the punctured case, we also know the following. Recall that a rigid indecomposable object in $\mathcal{C}(\mathbf{T})$ is *reachable* if it is a summand of some cluster tilting object, which is obtained from the canonical cluster tilting object by a sequence of mutations.

THEOREM 3.10 [BQ15]. *Let $\mathbf{A}^\times(\mathbf{S})$ be the set of tagged curves in $\mathbf{C}^\times(\mathbf{S})$ without self-intersections and $\mathcal{I}^\times(\mathbf{S})$ the set of reachable rigid indecomposable objects in $\mathcal{C}(\mathbf{T})$. Under a canonical bijection*

$$\varepsilon : \mathbf{A}^\times(\mathbf{S}) \rightarrow \mathcal{I}^\times(\mathbf{S}),$$

the tagged rotation ρ on $\mathbf{A}^\times(\mathbf{S})$ becomes the shift $[1]$ on $\mathcal{I}^\times(\mathbf{S})$.

3.4 Cluster(-tilting) exchange graphs

The *cluster(-tilting) exchange graph* $\text{CEG}(\mathcal{C})$ of a cluster category \mathcal{C} is the graph whose vertices are cluster tilting objects and whose edges are mutations. There are the following known results about connectedness of cluster exchange graphs.

THEOREM 3.11 [BMRRT06]. *If \mathcal{C} is the cluster category of an acyclic quiver, then $\text{CEG}(\mathcal{C})$ is connected.*

THEOREM 3.12 [BZ11]. *If \mathcal{C} is the cluster category from an unpunctured marked surface, then $\text{CEG}(\mathcal{C})$ is connected.*

4. Strings and tagged curves

4.1 Skewed-gentle algebras from admissible triangulations

Let \mathbf{T} be an admissible triangulation of \mathbf{S} , i.e. every puncture is in a self-folded triangle (see Lemma A.1 for the existence of \mathbf{T}), with the associated quiver with potential $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ and the cluster category $\mathcal{C}(\mathbf{T})$.

Let \mathbf{T}° be the subset of \mathbf{T} consisting of curves whose endpoints are in \mathbf{M} . Now we associate a biquiver $Q^{\mathbf{T}} = (Q_0^{\mathbf{T}}, Q_1^{\mathbf{T}}, Q_2^{\mathbf{T}})$ with potential $W^{\mathbf{T}}$ as follows:

- $Q_0^{\mathbf{T}} = \mathbf{T}^\circ$, i.e. curves in \mathbf{T} which are sides of non-self-folded triangles correspond to vertices in $Q_0^{\mathbf{T}}$;
- there is a solid arrow from i to j in $Q_1^{\mathbf{T}}$ whenever there is a non-self-folded triangle Δ in \mathbf{T} such that Δ has sides i and j with j following i in the clockwise orientation;
- there is a dashed loop at i in $Q_2^{\mathbf{T}}$, denoted by ε_i , whenever i is the remaining side of a self-folded triangle;
- each non-self-folded triangle in \mathbf{T} induces a unique three-cycle up to cyclic permutation. The potential $W^{\mathbf{T}}$ is the sum of all such three-cycles.

See the example in § 6. By construction, there are no loops in $Q_1^{\mathbf{T}}$, any arrow in $Q_2^{\mathbf{T}}$ is a loop and there is at most one loop in $Q_2^{\mathbf{T}}$ at each vertex. Hence the biquiver $Q^{\mathbf{T}}$ satisfies the conditions




FIGURE 7. Non-self-folded triangles with the corresponding quivers.




FIGURE 8. Self-folded triangles with the corresponding quivers.

on biquivers in § 2.1. Let $Z = \partial W^{\mathbf{T}} = \{\partial_a W^{\mathbf{T}} : a \in Q_1^{\mathbf{T}}\}$. We have the following straightforward observation.

LEMMA 4.1. *The set Z consists of $\beta\alpha$ for each pair $\alpha, \beta \in Q_1^{\mathbf{T}}$ such that they are from the same non-self-folded triangle in \mathbf{T} .*

PROPOSITION 4.2. *The pair $(Q^{\mathbf{T}}, Z)$ is skewed-gentle and the algebra $\Lambda^{\mathbf{T}} := \mathbf{k}Q^{\mathbf{T}}/(R)$ is a skewed-gentle algebra, where $R = Z \cup \{\varepsilon^2 = \varepsilon \mid \varepsilon \in Q_2^{\mathbf{T}}\}$.*

Proof. By Lemma 4.1, each element in Z is of the form $\beta\alpha$ for some α, β in $Q_1^{\mathbf{T}}$. Since every vertex $i \in Q_0^{\mathbf{T}}$ is a side of a non-self-folded triangle in \mathbf{T} , there are two possible cases.

(1) The curve i is a common side of two non-self-folded triangles in \mathbf{T} , see Figure 7. Then there is no dashed loop at i and there are at most two solid arrows α_1, α_2 ending at i and at most two solid arrows β_1, β_2 starting at i . By Lemma 4.1, $\beta_1\alpha_1 \in Z$, $\beta_2\alpha_2 \in Z$, $\beta_1\alpha_2 \notin Z$ and $\beta_2\alpha_1 \notin Z$ (if they exist).

(2) The curve i is a common side of a non-self-folded triangle and a self-folded triangle in \mathbf{T} , see Figure 8. Then there is a dashed loop at i and there is at most one solid arrow α ending at i and at most one solid arrow β starting at i . By Lemma 4.1, $\beta\alpha \in Z$ (if they exist).

Hence by Definition 2.1, $(Q^{\mathbf{T}}, Z)$ is skewed-gentle and the algebra $\Lambda^{\mathbf{T}} = \mathbf{k}Q^{\mathbf{T}}/(R)$ is a skewed-gentle algebra. \square

Let Q_0^{Sp} be the subset of $Q_0^{\mathbf{T}}$ consisting of vertices where there are dashed loops.

Remark 4.3. Comparing the constructions of $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ and $(Q^{\mathbf{T}}, W^{\mathbf{T}})$, one can obtain $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ (up to isomorphism) from $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ by splitting each vertex in Q_0^{Sp} into two vertices and removing all the dashed loops. More precisely:

- vertices in $Q_{\mathbf{T}}$ are indexed by elements in $Q_0^{\mathbf{T}} \cup \{i' \mid i \in Q_0^{\text{Sp}}\}$;
- arrows from i to j in $Q_{\mathbf{T}}$ are indexed by ${}^j\alpha^i$ induced by arrows $\alpha : \pi(i) \rightarrow \pi(j)$ in $Q_1^{\mathbf{T}}$, where π is the map on $Q_0^{\mathbf{T}} \cup \{i' \mid i \in Q_0^{\text{Sp}}\}$ sending i' to i and being the identity on $Q_0^{\mathbf{T}}$;
- the potential $W_{\mathbf{T}}$ is the sum of cycles that are obtained from cycles in $W^{\mathbf{T}}$ by replacing each α with ${}^j\alpha^i$ for possible i and j .

See the example in § 6.

Now we prove that the Jacobian algebra $\mathcal{P}(Q_{\mathbf{T}}, W_{\mathbf{T}})$ is isomorphic to the skewed-gentle algebra $\Lambda^{\mathbf{T}}$.

PROPOSITION 4.4. *There is an algebra isomorphism*

$$\varphi : \mathcal{P}(Q_{\mathbf{T}}, W_{\mathbf{T}}) \cong \Lambda^{\mathbf{T}}.$$

Proof. For each quiver, denote by e_i the trivial path associated with a vertex i . Noticing that $\varepsilon_i^2 - \varepsilon_i \in R$ for each $i \in Q_0^{\text{Sp}}$, a complete set of primitive orthogonal idempotents of $kQ^{\mathbf{T}}/(R^{\text{Sp}})$ is

$$\{e_i + (R^{\text{Sp}}) \mid i \in Q_0 \setminus Q_0^{\text{Sp}}\} \cup \{\varepsilon_i + (R^{\text{Sp}}), e_i - \varepsilon_i + (R^{\text{Sp}}) \mid i \in Q_0^{\text{Sp}}\},$$

where $R^{\text{Sp}} = \{\varepsilon_i^2 - \varepsilon_i \mid i \in Q_0^{\text{Sp}}\}$. Then using the recovery of $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ from $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ in Remark 4.3, there is an isomorphism φ of algebras from $kQ_{\mathbf{T}}$ to $kQ^{\mathbf{T}}/(R^{\text{Sp}})$ that sends ${}^j\alpha^i$ to $\varphi(e_j)\alpha\varphi(e_i)$, where

$$\varphi(e_i) = \begin{cases} e_i + (R^{\text{Sp}}) & \text{if } i \in Q_0^{\mathbf{T}} \setminus Q_0^{\text{Sp}}, \\ e_i - \varepsilon_i + (R^{\text{Sp}}) & \text{if } i \in Q_0^{\text{Sp}}, \\ \varepsilon_k + (R^{\text{Sp}}) & \text{if } i = k' \text{ for } k \in Q_0^{\text{Sp}}. \end{cases}$$

By [IL12, Theorem 5.7], the Jacobian algebra $\mathcal{P}(Q_{\mathbf{T}}, W_{\mathbf{T}})$ is isomorphic to $kQ_{\mathbf{T}}/\partial W_{\mathbf{T}}$. Then what is left to show is that $\varphi(\partial W_{\mathbf{T}}) = \partial W^{\mathbf{T}}$, which follows directly from the recovery of $W_{\mathbf{T}}$ from $W^{\mathbf{T}}$ in Remark 4.3. \square

Remark 4.5. Note that Propositions 4.2 and 4.4 imply that the Jacobian algebra $\mathcal{P}(Q_{\mathbf{T}}, W_{\mathbf{T}})$ is a skewed-gentle algebra for any admissible triangulation \mathbf{T} of a marked surface. This result was first announced by Labardini-Fragoso (cf. [Lab16]) and was first proved in [GLS16].

4.2 Correspondence

Denote by $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ the category of finite dimensional \mathbf{k} -linear representations of Q bounded by $R = \partial W^{\mathbf{T}} \cup \{\varepsilon^2 = \varepsilon \mid \varepsilon \in Q_2^{\mathbf{T}}\}$. By the equivalence (3.2) and Proposition 4.4, we have an equivalence

$$F_{\mathbf{T}} : \mathcal{C}(\mathbf{T})/(T_{\mathbf{T}}) \simeq \text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}}), \tag{4.1}$$

where $T_{\mathbf{T}}$ denotes the canonical cluster tilting object in $\mathcal{C}(\mathbf{T})$ (cf. § 3.1). So one can regard the set of indecomposable objects in $\mathcal{C}(\mathbf{T})$ as the union of the set of indecomposable representations in $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ and the set of indecomposable direct summands of $T_{\mathbf{T}}$. Note that indecomposable direct summands of $T_{\mathbf{T}}$ are indexed by curves in \mathbf{T} and hence also by tagged curves in \mathbf{T}^{\times} (which is the tagged version of \mathbf{T}). Thus, we can write

$$T_{\mathbf{T}} = \bigoplus_{(\gamma, \kappa) \in \mathbf{T}^{\times}} T_{(\gamma, \kappa)}. \tag{4.2}$$




FIGURE 9. The arc segments associated with letters.

DEFINITION 4.6. We call an indecomposable object X in $\mathcal{C}(\mathbf{T})$ a *string object* if either X is a direct summand of the canonical cluster tilting object $T_{\mathbf{T}}$, or $F_{\mathbf{T}}(X)$ is an indecomposable representation in $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ which is in the image of the injective map in Theorem 2.9. Denote the set of string objects in $\mathcal{C}(\mathbf{T})$ by $\mathfrak{S}(\mathbf{T})$.

To describe indecomposable representations in $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$, we shall use notions and notation stated in § 2. Recall that from the biquiver $Q^{\mathbf{T}}$, in § 2.2, we constructed a new biquiver $\widehat{Q} = (\widehat{Q}_0, \widehat{Q}_1, \widehat{Q}_2)$ by adding two new solid arrows $a_{i\pm}$ (whose terminal vertices are also new vertices) for each vertex $i \in Q_0^{\mathbf{T}}$. Recall that a letter is an arrow in \widehat{Q} or its inverse. For each $i \in Q_0^{\mathbf{T}} \subset \widehat{Q}_0$ the set $L(i) = \{l \in L \mid s(l) = i\}$ is divided into two disjoint subsets $L_{\theta}(i)$, $\theta \in \{\pm\}$, with linear orders satisfying certain conditions.

CONSTRUCTION 4.7. We associate an arc segment $\mathbf{a}(l)$ with each letter l as follows.

- For $l = \alpha$ with α an arrow from i to j in $Q_1^{\mathbf{T}} \subset \widehat{Q}_1$, α is induced from a triangle Δ in \mathbf{T} having i and j as sides with j following i in the clockwise orientation. Then we associate the arc segment (which is unique up to homotopy) in Δ starting at (a point in) i and ending at (a point in) j .
- For $l = \varepsilon$ with ε a dashed loop at i , we associate the arc segment in the self-folded triangle whose remaining side is i with the clockwise orientation.
- For $l = a_{i\theta}$ with $a_{i\theta}$ an arrow in $\widehat{Q}_1 \setminus Q_1^{\mathbf{T}}$, its associated segment is in the same triangle as the associated arc segments of other letters in $L_{\theta}(i)$. It starts at (a point in) i and ends at a point in $\mathbf{M} \cup \mathbf{P}$.
- The arc segment $\mathbf{a}(l^{-1})$ is the same as $\mathbf{a}(l)$ but with the opposite orientation.

See Figure 9 (cf. Figures 7 and 8), where the disjoint subsets $L_{\theta}(i)$ are given by $L_+(i) = \{\alpha_1^{-1} > a_{i+} > \beta_1\}$ and $L_-(i) = \{\alpha_2^{-1} > a_{i-} > \beta_2\}$ for the left picture and by $L_+(i) = \{\alpha^{-1} > a_{i+} > \beta\}$ and $L_-(i) = \{\varepsilon^{-1} > a_{i-} > \varepsilon\}$ for the right picture. Note that $\alpha, \beta, \alpha_1, \beta_1, \alpha_2, \beta_2$ might not exist.

By Construction 4.7, any arc segment associated with a letter in $L(i)$ starts at i . The following lemma gives some basic topological interpretation of notions about letters.




FIGURE 10. The orders.

LEMMA 4.8. For any two letters $l_1, l_2 \in L(i)$:

- (1) l_1 and l_2 are in the same subset $L_\theta(i)$ if and only if $\mathbf{a}(l_1)$ and $\mathbf{a}(l_2)$ are in the same triangle;
- (2) $l_1 > l_2$ if and only if they are of one of the forms in Figure 10(i) with $\mathbf{a}(l_i)$ being the segment of γ_i cut out by the triangle;
- (3) $l = l_i$ is punctured if and only if one of the endpoints of $\mathbf{a}(l)$ is a puncture.

Proof. For (1), recall that if $\gamma, \delta^{-1} \in L(i)$ for two solid arrows γ and δ then γ and δ^{-1} are in the same subset $L_\theta(i)$ if and only if $\gamma\delta \in Z$. Then by Lemma 4.1, γ and δ are from the same triangle. By Construction 4.7, $\mathbf{a}(\gamma)$ and $\mathbf{a}(\delta^{-1})$ are in the same triangle. Also by Construction 4.7, $\mathbf{a}(\varepsilon)$ and $\mathbf{a}(\varepsilon^{-1})$ are in the same triangle for a dashed loop ε and $\mathbf{a}(a_{i_\theta})$ is in the same triangle as the arc segments associated with other letters in $L_\theta(i)$. Hence we are done.

For (2), by definition we know two letters l_1 and l_2 are comparable if they are in the same subset $L_\theta(i)$. Then by (1) this is equivalent to $\mathbf{a}(l_1)$ and $\mathbf{a}(l_2)$ being in the same triangle and starting at the same curve i . Note that the forms in Figure 10(i) correspond to $\alpha^{-1} > \beta$ for $\beta\alpha \in Z$, $\alpha^{-1} > a_{i_\theta}$, $a_{i_\theta} > \beta$, $\varepsilon^{-1} > \varepsilon$, $a_{i_\theta} > \varepsilon$, and $\varepsilon^{-1} > a_{i_\theta}$, respectively. These give all the possible cases for the order.

For (3), l is punctured by definition if and only if l is of the form a_{i_θ} or $a_{i_\theta}^{-1}$ such that $L_{i_\theta} = \{\varepsilon^{-1} > a_{i_\theta} > \varepsilon\}$. Then by Construction 4.7, $\mathbf{a}(l)$ is in a self-folded triangle and hence it has to connect the puncture in this triangle. \square

Recall that a word is a sequence $\mathbf{m} = \omega_m \cdots \omega_1$ of letters in L such that for any $1 \leq j \leq m-1$, $\omega_j^{-1} \in L_\theta(i)$ and $\omega_{j+1} \in L_{\theta'}(i)$ for different $\theta, \theta' \in \{\pm\}$ and some $i \in Q_0^{\mathbf{T}}$. By Lemma 4.8(1), this condition is equivalent to $\mathbf{a}(\omega_j^{-1})$ and $\mathbf{a}(\omega_{j+1})$ starting at the same curve in \mathbf{T} , but being in the two triangles adjacent to this curve, respectively. Hence we can glue the two arc segments $\mathbf{a}(\omega_j)$ and $\mathbf{a}(\omega_{j+1})$ together to get a curve segment.

CONSTRUCTION 4.9. For each word $\mathbf{m} = \omega_m \cdots \omega_1$, we glue the corresponding arc segments $\mathbf{a}(\omega_1), \dots, \mathbf{a}(\omega_m)$ together in order to get a curve segment, denoted by $\mathbf{a}(\mathbf{m})$.

Recall that for two words $\mathbf{m} = \omega_m \cdots \omega_2 \omega_1$ and $\mathbf{t} = \nu_r \cdots \nu_2 \nu_1$, $\mathbf{m} > \mathbf{t}$ if and only if there is j such that $\omega_j \cdots \omega_1 = \nu_j \cdots \nu_1$ and $\omega_{j+1} > \nu_{j+1}$.

LEMMA 4.10. *Let $\mathbf{m}_1, \mathbf{m}_2$ be two words and $\gamma_1 = \mathbf{a}(\mathbf{m}_1), \gamma_2 = \mathbf{a}(\mathbf{m}_2)$. Then $\mathbf{m}_1 \geq \mathbf{m}_2$ if and only if γ_1 and γ_2 separate as in one of the forms in Figure 10 after they share the same curve segments from the start. The equality holds if and only if one of the forms in Figure 10(ii) occurs, and in this case $\gamma_1 \sim \gamma_2$.*

Proof. By definition, $\mathbf{m}_1 = \omega_m \cdots \omega_1 > \mathbf{m}_2 = \omega'_r \cdots \omega'_1$ if and only if there is a maximal integer $j \geq 0$ such that the first j letters (from right to left) of \mathbf{m}_1 and \mathbf{m}_2 are the same pointwise and $\omega_{j+1} > \omega'_{j+1}$. By Lemma 4.8(2), this is equivalent to $\mathbf{a}(\mathbf{m}_1)$ and $\mathbf{a}(\mathbf{m}_2)$ sharing the first j arc segments and their $(j + 1)$ th arc segments having one of the forms in Figure 10(i). Clearly $\mathbf{m}_1 = \mathbf{m}_2$ if and only if one of the forms in Figure 10(ii) occurs and hence $\gamma_1 \sim \gamma_2$. Thus the lemma holds. □

Recall that a word $\mathbf{m} = \omega_m \cdots \omega_1$ is maximal if and only if $\omega_1 = a_{i_\theta}^{-1}$ and $\omega_m = a_{j_{\theta'}}$ for some $i, j \in Q_0^{\mathbf{T}}$ and some $\theta, \theta' \in \{\pm\}$.

LEMMA 4.11. *The map $\mathbf{m} \mapsto \mathbf{a}(\mathbf{m})$ is a bijection from the set of maximal words to the set of curves (up to homotopy) in \mathbf{S} that are not in \mathbf{T} . Moreover, $\mathbf{a}(\mathbf{m}^{-1}) = \mathbf{a}(\mathbf{m})^{-1}$.*

Proof. Let $\mathbf{m} = \omega_m \cdots \omega_1$. Since \mathbf{m} is maximal, we have $\omega_1 = a_{i_\theta}^{-1}$ and $\omega_m = a_{j_{\theta'}}$ for some $i, j \in Q_0^{\mathbf{T}}$ and some $\theta, \theta' \in \{\pm\}$. By Construction 4.7, the endpoints of $\mathbf{a}(\mathbf{m})$ are in $\mathbf{M} \cup \mathbf{P}$. Because by Lemma 4.8(1), among the arc segments $\mathbf{a}(\omega_1), \dots, \mathbf{a}(\omega_m)$, no two adjacent arc segments are in the same triangle, the curve $\mathbf{a}(\mathbf{m})$ has minimal intersections with the curves in \mathbf{T}° . In particular, the intersection number of $\mathbf{a}(\mathbf{m})$ with \mathbf{T} is not zero. Hence $\mathbf{a}(\mathbf{m})$ is a curve in \mathbf{S} which is not in \mathbf{T} .

On the other hand, for a curve γ in $\mathbf{C}(\mathbf{S}) \setminus \mathbf{T}$, since we consider it up to homotopy, we can assume γ has minimal intersections with the curves in \mathbf{T} . Take the product, denoted by \mathbf{m}_γ , of letters corresponding to the arc segments of γ divided by its intersections with \mathbf{T}° in order. Then by Lemma 4.8(1), \mathbf{m}_γ is a word and clearly it is maximal. Moreover, the correspondence between arc segments and letters implies that $\mathbf{m}_{\mathbf{a}(\mathbf{m})} = \mathbf{m}$ and $\mathbf{a}(\mathbf{m}_\gamma) = \gamma$ up to homotopy. Therefore, $\mathbf{m} \mapsto \mathbf{a}(\mathbf{m})$ is the required bijection with $\mathbf{a}(\mathbf{m}^{-1}) = \mathbf{a}(\mathbf{m})^{-1}$. □

Recall from § 2.5 that a maximal word $\mathbf{m} = \omega_m \cdots \omega_1$ is called admissible if the following hold.

- (A1) For each $\omega_i = \varepsilon$ with ε a dashed loop, we have that $\omega_1^{-1} \cdots \omega_{i-1}^{-1} > \omega_m \cdots \omega_{i+1}$, and for each $\omega_i = \varepsilon^{-1}$ with ε a dashed loop, we have that $\omega_1^{-1} \cdots \omega_{i-1}^{-1} < \omega_m \cdots \omega_{i+1}$.
- (A2) If \mathbf{m} contains two punctured letters then $F(\mathbf{m})$ is not a proper power of $F(\mathbf{m}')$ for any maximal word \mathbf{m}' containing two punctured letters, where

$$F(\mathbf{m}) := F(\omega_m) \cdots F(\omega_1) F(\omega_2^{-1}) \cdots F(\omega_{m-1}^{-1}).$$

On the other hand, recall from Definition 3.1 that a pair (γ, κ) is called a tagged curve if the following hold.




FIGURE 11. A once-punctured monogon from a curve.

- (T1) The curve γ does not cut out a once-punctured monogon by a self-intersection (including endpoints), cf. Figure 2.
- (T2) If $\gamma(0), \gamma(1) \in \mathbf{P}$, then the completion $\bar{\gamma}$ is not a proper power of a closed curve in the sense of the multiplication in the fundamental group of \mathbf{S} .

LEMMA 4.12. *Let \mathbf{m} be a maximal word. Then \mathbf{m} satisfies (A1) if and only if $\mathbf{a}(\mathbf{m})$ satisfies (T1); \mathbf{m} satisfies (A2) if and only if $\mathbf{a}(\mathbf{m})$ satisfies (T2).*

Proof. Let $\mathbf{m} = \omega_m \cdots \omega_1$. Note that the curve $\mathbf{a}(\mathbf{m})$ does not satisfy (T1) if and only if it cuts out a once-punctured monogon as in Figure 2. This is equivalent to there being an arc segment $\mathbf{a}(\omega_i)$ of $\mathbf{a}(\mathbf{m})$ with $\omega_i = \varepsilon$ or ε^{-1} (for some dashed loop ε) which has the form as in Figure 11 with the right part being one of the forms in Figure 10. Let $\gamma_1 := \mathbf{a}(\omega_m \cdots \omega_{i+1})$ and $\gamma_2 := \mathbf{a}(\omega_1^{-1} \cdots \omega_{i-1}^{-1})$. If $\omega_i = \varepsilon$, then the orientation of γ is as shown in Figure 11.

By Lemma 4.10, $\omega_m \cdots \omega_{i+1} \geq \omega_1^{-1} \cdots \omega_{i-1}^{-1}$. Similarly, if $\omega_i = \varepsilon^{-1}$ then $\omega_m \cdots \omega_{i+1} \leq \omega_1^{-1} \cdots \omega_{i-1}^{-1}$. This implies that (T1) does not hold for $\mathbf{a}(\mathbf{m})$ if and only if (A1) does not hold for \mathbf{m} .

Now consider the condition (A2). Note that by Lemma 4.8(3) both of the endpoints of $\mathbf{a}(\mathbf{m})$ are punctures if and only if \mathbf{m} has two punctured letters ω_1 and ω_m . In this case, $F(\mathbf{m})$ (as a cycle) corresponds to the completion of $\mathbf{a}(\mathbf{m})$. Hence $\mathbf{a}(\mathbf{m})$ satisfies (T2) if and only if \mathbf{m} satisfies (A2). \square

Let $\mathbf{C}_0(\mathbf{S})$ be the subset of $\mathbf{C}(\mathbf{S}) \setminus \mathbf{T}$ consisting of curves satisfying the conditions (T1) and (T2). That is, $\mathbf{C}_0(\mathbf{S})$ consists of curves γ in $\mathbf{C}(\mathbf{S}) \setminus \mathbf{T}$ such that there exists a tagged curve (γ, κ) for some κ .

By Lemmas 4.11 and 4.12, we have the following.

LEMMA 4.13. *The map $\mathbf{m} \mapsto \mathbf{a}(\mathbf{m})$ is a bijection*

$$\mathbf{a}: \bar{\mathfrak{X}} \longrightarrow \mathbf{C}_0(\mathbf{S}), \tag{4.3}$$

where $\bar{\mathfrak{X}}$ is the set of admissible words up to inverse.

For each $\gamma \in \mathbf{C}_0(\mathbf{S})$, we denote by \mathbf{m}_γ the preimage of γ under the bijection (4.3). Recall from § 2.6 that the algebra $A_{\mathbf{m}_\gamma}$ is generated by the indeterminates x associated with punctured letters in \mathbf{m}_γ with $x^2 = x$. So by Lemma 4.8(3) indeterminates of $A_{\mathbf{m}_\gamma}$ are indexed by endpoints of γ that are punctures.

CONSTRUCTION 4.14. The one-dimensional $A_{\mathbf{m}_\gamma}$ -modules are classified in § 2.6. Using the notation there, each map κ gives a one-dimensional $A_{\mathbf{m}_\gamma}$ -module $N(\gamma, \kappa)$ as follows.

- (i) If neither of the endpoints of γ is a puncture, then $A_{m_\gamma} = \mathbf{k}$. Let $N_{(\gamma, \kappa)} = \mathbf{k}$.
- (ii) If exactly one endpoint of γ is a puncture, then $A_{m_\gamma} = \mathbf{k}[x_a]/(x_a^2 - x_a)$, where $a \in \{0, 1\}$ with $\gamma(a) \in \mathbf{P}$. Let $N_{(\gamma, \kappa)} = \mathbf{k}_{\kappa(a)}$.
- (iii) If both of the endpoints of γ are punctures, then $A_{m_\gamma} = \mathbf{k}\langle x_0, x_1 \rangle / (x_0^2 - x_0, x_1^2 - x_1)$. Let $N_{(\gamma, \kappa)} = \mathbf{k}_{\kappa(0), \kappa(1)}$.

Thus, for each tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S}) \setminus \mathbf{T}^\times$, where \mathbf{T}^\times is the tagged version of \mathbf{T} (see §3.2), there is an associated indecomposable representation

$$M_{(\gamma, \kappa)}^{\mathbf{T}} := M^{\mathbf{T}}(\mathbf{m}_\gamma, N_{(\gamma, \kappa)}) \tag{4.4}$$

in $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ by Construction 2.7 and Theorem 2.9. For $(\gamma, \kappa) \in \mathbf{T}^\times$, let $M_{(\gamma, \kappa)}^{\mathbf{T}} = 0$.

DEFINITION 4.15. Define a map $X^{\mathbf{T}}$ from $\mathbf{C}^\times(\mathbf{S})$ to the set $\mathfrak{S}(\mathbf{T})$ of string objects in $\mathcal{C}(\mathbf{T})$ by

$$X_{(\gamma, \kappa)}^{\mathbf{T}} = \begin{cases} M_{(\gamma, \kappa)}^{\mathbf{T}} & \text{if } (\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S}) \setminus \mathbf{T}^\times, \\ T_{(\gamma, \kappa)} & \text{if } (\gamma, \kappa) \in \mathbf{T}^\times. \end{cases}$$

THEOREM 4.16. *The map $X^{\mathbf{T}}$ is a bijection.*

Proof. It is sufficient to prove that $X^{\mathbf{T}}$ is a bijection from the set $\mathbf{C}^\times(\mathbf{S}) \setminus \mathbf{T}^\times$ to the set of indecomposable representations $M^{\mathbf{T}}(\mathbf{m}, N)$ with $\mathbf{m} \in \bar{\mathfrak{X}}$ and $\dim_{\mathbf{k}} N = 1$. This follows from the bijection (4.3) in Lemma 4.13 and the description of the one-dimensional A_{m_γ} -modules in Construction 4.14. \square

4.3 Flips and mutations

We study \diamond -flips of an admissible triangulation \mathbf{T} in this subsection. Recall that the function $\pi_{\mathbf{T}}$ on \mathbf{T} is defined as follows: if γ is the folded side of a self-folded triangle in \mathbf{T} , then $\pi_{\mathbf{T}}(\gamma)$ is the corresponding remaining side; otherwise, $\pi_{\mathbf{T}}(\gamma) = \gamma$.

DEFINITION 4.17 [FST08, Definition 9.11]. Let \mathbf{T} be an admissible triangulation of \mathbf{S} . The \diamond -flip $f_i(\mathbf{T})$ associated with a curve $i \in \mathbf{T}^o$ is the unique admissible triangulation that shares all curves with \mathbf{T} except for the curves j satisfying $\pi_{\mathbf{T}}(j) = i$.

Note that there are two types of \diamond -flips: when i is not a side of a self-folded triangle, the corresponding \diamond -flip is an ordinary flip, and when i is the remaining side of a self-folded triangle, the corresponding \diamond -flip is a combination of two ordinary flips occurring inside a once-punctured digon (see Figure A.1). Recall that an ordinary flip of a triangulation \mathbf{T}' is a new triangulation \mathbf{T}'' which shares all curves with \mathbf{T}' except for one.

The \diamond -flips of an admissible triangulation \mathbf{T} are indexed by the vertices of the quiver $Q^{\mathbf{T}}$. Define the mutation $(Q', W') = \mu_i(Q^{\mathbf{T}}, W^{\mathbf{T}})$ at a vertex $i \in Q_0^{\mathbf{T}}$ to be (Q'_0, Q'_1, W') $= \mu_i(Q_0, Q_1, W)$ in the sense of [DWZ08] with $Q'_2 = Q_2$. By [FST08, Lab09a],

$$\mu_i(Q^{\mathbf{T}}, W^{\mathbf{T}}) \simeq (Q^{f_i(\mathbf{T})}, W^{f_i(\mathbf{T})})$$

for each $i \in \mathbf{T}^o = Q_0^{\mathbf{T}}$, where f_i is the \diamond -flip associated with i . Then by [KY11], there is an equivalence $\tilde{\mu}_i : \mathcal{C}(\mathbf{T}) \simeq \mathcal{C}(\mathbf{T}')$.

For each tagged curve (γ, κ) , the corresponding object $X_{(\gamma, \kappa)}^{\mathbf{T}}$ in $\mathcal{C}(\mathbf{T})$ is given by the associated representation $M_{(\gamma, \kappa)}^{\mathbf{T}}$ in $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ through the equivalence $F_{\mathbf{T}} : \mathcal{C}(\mathbf{T}) / (T_{\mathbf{T}}) \rightarrow \text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$. Then the equivalence $\tilde{\mu}_i$ should be compatible with some mutation of

representations. However, there is not a well-defined mutation on $\text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$. Instead, we shall use decorated representations and their mutations introduced in [DWZ08]. A decorated representation of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ is a pair (M, V) , where $M \in \text{rep}(Q^{\mathbf{T}}, W^{\mathbf{T}})$ is a usual representation of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ and V is a representation of $(Q_0^{\mathbf{T}}, Q_2^{\mathbf{T}})$ bounded by $\{\varepsilon^2 - \varepsilon \mid \varepsilon \in Q_2^{\mathbf{T}}\}$.

CONSTRUCTION 4.18. Let (γ, κ) be a tagged curve in $\mathbf{C}^\times(\mathbf{S})$. If $(\gamma, \kappa) \notin \mathbf{T}^\times$, define $V_{(\gamma, \kappa)}^{\mathbf{T}} = 0$; if $(\gamma, \kappa) \in \mathbf{T}^\times$, define $V_{(\gamma, \kappa)}^{\mathbf{T}}$ by:

- $V_j = 0$ and $V_{\varepsilon_j} = 0$, for $j \neq \pi_{\mathbf{T}}(\gamma)$;
- $V_{\pi_{\mathbf{T}}(\gamma)} = \mathbf{k}$;
- $V_{\varepsilon_{\pi_{\mathbf{T}}(\gamma)}} = 1 - \kappa(a)$ if there exists a (unique) a with $\gamma(a) \in \mathbf{P}$.

This construction, together with (4.4), gives a decorated representation $(M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}})$ associated with each tagged curve (γ, κ) .

Let (γ, κ) be a tagged curve in $\mathbf{C}^\times(\mathbf{S})$ and $(M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}})$ the corresponding decorated representation. Construct $\mu_i(M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}})$ as in Appendix B. Now we prove that the map $X^{\mathbf{T}}$ from tagged curves to string objects is independent of the choice of the admissible triangulation \mathbf{T} .

THEOREM 4.19. For any two admissible triangulations \mathbf{T} and \mathbf{T}' , there is an equivalence $\Theta: \mathcal{C}(\mathbf{T}) \simeq \mathcal{C}(\mathbf{T}')$ such that $\Theta(X_{(\gamma, \kappa)}^{\mathbf{T}}) \cong X_{(\gamma, \kappa)}^{\mathbf{T}'}$, for every tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})$.

Proof. By Lemma A.2, any two admissible triangulations are connected by a sequence of \diamond -flips. Then using induction, it is sufficient to consider the case of $\mathbf{T}' = f_i(\mathbf{T})$, a \diamond -flip of \mathbf{T} . Recall that there is an equivalence $\tilde{\mu}_i: \mathcal{C}(\mathbf{T}) \simeq \mathcal{C}(\mathbf{T}')$.

We claim that for any tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})$,

$$\mu_i(M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}}) \cong (M_{(\gamma, \kappa)}^{\mathbf{T}'}, V_{(\gamma, \kappa)}^{\mathbf{T}'}). \tag{4.5}$$

Indeed, since $M_{(\gamma, \kappa)}^{\mathbf{T}}$ and $V_{(\gamma, \kappa)}^{\mathbf{T}}$ are constructed locally, we only need to prove that for each segment of γ crossing i , the corresponding decorated representations (M, V) of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ and (M', V') of $(Q^{f_i(\mathbf{T})}, W^{f_i(\mathbf{T})})$ satisfy $\mu_i(M, V) \cong (M', V')$. We list all the possible cases in Tables B.1 and B.2 in Appendix B for the first and second types of \diamond -flips, respectively, up to symmetry. And in the same row of each case, we list the corresponding decorated representations, using Constructions 2.7 and 4.18. Then one can check (4.5) on a case by case basis.

Let $F^{\mathbf{T}}$ denote the equivalence (4.1). Consider the map $\Phi_{\mathbf{T}}$ from the set of (isoclasses of) objects in $\mathcal{C}(\mathbf{T})$ to the set of (isoclasses of) decorated representations of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ defined as follows. For any indecomposable object $X \in \mathcal{C}(\mathbf{T})$ which is not isomorphic to a direct summand of $T_{\mathbf{T}}$, define $\Phi_{\mathbf{T}}(X) \cong (F_{\mathbf{T}}(X), 0)$; for any tagged curve $(\gamma, \kappa) \in \mathbf{T}^\times$, define $\Phi_{\mathbf{T}}(T_{(\gamma, \kappa)}) = (0, V_{(\gamma, \kappa)}^{\mathbf{T}})$. By definition, for any tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})$, $\Phi_{\mathbf{T}}(X_{(\gamma, \kappa)}^{\mathbf{T}}) \cong (M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}})$. Hence by (4.5), we have

$$\mu_i(\Phi_{\mathbf{T}}(X_{(\gamma, \kappa)}^{\mathbf{T}})) = \Phi_{\mathbf{T}'}(X_{(\gamma, \kappa)}^{\mathbf{T}'}).$$

By [Pla11, Proposition 4.1], $\Phi_{\mathbf{T}}$ is a bijection and $\mu_i \Phi_{\mathbf{T}}(X) \simeq \Phi_{\mathbf{T}'} \tilde{\mu}_i(X)$ for any indecomposable object X in $\mathcal{C}(\mathbf{T})$. Hence $\tilde{\mu}_i(X_{(\gamma, \kappa)}^{\mathbf{T}}) \cong X_{(\gamma, \kappa)}^{\mathbf{T}'}$, as required. \square

5. Homological interpretation of marked surfaces


5.1 AR-translation and AR-triangles

Note that we have chosen an admissible triangulation \mathbf{T} and have a bijection $X^{\mathbf{T}} : \mathbf{C}^{\times}(\mathbf{S}) \rightarrow \mathfrak{S}(\mathbf{T})$ from the set of tagged curves to the set of string objects in the cluster category $\mathcal{C}(\mathbf{T})$. The tagged rotation (cf. Definition 3.2 and Figure 3) ρ is a permutation on $\mathbf{C}^{\times}(\mathbf{S})$ while the shift functor $[1]$ in $\mathcal{C}(\mathbf{T})$ gives a permutation on the set $\mathfrak{S}(\mathbf{T})$. We will give a straightforward proof of Theorem 3.10, with a slight generalization to tagged curves.

For a curve γ in $\mathbf{C}(\mathbf{S})$ with $\gamma(1) \in \mathbf{M}$, denote by $[+]\gamma$ the curve obtained from γ by moving $\gamma(1)$ along the boundary anticlockwise to the next marked point; dually, for a curve γ in $\mathbf{C}(\mathbf{S})$ with $\gamma(0) \in \mathbf{M}$, denote by $\gamma[+]$ the curve obtained from γ by moving $\gamma(0)$ along the boundary anticlockwise to the next marked point. We first show the following lemma, where \mathbf{m}_{γ} is the word associated with γ defined by the bijection (4.3) and $[+]\mathbf{m}$ and $\mathbf{m}[+]$ are defined in §2.4. Recall that the set $\mathbf{C}_0(\mathbf{S})$ consists of curves γ in $\mathbf{C}(\mathbf{S}) \setminus \mathbf{T}$ such that there exists a tagged curve (γ, κ) for some κ .

LEMMA 5.1. *If γ is a curve in $\mathbf{C}_0(\mathbf{S})$ with $\gamma(1) \in \mathbf{M}$ such that $[+]\gamma$ is in $\mathbf{C}_0(\mathbf{S})$, then $[+]\mathbf{m}_{\gamma}$ exists and $[+]\mathbf{m}_{\gamma} = \mathbf{m}_{[+]\gamma}$. Dually, if γ is a curve in $\mathbf{C}_0(\mathbf{S})$ with $\gamma(0) \in \mathbf{M}$ such that $\gamma[+]$ is in $\mathbf{C}_0(\mathbf{S})$, then $\mathbf{m}_{\gamma}[+]$ exists and $\mathbf{m}_{\gamma}[+] = \mathbf{m}_{\gamma[+]}$.*

Proof. We only prove the first assertion. By construction, γ and $[+]\gamma$ start at the same point, go through the same way at the beginning and then separate in a triangle as one of the following two forms, where δ is the boundary segment from $\gamma(1)$ to $[+]\gamma(1)$ anticlockwise:



By Lemma 4.10, $\mathbf{m}_{\gamma} > \mathbf{m}_{[+]\gamma}$. Moreover, since by construction γ , $[+]\gamma$ and δ enclose a contractible triangle in the surface (i.e. a triangle which is homotopic to a point), by Lemma 4.10 again there is no curve $\gamma' \in \mathbf{C}_0(\mathbf{S})$ starting at $\gamma(0) = [+]\gamma(0)$ such that $\mathbf{m}_{\gamma} > \mathbf{m}_{\gamma'} > \mathbf{m}_{[+]\gamma}$. Therefore, the bijection (4.3) between curves in $\mathbf{C}_0(\mathbf{S})$ and words in $\bar{\mathfrak{X}}$ implies that $\mathbf{m}_{[+]\gamma}$ is the successor of \mathbf{m}_{γ} , i.e. $[+]\mathbf{m}_{\gamma} = \mathbf{m}_{[+]\gamma}$. \square

THEOREM 5.2. *Under the bijection $X^{\mathbf{T}} : \mathbf{C}^{\times}(\mathbf{S}) \rightarrow \mathfrak{S}(\mathbf{T})$, the tagged rotation ρ on $\mathbf{C}^{\times}(\mathbf{S})$ becomes the shift $[1]$ on the set $\mathfrak{S}(\mathbf{T})$, i.e. we have the following commutative diagram.*

$$\begin{array}{ccc} \mathbf{C}^{\times}(\mathbf{S}) & \xrightarrow{X^{\mathbf{T}}} & \mathfrak{S}(\mathbf{T}) \\ \downarrow \rho & & \downarrow [1] \\ \mathbf{C}^{\times}(\mathbf{S}) & \xrightarrow{X^{\mathbf{T}}} & \mathfrak{S}(\mathbf{T}) \end{array}$$

In particular, restricting to $\mathbf{A}^{\times}(\mathbf{S})$, we get Theorem 3.10.

Proof. Let $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})$ such that neither (γ, κ) nor $\rho(\gamma, \kappa)$ is in \mathbf{T}^\times . So both γ and $\rho(\gamma)$ are in $\mathbf{C}_0(\mathbf{S})$. Using Theorem 2.11, we show that $\tau M_{(\gamma, \kappa)}^\mathbf{T} = M_{\rho(\gamma, \kappa)}^\mathbf{T}$, where there are three cases, as follows.

(1) If both $\gamma(0)$ and $\gamma(1)$ are in \mathbf{M} , then at least one of $[+]\gamma$ and $\gamma[+]$ is in $\mathbf{C}_0(\mathbf{S})$. This is because when they are both in \mathbf{T} , one of (γ, κ) and $\rho(\gamma, \kappa)$ is forced to be in \mathbf{T} , which contradicts our assumption. Then we deduce that $\mathbf{m}_{\rho(\gamma)} = [+]\mathbf{m}_\gamma[+]$ by Lemma 5.1. On the other hand, \mathbf{m}_γ contains no punctured letters by Lemma 4.8(3). Therefore,

$$\tau M_{(\gamma, \kappa)}^\mathbf{T} = M^\mathbf{T}([+]\mathbf{m}_\gamma[+], \mathbf{k}) = M_{\rho(\gamma, \kappa)}^\mathbf{T},$$

where $\kappa = \emptyset$.

(2) If exactly one of $\gamma(1)$ and $\gamma(0)$ is a puncture, assume that $\gamma(0) \in \mathbf{P}$ and $\gamma(1) \in \mathbf{M}$ without loss of generality. Then $\rho(\gamma) = [+]\gamma$, $\mathbf{m}_{\rho(\gamma)} = [+]\mathbf{m}_\gamma$, and

$$\tau M_{(\gamma, \kappa)}^\mathbf{T} = M^\mathbf{T}([+]\mathbf{m}_\gamma, \mathbf{k}_{1-\kappa(0)}) = M_{\rho(\gamma, \kappa)}^\mathbf{T}.$$

Note that the part $\mathbf{k}_{1-\kappa(0)}$ is constructed in Construction 4.14, which is determined by the tagging.

(3) If both $\gamma(0)$ and $\gamma(1)$ are in \mathbf{P} , then $\rho(\gamma) = \gamma$ and

$$\tau M_{(\gamma, \kappa)}^\mathbf{T} = M^\mathbf{T}(\mathbf{m}_\gamma, \mathbf{k}_{1-\kappa(0), 1-\kappa(1)}) = M_{\rho(\gamma, \kappa)}^\mathbf{T}.$$


By [KR07, Lemma in §3.5], the shift [1] in the triangulated category $\mathcal{C}(\mathbf{T})$ gives the AR-translation τ in $\text{rep}(Q^\mathbf{T}, W^\mathbf{T})$. Then we have

$$X_{(\gamma, \kappa)}^\mathbf{T}[1] = X_{\rho(\gamma, \kappa)}^\mathbf{T}, \tag{5.1}$$

for $(\gamma, \kappa) \notin \mathbf{T}^\times \cup \rho^{-1}(\mathbf{T}^\times)$. Furthermore, $M_{(\gamma, \kappa)}^\mathbf{T}$ is a projective representation for $(\gamma, \kappa) \in \rho^{-1}(\mathbf{T}^\times)$ and $P_{(\gamma, \kappa)}[1] = T_{(\gamma, \kappa)}$ for any $(\gamma, \kappa) \in \mathbf{T}^\times$, where $P_{(\gamma, \kappa)}$ is the projective representation associated with the primitive idempotent indexed by (γ, κ) . In particular,

$$\{X_{(\gamma, \kappa)}^\mathbf{T}[1] \mid (\gamma, \kappa) \in \rho^{-1}(\mathbf{T}^\times)\} = \{X_{(\gamma, \kappa)}^\mathbf{T} \mid (\gamma, \kappa) \in \mathbf{T}^\times\}.$$

To finish the proof, we only need to show that $X_{\rho^{\pm 1}(\gamma, \kappa)} = T_{(\gamma, \kappa)}[\pm 1]$ for $(\gamma, \kappa) \in \mathbf{T}^\times$. There are two cases and we only deal with $\rho^{-1}(\gamma, \kappa)$. Note that $\rho(\gamma, \kappa)$ intersects (γ, κ) and the following diagrams show the local situation near this intersection.



This enables us to deduce that $M_{\rho^{-1}(\gamma, \kappa)}^\mathbf{T}$ is not a projective presentation different from $P_{(\gamma, \kappa)}$. So $M_{\rho^{-1}(\gamma, \kappa)}^\mathbf{T} = P_{(\gamma, \kappa)}$ and $X_{\rho^{-1}(\gamma, \kappa)}[1] = T_{(\gamma, \kappa)}$. \square




FIGURE 12. Cutting: five cases.

Remark 5.3. Using the description of AR-sequences in [Gei99, § 5.4] and their relation with AR-triangles in [KZ08, Proposition 4.7], we can describe the AR-triangle ending at $X_{(\gamma, \kappa)}^{\mathbf{T}}$, where (γ, κ) is a tagged curve in $\mathbf{C}^{\times}(\mathbf{S})$ that does not connect two punctures. In the case when (γ, κ) does connect two punctures, the middle term of the AR-triangle is not a string object (and hence we do not have a description).

In the following, let $\bar{\delta}$ be the completion of a curve δ which connects a marked point in \mathbf{M} and a puncture in \mathbf{P} (see the top picture of Figure 1 for the induced orientation). Let $X = X^{\mathbf{T}}$ and

$$X_{(\bar{\delta}, \emptyset)} := X_{(\delta, \emptyset)} \oplus X_{(\delta, \kappa')},$$

where $\kappa'(t) = 1$ for t with $\delta(t) \in \mathbf{P}$.

- If both $\gamma(0)$ and $\gamma(1)$ are in \mathbf{M} (which implies $\kappa = \emptyset$), the AR-triangle ending at $X_{(\gamma, \emptyset)}$ is

$$X_{\rho(\gamma, \emptyset)} \rightarrow X_{(\gamma_{[+], \emptyset})} \oplus X_{([+] \gamma, \emptyset)} \rightarrow X_{(\gamma, \emptyset)} \rightarrow.$$

- If exactly one of $\gamma(0)$ and $\gamma(1)$ is in \mathbf{P} , the AR-triangle ending at $X_{(\gamma, \kappa)}$ is

$$X_{\rho(\gamma, \kappa)} \rightarrow X_{([+] \bar{\gamma}, \emptyset)} \rightarrow X_{(\gamma, \kappa)} \rightarrow.$$

5.2 Cutting and Calabi–Yau reductions

Given a tagged curve $(\gamma, \kappa) \in \mathbf{C}^{\times}(\mathbf{S})$ without self-intersections (i.e. $\text{Int}((\gamma, \kappa), (\gamma, \kappa)) = 0$), let $\mathbf{S}/(\gamma, \kappa)$ be the marked surface obtained from \mathbf{S} by cutting along γ . More precisely, there are five cases as listed below.

- (1) If γ connects two different marked points $M_1, M_2 \in \mathbf{M}$, then the resulting surface is shown as in the first column of Figure 12.




FIGURE 13. The non-trivial bijections for cutting.

- (2) If γ is a loop based on a marked point $M \in \mathbf{M}$, then the resulting surface is shown as in the second column of Figure 12.
- (3) If γ connects a marked point $M \in \mathbf{M}$ and a puncture $P \in \mathbf{P}$, then the resulting surface is shown as in the third column of Figure 12.
- (4) If γ connects two different punctures $P_1, P_2 \in \mathbf{P}$, then the resulting surface is shown as in the fourth column of Figure 12.
- (5) If γ is a loop based on a puncture $P \in \mathbf{P}$, then the resulting surface is shown as in the fifth column of Figure 12.

There is a canonical bijection between the tagged curves in \mathbf{S} that do not intersect (γ, κ) and the tagged curves in $\mathbf{S}/(\gamma, \kappa)$. This bijection is straightforward to see for cases (1), (2), and (5), while there are several non-obvious correspondences between the tagged curves in the cases (3) and (4), which have been shown in Figure 13 (up to tagging). It is easy to check that this bijection preserves the intersection numbers.

Let \mathbf{R} be a subset of an admissible triangulation \mathbf{T} . We define \mathbf{S}/\mathbf{R} to be the marked surface obtained from \mathbf{S} by cutting successively along each tagged curve in \mathbf{R} . Clearly the new marked surface is independent of the choice of orders of tagged curves in \mathbf{R} and it inherits an admissible triangulation $\mathbf{T} \setminus \mathbf{R}$ from \mathbf{S} . Denote by $\mathbf{C}^\times(\mathbf{S})_{\mathbf{R}}$ the set of tagged curves (γ, κ) in $\mathbf{C}^\times(\mathbf{S}) \setminus \mathbf{R}$ that do not intersect the tagged curves in \mathbf{R} . By induction, there is a canonical bijection from $\mathbf{C}^\times(\mathbf{S})_{\mathbf{R}}$ to $\mathbf{C}^\times(\mathbf{S}/\mathbf{R})$. For each tagged curve $(\gamma, \kappa) \in \mathbf{C}^\times(\mathbf{S})_{\mathbf{R}}$, we still use the notation (γ, κ) to denote its image under this bijection.

On the other hand, the object $R = \bigoplus_{(\gamma, \kappa) \in \mathbf{R}} X_{(\gamma, \kappa)}^{\mathbf{T}}$ is a direct summand of the canonical cluster tilting object $T_{\mathbf{T}}$ (cf. (4.2)) in $\mathcal{C}(\mathbf{T})$. Then the Calabi–Yau reduction ${}^\perp R[1]/(R)$ is a 2-Calabi–Yau category with a cluster tilting object $T_{\mathbf{T}} \setminus R$ (see [IY08, § 4]). The following lemma will be used in the proof of the main result in the next subsection. Indeed, this generalizes a result in [MP14] on the relation between Calabi–Yau reduction and cutting to the punctured case.

LEMMA 5.4. Let \mathbf{R} be a subset of an admissible triangulation \mathbf{T} and $R = \bigoplus_{(\gamma, \kappa) \in \mathbf{R}} X_{(\gamma, \kappa)}^{\mathbf{T}}$. Then there is a canonical triangle equivalence $\xi: {}^{\perp}R[1]/(R) \simeq \mathcal{C}(\mathbf{T} \setminus \mathbf{R})$ satisfying

$$\xi(X_{(\gamma, \kappa)}^{\mathbf{T}}) \cong X_{(\gamma, \kappa)}^{\mathbf{T} \setminus \mathbf{R}} \tag{5.2}$$

for each $(\gamma, \kappa) \in \mathbf{C}^{\times}(\mathbf{S})_{\mathbf{R}}$.

Proof. Notice that the corresponding biquiver with potential $(Q^{\mathbf{T} \setminus \mathbf{R}}, W^{\mathbf{T} \setminus \mathbf{R}})$ can be obtained from the biquiver with potential $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ by deleting the vertices corresponding to the tagged curves in \mathbf{R} . By [Kel13, Theorem 7.4], the canonical projection $\pi: \Lambda^{\mathbf{T}} \rightarrow \Lambda^{\mathbf{T} \setminus \mathbf{R}}$ induces the required equivalence $\xi: {}^{\perp}\mathcal{R}[1]/\mathcal{R} \simeq \mathcal{C}(\mathbf{S}/\mathbf{R})$. Furthermore, for each tagged curve $(\gamma, \kappa) \in \mathbf{C}^{\times}(\mathbf{S})_{\mathbf{R}}$, the support of the representation $M_{(\gamma, \kappa)}^{\mathbf{T}}$ does not contain the vertices corresponding to tagged curves in \mathbf{R} . Hence it is preserved by the projection π . Thus we deduce that (5.2) holds. \square

5.3 Intersection numbers

THEOREM 5.5. Let (γ_1, κ_1) and (γ_2, κ_2) be two tagged curves in $\mathbf{C}^{\times}(\mathbf{S})$. Then

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}})$$

for any admissible triangulation \mathbf{T} of \mathbf{S} .

Proof. The proof is given in § 7. \square

5.4 Connectedness of cluster exchange graphs

We apply our main result to study the exchange graph $\text{CEG}(\mathcal{C}(\mathbf{T}))$ of the cluster category $\mathcal{C}(\mathbf{T})$.

COROLLARY 5.6. The correspondence $X^{\mathbf{T}}$ in Theorem 4.16 induces bijections:

- (i) between tagged curves without self-intersections in \mathbf{S} and indecomposable rigid objects in $\mathcal{C}(\mathbf{T})$;
- (ii) between tagged triangulations of \mathbf{S} and cluster tilting objects in $\mathcal{C}(\mathbf{T})$.

Moreover, under the last bijection, flip of tagged triangulations is compatible with mutation of cluster tilting objects.

Proof. By Remark 2.10, for any indecomposable representation M of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$, if there is no tagged curve (γ, κ) such that $M \cong M_{(\gamma, \kappa)}^{\mathbf{T}}$, then $\text{Hom}_{\Lambda^{\mathbf{T}}}(M, \tau M) \neq 0$. This implies that $\text{Ext}_{\mathcal{C}(\mathbf{T})}^1(M, M) \neq 0$. Hence M is not a rigid object in $\mathcal{C}(\mathbf{T})$. Thus, all rigid objects are string objects and hence the proposition follows from Theorem 5.5. \square

THEOREM 5.7. The cluster exchange graph $\text{CEG}(\mathcal{C}(\mathbf{T}))$ is connected and $\text{CEG}(\mathcal{C}(\mathbf{T})) \cong \text{EG}^{\times}(\mathbf{S})$. In particular, each rigid object in $\mathcal{C}(\mathbf{T})$ is reachable.

Proof. The isomorphism $\text{EG}^{\times}(\mathbf{S}) \cong \text{CEG}(\mathcal{C}(\mathbf{T}))$ of graphs follows directly from Corollary 5.6. The connectedness of $\text{EG}^{\times}(\mathbf{S})$ is proved in [FST08, Proposition 7.10]. \square





FIGURE 14. An example.


6. An example

Let \mathbf{S} be a disk with three marked points on the boundary and two punctures in the interior. The corresponding cluster category $\mathcal{C}(\mathbf{T})$ is the classical cluster category of type \tilde{D}_5 . Let \mathbf{T} be the admissible triangulation shown in the top left picture of Figure 14.

Then the associated biquiver with potential $(Q^{\mathbf{T}}, W^{\mathbf{T}})$ introduced in § 4.1 is as shown in the following diagram with $W^{\mathbf{T}} = cba$.



And the associated quiver with potential $(Q_{\mathbf{T}}, W_{\mathbf{T}})$ introduced in § 3.2 is as shown in the following diagram with $W_{\mathbf{T}} = c^{3 \rightarrow 1}ba^{1 \rightarrow 2} + c^{3 \rightarrow 1'}ba^{1' \rightarrow 2}$ (where the terms $c^{3 \rightarrow 1'}ba^{1 \rightarrow 2}$ and $c^{3 \rightarrow 1}ba^{1' \rightarrow 2}$ do not appear in $W_{\mathbf{T}}$ because they are not cycles).



We have $Z = \partial W^{\mathbf{T}} = \{ba, cb, ac\}$. Then the biquiver $Q^{\mathbf{T}}$ with Z is precisely the one in Example 2.2. Use the choice of disjoint subsets given in Example 2.3:

- $L_+(1) = \{c^{-1} > a_{1+} > a\}$ and $L_-(1) = \{\varepsilon_1^{-1} > a_{1-} > \varepsilon_1\}$;
- $L_+(2) = \{a^{-1} > a_{2+} > b\}$ and $L_-(2) = \{a_{2-}\}$;
- $L_+(3) = \{b^{-1} > a_{3+} > c\}$ and $L_-(3) = \{a_{3-} > d\}$;
- $L_+(4) = \{d^{-1} > a_{4+}\}$ and $L_-(4) = \{\varepsilon_4^{-1} > a_{4-} > \varepsilon_4\}$.

Using Construction 4.7, the arc segments corresponding to direct letters are shown in the top pictures of Figure 14, and the inverse of letters corresponds to the reverse of direction of arc segments.

Consider the tagged curves (γ_1, κ_1) and (γ_2, κ_2) shown in the bottom left figure of Figure 14, where $\kappa_1(0) = 0$ and $\kappa_2(0) = 1$. The rotations $\rho(\gamma_1, \kappa_1)$ and $\rho(\gamma_2, \kappa_2)$ are shown in the bottom right picture of Figure 14. The intersection number between (γ_1, κ_1) and (γ_2, κ_2) is

$$\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = 1.$$

On the representation side, since $\rho(\gamma_1, \kappa_1) \in \mathbf{T}^\times$, we have $M_{\rho(\gamma_1, \kappa_1)}^{\mathbf{T}} = 0$ by Construction 4.14. Hence

$$\begin{aligned} \dim \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}}) &= \dim \text{Hom}_{\Lambda^{\mathbf{T}}}(M_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, M_{\rho(\gamma_2, \kappa_2)}^{\mathbf{T}}) \\ &\quad + \dim \text{Hom}_{\Lambda^{\mathbf{T}}}(M_{(\gamma_2, \kappa_2)}^{\mathbf{T}}, M_{\rho(\gamma_1, \kappa_1)}^{\mathbf{T}}) \\ &= \dim \text{Hom}_{\Lambda^{\mathbf{T}}}(M_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, M_{\rho(\gamma_2, \kappa_2)}^{\mathbf{T}}). \end{aligned}$$

The words corresponding to γ_1 and $\rho(\gamma_2)$ are

$$\mathbf{m}_{\gamma_1} = a_{4+} a_{4-}^{-1} \quad \text{and} \quad \mathbf{m}_{\rho(\gamma_2)} = a_{3-} c^{-1} \varepsilon_1^{-1} c d^{-1} a_{4-}^{-1},$$

respectively. These two admissible words are precisely the two given in Example 2.12 and we have $H^{\mathbf{m}_{\gamma_1}, \mathbf{m}_{\rho(\gamma_2)}}$ contains only one element, which is $J = ((0, 2), (0, 2))$ containing one punctured letter a_{4-}^{-1} . Thus, we have

$$\begin{aligned} \dim \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}}) &= \dim_{\mathbf{k}} \text{Hom}_{\Lambda^{\mathbf{T}}}(M_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, M_{\rho(\gamma_2, \kappa_2)}^{\mathbf{T}}) \\ &= \dim_{\mathbf{k}} \text{Hom}_{A_J}(\mathbf{k}_{\kappa_1(0)}, \mathbf{k}_{1-\kappa_2(0)}) \\ &= 1. \end{aligned}$$

7. Proof of Theorem 5.5

7.1 Adding marked points

Recall that we fix an admissible triangulation \mathbf{T} of \mathbf{S} and two tagged curves (γ_1, κ_1) and (γ_2, κ_2) in $\mathbf{C}^\times(\mathbf{S})$.

For technical reasons, we add some new marked points on the boundary of \mathbf{S} as follows. Let E be the set of marked points P in M that are endpoints of γ_1 or γ_2 . For each $P \in E$, we add two marked points on each side of P on the boundary component containing P , denoted by P'_l, P''_l , and P'_r, P''_r , respectively, see Figure 15.

Let \mathbf{S}' be the new marked surface obtained from \mathbf{S} by adding these new marked points. For each $P \in E$, let \mathbf{R}_P be the set of tagged curves as in Figure 15, where the right picture is for the case that P is a unique marked point on a boundary component. Take \mathbf{R} to be the disjoint union of \mathbf{R}_P for $P \in E$. Note that no two tagged curves in \mathbf{R} cross each other and the cutting \mathbf{S}'/\mathbf{R} is canonically homeomorphic to \mathbf{S} . Hence $\mathbf{T} \cup \mathbf{R}$ is an admissible triangulation of \mathbf{S}' .




FIGURE 15. Adding marked points.




FIGURE 16. A partial triangulation associated with new marked points.

LEMMA 7.1. *Under the notation above, we have*

$$\dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}}) = \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T} \cup \mathbf{R})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T} \cup \mathbf{R}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T} \cup \mathbf{R}}).$$

Proof. Let $R = \bigoplus_{(\gamma, \kappa) \in \mathbf{R}} X_{(\gamma, \kappa)}^{\mathbf{T} \cup \mathbf{R}}$, a direct summand of $T_{\mathbf{T} \cup \mathbf{R}}$ in $\mathcal{C}(\mathbf{T} \cup \mathbf{R})$. By Lemma 5.4, there is an equivalence $\xi: {}^\perp R[1]/(R) \simeq \mathcal{C}(\mathbf{T})$ sending $X_{(\gamma_i, \kappa_i)}^{\mathbf{T} \cup \mathbf{R}}$ to $X_{(\gamma_i, \kappa_i)}^{\mathbf{T}}$, $i = 1, 2$ (noting that any tagged curve in \mathbf{R} does not cross (γ_i, κ_i)). Hence,

$$\text{Ext}_{\mathcal{C}(\mathbf{T})}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}}) \cong \text{Ext}_{{}^\perp R[1]/(R)}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T} \cup \mathbf{R}}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T} \cup \mathbf{R}}).$$

By [IY08, Lemma 4.8], for any two objects $X_1, X_2 \in {}^\perp R[1]$, there is an isomorphism

$$\text{Ext}_{{}^\perp R[1]/(R)}^1(X_1, X_2) \cong \text{Ext}_{\mathcal{C}(\mathbf{T} \cup \mathbf{R})}^1(X_1, X_2).$$

Therefore, we get the equality, as required. □

Now we consider another admissible triangulation of \mathbf{S}' . For each $P \in E$, let \mathbf{R}'_P be the set of tagged curves as in Figure 16 and \mathbf{R}' the disjoint union of \mathbf{R}'_P . By Lemma A.1, we can extend \mathbf{R}' to an admissible triangulation \mathbf{T}' of \mathbf{S}' . Due to Theorem 4.19, there is an equivalence $\Theta: \mathcal{C}(\mathbf{T} \cup \mathbf{R}) \simeq \mathcal{C}(\mathbf{T}')$ such that $\Theta(X_{(\gamma_i, \kappa_i)}^{\mathbf{T} \cup \mathbf{R}}) \cong X_{(\gamma_i, \kappa_i)}^{\mathbf{T}'}$, for $i = 1, 2$.

Therefore, this equivalence together with Lemma 7.1 implies that it is sufficient to prove $\text{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) = \dim_{\mathbf{k}} \text{Ext}_{\mathcal{C}(\mathbf{T}')}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}'}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}'})$.

Without loss of generality, fix representatives of γ_1 and γ_2 with minimal intersections with \mathbf{T}' and with each other. We further require that any intersection $\gamma_1 \cap \gamma_2$ does not intersect \mathbf{T}' .

7.2 Normal intersections in the interior

We will use the notion of int-pairs from § 2.6. Recall that for any two admissible words \mathbf{m} and \mathbf{r} , we use $H^{\mathbf{m}, \mathbf{r}}$ to denote the set of int-pairs from \mathbf{m} to \mathbf{r} . For $v = 0, 1, 2$, let $H_v^{\mathbf{m}, \mathbf{r}}$ be the subset of $H^{\mathbf{m}, \mathbf{r}}$ consisting of int-pairs that contain v punctured letters. Then $H^{\mathbf{m}, \mathbf{r}} = H_0^{\mathbf{m}, \mathbf{r}} \cup H_1^{\mathbf{m}, \mathbf{r}} \cup H_2^{\mathbf{m}, \mathbf{r}}$.





FIGURE 17. An interior intersection.

LEMMA 7.2. Let (γ_1, κ_1) and (γ_2, κ_2) be the two tagged curves in the theorem. There is a bijection between $\gamma_1 \cap \gamma_2$ and the disjoint union $H_0^{m_{\gamma_1}, m_{\rho(\gamma_2)}} \cup H_0^{m_{\gamma_2}, m_{\rho(\gamma_1)}}$.

Proof. Consider the triangle Δ_I that contains an intersection I in $\gamma_1 \cap \gamma_2$. Since the triangulation \mathbf{T}' contains \mathbf{R}' which is constructed as in Figure 16, the following situations do not occur:



Therefore, we can deduce that γ_1 and γ_2 share at least one curve in \mathbf{T}' . Hence the curve segments of γ_1 and γ_2 near I has the form in Figure 17 with four possible right (respectively left) parts, where $\{a, b\} = \{1, 2\}$.

We will prove that such an intersection induces an int-pair in $H_0^{m_{\gamma_b}, m_{\rho(\gamma_a)}}$. Let

$$\mathbf{m} = \mathbf{m}_{\gamma_b} = \omega_m \cdots \omega_1 \quad \text{and} \quad \boldsymbol{\tau} = \mathbf{m}_{\gamma_a} = \nu_r \cdots \nu_1.$$

Set $\boldsymbol{\tau}' = \mathbf{m}_{\rho(\gamma_a)}$. We say that an arc segment is in \mathbf{R}' if it is in a triangle formed by curves in \mathbf{R}' and boundary segments, and we say that a letter is from \mathbf{R}' if its corresponding arc segment is in \mathbf{R}' . Note that γ_a and $\rho(\gamma_a)$ intersect the triangulation \mathbf{T}' in the same way except for the parts near endpoints of γ_a in \mathbf{M} , where $\rho(\gamma_a)$ intersects one more curve in \mathbf{R}' than γ_a as in Figure 18. Then $\boldsymbol{\tau}' = x\nu_{r-1} \cdots \nu_2y$, where

$$x = \begin{cases} \nu_r, & \gamma_a(1) \in \mathbf{P} \\ \nu'_{r+1}\nu'_r, & \gamma_a(1) \in \mathbf{M} \end{cases} \quad \text{and} \quad y = \begin{cases} \nu_r, & \gamma_a(0) \in \mathbf{P} \\ \nu'_1\nu'_0, & \gamma_a(0) \in \mathbf{M}, \end{cases}$$

where ν'_0, ν'_1, ν'_r , and ν'_{r+1} are letters corresponding to arc segments of $\rho(\gamma_a)$ in \mathbf{R}' . By Lemma 4.8(1) a letter in $(\boldsymbol{\tau}')^{\pm 1}$ from \mathbf{R}' is not smaller than any letter in $\mathbf{m}^{\pm 1}$ or $\boldsymbol{\tau}^{\pm 1}$ from \mathbf{R}' (cf. also Figure 18). In particular, $(\boldsymbol{\tau}')^{\pm 1}$ and $\mathbf{m}^{\pm 1}/\boldsymbol{\tau}^{\pm 1}$ do not share letters from \mathbf{R}' .

Without loss of generality, we assume that both of the orientations of γ_a and γ_b are from right to left in Figure 17. Then the curve segments in the middle part correspond to the subwords




FIGURE 18. The rotation of γ_a .

$\mathbf{m}_{(i,j)}$ and $\mathbf{r}_{(h,l)}$ for some $0 < i < j < m$, $0 < h < l < r$ and the arc segments in the left (respectively right) part correspond to ω_j and ν_l (respectively ω_i and ν_h). By Lemma 4.8(2), we have $\omega_i^{-1} < \nu_h^{-1}$ and $\omega_j < \nu_l$. Thus $J_I := ((i, j), (h, l))$ is an int-pair in $H_0^{\mathbf{m}, \mathbf{r}}$. Moreover, it is clear that any arc segment of γ_a which connects a marked point in M does not appear in Figure 17. Therefore $\nu_l \mathbf{r}_{(h,l)} \nu_j$ does not have letters from \mathbf{R}' and thus it is also a subword of \mathbf{r}' . Hence J_I is also an int-pair in $H_0^{\mathbf{m}, \mathbf{r}'}$ and we obtain a map

$$J: \gamma_1 \cap \gamma_2 \rightarrow H_0^{\mathbf{m}_{\gamma_1}, \mathbf{m}_{\rho(\gamma_2)}} \cup H_0^{\mathbf{m}_{\gamma_2}, \mathbf{m}_{\rho(\gamma_1)}} \\ I \mapsto J_I.$$

Clearly, different intersections correspond to different int-pairs. Hence the map J is injective. So what is left to show is that the map J is surjective.

Let $J_0 = ((i, j), (h, l))$ be an int-pair in $H_0^{\mathbf{m}, \mathbf{r}'}$. Without loss of generality, we assume that $\mathbf{m}_{(i,j)} = \mathbf{r}'_{(h,l)}$ with $\omega_i^{-1} < \nu'_h$ and $\omega_j < \nu'_l$, where $\nu'_a = \nu_a$ if $1 < a < r$. Since \mathbf{m} and \mathbf{r}' do not share letters from \mathbf{R}' and $\mathbf{m}_{(i,j)} = \mathbf{r}'_{(h,l)}$ contains no punctured letters, we have $\mathbf{r}'_{(h,l)} = \mathbf{r}_{(h,l)}$ and the letters ω_i , ν'_h , ω_j , and ν'_l exist. Since ω_i^{-1} and ν'_h are comparable, by Lemma 4.8(1) their corresponding arc segments are in the same triangle. Hence if ν'_h is from \mathbf{R}' , then so is ω_i^{-1} . This is a contradiction because ν'_h is a letter in $(\mathbf{r}')^{-1}$ and ω_i^{-1} is in \mathbf{m}^{-1} . Hence neither ν'_h nor ω_i is from \mathbf{R}' . Similarly, neither ν'_l nor ω_j is from \mathbf{R}' . In particular, their corresponding arc segments do not connect to a marked point in \mathbf{M} and we have $\nu'_h = \nu_h$ and $\nu'_l = \nu_l$. Then the curve segments corresponding to $\omega_j \mathbf{m}_{(i,j)} \omega_i$ and $\nu_l \mathbf{r}_{(h,l)} \nu_h = \nu'_l \mathbf{r}'_{(h,l)} \nu'_h$ are of the form in Figure 17 (note that the left/right parts are the cases in Figure 10(1), where neither γ_i connect to marked points in \mathbf{M}). By construction, we see that J_0 is in the image of the map J above, as required. \square

For an int-pair J without punctured letters, since $A_J \cong \mathbf{k}$, the unique one-dimensional A_J -module is \mathbf{k} . Then $N_{(\gamma_i, \kappa_i)} \cong \mathbf{k}$ as A_J -modules (cf. Notation 2.13). So we have the following consequence:

$$\text{Int}(\gamma_1, \gamma_2) = \sum_{\{a,b\}=\{1,2\}} \sum_{J \in H_0^{\mathbf{m}_{\gamma_a}, \mathbf{m}_{\rho(\gamma_b)}}} \dim_{\mathbf{k}} \text{Hom}_{A_J}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}). \quad (7.1)$$

7.3 Tagged intersections at the ends

Recall that $\mathfrak{P} = \mathfrak{P}(\gamma_1, \gamma_2) = \{(t_1, t_2) \in \{0, 1\}^2 \mid \gamma_1(t_1) = \gamma_2(t_2) \in \mathbf{P}\}$ is the set of intersections between γ_1 and γ_2 at \mathbf{P} . Let

$$\mathfrak{P}_1 = \{(t_1, t_2) \in \mathfrak{P} \mid \gamma_1|_{t_1 \rightarrow (1-t_1)} \approx \gamma_2|_{t_2 \rightarrow (1-t_2)}\}$$




FIGURE 19. A punctured intersection.

and

$$\mathfrak{T}_1 = \{(t_1, t_2) \in \mathfrak{P}_1 \mid \kappa_1(t_1) \neq \kappa_2(t_2)\}.$$

There is an analogue result of Lemma 7.2 for \mathfrak{P}_1 .

LEMMA 7.3. *There is a bijection between \mathfrak{P}_1 and the disjoint union $H_1^{m_{\gamma_1}, m_{\rho(\gamma_2)}} \cup H_1^{m_{\gamma_2}, m_{\rho(\gamma_1)}}$.*

Proof. Each intersection in \mathfrak{P}_1 has the form in Figure 19 with four possible right parts and $\{a, b\} = \{1, 2\}$. Then the required bijection follows from a proof similar to that of Lemma 7.2. \square

For an intersection $I = (t_1, t_2) \in \mathfrak{P}_1$, let J_I be the corresponding int-pair in $H_1^{m_{\gamma_1}, m_{\rho(\gamma_2)}} \cup H_1^{m_{\gamma_2}, m_{\rho(\gamma_1)}}$. Then we have the associated algebra $A_{J_I} = \mathbf{k}[x]/(x^2 - x)$ and $N_{(\gamma_a, \kappa_a)} \cong \mathbf{k}_{\kappa_a(t_a)}$ and $N_{\rho(\gamma_b, \kappa_b)} \cong \mathbf{k}_{1-\kappa_b(t_b)}$ as A_{J_I} -modules (cf. Notation 2.13), for $\{a, b\} = \{1, 2\}$. Using the formula (2.1), we have

$$\dim_{\mathbf{k}} \text{Hom}_{A_{J_I}}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}) = \begin{cases} 1 & \text{for } (t_1, t_2) \in \mathfrak{T}_1, \\ 0 & \text{for } (t_1, t_2) \notin \mathfrak{T}_1. \end{cases}$$

Hence, we obtain a consequence of Lemma 7.3:

$$|\mathfrak{T}_1| = \sum_{\{a,b\}=\{1,2\}} \sum_{J \in H_1^{m_{\gamma_a}, m_{\rho(\gamma_b)}}} \dim_{\mathbf{k}} \text{Hom}_{A_J}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}). \tag{7.2}$$

Let

$$\mathfrak{P}_2 = \{(t_1, t_2) \in \mathfrak{P} \mid \gamma_1|_{t_1 \rightarrow (1-t_1)} \sim \gamma_2|_{t_2 \rightarrow (1-t_2)}, \gamma_1(1-t_1) = \gamma_2(1-t_2) \in \mathbf{P}\}.$$

Observe that for each $(t_1, t_2) \in \mathfrak{P}_2$, $(1-t_1, 1-t_2)$ is also in \mathfrak{P}_2 . We call them twin intersections (at punctures). Clearly, there is at most one pair of twin intersections in \mathfrak{P}_2 . Let

$$\mathfrak{T}_2 = \{(t_1, t_2) \in \mathfrak{P}_2 \mid \kappa_1(t_1) \neq \kappa_2(t_2), \kappa_1(1-t_1) \neq \kappa_2(1-t_2)\}.$$

Suppose that there is a (unique) pair of twin intersections (t_1, t_2) and $(1-t_1, 1-t_2)$ in \mathfrak{P}_2 . Then both the endpoints of γ_i are in \mathbf{P} and thus $\gamma_i = \rho(\gamma_i)$. Reversing one of γ_i if necessary, assume that $\gamma_1 \sim \gamma_2$. So $m_{\gamma_1} = m_{\gamma_2} = m_{\rho(\gamma_1)} = m_{\rho(\gamma_2)}$. Then this pair of twin intersections induces the int-pairs $J_{1,2} = (m_{\gamma_1}, m_{\rho(\gamma_2)})$ in $H_2^{m_{\gamma_1}, m_{\rho(\gamma_2)}}$ and $J_{2,1} = (m_{\gamma_2}, m_{\rho(\gamma_1)})$ in $H_2^{m_{\gamma_2}, m_{\rho(\gamma_1)}}$,

which are the only ones with two punctured letters. In this case, for any $\{a, b\} = \{1, 2\}$ we have $A_{J_{a,b}} \cong \mathbf{k}\langle x, y \rangle / (x^2 - x, y^2 - y)$ and $N_{(\gamma_a, \kappa_a)} \cong \mathbf{k}_{\kappa_a(t_a), \kappa_a(1-t_a)}$ and $N_{\rho(\gamma_b, \kappa_b)} \cong \mathbf{k}_{1-\kappa_b(t_b), 1-\kappa_b(1-t_b)}$ as $A_{J_{a,b}}$ -modules (cf. Notation 2.13). Using the formula (2.2), we have

$$\dim_{\mathbf{k}} \operatorname{Hom}_{A_{J_{a,b}}}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}) = \begin{cases} 1 & \text{for } (t_1, t_2) \in \mathfrak{I}_2, \\ 0 & \text{for } (t_1, t_2) \notin \mathfrak{I}_2, \end{cases}$$

and hence

$$|\mathfrak{I}_2| = \sum_{\{a,b\}=\{1,2\}} \sum_{J \in H_2^{m\gamma_a, m\rho(\gamma_b)}} \dim_{\mathbf{k}} \operatorname{Hom}_{A_J}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}). \tag{7.3}$$

7.4 Summary

By definition, we have

$$|\mathfrak{I}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2))| = |\mathfrak{I}_1| + |\mathfrak{I}_2|. \tag{7.4}$$

Combining (7.1), (7.2), (7.3), and (7.4), we have

$$\begin{aligned} \operatorname{Int}((\gamma_1, \kappa_1), (\gamma_2, \kappa_2)) &= \sum_{\{a,b\}=\{1,2\}} \sum_{J \in H^{m\gamma_a, m\rho(\gamma_b)}} \dim_{\mathbf{k}} \operatorname{Hom}_{A_J}(N_{(\gamma_a, \kappa_a)}, N_{\rho(\gamma_b, \kappa_b)}) \\ &= \sum_{\{a,b\}=\{1,2\}} \dim_{\mathbf{k}} \operatorname{Hom}_{\Lambda^{\mathbf{T}'}}(M_{(\gamma_a, \kappa_a)}^{\mathbf{T}'}, M_{\rho(\gamma_b, \kappa_b)}^{\mathbf{T}'}) \\ &= \sum_{\{a,b\}=\{1,2\}} \dim_{\mathbf{k}} \operatorname{Hom}_{\Lambda^{\mathbf{T}'}}(M_{(\gamma_a, \kappa_a)}^{\mathbf{T}'}, \tau M_{(\gamma_b, \kappa_b)}^{\mathbf{T}'}) \\ &= \dim_{\mathbf{k}} \operatorname{Ext}_{\mathcal{C}(\mathbf{T}')}^1(X_{(\gamma_1, \kappa_1)}^{\mathbf{T}'}, X_{(\gamma_2, \kappa_2)}^{\mathbf{T}'}). \end{aligned}$$

Here, the second equality follows from Theorem 2.14, the third one follows from the fact that $M_{\rho(\gamma_b, \kappa_b)}^{\mathbf{T}'} = \tau M_{\gamma_b, \kappa_b}^{\mathbf{T}'}$ (Theorem 5.2), and the last one follows from [Pal08, Lemma 3.3]. This finishes the proof. \square

ACKNOWLEDGEMENTS

The authors would like to thank the algebra group of NTNU, especially Aslak Bakke Buan and Idun Reiten, for their enthusiastic help. Further thanks are due to Daniel Labardini-Fragoso, Bangming Deng, Jie Xiao and Bin Zhu for their helpful discussions and comments. The authors would also like to thank the anonymous referees for many comments and suggestions that improved the presentation of the paper.

Appendix A. Admissible triangulations

In this section, we give some results about admissible triangulations which will be used in the paper.

LEMMA A.1. *There is an admissible triangulation, i.e. every puncture is in a self-folded triangle, of any marked surface with non-empty boundary.*

Proof. If the surface \mathbf{S} does not admit any puncture, then any triangulation is admissible. Now let P be a puncture in \mathbf{S} . Use induction on the rank n , starting from the trivial case when $n = 1$ and \mathbf{S} is a once-punctured monogon. Now suppose $n \geq 2$. Consider the curve α with $\operatorname{Int}(\alpha, \alpha) = 0$ and $\alpha(0) = \alpha(1) = M$, which encloses a disk with the puncture P . Cutting along α




FIGURE A.1. A \diamond -flip.

we obtain a surface \mathbf{S}/α (cf. the second column of Figure 12) whose rank is $n - 1$. By inductive assumption we deduce that \mathbf{S}/α , and hence \mathbf{S} admits an admissible triangulation. \square

LEMMA A.2. Any two admissible triangulations are connected by a sequence of \diamond -flips.

Proof. Let \mathbf{T}_i , $i = 1, 2$, be two admissible triangulations. Use induction on the rank n of the marked surface \mathbf{S} , starting with the trivial case when $n = 1$.

Consider a puncture P , which is connected to exactly one marked point M_i in \mathbf{T}_i . If $PM_1 \sim PM_2$, we can delete the self-folded triangles containing P from \mathbf{T}_i and reduce to the case with a smaller rank.

Now suppose that $PM_1 \not\sim PM_2$. Freeze the self-folded triangle in \mathbf{T}_1 containing PM_1 . By inductive assumption for the remaining surface, we can flip \mathbf{T}_1 to a triangulation \mathbf{T}'_1 , with local picture as in the left picture of Figure A.1 with $A = M_2$, $B = M_1$, and the curve $PA \sim PM_2$. Then by one \diamond -flip we can locally flip \mathbf{T}'_1 to another triangulation \mathbf{T}''_1 such that it contains the curve PM_2 in \mathbf{T}_2 , which becomes the $PM_1 \sim PM_2$ case above. \square

Appendix B. Explicit version of Derksen *et al.*'s mutations of decorated representations for biquivers with potential

Let (γ, κ) be a tagged curve in $\mathbf{C}^\times(\mathbf{S})$ and $(M, V) = (M_{(\gamma, \kappa)}^{\mathbf{T}}, V_{(\gamma, \kappa)}^{\mathbf{T}})$ be the corresponding decorated representation of $(Q^{\mathbf{T}}, W^{\mathbf{T}})$, defined in Construction 4.18. For a vertex $i \in Q_0^{\mathbf{T}}$, construct $\mu_i(M, V) = (M', V')$ as follows, where we use \hookrightarrow to denote the canonical inclusion and \twoheadrightarrow the canonical projection.

If there is no dashed loop at i , the subquivers of $Q^{\mathbf{T}}$ and $\mu_i(Q^{\mathbf{T}})$ consisting of all arrows adjacent to i are shown in the second row in Table B.1. Construct $\mu_i(M, V) = (M', V')$ as follows.

- For any $j \neq i$, set $M'_j = M_j$ and $V'_j = V_j$.
- Define

$$M'_i = \frac{\ker M_{\gamma_1} \oplus \ker M_{\gamma_2}}{\text{Im} \begin{pmatrix} M_{\beta_1} \\ M_{\beta_2} \end{pmatrix}} \oplus \text{Im} M_{\gamma_1} \oplus \text{Im} M_{\gamma_2} \oplus \frac{\ker(M_{\alpha_1} \ M_{\alpha_2})}{\text{Im} M_{\gamma_1} \oplus \text{Im} M_{\gamma_2}} \oplus V_i$$

and

$$V'_i = \frac{\ker M_{\beta_1} \cap \ker M_{\beta_2}}{\ker M_{\beta_1} \cap \ker M_{\beta_2} \cap (\text{Im} M_{\alpha_1} + \text{Im} M_{\alpha_2})}.$$

- For any arrow $a \in Q_1^{\mathbf{T}}$ not incident with i , set $M'_a = M_a$.
- For any arrow $\varepsilon \in Q_2^{\mathbf{T}}$, set $M'_\varepsilon = M_\varepsilon$ and $V'_\varepsilon = V_\varepsilon$.

TABLE B.1. The first type of \diamond -flips.

	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & \mathbf{k} \\ \uparrow & \mathbf{k} & \downarrow \\ 0 & \leftarrow & 0 \end{array}$	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & \mathbf{k} \\ 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \end{array}$
	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & 0 \\ \uparrow & \mathbf{k} & \downarrow \\ 0 & \leftarrow & 0 \end{array}$	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \\ 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \end{array}$
	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & 0 \\ \uparrow & \mathbf{k} & \downarrow \\ 0 & \leftarrow & \mathbf{k} \end{array}$	$\begin{array}{ccc} \mathbf{k} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \\ 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & \mathbf{k} \end{array}$
	$\begin{array}{ccc} 0 & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & \mathbf{k} \\ \uparrow & \mathbf{k} & \downarrow \\ 0 & \leftarrow & 0 \end{array}$	$\begin{array}{ccc} 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & \mathbf{k} \\ 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \end{array}$
	$\begin{array}{ccc} 0 & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & \mathbf{k} \\ \uparrow & \mathbf{k} & \downarrow \\ \mathbf{k} & \leftarrow & 0 \end{array}$	$\begin{array}{ccc} 0 & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & \mathbf{k} \\ \mathbf{k} & \begin{array}{c} \xrightarrow{\quad} \\ \leftarrow \end{array} & 0 \end{array}$
	$\begin{array}{ccc} 0 & \begin{array}{c} \downarrow \\ \leftarrow \end{array} & 0 \\ \uparrow & \mathbf{k} & \downarrow \\ 0 & \leftarrow & 0 \end{array}$	$\begin{array}{l} M' = 0 \\ V'_j = \delta_{ij} \mathbf{k} \end{array}$

- Define $M'_{[\beta_2 \alpha_1]} = M_{\beta_2} M_{\alpha_1}$ and $M'_{[\beta_1 \alpha_2]} = M_{\beta_1} M_{\alpha_2}$.
- The map $M'_{\alpha_x^*} : M'_i \rightarrow M'_{s(\alpha_x)}$, is given by the canonical inclusion $\text{Im } M_{\gamma_x} \hookrightarrow M_{s(\alpha_x)}$, and the composition

$$\frac{\ker(M_{\alpha_1} \ M_{\alpha_2})}{\text{Im } M_{\gamma_1} \oplus \text{Im } M_{\gamma_2}} \xrightarrow{a} \ker(M_{\alpha_1} \ M_{\alpha_2}) \hookrightarrow M_{s(\alpha_1)} \oplus M_{s(\alpha_2)} \twoheadrightarrow M_{s(\alpha_x)},$$

where a is a right inverse of the canonical projection

$$\ker(M_{\alpha_1} \ M_{\alpha_2}) \twoheadrightarrow \frac{(\ker(M_{\alpha_1} \ M_{\alpha_2}))}{(\text{Im } M_{\gamma_1} \oplus \text{Im } M_{\gamma_2})}.$$

TABLE B.2. The second type of \diamond -flips.

		$\begin{aligned} M' &= 0 \\ V'_j &= \delta_{ij} \mathbf{k} \\ V'_{\varepsilon_i} &= 1 - \kappa \end{aligned}$

– The map $M'_{\beta_x} : M'_{t(\beta_x)} \rightarrow M'_i$, is given by the map $M_{t(\beta_x)} \xrightarrow{M_{\gamma_x}} \text{Im } M_{\gamma_x}$ and the composition

$$M_{t(\beta_x)} \hookrightarrow M_{t(\beta_1)} \oplus M_{t(\beta_2)} \xrightarrow{b} \ker M_{\gamma_1} \oplus \ker M_{\gamma_2} \twoheadrightarrow \frac{\ker M_{\gamma_1} \oplus \ker M_{\gamma_2}}{\text{Im} \begin{pmatrix} M_{\beta_1} \\ M_{\beta_2} \end{pmatrix}},$$

where b is a left inverse of the inclusion $\ker M_{\gamma_1} \oplus \ker M_{\gamma_2} \hookrightarrow M_{t(\beta_1)} \oplus M_{t(\beta_2)}$.

If there is a dashed loop ε_i at i , the subquivers of $Q^{\mathbf{T}}$ and $\mu_i(Q^{\mathbf{T}})$ consisting of all arrows adjacent to i are shown in the second row in Table B.2. Construct $\mu_i(M, V) = (M', V')$ as follows.

- For any $j \neq i$, set $M'_j = M_j$ and $V'_j = V_j$.
- Define

$$M'_i = \frac{\ker M_\gamma}{\operatorname{Im} M_\beta M_{\varepsilon_i}} \oplus \frac{\ker M_\gamma}{\operatorname{Im} M_\beta(1 - M_{\varepsilon_i})} \oplus \operatorname{Im}(M_\gamma)^{\oplus 2} \oplus \frac{\ker M_{\varepsilon_i} M_\alpha \oplus \ker(1 - M_{\varepsilon_i}) M_\alpha}{\operatorname{Im} M_\gamma} \oplus V_i$$

and

$$V'_i = \frac{\ker M_\beta M_{\varepsilon_i}}{\ker M_\beta M_{\varepsilon_i} \cap \operatorname{Im} M_{\varepsilon_i} M_\alpha} \oplus \frac{\ker M_\beta(1 - M_{\varepsilon_i})}{\ker M_\beta(1 - M_{\varepsilon_i}) \cap \operatorname{Im}(1 - M_{\varepsilon_i}) M_\alpha}.$$

- The map V'_{ε_i} is given by the identity on the first summand.
- For any arrow $a \in Q_1^{\mathbf{T}}$ not incident with i , set $M'_\alpha = M_\alpha$.
- For any arrow $\varepsilon \in Q_2^{\mathbf{T}}$ not incident with i , set $M'_\varepsilon = M_\varepsilon$ and $V'_\varepsilon = V_\varepsilon$.
- Define $M'_{[\beta\alpha]} = M_\beta M_{\varepsilon_i} M_\alpha$.
- The map $M'_{\alpha^*} : M'_i \rightarrow M'_{s(\alpha)}$ is given by the map $(\operatorname{Im} M_\gamma)^{\oplus 2} \xrightarrow{(\iota, \iota)} M_{s(\alpha_x)}$ and the composition

$$\frac{\ker M_{\varepsilon_i} M_\alpha \oplus \ker(1 - M_{\varepsilon_i}) M_\alpha}{\operatorname{Im} M_\gamma} \xrightarrow{a} \ker M_{\varepsilon_i} M_\alpha \oplus \ker(1 - M_{\varepsilon_i}) M_\alpha \hookrightarrow M_{s(\alpha)} \oplus M_{s(\alpha)},$$

where ι is the inclusion and a is a right inverse of the projections $\ker M_{\varepsilon_i} M_\alpha \oplus \ker(1 - M_{\varepsilon_i}) M_\alpha \rightarrow (\ker M_{\varepsilon_i} M_\alpha \oplus \ker(1 - M_{\varepsilon_i}) M_\alpha) / (\operatorname{Im} M_\gamma)$.

- The map $M'_{\beta^*} : M'_{t(\beta)} \rightarrow M'_i$ is given by the map $M_{t(\beta)} \xrightarrow{(M_\gamma, M_\gamma)^\iota} (\operatorname{Im} M_\gamma)^{\oplus 2}$ and the composition

$$M_{t(\beta)} \xrightarrow{b} \ker M_\gamma \xrightarrow{(\pi, \pi')} \frac{\ker M_\gamma}{\operatorname{Im} M_\beta M_{\varepsilon_i}} \oplus \frac{\ker M_\gamma}{\operatorname{Im} M_\beta(1 - M_{\varepsilon_i})},$$

where b is a left inverse of the inclusion $\ker M_\gamma \hookrightarrow M_{t(\beta)}$ and π, π' are the projections.

- The map M'_{ε_i} is given by the identity on $(\ker M_\gamma) / (\operatorname{Im} M_\beta M_{\varepsilon_i})$, the identity on $(\ker M_{\varepsilon_i} M_\alpha) / (\operatorname{Im} M_\gamma)$, the map $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} : \operatorname{Im}(M_\gamma)^{\oplus 2} \rightarrow \operatorname{Im}(M_\gamma)^{\oplus 2}$, and $V_{\varepsilon_i} : V_i \rightarrow V_i$.

By [DWZ08, Corollary 10.12], $\mu_i(M, V)$ is a decorated representation of $\mu_i(Q^{\mathbf{T}}, W^{\mathbf{T}})$ in both cases.

Remark B.1. The first mutation formula above for (M, V) is an explicit version of Derksen *et al.*'s mutation of decorated representations [DWZ08]. The second mutation formula above is the composition of two Derksen *et al.*'s mutations of decorated representations.

REFERENCES

ACCERV13 M. Alim, S. Cecotti, C. Córdova, S. Espahbodi, A. Rastogi and C. Vafa, *BPS quivers and spectra of complete $\mathcal{N} = 2$ quantum field theories*, Comm. Math. Phys. **323** (2013), 1185–1227.

Ami09 C. Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, Ann. Inst. Fourier (Grenoble) **59** (2009), 2525–2590.

ABCP12 I. Assem, T. Brüstle, G. Charbonneau-Jodoin and P. G. Plamondon, *Gentle algebras arising from surface triangulations*, Algebra Number Theory **4** (2012), 201–229.

Bon92 V. M. Bondarenko, *Representations of bundles of semichained sets and their applications*, St. Petersburg Math. J. **3** (1992), 973–996.

BQ15 T. Brüstle and Y. Qiu, *Tagged mapping class groups: Auslander–Reiten translation*, Math. Z. **279** (2015), 1103–1120.

- BZ11 T. Brüstle and J. Zhang, *On the cluster category of a marked surface without punctures*, Algebra Number Theory **5** (2011), 529–566.
- BMRRT06 A. B. Buan, R. Marsh, M. Reineke, I. Reiten and G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), 572–618.
- CCS06 P. Caldero, F. Chapoton and R. Schiffler, *Quivers with relations arising from clusters (A_n case)*, Trans. Amer. Math. Soc. **358** (2006), 1347–1364.
- CS13 I. Canakci and R. Schiffler, *Snake graph calculus and cluster algebras from surfaces*, J. Algebra **382** (2013), 240–281.
- CS15a I. Canakci and R. Schiffler, *Snake graph calculus and cluster algebras from surfaces II: Self-crossing snake graphs*, Math. Z. **281** (2015), 55–102.
- CS15b I. Canakci and R. Schiffler, *Snake graph calculus and cluster algebras from surfaces III: Band graphs and snake rings*, Preprint (2015), [arXiv:1506.01742](https://arxiv.org/abs/1506.01742).
- CS14 I. Canakci and S. Schroll, *Extensions in Jacobian algebras and cluster categories of marked surfaces*, Preprint (2014), [arXiv:1408.2074](https://arxiv.org/abs/1408.2074).
- Cra89 W. Crawley-Boevey, *Functorial filtrations II: Clans and the Gelfand problem*, J. Lond. Math. Soc. (2) **40** (1989), 9–30.
- Den00 B. Deng, *On a problem of Nazarova and Roiter*, Comment. Math. Helv. **75** (2000), 368–409.
- DWZ08 H. Derksen, J. Weyman and A. Zelevinsky, *Quivers with potentials and their representations I: Mutations*, Selecta Math. (N.S.) **14** (2008), 59–119.
- FST12 A. Felikson, M. Shapiro and P. Tumarkin, *Skew-symmetric cluster algebras of finite mutation type*, J. Eur. Math. Soc. (JEMS) **14** (2012), 1135–1180.
- FG06 V. V. Fock and A. B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. **103** (2006), 1–211.
- FG09 V. V. Fock and A. B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), 865–930.
- FST08 S. Fomin, M. Shapiro and D. Thurston, *Cluster algebras and triangulated surfaces, part I: Cluster complexes*, Acta Math. **201** (2008), 83–146.
- FZ02 S. Fomin and A. Zelevinsky, *Cluster algebras I: Foundations*, J. Amer. Math. Soc. **15** (2002), 497–529.
- Gei99 C. Geiß, *Maps between representations of clans*, J. Algebra **218** (1999), 131–164.
- GdlP99 C. Geiß and J. A. de la Peña, *Auslander–Reiten components for clans*, Boll. Soc. Mat. Mexicana **5** (1999), 307–326.
- GLS16 C. Geiß, D. Labardini-Fragoso and J. Schröer, *The representation type of Jacobian algebras*, Adv. Math. **290** (2016), 364–452.
- IL12 G. Irelli and D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces, part III: Tagged triangulations and cluster monomials*, Compos. Math. **148** (2012), 1833–1866.
- IY08 O. Iyama and Y. Yoshino, *Mutations in triangulated categories and rigid Cohen–Macaulay modules*, Invent. Math. **172** (2008), 117–168.
- Kel12 B. Keller, *Cluster algebras and derived categories*, in *Derived categories in algebraic geometry*, EMS Series of Congress Reports (European Mathematical Society, Zürich, 2012), 123–183.
- Kel13 B. Keller, *Deformed Calabi–Yau completions*, J. reine angew. Math. **2013** (2013), 125–180.
- KR07 B. Keller and I. Reiten, *Cluster tilted algebras are Gorenstein and stably Calabi–Yau*, Adv. Math. **211** (2007), 123–151.
- KY11 B. Keller and D. Yang, *Derived equivalences from mutations of quivers with potential*, Adv. Math. **226** (2011), 2118–2168.

- KZ08 S. Koenig and B. Zhu, *From triangulated categories to abelian categories: cluster tilting in a general framework*, *Math. Z.* **258** (2008), 143–160.
- Lab09a D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces*, *Proc. Lond. Math. Soc.* (3) **98** (2009), 797–839.
- Lab09b D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces, Part II: Arc representations*, Preprint (2009), [arXiv:0909.4100](https://arxiv.org/abs/0909.4100).
- Lab16 D. Labardini-Fragoso, *On triangulations, quivers with potentials and mutations*, in *Mexican mathematicians abroad: recent contributions*, Contemporary Mathematics, vol. 657 (American Mathematical Society, Providence, RI, 2016), 103–127.
- MP14 R. J. Marsh and Y. Palu, *Coloured quivers for rigid objects and partial triangulations: the unpunctured case*, *Proc. Lond. Math. Soc.* (3) **108** (2014), 411–440.
- Mil16 M. Mills, *Maximal green sequences for quivers of finite mutation type*, Preprint (2016), [arXiv:1606.03799](https://arxiv.org/abs/1606.03799).
- MSW11 G. Musiker, R. Schiffler and L. Williams, *Positivity for cluster algebras from surfaces*, *Adv. Math.* **227** (2011), 2241–2308.
- MSW13 G. Musiker, R. Schiffler and L. Williams, *Bases for cluster algebras from surfaces*, *Compos. Math.* **149** (2013), 217–263.
- MW13 G. Musiker and L. Williams, *Matrix formulae and skein relations for cluster algebras from surfaces*, *Int. Math. Res. Not. IMRN* **2013** (2012), 2891–2944.
- Pal08 Y. Palu, *Cluster characters for 2-Calabi–Yau triangulated categories*, *Ann. Inst. Fourier (Grenoble)* **58** (2008), 2221–2248.
- Pla11 P.-G. Plamondon, *Cluster algebras via cluster categories with infinite dimensional morphism spaces*, *Compos. Math.* **147** (2011), 1921–1954.
- Qiu16 Y. Qiu, *Decorated marked surfaces: spherical twists versus braid twists*, *Math. Ann.* **365** (2016), 595–633.
- QZ14 Y. Qiu and Y. Zhou, *Decorated marked surfaces II: Intersection numbers and dimensions of Hom's*, Preprint (2014), [arXiv:1411.4003](https://arxiv.org/abs/1411.4003).
- Sch08 R. Schiffler, *A geometric model for cluster categories of type D_n* , *J. Algebraic Combin.* **27** (2008), 1–21.
- ZZZ13 J. Zhang, Y. Zhou and B. Zhu, *Cotorsion pairs in the cluster category of a marked surface*, *J. Algebra* **391** (2013), 209–226.

Yu Qiu yu.qiu@bath.edu

Department of Mathematics, Chinese University of Hong Kong,
Shatin, N.T., Hong Kong

Yu Zhou yu.zhou@ntnu.no

Department of Mathematical Sciences,
Norwegian University of Science and Technology, N-7491,
Trondheim, Norway