

## Original Article

\*These authors contributed equally to this work.

**Cite this article:** Zhao W, Zhu Dao-min, Li Q, Xu X, Zhang Y, Zhang C, Zhu J, Yu Y (2023). Brain function mediates the association between low vitamin D and neurocognitive status in female patients with major depressive disorder. *Psychological Medicine* **53**, 4032–4045. <https://doi.org/10.1017/S0033291722000708>

Received: 5 August 2021  
Revised: 15 February 2022  
Accepted: 22 February 2022  
First published online: 1 April 2022

**Key words:**

Brain function; cognition; fMRI; gender; major depressive disorder; vitamin D

**Author for correspondence:**

Yongqiang Yu,  
E-mail: [cjr.yuyongqiang@vip.163.com](mailto:cjr.yuyongqiang@vip.163.com);  
Jiajia Zhu,  
E-mail: [zhujiajiagraduate@163.com](mailto:zhujiajiagraduate@163.com)

# Brain function mediates the association between low vitamin D and neurocognitive status in female patients with major depressive disorder

Wenming Zhao<sup>1,2,3,\*</sup>, Dao-min Zhu<sup>4,5,6,\*</sup>, Qian Li<sup>1,2,3</sup>, Xiaotao Xu<sup>1,2,3</sup>, Yu Zhang<sup>4,5,6</sup>, Cun Zhang<sup>1,2,3</sup>, Jiajia Zhu<sup>1,2,3</sup> and Yongqiang Yu<sup>1,2,3</sup>

<sup>1</sup>Department of Radiology, The First Affiliated Hospital of Anhui Medical University, Hefei 230022, China; <sup>2</sup>Research Center of Clinical Medical Imaging, Anhui Province, Hefei 230032, China; <sup>3</sup>Anhui Provincial Institute of Translational Medicine, Hefei 230032, China; <sup>4</sup>Department of Sleep Disorders, Affiliated Psychological Hospital of Anhui Medical University, Hefei 230022, China; <sup>5</sup>Hefei Fourth People's Hospital, Hefei 230022, China and <sup>6</sup>Anhui Mental Health Center, Hefei 230022, China

**Abstract**

**Background.** Vitamin D is engaged in various neural processes, with low vitamin D linked to depression and cognitive dysfunction. There are gender differences in depression and vitamin D level. However, the relationship between depression, gender, vitamin D, cognition, and brain function has yet to be determined.

**Methods.** One hundred and twenty-two patients with major depressive disorder (MDD) and 119 healthy controls underwent resting-state functional MRI and fractional amplitude of low-frequency fluctuations (fALFF) was calculated to assess brain function. Serum concentration of vitamin D (SCVD) and cognition (i.e. prospective memory and sustained attention) were also measured.

**Results.** We found a significant group-by-gender interaction effect on SCVD whereby MDD patients showed a reduction in SCVD relative to controls in females but not males. Concurrently, there was a female-specific association of SCVD with cognition and MDD-related fALFF alterations in widespread brain regions. Remarkably, MDD- and SCVD-related fALFF changes mediated the relation between SCVD and cognition in females.

**Conclusion.** Apart from providing insights into the neural mechanisms by which low vitamin D contributes to cognitive impairment in MDD in a gender-dependent manner, these findings might have clinical implications for assignment of female patients with MDD and cognitive dysfunction to adjuvant vitamin D supplementation therapy, which may ultimately advance a precision approach to personalized antidepressant choice.

**Introduction**

Major depressive disorder (MDD) is the single largest contributor to global disability, affecting as many as 300 million people worldwide (World Health Organization, 2017). Despite great efforts, our current understanding of its etiology is still limited given its complex symptomatology and the mixed findings regarding MDD-related brain abnormalities (Zhuo et al., 2019) and antidepressant treatment effects (Cipriani et al., 2018). Mounting evidence demonstrates a bidirectional relationship of dietary profiles with neurobiology and psychiatry, giving rise to the nascent field of nutritional neuroscience (Kiecolt-Glaser, Jaremka, & Hughes, 2014). In this framework, several nutrients (e.g. vitamins, omega-3 polyunsaturated fatty acids, and minerals) have gained considerable attention in depression (Mikkelsen, Stojanovska, & Apostolopoulos, 2016; Parker & Brotchie, 2011; Parker, Brotchie, & Graham, 2017; Wang, Um, Dickerman, & Liu, 2018; Wani, Bhat, & Ara, 2015). Remarkably, vitamin D has been suggested to play a prominent role in the pathogenesis of depression and its treatment (Parker et al., 2017).

The involvement of vitamin D in the development and functioning of the central nervous system has been studied over the last two decades. It is well established that vitamin D is engaged in various neural processes such as the growth and development of neurons as well as the synthesis, release and regulation of neurotransmitters (Eyles, Burne, & McGrath, 2013). Active form of vitamin D can be produced by the brain and vitamin D receptors are widely distributed across distinct brain systems (e.g. the prefrontal cortex and limbic system) (Eyles, Smith, Kinobe, Hewison, & McGrath, 2005), providing a mechanistic account for the role of vitamin D in emotion, cognition, and neuropsychiatric diseases (Di Somma et al., 2017; Eyles et al., 2013; Schlogl & Holick, 2014). In parallel, a large number of clinical studies have

documented an intimate association between low serum concentration of vitamin D (SCVD) and depression (Aghajafari, Letourneau, Mahinpey, Cosic, & Giesbrecht, 2018; Anglin, Samaan, Walter, & McDonald, 2013; Ju, Lee, & Jeong, 2013) along with a beneficial effect of vitamin D supplementation on this disease (Cheng, Huang, & Huang, 2020; Gowda, Mutowo, Smith, Wluka, & Renzaho, 2015). While preliminary, our prior work found that total intracranial volume mediated the relationship between SCVD and depressive symptoms in patients with MDD (Zhu et al., 2019). Nonetheless, the neural substrates underlying the association between vitamin D and depression are rather complex and need further investigation.

There is a significant gender difference in depression characterized by the fact that females are more likely to develop MDD than males (Ferrari et al., 2013; Seedat et al., 2009), yet the exact mechanisms contributing to this gender difference remain poorly understood. Several possible explanations have been proposed, including gender differences in (1) the monoamine systems (Ngun, Ghahramani, Sanchez, Bocklandt, & Vilain, 2011) whose dysfunction may be causally related to MDD (Booij, Van der Does, & Riedel, 2003; Ruhe, Mason, & Schene, 2007), (2) hypothalamus-pituitary-adrenal axis function responsible for stress responses (Bangasser & Valentino, 2014; Fernandez-Guasti, Fiedler, Herrera, & Handa, 2012; Zagni, Simoni, & Colombo, 2016), and/or (3) the levels of gonadal steroid hormones such as estrogens (Fernandez-Guasti et al., 2012; Solomon & Herman, 2009). It is now also clear that there is a link between vitamin D and production/release of gonadal hormones (Kinuta et al., 2000; Lorenzen et al., 2017). Meanwhile, population-based studies have reported consistent findings of lower SCVD in females than males (Choi et al., 2020; Milaneschi et al., 2010; Rhee, Lee, & Ahn, 2020; Song, Kim, Rhee, Youm, & Kim, 2016; Toffanello et al., 2014b) but conflicting findings of gender-dependent associations between SCVD and depressive symptoms (Milaneschi et al., 2010; Rhee et al., 2020; Toffanello et al., 2014b). Despite these previous findings, however, whether and how SCVD relates to MDD in a gender-dependent manner is still an open question.

Depression involves deficits in a variety of cognitive domains including attention, memory, learning, and executive function (Bortolato et al., 2016; Pan et al., 2019). Cognitive deficits are thought to be a core domain of depressive psychopathology and a principal mediator of psychosocial functioning in MDD (Pan et al., 2019). The persistence of cognitive deficits after remission of depressive symptoms has been proved to contribute to the failure in achieving full functional recovery in patients suffering from MDD (Bortolato et al., 2016). Prospective memory (PM) is a multi-phase process entailing intention encoding, maintenance, and retrieval (Cona, Scarpazza, Sartori, Moscovitch, & Bisiacchi, 2015). This complex process involves a plurality of neural networks (Cona et al., 2015; Gonneaud et al., 2014). Likewise, sustained attention is a multicomponent mental faculty supported by distributed neural networks (Fortenbaugh, DeGutis, & Esterman, 2017; Langner & Eickhoff, 2013). There is mounting evidence that cognitive deficits such as PM and sustained attention seem to be typical features of depression (Han et al., 2012; McFarland & Vasterling, 2018; Rock, Roiser, Riedel, & Blackwell, 2014; Zhou et al., 2017). Interestingly, strong evidence has revealed that vitamin D deficiency is associated with worse cognitive abilities in healthy adults (Mayne & Burne, 2019) and low vitamin D gives rise to cognitive decline in the elderly population (Toffanello et al., 2014a). Moreover, Pettersen observed that vitamin D supplementation

significantly improved cognitive performance, particularly among individual who had lower vitamin D status at the baseline (Pettersen, 2017). Together, these previous reports have led to an assumption of a potential association between depression, vitamin D, and cognitive disturbances. Nevertheless, whether depression and vitamin D influence cognition independently or synergistically with each other as well as the underlying neural mechanisms have received scant attention with a few exceptions (Lerner, Sharony, & Miodownik, 2018; Roy, 2021).

Resting-state functional magnetic resonance imaging (rs-fMRI) has emerged as a non-invasive neuroimaging technique to assess spontaneous brain activity based on the blood-oxygen-level-dependent (BOLD) signal (Biswal, Yetkin, Haughton, & Hyde, 1995). As the most commonly used metrics derived from rs-fMRI, amplitude of low-frequency fluctuations (ALFF) (Zang et al., 2007) and its normalized version fractional ALFF (fALFF) (Zou et al., 2008) measure the low-frequency oscillation intensity of BOLD time courses and reflect local neural activity strength. Taking advantage of effectively suppressing non-specific signal components and showing improved sensitivity and specificity relative to ALFF, fALFF has been widely applied to the research of MDD and has enjoyed significant success in unraveling local neural activity abnormalities in this condition (Guo et al., 2013; Lai & Wu, 2015; Qiu et al., 2019; Zhu et al., 2020). In regard to the vitamin D-brain relation, earlier structural MRI research has established a link between SCVD and brain morphology (i.e. volume and thickness) in both healthy and depressed individuals (Ali et al., 2020; Annweiler, Annweiler, Montero-Odasso, Bartha, & Beauchet, 2014; Foucault et al., 2019; Karakis et al., 2016; Plozer et al., 2015; Zhu et al., 2019), leaving the relationship between SCVD and brain function largely unknown. Thus, it would be of great interest to examine the association between SCVD and fALFF modulated by MDD.

Here, we sought to determine the relationship between gender, SCVD, cognition, and fALFF in a large sample of MDD patients and healthy subjects. Building on previous literature, we hypothesized (1) a group by gender interaction for SCVD, (2) a gender-specific association of SCVD with cognition and MDD-related fALFF alterations, and (3) a mediative role of MDD- and SCVD-related fALFF changes in accounting for the relation between SCVD and cognition in a gender-specific fashion.

## Materials and methods

### Study design and setting

A cross-sectional design was employed in this study. Patients were recruited consecutively from Affiliated Psychological Hospital of Anhui Medical University between November 2017 and January 2020. We also enrolled healthy controls (HC) from the local community via poster advertisements.

### Participants

In this study, a total of 244 right-handed subjects were enrolled, including 122 MDD patients and 122 gender- and age-matched HC. Two well-trained clinical psychiatrists confirmed the diagnoses of depression using the MINI-International Neuropsychiatric Interview (M.I.N.I.) in accordance with the International Classification of Diseases (ICD-10) criteria (Sheehan et al., 1998). HC were carefully screened to confirm an absence of any psychiatric illness using the M.I.N.I. The exclusion criteria for

all participants included (1) the presence of other psychiatric disorders such as substance-induced mood disorder, bipolar disorder, anxiety disorders, schizophrenia, substance abuse, or dependence; (2) a history of significant physical or neurological diseases; (3) a history of head injury with loss of consciousness; (4) contraindications for MRI such as pregnancy. Additional exclusion criterion for HC was a family history of major neurological or psychiatric illnesses among their first-degree relatives. In addition, participants with SCVD values greater than mean  $+3 \times$  standard deviation (s.d.) or smaller than mean  $-3 \times$  s.d. (i.e. extreme outliers,  $n = 3$ ) were excluded, resulting in the final sample of 241 participants (122 patients with MDD and 119 HC). Moreover, regular antidepressant medications were received by all patients with MDD, including selective serotonin reuptake inhibitors, serotonin-norepinephrine reuptake inhibitors, and noradrenergic and specific serotonergic antidepressants. This study was approved by the ethics committee of The First Affiliated Hospital of Anhui Medical University. Written informed consent was obtained from all participants after being given a complete description of the study.

### Sample size definition

Since there are no studies directly exploring the relationship between vitamin D and fALFF in patients with MDD, we reviewed previous literature on group comparisons in fALFF and listed the effect sizes (Cohen's  $d$ ) of fALFF changes in online Supplementary Table S1. We found that Cohen's  $d$  of fALFF changes in these previous studies was 0.54–1.56. We performed a statistical power analysis to estimate the required sample size using the software G\*Power (Faul, Erdfelder, Lang, & Buchner, 2007) with the following parameters: test family =  $t$  tests, statistical test = difference between two independent groups, minimum effect size  $d = 0.54$ , significance criterion  $\sigma = 0.05$ , statistical power  $1 - \beta = 0.8$ . The statistical power analysis demonstrated that the required sample size to detect the most subtle inter-group difference (minimum  $d = 0.54$ ) is 55 for each group. Therefore, our sample of 122 patients with MDD and 119 HC used in this study provided sufficient statistical power for analysis.

### Demographic and clinical variables

The demographic and clinical data including gender, age, educational level, body mass index (BMI), illness duration, onset age, episode number, and antidepressant medications were collected by the trained research staff. We used the 24-item Hamilton Rating Scale for Depression (HAMD) (Williams, 1988) and the 14-item Hamilton Rating Scale for Anxiety (HAMA) (Thompson, 2015) to assess the severity of depression and anxiety symptoms. Notably, we used the Chinese versions of the HAMD and HAMA, which have been documented to show great reliability and validity (Zheng *et al.*, 1988) and have been widely applied in the Chinese population (Guo *et al.*, 2015; Lai *et al.*, 2021; Shen *et al.*, 2020; Tong *et al.*, 2021).

The cognitive assessment was done by a clinical psychiatrist trained in neuropsychological testing. PM is defined as the ability to remember to carry out an intended action after a delay without any explicit instruction to do so (Einstein & McDaniel, 1990; McDaniel & Einstein, 2011). PM is typically classified into event-based prospective memory (EBPM; i.e. the ability to remember to carry out an intended action at the occurrence of a certain event) and time-based prospective memory (TBPM; i.e. the ability to

remember to carry out an intended action at a certain time). The schematic representation of the EBPM and TBPM tests is shown in online Supplementary Fig. S1. A computerized version of the Continuous Performance Task-Identical Pairs (CPT-IP) (Cornblatt, Risch, Faris, Friedman, & Erlenmeyer-Kimling, 1988) was used to measure sustained attention. The schematic representation of the CPT-IP design is shown in online Supplementary Fig. S2. The main outcome variables of interest are CPT-IP-2, -3, and -4 reflecting sustained attention scores at increasing memory loads on digit span.

### Serum concentration measurement of vitamin D

After an overnight fasting period, peripheral venous blood samples (2 ml) were collected from all participants in the morning of MRI scanning. Samples were sent to the Department of Clinical Laboratory, Affiliated Psychological Hospital of Anhui Medical University immediately for centrifugation and serum was separated. Vitamin D [25(OH)D] was measured in serum using a chemiluminescence immunoassay technique in a fully automated Maglumi 1000 analyzer (SNIBE Co., Ltd., China). Internal quality control provided by the manufacturer was used to assure quality. SCVD was stratified as follows: 30–100 ng/ml (75–250 nmol/l) as sufficiency, 20–30 ng/ml (50–75 nmol/l) as insufficiency, <20 ng/ml (50 nmol/l) as deficiency (Ringe & Kipshoven, 2012).

### Image acquisition

MRI data were acquired using a 3.0-Tesla MR system (Discovery MR750w, General Electric, Milwaukee, WI, USA) with a 24-channel head coil. During scanning, tight but comfortable foam and earplugs were used to minimize head movement and scanner noise. All subjects were instructed to relax, keep their eyes closed but not fall asleep, think of nothing in particular, and move as little as possible. All participants underwent a high-resolution three-dimensional T1-weighted brain volume sequence with the following parameters: repetition time (TR) = 8.5 ms; echo time (TE) = 3.2 ms; inversion time (TI) = 450 ms; flip angle (FA) = 12°; field of view (FOV) = 256 mm  $\times$  256 mm; matrix size = 256  $\times$  256; slice thickness = 1 mm, no gap; voxel size = 1 mm  $\times$  1 mm  $\times$  1 mm; 188 sagittal slices; and acquisition time = 296 s. Resting-state BOLD fMRI data were acquired using a gradient-echo single-shot echo planar imaging sequence with the following parameters: TR = 2000 ms; TE = 30 ms; FA = 90°; FOV = 220 mm  $\times$  220 mm; matrix size = 64  $\times$  64; slice thickness = 3 mm, slice gap = 1 mm; 35 interleaved axial slices; 185 volumes; and acquisition time = 370 s. Routine T2-weighted images were also collected to exclude any organic brain abnormality. None of the participants was excluded for visually inspected imaging artifacts.

### fMRI data preprocessing and fALFF analysis

Resting-state fMRI data were preprocessed using Statistical Parametric Mapping software (SPM12, <http://www.fil.ion.ucl.ac.uk/spm>) and Data Processing & Analysis for Brain Imaging (DPABI, <http://rfmri.org/dpabi>) (Yan, Wang, Zuo, & Zang, 2016). The first 10 volumes for each subject were discarded, and the remaining volumes were corrected for the acquisition time delay between slices. Realignment was then performed to correct the motion between time points. Head motion parameters were computed by estimating the translation in each direction and

the angular rotation on each axis for each volume. All data were within the defined motion thresholds (i.e. translational or rotational motion parameters  $<2.0$  mm or  $2.0^\circ$ ). We also calculated FD, which indexes the volume-to-volume changes in head position. Several nuisance covariates (the estimated motion parameters based on the Friston-24 model, the linear drift, the white matter signal, the cerebrospinal fluid signal, and the spike volumes with  $FD > 0.5$ ) were regressed out from the data. In the normalization step, individual structural images were firstly co-registered with the mean functional image; the transformed structural images were then segmented and normalized to the Montreal Neurological Institute (MNI) space using a high-level nonlinear warping algorithm, i.e. the diffeomorphic anatomical registration through exponentiated Lie algebra technique (Ashburner, 2007). Finally, each filtered functional volume was spatially normalized to MNI space using the deformation parameters estimated during the above step and resampled into a 3 mm isotropic voxel. After spatial normalization, all data sets were smoothed with a 6 mm full-width at half-maximum Gaussian kernel.

The BOLD time course of each voxel obtained from the preprocessed data was transformed to a frequency domain via a Fast Fourier Transform (FFT), and then the power spectrum was obtained. fALFF was defined as the ratio of the power spectrum in the low-frequency band (0.01–0.1 Hz) to that in the entire frequency range (Zou et al., 2008). For the purpose of standardization, the fALFF value of each voxel was divided by the global mean fALFF value, yielding a standardized fALFF map per subject.

### Statistical methods

The statistical analyses of demographic and clinical data were performed using the SPSS 23.0 software package (SPSS, Chicago, IL, USA). Two-sample *t* tests were used to compare age, years of education, BMI, FD, HAMD, HAMA, EBPM, TBPM, and CPT-IP between patients with MDD and HC. Pearson  $\chi^2$  test was used to test group differences in gender. A threshold of  $p < 0.05$  was considered statistically significant (two-tailed).

As our primary focus was set on the group  $\times$  gender interaction effect on SCVD, we employed a general linear model to test this effect with age and education as nuisance covariates. In case of significant interaction, we conducted post hoc pairwise comparisons in SCVD between patients and HC in females and males, respectively. For a gender showing significant group differences, subsequent analyses examining the associations of SCVD with clinical variables and neuroimaging data were performed in this gender. With respect to clinical variables, we tested their associations with SCVD using partial correlation analyses controlling for age and education. Multiple comparison correction was performed using the false discovery rate (FDR) method with a corrected significance level of  $p < 0.05$ . For neuroimaging data, a voxel-wise two-sample *t* test was initially used to test fALFF differences between patients and HC controlling for age, years of education, and FD. For the voxel-based analysis, multiple comparisons were corrected using the cluster-level family-wise error (FWE) method, resulting in a cluster defining threshold of  $p = 0.001$  and a corrected cluster significance of  $p < 0.05$ . Then, fALFF value of each cluster with a significant group difference was extracted and used for region of interest (ROI)-based correlation analyses with SCVD. To assess their clinical relevance, we further examined the associations of the SCVD-sensitive fALFF with clinical variables. For these ROI-based analyses, age,

education, and FD were included as nuisance covariates and multiple comparisons were corrected by the FDR method.

To investigate the potential relationship among SCVD, neuroimaging parameters, and clinical variables, we tested the SCVD-fALFF-clinical variables mediation model where fALFF mediated the relation between SCVD and clinical variables. The mediation analyses were performed using the PROCESS macro (<http://www.processmacro.org/>) available for SPSS (Hayes, 2009; Hayes, 2014). Only variables that showed a significant correlation with others were considered independent, dependent, or mediating variables in the mediation analyses. Age, education, and FD were included as covariates of no interest. Based on 5000 bootstrap realizations, the significance of mediation effects was assessed by the bootstrap 95% confidence interval (CI) in the way a significant indirect effect is indicated when the bootstrap 95% CI does not include zero.

### Validation analysis

Given a significant effect of BMI on SCVD (Wortsman, Matsuoaka, Chen, Lu, & Holick, 2000), we performed the above analyses after additionally controlling for BMI. To test the specificity of the results, we also repeated our analyses in the gender showing no group effect on SCVD.

## Results

### Group $\times$ gender interaction on SCVD

In favor of our hypothesis, we observed a significant interaction effect ( $F = 6.050$ ,  $p = 0.015$ ) of group  $\times$  gender on SCVD (online Supplementary Fig. S3). Post hoc analyses revealed that patients with MDD showed a significant SCVD reduction relative to HC in females ( $p < 0.001$ ) but not males ( $p > 0.05$ ), suggesting a gender-specific effect. Accordingly, we focused our subsequent analyses on females. Demographic and clinical data of females are listed in Table 1 and described in the online Supplementary materials. Among 82 female patients, 11 (13.41%) were classified as vitamin D insufficiency and 70 (85.37%) as deficiency. In addition, demographic and clinical data of all subjects are shown in online Supplementary Table S2.

### Associations between SCVD and cognition in females

We found significant positive correlations of SCVD with EBPM [partial correlation coefficient ( $pr$ ) = 0.267,  $p < 0.001$ ], TBPM ( $pr = 0.355$ ,  $p < 0.001$ ), and CPT-IP-2 ( $pr = 0.215$ ,  $p = 0.006$ ) in females ( $p < 0.05$ , FDR corrected). No significant correlations were found between SCVD and clinical symptoms (HAMD and HAMA) in female patients with MDD (online Supplementary Table S3).

### Associations between SCVD and fALFF in females

Compared with female HC, female patients with MDD exhibited increased fALFF of the right middle temporal gyrus (MTG), left precuneus, right angular gyrus (ANG), right precuneus, bilateral middle cingulate cortex (MCC), right cuneus and left middle occipital gyrus, and reduced fALFF in the left supramarginal gyrus and left middle frontal gyrus (MFG) ( $p < 0.05$ , cluster-level FWE corrected) (Table 2 and online Supplementary Fig. S4).

The significant correlations between SCVD and MDD-related fALFF alterations in females are illustrated in Fig. 1. SCVD was

**Table 1.** Demographic and clinical data of females

Characteristics	MDD ( <i>n</i> = 82)	HC ( <i>n</i> = 82)	Statistics	<i>p</i> Value
Age (years)	44.35 ± 10.28 (21–62)	44.24 ± 12.75 (21–62)	<i>t</i> = 0.061	0.952
Education (years)	8.45 ± 3.76 (0–16)	11.15 ± 4.63 (3–20)	<i>t</i> = −4.100	<0.001
BMI (kg/m <sup>2</sup> )	22.39 ± 3.66 (13.93–32.44)	22.93 ± 2.67 (15.98–31.22)	<i>t</i> = −1.076	0.284
HAMD	30.06 ± 11.25 (1–52)	1.49 ± 2.97 (0–19)	<i>t</i> = 22.244	<0.001
HAMA	20.68 ± 7.40 (2–35)	1.63 ± 3.31 (0–21)	<i>t</i> = 21.287	<0.001
EBPM	2.28 ± 2.55 (0–8)	5.49 ± 2.82 (0–8)	<i>t</i> = −7.638	<0.001
TBPM	1.89 ± 2.27 (0–6)	5.30 ± 1.45 (0–6)	<i>t</i> = −11.482	<0.001
CPT-IP-2	2.07 ± 1.09 (−0.12 to 4.24)	3.11 ± 0.93 (1.11–4.24)	<i>t</i> = −6.559	<0.001
CPT-IP-3	1.50 ± 0.97 (−0.26 to 4.24)	2.29 ± 1.08 (0.28–4.24)	<i>t</i> = −4.954	<0.001
CPT-IP-4	0.85 ± 0.69 (−0.41 to 3.12)	1.24 ± 0.83 (−0.44 to 3.34)	<i>t</i> = −3.273	0.001
SCVD (nmol/l)	38.87 ± 12.05 (15.95–78.36)	50.91 ± 15.83 (21.50–86.25)	<i>t</i> = −5.940	<0.001
FD (mm)	0.13 ± 0.09 (0.05–0.60)	0.13 ± 0.07 (0.06–0.40)	<i>t</i> = −0.003	0.998
Illness duration (months)	62.74 ± 71.44 (0.30–306)	–	–	–
Onset age (years)	38.95 ± 10.46 (19–55)	–	–	–
Episode number	2.52 ± 2.40 (1–21)	–	–	–
Antidepressant medications		–	–	–
SSRIs	58	–	–	–
SNRIs	19	–	–	–
NaSSA	5	–	–	–

MDD, major depressive disorder; HC, healthy controls; BMI, body mass index; HAMD, Hamilton Rating Scale for Depression; HAMA, Hamilton Rating Scale for Anxiety; EBPM, event-based prospective memory; TBPM, time-based prospective memory; CPT-IP, Continuous Performance Task-Identical Pairs; SCVD, serum concentration of vitamin D; FD, frame-wise displacement; SSRIs, selective serotonin reuptake inhibitors; SNRIs, serotonin norepinephrine reuptake inhibitors; NaSSA, noradrenergic and specific serotonergic antidepressant. Data are expressed as means ± standard deviations. Numbers in parentheses are the range.

**Table 2.** Brain regions showing group fALFF differences in females

Brain regions	Cluster size (voxels)	Peak <i>t</i> values	Coordinates in MNI ( <i>x</i> , <i>y</i> , <i>z</i> )
MDD > HC			
Right middle temporal gyrus	29	4.075	57, −66, −3
Left precuneus	38	4.338	−12, −63, 54
Right angular gyrus	43	5.772	45, −51, 30
Right precuneus	45	4.741	6, −54, 42
Bilateral middle cingulate cortex	48	4.738	6, −33, 33
Right cuneus	75	4.855	18, −84, 36
Left middle occipital gyrus	166	5.475	−24, −72, 39
MDD < HC			
Left supramarginal gyrus	32	−4.818	−63, −33, 36
Left middle frontal gyrus	101	−4.728	−39, 48, 30

fALFF, fractional amplitude of low-frequency fluctuations; MDD, major depressive disorder; HC, healthy controls; MNI, Montreal Neurological Institute.

negatively correlated with fALFF of the right MTG ( $pr = -0.215$ ,  $p = 0.006$ , Fig. 1a), left precuneus ( $pr = -0.265$ ,  $p < 0.001$ , Fig. 1b), right ANG ( $pr = -0.207$ ,  $p = 0.009$ , Fig. 1c), right precuneus ( $pr = -0.228$ ,  $p = 0.004$ , Fig. 1d), bilateral MCC ( $pr = -0.201$ ,  $p = 0.011$ , Fig. 1e), and right cuneus ( $pr = -0.169$ ,  $p = 0.032$ , Fig. 1f), and positively correlated with fALFF of the left MFG ( $pr = 0.248$ ,  $p = 0.002$ , Fig. 1g) in females ( $p < 0.05$ , FDR corrected).

#### Associations between SCVD, fALFF, and cognition in females

EBPM was negatively correlated with fALFF of the right MTG ( $pr = -0.194$ ,  $p = 0.014$ , Fig. 2a), right ANG ( $pr = -0.232$ ,  $p = 0.003$ , Fig. 2b), right precuneus ( $pr = -0.177$ ,  $p = 0.025$ , Fig. 2c), bilateral MCC ( $pr = -0.183$ ,  $p = 0.020$ , Fig. 2d), and right cuneus ( $pr = -0.203$ ,  $p = 0.010$ , Fig. 2e), and positively correlated with fALFF of the left MFG ( $pr = 0.227$ ,  $p = 0.004$ , Fig. 2f) in females

( $p < 0.05$ , FDR corrected). To summarize individual differences in fALFF, a principal component analysis (PCA) was performed to identify latent components underlying the EBPM-related fALFF. Based on the Kaiser–Guttman criterion, components with an eigenvalue  $< 1.5$  were removed. As a consequence, only the first fALFF component that accounted for 41.02% of the variance was retained and extracted for subsequent mediation analysis. In the mediation analysis model, all paths were reported as unstandardized ordinary least squares regression coefficients, namely, total effect of  $X$  on  $Y$  ( $c$ ) = indirect effect of  $X$  on  $Y$  through  $M$  ( $a \times b$ ) + direct effect of  $X$  on  $Y$  ( $c'$ ). We found that the relationship between SCVD and EBPM was significantly mediated by the first fALFF component (indirect effect = 0.0056, *s.e.* = 0.0038, 95% CI 0.0003–0.0154) in females (Fig. 2g).

TBPM was negatively correlated with fALFF of the right MTG ( $pr = -0.336$ ,  $p < 0.001$ , Fig. 3a), left precuneus ( $pr = -0.278$ ,  $p < 0.001$ , Fig. 3b), right ANG ( $pr = -0.269$ ,  $p = 0.001$ , Fig. 3c), right precuneus ( $pr = -0.173$ ,  $p = 0.028$ , Fig. 3d), bilateral MCC ( $pr = -0.295$ ,  $p < 0.001$ , Fig. 3e), and right cuneus ( $pr = -0.321$ ,  $p < 0.001$ , Fig. 3f), and positively correlated with fALFF of the left MFG ( $pr = 0.319$ ,  $p < 0.001$ , Fig. 3g) in females ( $p < 0.05$ , FDR corrected). PCA revealed that the first fALFF component accounted for 42.12% of the variance. Then, we found that the relationship between SCVD and TBPM was significantly mediated by the first fALFF component (indirect effect = 0.0067, *s.e.* = 0.0038, 95% CI 0.0014–0.0159) in females (Fig. 3h).

CPT-IP-2 was negatively correlated with fALFF of the right MTG ( $pr = -0.214$ ,  $p = 0.006$ , Fig. 4a), left precuneus ( $pr = -0.301$ ,  $p < 0.001$ , Fig. 4b), right ANG ( $pr = -0.218$ ,  $p = 0.006$ , Fig. 4c), right precuneus ( $pr = -0.204$ ,  $p = 0.010$ , Fig. 4d), bilateral MCC ( $pr = -0.315$ ,  $p < 0.001$ , Fig. 4e), and right cuneus ( $pr = -0.279$ ,  $p < 0.001$ , Fig. 4f) in females ( $p < 0.05$ , FDR corrected). PCA revealed that the first fALFF component accounted for 48.14% of the variance. Then, we found that the relationship between SCVD and CPT-IP-2 was significantly mediated by the first fALFF component (indirect effect = 0.0040, *s.e.* = 0.0020, 95% CI 0.0011–0.0089) in females (Fig. 4g).

There were no significant correlations of fALFF with other cognitive variables (CPT-IP-3 and CPT-IP-4) in females or with clinical symptoms (HAMD and HAMA) in female patients with MDD (online Supplementary Table S4).

### Validation analysis

After additionally adjusting for BMI, the interaction effect of group  $\times$  gender on SCVD was still significant ( $F = 6.066$ ,  $p = 0.015$ ) and the correlations of SCVD with clinical variables and fALFF remained unchanged (online Supplementary Tables S5 and S6), suggesting no effect of BMI on our findings. In males, no significant correlations were found between SCVD and cognition (online Supplementary Table S7), between SCVD and fALFF (online Supplementary Table S8), and between fALFF and cognition (online Supplementary Table S9), indicating female specificity of our results.


### Discussion

To the best of our knowledge, this is the first study to examine the relationship between gender, SCVD, cognition, and fALFF in a large sample of MDD patients and HC. Three main findings were observed in the present study. First, we found a significant group-by-gender interaction effect on SCVD whereby MDD

patients showed a reduction in SCVD relative to HC in females but not males. Second, there was a female-specific association of SCVD with cognition (i.e. PM and sustained attention) and MDD-related fALFF alterations in widespread brain regions. Finally, MDD- and SCVD-related fALFF changes mediated the relation between SCVD and cognition in females.

A large body of evidence from both cross-sectional and longitudinal studies has suggested a close relationship between circulating levels of vitamin D and depression (Briggs et al., 2019; Milaneschi et al., 2014; Wong, Chin, & Ima-Nirwana, 2018). In line with these previous reports, our data showed that most female patients with MDD exhibited insufficient or deficient circulating vitamin D that was markedly lower than HC, raising the possibility that low vitamin D may have a pivotal role in the pathophysiology of MDD. Moreover, extensive research has evidenced that vitamin D supplementation has the potential for reducing depressive symptoms and/or preventing depression (Alghamdi et al., 2020; Casseb, Kaster, & Rodrigues, 2019; de Koning et al., 2015). Despite these promising findings, further mechanistic studies might help advance the clinical translation of vitamin D supplementation by identifying biological pathways through which vitamin D exerts its effect on depression.

Consistent with prior epidemiological research demonstrating gender differences in vitamin D status (Choi et al., 2020), we found that females had lower SCVD than males both in MDD patients and HC. Factors contributing to lower vitamin D in females are insufficient sunlight exposure, higher BMI, more fat tissue, and more sedentary life (Sezgin, Ozturk, Turkal, & Caykara, 2019). For instance, female urban dwellers tend to adhere to the traditional Chinese aesthetic opinion that values paler skin, such that females often intentionally avoid sunshine or use sun cream when they are outdoors (Yan, Zhang, Cheng, Wang, & Qin, 2019). Using sun cream may exponentially suppress vitamin D synthesis in the skin (Fauschou et al., 2012). By contrast, males prefer to do outdoor activities, increasing their sun exposure and vitamin D synthesis. Moreover, it is well known that females typically exceed males in subcutaneous fat (Eisner et al., 2010), and subcutaneous adipose tissue may store large amounts of vitamin D (Didriksen, Burild, Jakobsen, Fuskevåg, & Jorde, 2015). Therefore, the greater subcutaneous fat in females takes up more vitamin D produced by the skin, resulting in fewer vitamin D molecules entering the blood circulation in females. It is also apparent that females are more likely to develop depression than males (Ferrari et al., 2013; Seedat et al., 2009). Based on these phenomena, one may speculate that lower vitamin D in females might render them more susceptible to genetic and environmental risk factors for MDD. Crucially, our current observation that MDD patients showed a reduction in SCVD relative to HC in females rather than males emphasizes a specific involvement of vitamin D in the pathogenesis of depression in females. The reasons for this are unknown and seem to be multifaceted. Although speculative, a potential explanation is that lower vitamin D may lead to altered gonadal hormone levels that may in turn contribute to the development of depression given an intimate association between vitamin D and production/release of gonadal hormones (Kinuta et al., 2000; Lorenzen et al., 2017). This notion is partially supported by some investigations showing that (1) females are more likely to experience mood disturbances and depression during times of hormonal flux such as puberty, perimenstrual, postpartum, and perimenopausal periods (Ahokas, Kaukoranta, Wahlbeck, & Aito, 2001; Parker & Brotchie, 2004; Solomon & Herman, 2009), (2) the incidence of




**Fig. 1.** Associations between SCVD and fALFF in females. (a–g) On the left, SCVD was correlated with fALFF of brain regions showing group differences; on the right, scatter plots show the correlations between SCVD and fALFF.  $^\#p < 0.05$ , false discovery rate correction for multiple comparisons. The warm color represents increased fALFF and the cold color denotes decreased fALFF in patients with MDD. SCVD, serum concentration of vitamin D; fALFF, fractional amplitude of low-frequency fluctuations; MTG, middle temporal gyrus; ANG, angular gyrus; MCC, middle cingulate cortex; MFG, middle frontal gyrus; L, left; R, right; MDD, major depressive disorder; HC, healthy controls;  $pr$ , partial correlation coefficient.

depression in females during menopause appears to be similar to that in males (Bebbington et al., 1998), and (3) hormone replacement therapy during the perimenopausal period can effectively prevent postmenopausal depression (Gordon & Girdler, 2014).

It is widely accepted that cognitive impairment is a characteristic of MDD (Bortolato et al., 2016; Pan et al., 2019), with persistence of cognitive deficits precluding patients from achieving complete functional recovery (Bortolato et al., 2016). Parallel empirical evidence has linked vitamin D deficiency to poorer cognitive performance in normal adults (Annweiler et al., 2013;

Balton et al., 2012) and has suggested low vitamin D as a significant predictor of cognitive worsening in older adults (Llewellyn, Lang, Langa, & Melzer, 2011; Miller et al., 2015; Pavlovic et al., 2018; Slinin et al., 2012). Besides, a randomized trial revealed that high dose vitamin D supplementation enhanced cognition in healthy adults (Pettersen, 2017). These previous findings jointly work to establish a possible causal relationship between vitamin D changes and cognitive alterations in healthy populations. Coherent with previous reports (Han et al., 2012; McFarland & Vasterling, 2018; Rock et al., 2014; Zhou et al., 2017), we found




**Fig. 2.** Associations between SCVD, fALFF, and EBPM in females. (a–f) Scatter plots show correlations between SCVD-related fALFF and EBPM. (g) Graphical representation of the mediation analysis between SCVD and EBPM with first fALFF component as the mediator: estimates of the mediated ( $a \times b$ ), direct ( $c'$ ), and total ( $c$ ) effects.  $^{\#}p < 0.05$ , false discovery rate correction for multiple comparisons;  $^{**}p < 0.01$ ;  $^{***}p < 0.001$ . SCVD, serum concentration of vitamin D; fALFF, fractional amplitude of low-frequency fluctuations; EBPM, event-based prospective memory; MTG, middle temporal gyrus; ANG, angular gyrus; MCC, middle cingulate cortex; MFG, middle frontal gyrus; MDD, major depressive disorder; HC, healthy controls;  $pr$ , partial correlation coefficient; SE, standard error; CI, confidence interval.

that female MDD patients exhibited cognitive dysfunction in PM and sustained attention, which was associated with lower SCVD. Apart from complementing and extending previous literature by identifying a similar vitamin D–cognition association in a psychiatric condition, this finding may represent an informative clue yielding insights into real-world clinical practice by assignment of female patients with MDD and cognitive dysfunction to adjuvant vitamin D supplementation therapy, which may ultimately advance a precision approach to personalized antidepressant choice.

Female patients with MDD presented with local neural activity abnormalities characterized by altered fALFF in widespread areas of the brain that are distributed across the frontal, temporal, parietal, occipital, and limbic regions. Several previous rs-fMRI

studies have reported fALFF differences between MDD patients and healthy subjects (Huang et al., 2017; Li, Rossbach, Zhang, Liu, & Zhang, 2018; Liu et al., 2013; Liu et al., 2017; Shen et al., 2014; Wang et al., 2012), with some endorsing our finding of increased fALFF in the cuneus and decreased fALFF in the supra-marginal gyrus and MFG in patients (Huang et al., 2017; Liu et al., 2013; Wang et al., 2012), and others contrasting with our observation of increased fALFF in the precuneus, ANG, and MTG in patients (Li et al., 2018; Liu et al., 2017; Shen et al., 2014; Wang et al., 2012). As our group comparison analysis focused solely on females, we identified local neural activity alterations specific to female MDD patients, which may in part explain the commonalities and differences with prior research in mixed patients.






**Fig. 3.** Associations between SCVD, fALFF, and TBPM in females. (a–g) Scatter plots show correlations between SCVD-related fALFF and TBPM. (h) Graphical representation of the mediation analysis between SCVD and TBPM with first fALFF component as the mediator: estimates of the mediated (a × b), direct (c'), and total (c) effects. <sup>#</sup> $p < 0.05$ , false discovery rate correction for multiple comparisons; \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$ . SCVD, serum concentration of vitamin D; fALFF, fractional amplitude of low-frequency fluctuations; TBPM, time-based prospective memory; MTG, middle temporal gyrus; ANG, angular gyrus; MCC, middle cingulate cortex; MFG, middle frontal gyrus; MDD, major depressive disorder; HC, healthy controls;  $pr$ , partial correlation coefficient; s.e., standard error; CI, confidence interval.

The relationship between vitamin D and the brain has been well established, as characterized by the critical involvement of vitamin D in the growth and development of neurons as well as the synthesis, release, and regulation of neurotransmitters (Patrick & Ames, 2015) and the broad spatial distribution of vitamin D receptors across the entire brain (Eyles et al., 2005). Recently, such association has attracted intense interest from researchers in the field of neuroimaging. For example, a wealth of evidence from brain structural MRI studies suggests that healthy subjects have SCVD that is linked to different brain morphological features including global and regional volume and regional cortical thickness (Ali et al., 2020; Annweiler et al., 2014; Foucault et al., 2019; Karakis et al., 2016; Plozer et al.,

2015). In patients with MDD, our previous research revealed that SCVD was correlated with total intracranial volume, which further mediated the association between SCVD and depressive symptoms (Zhu et al., 2019). However, these past investigations place less emphasis on brain function. It is generally assumed that functional measures (e.g. fALFF) are a more specific index of state-related changes in brain function, perhaps more closely tracking the onset and offset of symptom expression, whereas structural measures are more stable and thus may relate to more enduring illness characteristics. Leveraging fALFF, the present study found that most of the MDD-related brain functional abnormalities were correlated with SCVD in females. These findings might have clinical implications for informing a novel



**Fig. 4.** Associations between SCVD, fALFF, and CPT-IP-2 in females. (a–f) Scatter plots show correlations between SCVD-related fALFF and CPT-IP-2. (g) Graphical representation of the mediation analysis between SCVD and CPT-IP-2 with first fALFF component as the mediator: estimates of the mediated (a × b), direct (c'), and total (c) effects. #*p* < 0.05, false discovery rate correction for multiple comparisons; \*\**p* < 0.01. SCVD, serum concentration of vitamin D; fALFF, fractional amplitude of low-frequency fluctuations; CPT-IP, Continuous Performance Task-Identical Pairs; MTG, middle temporal gyrus; ANG, angular gyrus; MCC, middle cingulate cortex; MDD, major depressive disorder; HC, healthy controls; *pr*, partial correlation coefficient; SE, standard error; CI, confidence interval.

conceptualization that vitamin D supplementation may reverse MDD-related functional alterations in a gender-dependent way.

Of note, we found that the MDD- and SCVD-related fALFF alterations were associated with PM and sustained attention in females. Further mediation analyses revealed that these fALFF alterations mediated the relation between SCVD and cognition. The affected brain regions are a widely distributed set of areas within the default mode network (the precuneus, ANG, and MTG), executive control network (the MFG), and visual network (the cuneus), in support of the view that complex cognitive functions are assumed to rely on multiple neural processes arising from functional integration and segregation of distinct brain systems. Our data may shed important light on the potential neurobiological mechanisms whereby lower SCVD contributes to cognitive impairment in female patients suffering from MDD.

Several limitations should be considered in this study. First, our chronic and antidepressant-medicated MDD patients may introduce various confounding factors due to antidepressant medication and illness duration. Future studies would benefit from recruiting a sample of first-episode, medication-naive patients with MDD. Second, patients with MDD and HC were not matched in terms of educational level in this study. Although education was considered a nuisance covariate, we cannot completely exclude its potential effects on our results. Third, causal relationship cannot be inferred from this cross-sectional design. Longitudinal studies with intervention targeted toward elevating SCVD in MDD patients are warranted to establish the direction of causality. Fourth, the diagnoses of depression were determined by two well-trained clinical psychiatrists using the M.I.N.I. according to the ICD-10 criteria and the severity of

depression symptoms was assessed using the HAMD. However, the gold standard interview, such as Composite International Diagnostic Interview (CIDI), is the most preferred. Finally, we failed to collect more relevant information about participants' lifestyle profiles. Further analysis of these data may facilitate the interpretation of our findings.

In conclusion, our data revealed a plausible association between low vitamin D, brain dysfunction, and cognitive impairment in female patients with MDD. These results could help elucidate the neural mechanisms by which low vitamin D contributes to cognitive impairment in MDD in a gender-dependent manner. More generally, these findings might have clinical implications for assignment of female patients with MDD and cognitive dysfunction to adjuvant vitamin D supplementation therapy, which may ultimately advance a precision approach to personalized antidepressant choice.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291722000708>

**Acknowledgements.** We are grateful to study participants for volunteering their time. We greatly appreciate the Chinese psychiatrists and psychologists who have developed the Chinese versions of the HAMD and HAMA, the MINI-International Neuropsychiatric Interview (M.I.N.I.), and the International Classification of Diseases (ICD-10) criteria.

**Financial support.** The study was supported by the National Natural Science Foundation of China (grant numbers: 81801679, 82071905, and 81771817).

**Conflict of interest.** None.

## References

- Aghajafari, F., Letourneau, N., Mahinpey, N., Cosic, N., & Giesbrecht, G. (2018). Vitamin D deficiency and antenatal and postpartum depression: A systematic review. *Nutrients*, *10*(4), 478. doi: 10.3390/nu10040478.
- Ahokas, A., Kaukoranta, J., Wahlbeck, K., & Aito, M. (2001). Estrogen deficiency in severe postpartum depression: Successful treatment with sublingual physiologic 17beta-estradiol: A preliminary study. *The Journal of Clinical Psychiatry*, *62*(5), 332–336. doi: 10.4088/jcp.v62n0504.
- Alghamdi, S., Alsulami, N., Khoja, S., Alsufiani, H., Tayeb, H. O., & Tarazi, F. I. (2020). Vitamin D supplementation ameliorates severity of major depressive disorder. *Journal of Molecular Neuroscience: MN*, *70*(2), 230–235. doi:10.1007/s12031-019-01461-2.
- Ali, P., Labriffe, M., Navasiolava, N., Custaud, M. A., Dinomais, M., & Annweiler, C. (2020). Vitamin D concentration and focal brain atrophy in older adults: A voxel-based morphometric study. *Annals of Clinical and Translational Neurology*, *7*(4), 554–558. doi: 10.1002/acn3.50997.
- Anglin, R. E., Samaan, Z., Walter, S. D., & McDonald, S. D. (2013). Vitamin D deficiency and depression in adults: Systematic review and meta-analysis. *The British Journal of Psychiatry: The Journal of Mental Science*, *202*, 100–107. doi: 10.1192/bjp.bp.111.106666.
- Annweiler, C., Annweiler, T., Montero-Odasso, M., Bartha, R., & Beauchet, O. (2014). Vitamin D and brain volumetric changes: Systematic review and meta-analysis. *Maturitas*, *78*(1), 30–39. doi: 10.1016/j.maturitas.2014.02.013.
- Annweiler, C., Montero-Odasso, M., Llewellyn, D. J., Richard-Devantoy, S., Duque, G., & Beauchet, O. (2013). Meta-analysis of memory and executive dysfunctions in relation to vitamin D. *Journal of Alzheimer's Disease: JAD*, *37*(1), 147–171. doi: 10.3233/JAD-130452.
- Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage*, *38*(1), 95–113. doi: 10.1016/j.neuroimage.2007.07.007.
- Balton, C., Griffith, L. E., Strifler, L., Henderson, M., Patterson, C., Heckman, G., ... Raina, P. (2012). Vitamin D, cognition, and dementia: A systematic review and meta-analysis. *Neurology*, *79*(13), 1397–1405. doi: 10.1212/WNL.0b013e31826c197f.
- Bangasser, D. A., & Valentino, R. J. (2014). Sex differences in stress-related psychiatric disorders: Neurobiological perspectives. *Frontiers in Neuroendocrinology*, *35*(3), 303–319. doi: 10.1016/j.yfrne.2014.03.008.
- Bebbington, P. E., Dunn, G., Jenkins, R., Lewis, G., Brugha, T., Farrell, M., & Meltzer, H. (1998). The influence of age and sex on the prevalence of depressive conditions: Report from the National Survey of Psychiatric Morbidity. *Psychological Medicine*, *28*(1), 9–19. doi: 10.1017/s0033291797006077.
- Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magnetic Resonance in Medicine*, *34*(4), 537–541. doi: 10.1002/mrm.1910340409.
- Booij, L., Van der Does, A. J., & Riedel, W. J. (2003). Monoamine depletion in psychiatric and healthy populations: Review. *Molecular Psychiatry*, *8*(12), 951–973. doi: 10.1038/sj.mp.4001423.
- Bortolato, B., Miskowiak, K. W., Kohler, C. A., Maes, M., Fernandes, B. S., Berk, M., & Carvalho, A. F. (2016). Cognitive remission: A novel objective for the treatment of major depression? *BMC Medicine*, *14*, 9. doi: 10.1186/s12916-016-0560-3.
- Briggs, R., McCarroll, K., O'Halloran, A., Healy, M., Kenny, R. A., & Laird, E. (2019). Vitamin D deficiency is associated with an increased likelihood of incident depression in community-dwelling older adults. *Journal of the American Medical Directors Association*, *20*(5), 517–523. doi: 10.1016/j.jamda.2018.10.006.
- Casseb, G. A. S., Kaster, M. P., & Rodrigues, A. L. S. (2019). Potential role of vitamin D for the management of depression and anxiety. *CNS Drugs*, *33* (7), 619–637. doi: 10.1007/s40263-019-00640-4.
- Cheng, Y. C., Huang, Y. C., & Huang, W. L. (2020). The effect of vitamin D supplement on negative emotions: A systematic review and meta-analysis. *Depression and Anxiety*, *37*(6), 549–564. doi: 10.1002/da.23025.
- Choi, J. H., Lee, B., Lee, J. Y., Kim, C. H., Park, B., Kim, D. Y., ... Park, D. Y. (2020). Relationship between sleep duration, sun exposure, and serum 25-hydroxyvitamin D status: A cross-sectional study. *Scientific Reports*, *10*(1), 4168. doi: 10.1038/s41598-020-61061-8.
- Cipriani, A., Furukawa, T. A., Salanti, G., Chaimani, A., Atkinson, L. Z., Ogawa, Y., ... Geddes, J. R. (2018). Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *The Lancet*, *391*(10128), 1357–1366. doi: 10.1016/s0140-6736(17)32802-7.
- Cona, G., Scarpazza, C., Sartori, G., Moscovitch, M., & Bisiacchi, P. S. (2015). Neural bases of prospective memory: A meta-analysis and the 'Attention to Delayed Intention' (AtoDI) model. *Neuroscience and Biobehavioral Reviews*, *52*, 21–37. doi: 10.1016/j.neubiorev.2015.02.007.
- Cornblatt, B. A., Risch, N. J., Faris, G., Friedman, D., & Erlenmeyer-Kimling, L. (1988). The Continuous Performance Test, identical pairs version (CPT-IP): I. New findings about sustained attention in normal families. *Psychiatry Research*, *26*(2), 223–238. doi: 10.1016/0165-1781(88)90076-5.
- de Koning, E. J., van Schoor, N. M., Penninx, B. W., Elders, P. J., Heijboer, A. C., Smit, J. H., ... Lips, P. (2015). Vitamin D supplementation to prevent depression and poor physical function in older adults: Study protocol of the D-Vitaal study, a randomized placebo-controlled clinical trial. *BMC Geriatrics*, *15*, 151. doi: 10.1186/s12877-015-0148-3.
- Didriksen, A., Burild, A., Jakobsen, J., Fuskevåg, O. M., & Jorde, R. (2015). Vitamin D3 increases in abdominal subcutaneous fat tissue after supplementation with vitamin D3. *European Journal of Endocrinology*, *172*(3), 235–241. doi: 10.1530/eje-14-0870.
- Di Somma, C., Scarano, E., Barrea, L., Zhukouskaya, V. V., Savastano, S., Mele, C., ... Marzullo, P. (2017). Vitamin D and neurological diseases: An endocrine view. *International Journal of Molecular Sciences*, *18*(11), 2482. doi: 10.3390/ijms18112482.
- Einstein, G. O., & McDaniel, M. A. (1990). Normal aging and prospective memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *16*(4), 717–726. doi: 10.1037//0278-7393.16.4.717.
- Eisner, B. H., Zargooshi, J., Berger, A. D., Cooperberg, M. R., Doyle, S. M., Sheth, S., & Stoller, M. L. (2010). Gender differences in subcutaneous and perirenal fat distribution. *Surgical and Radiologic Anatomy: SRA*, *32*(9), 879–882. doi: 10.1007/s00276-010-0692-7.
- Eyles, D. W., Burne, T. H., & McGrath, J. J. (2013). Vitamin D, effects on brain development, adult brain function and the links between low levels of

- vitamin D and neuropsychiatric disease. *Frontiers in Neuroendocrinology*, 34(1), 47–64. doi: 10.1016/j.yfrne.2012.07.001.
- Eyles, D. W., Smith, S., Kinobe, R., Hewison, M., & McGrath, J. J. (2005). Distribution of the vitamin D receptor and 1 alpha-hydroxylase in human brain. *Journal of Chemical Neuroanatomy*, 29(1), 21–30. doi: 10.1016/j.jchemneu.2004.08.006.
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G\*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17695343>.
- Faurschou, A., Beyer, D. M., Schmedes, A., Bogh, M. K., Philipsen, P. A., & Wulf, H. C. (2012). The relation between sunscreen layer thickness and vitamin D production after ultraviolet B exposure: A randomized clinical trial. *The British Journal of Dermatology*, 167(2), 391–395. doi: 10.1111/j.1365-2133.2012.11004.x.
- Fernandez-Guasti, A., Fiedler, J. L., Herrera, L., & Handa, R. J. (2012). Sex, stress, and mood disorders: At the intersection of adrenal and gonadal hormones. *Hormone and Metabolic Research = Hormon- und Stoffwechselforschung = Hormones et métabolisme*, 44(8), 607–618. doi: 10.1055/s-0032-1312592.
- Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J., ... Whiteford, H. A. (2013). Burden of depressive disorders by country, sex, age, and year: Findings from the global burden of disease study 2010. *PLoS Medicine*, 10(11), e1001547. doi: 10.1371/journal.pmed.1001547.
- Fortenbaugh, F. C., DeGutis, J., & Esterman, M. (2017). Recent theoretical, neural, and clinical advances in sustained attention research. *Annals of the New York Academy of Sciences*, 1396(1), 70–91. doi: 10.1111/nyas.13318.
- Foucault, G., Duval, G. T., Simon, R., Beauchet, O., Dinomais, M., & Annweiler, C. (2019). Serum vitamin D and cingulate cortex thickness in older adults: Quantitative MRI of the brain. *Current Alzheimer Research*, 16(11), 1063–1071. doi: 10.2174/1567205016666191113124356.
- Gonneaud, J., Rauchs, G., Groussard, M., Landeau, B., Mezenge, F., de La Sayette, V., ... Desgranges, B. (2014). How do we process event-based and time-based intentions in the brain? An fMRI study of prospective memory in healthy individuals. *Human Brain Mapping*, 35(7), 3066–3082. doi: 10.1002/hbm.22385.
- Gordon, J. L., & Girdler, S. S. (2014). Hormone replacement therapy in the treatment of perimenopausal depression. *Current Psychiatry Reports*, 16(12), 517. doi: 10.1007/s11920-014-0517-1.
- Gowda, U., Mutowo, M. P., Smith, B. J., Wluka, A. E., & Renzaho, A. M. (2015). Vitamin D supplementation to reduce depression in adults: Meta-analysis of randomized controlled trials. *Nutrition*, 31(3), 421–429. doi: 10.1016/j.nut.2014.06.017.
- Guo, T., Xiang, Y. T., Xiao, L., Hu, C. Q., Chiu, H. F., Ungvari, G. S., ... Wang, G. (2015). Measurement-based care versus standard care for major depression: A randomized controlled trial with blind raters. *The American Journal of Psychiatry*, 172(10), 1004–1013. doi: 10.1176/appi.ajp.2015.14050652.
- Guo, W., Liu, F., Zhang, J., Zhang, Z., Yu, L., Liu, J., ... Xiao, C. (2013). Dissociation of regional activity in the default mode network in first-episode, drug-naive major depressive disorder at rest. *Journal of Affective Disorders*, 151(3), 1097–1101. doi: 10.1016/j.jad.2013.09.003.
- Han, G., Klimes-Dougan, B., Jepsen, S., Ballard, K., Nelson, M., Hour, A., ... Cullen, K. (2012). Selective neurocognitive impairments in adolescents with major depressive disorder. *Journal of Adolescence*, 35(1), 11–20. doi: 10.1016/j.adolescence.2011.06.009.
- Hayes, A. F. (2009). Beyond Baron and Kenny: Statistical mediation analysis in the new millennium. *Communication Monographs*, 76(4), 408–420. doi: 10.1080/03637750903310360.
- Hayes, A. F. (2014). Introduction to mediation, moderation, and conditional process analysis: A regression-based approach. *Journal of Educational Measurement*, 51(3), 335–337.
- Huang, M., Lu, S., Yu, L., Li, L., Zhang, P., Hu, J., ... Xu, Y. (2017). Altered fractional amplitude of low frequency fluctuation associated with cognitive dysfunction in first-episode drug-naive major depressive disorder patients. *BMC Psychiatry*, 17(1), 11. doi: 10.1186/s12888-016-1190-1.
- Ju, S. Y., Lee, Y. J., & Jeong, S. N. (2013). Serum 25-hydroxyvitamin D levels and the risk of depression: A systematic review and meta-analysis. *The Journal of Nutrition, Health & Aging*, 17(5), 447–455. doi: 10.1007/s12603-012-0418-0.
- Karakis, I., Pase, M. P., Beiser, A., Booth, S. L., Jacques, P. F., Rogers, G., ... Seshadri, S. (2016). Association of serum vitamin D with the risk of incident dementia and subclinical indices of brain aging: The Framingham heart study. *Journal of Alzheimer's Disease: JAD*, 51(2), 451–461. doi: 10.3233/JAD-150991.
- Kiecolt-Glaser, J. K., Jaremka, L. M., & Hughes, S. (2014). Psychiatry and social nutritional neuroscience. *World Psychiatry*, 13(2), 151–152. doi: 10.1002/wps.20127.
- Kinuta, K., Tanaka, H., Moriwake, T., Aya, K., Kato, S., & Seino, Y. (2000). Vitamin D is an important factor in estrogen biosynthesis of both female and male gonads. *Endocrinology*, 141(4), 1317–1324. doi: 10.1210/endo.141.4.7403.
- Lai, C. H., & Wu, Y. T. (2015). The patterns of fractional amplitude of low-frequency fluctuations in depression patients: The dissociation between temporal regions and fronto-parietal regions. *Journal of Affective Disorders*, 175, 441–445. doi: 10.1016/j.jad.2015.01.054.
- Lai, W. T., Deng, W. F., Xu, S. X., Zhao, J., Xu, D., Liu, Y. H., ... Rong, H. (2021). Shotgun metagenomics reveals both taxonomic and tryptophan pathway differences of gut microbiota in major depressive disorder patients. *Psychological Medicine*, 51(1), 90–101. doi: 10.1017/S0033291719003027.
- Langner, R., & Eickhoff, S. B. (2013). Sustaining attention to simple tasks: A meta-analytic review of the neural mechanisms of vigilant attention. *Psychological Bulletin*, 139(4), 870–900. doi: 10.1037/a0030694.
- Lerner, P. P., Sharony, L., & Miodownik, C. (2018). Association between mental disorders, cognitive disturbances and vitamin D serum level: Current state. *Clinical Nutrition ESPEN*, 23, 89–102. doi: 10.1016/j.clnesp.2017.11.011.
- Li, G., Rossbach, K., Zhang, A., Liu, P., & Zhang, K. (2018). Resting-state functional changes in the precuneus within first-episode drug-naive patients with MDD. *Neuropsychiatric Disease and Treatment*, 14, 1991–1998. doi: 10.2147/NDT.S168060.
- Liu, C. H., Ma, X., Yuan, Z., Song, L. P., Jing, B., Lu, H. Y., ... Wang, C. Y. (2017). Decreased resting-state activity in the precuneus is associated with depressive episodes in recurrent depression. *The Journal of Clinical Psychiatry*, 78(4), e372–e382. doi: 10.4088/JCP.15m10022.
- Liu, F., Guo, W., Liu, L., Long, Z., Ma, C., Xue, Z., ... Chen, H. (2013). Abnormal amplitude low-frequency oscillations in medication-naive, first-episode patients with major depressive disorder: A resting-state fMRI study. *Journal of Affective Disorders*, 146(3), 401–406. doi: 10.1016/j.jad.2012.10.001.
- Llewellyn, D. J., Lang, I. A., Langa, K. M., & Melzer, D. (2011). Vitamin D and cognitive impairment in the elderly U.S. population. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 66(1), 59–65. doi: 10.1093/gerona/glq185.
- Lorenzen, M., Boisen, I. M., Mortensen, L. J., Lanske, B., Juul, A., & Blomberg Jensen, M. (2017). Reproductive endocrinology of vitamin D. *Molecular and Cellular Endocrinology*, 453, 103–112. doi: 10.1016/j.mce.2017.03.023.
- Mayne, P. E., & Burne, T. H. J. (2019). Vitamin D in synaptic plasticity, cognitive function, and neuropsychiatric illness. *Trends in Neurosciences*, 42(4), 293–306. doi: 10.1016/j.tins.2019.01.003.
- McDaniel, M. A., & Einstein, G. O. (2011). The neuropsychology of prospective memory in normal aging: A componential approach. *Neuropsychologia*, 49(8), 2147–2155. doi: 10.1016/j.neuropsychologia.2010.12.029.
- McFarland, C. P., & Vasterling, J. J. (2018). Prospective memory in depression: Review of an emerging field. *Archives of Clinical Neuropsychology*, 33(7), 912–930. doi: 10.1093/arclin/axx118.
- Mikkelsen, K., Stojanovska, L., & Apostolopoulos, V. (2016). The effects of vitamin B in depression. *Current Medicinal Chemistry*, 23(38), 4317–4337. doi: 10.2174/0929867323666160920110810.
- Milaneschi, Y., Hoogendijk, W., Lips, P., Heijboer, A. C., Schoevers, R., van Hemert, A. M., ... Penninx, B. W. (2014). The association between low vitamin D and depressive disorders. *Molecular Psychiatry*, 19(4), 444–451. doi: 10.1038/mp.2013.36.
- Milaneschi, Y., Shardell, M., Corsi, A. M., Vazzana, R., Bandinelli, S., Guralnik, J. M., & Ferrucci, L. (2010). Serum 25-hydroxyvitamin D and depressive symptoms in older women and men. *The Journal of Clinical*

- Endocrinology and Metabolism*, 95(7), 3225–3233. doi: 10.1210/jc.2010-0347.
- Miller, J. W., Harvey, D. J., Beckett, L. A., Green, R., Farias, S. T., Reed, B. R., ... DeCarli, C. (2015). Vitamin D status and rates of cognitive decline in a multiethnic cohort of older adults. *JAMA Neurology*, 72(11), 1295–1303. doi: 10.1001/jamaneurol.2015.2115.
- Ngun, T. C., Ghahramani, N., Sanchez, F. J., Bocklandt, S., & Vilain, E. (2011). The genetics of sex differences in brain and behavior. *Frontiers in Neuroendocrinology*, 32(2), 227–246. doi: 10.1016/j.yfrne.2010.10.001.
- Pan, Z., Park, C., Brietzke, E., Zuckerman, H., Rong, C., Mansur, R. B., ... McIntyre, R. S. (2019). Cognitive impairment in major depressive disorder. *CNS Spectrums*, 24(1), 22–29. doi: 10.1017/S1092852918001207.
- Parker, G., & Brotchie, H. (2011). 'D' for depression: Any role for vitamin D? 'Food for thought' II. *Acta Psychiatrica Scandinavica*, 124(4), 243–249. doi: 10.1111/j.1600-0447.2011.01705.x.
- Parker, G. B., Brotchie, H., & Graham, R. K. (2017). Vitamin D and depression. *Journal of Affective Disorders*, 208, 56–61. doi: 10.1016/j.jad.2016.08.082.
- Parker, G. B., & Brotchie, H. L. (2004). From diathesis to dimorphism. *Journal of Nervous & Mental Disease*, 192(3), 210–216. doi: 10.1097/01.nmd.0000116464.60500.63.
- Patrick, R. P., & Ames, B. N. (2015). Vitamin D and the omega-3 fatty acids control serotonin synthesis and action, part 2: Relevance for ADHD, bipolar disorder, schizophrenia, and impulsive behavior. *FASEB Journal*, 29(6), 2207–2222. doi: 10.1096/fj.14-268342.
- Pavlovic, A., Abel, K., Barlow, C. E., Farrell, S. W., Weiner, M., & DeFina, L. F. (2018). The association between serum vitamin d level and cognitive function in older adults: Cooper Center Longitudinal Study. *Preventive Medicine*, 113, 57–61. doi: 10.1016/j.ypmed.2018.05.010.
- Pettersen, J. A. (2017). Does high dose vitamin D supplementation enhance cognition? A randomized trial in healthy adults. *Experimental Gerontology*, 90, 90–97. doi: 10.1016/j.exger.2017.01.019.
- Plozer, E., Altbacker, A., Darnai, G., Perlaki, G., Orsi, G., Nagy, S. A., ... Janszky, J. (2015). Intracranial volume inversely correlates with serum 25 (OH)D level in healthy young women. *Nutritional Neuroscience*, 18(1), 37–40. doi: 10.1179/1476830514Y.0000000109.
- Qiu, H., Li, X., Luo, Q., Li, Y., Zhou, X., Cao, H., ... Sun, M. (2019). Alterations in patients with major depressive disorder before and after electroconvulsive therapy measured by fractional amplitude of low-frequency fluctuations (fALFF). *Journal of Affective Disorders*, 244, 92–99. doi: 10.1016/j.jad.2018.10.099.
- Rhee, S. J., Lee, H., & Ahn, Y. M. (2020). Serum vitamin D concentrations are associated with depressive symptoms in men: The sixth Korea National Health and Nutrition Examination Survey 2014. *Frontiers in Psychiatry*, 11, 756. doi: 10.3389/fpsy.2020.00756.
- Ringe, J. D., & Kipshoven, C. (2012). Vitamin D-insufficiency: An estimate of the situation in Germany. *Dermato-Endocrinology*, 4(1), 72–80. doi: 10.4161/derm.19829.
- Rock, P. L., Roiser, J. P., Riedel, W. J., & Blackwell, A. D. (2014). Cognitive impairment in depression: A systematic review and meta-analysis. *Psychological Medicine*, 44(10), 2029–2040. doi: 10.1017/S0033291713002535.
- Roy, N. M. (2021). Impact of vitamin D on neurocognitive function in dementia, depression, schizophrenia and ADHD. *Frontiers in Bioscience*, 26(3), 566–611. doi: 10.2741/4908.
- Ruhe, H. G., Mason, N. S., & Schene, A. H. (2007). Mood is indirectly related to serotonin, norepinephrine and dopamine levels in humans: A meta-analysis of monoamine depletion studies. *Molecular Psychiatry*, 12(4), 331–359. doi: 10.1038/sj.mp.4001949.
- Schlogl, M., & Holick, M. F. (2014). Vitamin D and neurocognitive function. *Clinical Interventions in Aging*, 9, 559–568. doi: 10.2147/CIA.S51785.
- Seedat, S., Scott, K. M., Angermeyer, M. C., Berglund, P., Bromet, E. J., Brugha, T. S., ... Kessler, R. C. (2009). Cross-national associations between gender and mental disorders in the World Health Organization World Mental Health Surveys. *Archives of General Psychiatry*, 66(7), 785–795. doi: 10.1001/archgenpsychiatry.2009.36.
- Sezgin, G., Ozturk, G., Turkal, R., & Caykara, B. (2019). Vitamin D levels of outpatients admitted to a university hospital in the Marmara region of Turkey over 3 years. *Journal of Medical Biochemistry*, 38(2), 181–187. doi: 10.2478/jomb-2018-0027.
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., ... Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*, 59(Suppl 20), 22–33; quiz 34–57. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9881538>.
- Shen, T., Qiu, M., Li, C., Zhang, J., Wu, Z., Wang, B., ... Peng, D. (2014). Altered spontaneous neural activity in first-episode, unmedicated patients with major depressive disorder. *Neuroreport*, 25(16), 1302–1307. doi: 10.1097/WNR.0000000000000263.
- Shen, Y., Wei, Y., Yang, X. N., Zhang, G., Du, X., Jia, Q., ... Zhang, X. Y. (2020). Psychotic symptoms in first-episode and drug naive patients with major depressive disorder: Prevalence and related clinical factors. *Depression and Anxiety*, 37(8), 793–800. doi: 10.1002/da.23026.
- Slinin, Y., Paudel, M., Taylor, B. C., Ishani, A., Rossom, R., Yaffe, K., ... Ensrud, K. E. (2012). Association between serum 25(OH) vitamin D and the risk of cognitive decline in older women. *The Journals of Gerontology: Series A, Biological Sciences and Medical Sciences*, 67(10), 1092–1098. doi: 10.1093/gerona/gls075.
- Solomon, M. B., & Herman, J. P. (2009). Sex differences in psychopathology: Of gonads, adrenals and mental illness. *Physiology & Behavior*, 97(2), 250–258. doi: 10.1016/j.physbeh.2009.02.033.
- Song, B. M., Kim, H. C., Rhee, Y., Youm, Y., & Kim, C. O. (2016). Association between serum 25-hydroxyvitamin D concentrations and depressive symptoms in an older Korean population: A cross-sectional study. *Journal of Affective Disorders*, 189, 357–364. doi: 10.1016/j.jad.2015.09.043.
- Thompson, E. (2015). Hamilton Rating Scale for Anxiety (HAM-A). *Occupational Medicine*, 65(7), 601. doi: 10.1093/occmed/kqv054.
- Toffanello, E. D., Coin, A., Perissinotto, E., Zambon, S., Sarti, S., Veronese, N., ... Sergi, G. (2014a). Vitamin D deficiency predicts cognitive decline in older men and women: The Pro.V.A. Study. *Neurology*, 83(24), 2292–2298. doi: 10.1212/WNL.0000000000001080.
- Toffanello, E. D., Sergi, G., Veronese, N., Perissinotto, E., Zambon, S., Coin, A., ... Manzano, E. (2014b). Serum 25-hydroxyvitamin d and the onset of late-life depressive mood in older men and women: The Pro.V.A. study. *The Journals of Gerontology: Series A, Biological Sciences and Medical Sciences*, 69(12), 1554–1561. doi: 10.1093/gerona/glu081.
- Tong, P., Bo, P., Shi, Y., Dong, L., Sun, T., Gao, X., & Yang, Y. (2021). Clinical traits of patients with major depressive disorder with comorbid borderline personality disorder based on propensity score matching. *Depression and Anxiety*, 38(1), 100–106. doi: 10.1002/da.23122.
- Wang, J., Um, P., Dickerman, B. A., & Liu, J. (2018). Zinc, magnesium, selenium and depression: A review of the evidence, potential mechanisms and implications. *Nutrients*, 10(5), 584. doi: 10.3390/nu10050584.
- Wang, L., Dai, W., Su, Y., Wang, G., Tan, Y., Jin, Z., ... Si, T. (2012). Amplitude of low-frequency oscillations in first-episode, treatment-naive patients with major depressive disorder: A resting-state functional MRI study. *PLoS ONE*, 7(10), e48658. doi: 10.1371/journal.pone.0048658.
- Wani, A. L., Bhat, S. A., & Ara, A. (2015). Omega-3 fatty acids and the treatment of depression: A review of scientific evidence. *Integrative Medicine Research*, 4(3), 132–141. doi: 10.1016/j.imr.2015.07.003.
- Williams, J. B. (1988). A structured interview guide for the Hamilton Depression Rating Scale. *Archives of General Psychiatry*, 45(8), 742–747. doi: 10.1001/archpsyc.1988.01800320058007.
- Wong, S. K., Chin, K. Y., & Ima-Nirwana, S. (2018). Vitamin D and depression: The evidence from an indirect clue to treatment strategy. *Current Drug Targets*, 19(8), 888–897. doi: 10.2174/1389450118666170913161030.
- World Health Organization. (2017). *Depression and other common mental disorders: Global health estimates*. Geneva, World Health Organization. Retrieved from [https://www.who.int/mental\\_health/management/depression/prevalence\\_global\\_health\\_estimates/en/](https://www.who.int/mental_health/management/depression/prevalence_global_health_estimates/en/).
- Wortsman, J., Matsuoka, L. Y., Chen, T. C., Lu, Z., & Holick, M. F. (2000). Decreased bioavailability of vitamin D in obesity. *The American Journal of Clinical Nutrition*, 72(3), 690–693. doi: 10.1093/ajcn/72.3.690.
- Yan, C. G., Wang, X. D., Zuo, X. N., & Zang, Y. F. (2016). DPABI: Data processing & analysis for (resting-state) brain imaging. *Neuroinformatics*, 14(3), 339–351. doi: 10.1007/s12021-016-9299-4.
- Yan, X., Zhang, N., Cheng, S., Wang, Z., & Qin, Y. (2019). Gender differences in vitamin D status in China. *Medical Science Monitor: International*

- Medical Journal of Experimental and Clinical Research*, 25, 7094–7099. doi: 10.12659/MSM.916326.
- Zagni, E., Simoni, L., & Colombo, D. (2016). Sex and gender differences in central nervous system-related disorders. *Neuroscience Journal*, 2016, 2827090. doi: 10.1155/2016/2827090.
- Zang, Y. F., He, Y., Zhu, C. Z., Cao, Q. J., Sui, M. Q., Liang, M., ... Wang, Y. F. (2007). Altered baseline brain activity in children with ADHD revealed by resting-state functional MRI. *Brain & Development*, 29(2), 83–91. doi: 10.1016/j.braindev.2006.07.002.
- Zheng, Y. P., Zhao, J. P., Phillips, M., Liu, J. B., Cai, M. F., Sun, S. Q., & Huang, M. F. (1988). Validity and reliability of the Chinese Hamilton Depression Rating Scale. *The British Journal of Psychiatry: The Journal of Mental Science*, 152, 660–664. doi: 10.1192/bjp.152.5.660.
- Zhou, F. C., Wang, Y. Y., Zheng, W., Zhang, Q., Ungvari, G. S., Ng, C. H., ... Xiang, Y. T. (2017). Prospective memory deficits in patients with depression: A meta-analysis. *Journal of Affective Disorders*, 220, 79–85. doi: 10.1016/j.jad.2017.05.042.
- Zhu, D. M., Zhang, C., Yang, Y., Zhang, Y., Zhao, W., Zhang, B., ... Yu, Y. (2020). The relationship between sleep efficiency and clinical symptoms is mediated by brain function in major depressive disorder. *Journal of Affective Disorders*, 266, 327–337. doi: 10.1016/j.jad.2020.01.155.
- Zhu, D.-M., Zhao, W., Zhang, B., Zhang, Y., Yang, Y., Zhang, C., ... Yu, Y. (2019). The relationship between serum concentration of vitamin D, total intracranial volume, and severity of depressive symptoms in patients with major depressive disorder. *Frontiers in Psychiatry*, 10, 322. 10.3389/fpsyt.2019.00322.
- Zhuo, C., Li, G., Lin, X., Jiang, D., Xu, Y., Tian, H., ... Song, X. (2019). The rise and fall of MRI studies in major depressive disorder. *Translational Psychiatry*, 9(1), 335. doi: 10.1038/s41398-019-0680-6.
- Zou, Q. H., Zhu, C. Z., Yang, Y., Zuo, X. N., Long, X. Y., Cao, Q. J., ... Zang, Y. F. (2008). An improved approach to detection of amplitude of low-frequency fluctuation (ALFF) for resting-state fMRI: Fractional ALFF. *Journal of Neuroscience Methods*, 172(1), 137–141. doi: 10.1016/j.jneumeth.2008.04.012.