

Anaerobic cultures showed that species of *Bacteroides* were the predominant bacteria present during both restricted and *ad lib.* feeding. These appeared in Gram-films as two pleomorphic forms—mainly coccal in the presence of ciliates and rod-like during *ad lib.* intake.

Similar observations were made when two steers were changed from *ad lib.* to restricted intake and given an inoculum of rumen ciliates.

REFERENCE

Eadie, J. M., Hobson, P. N. & Mann, S. O. (1967). *Anim. Prod.* **9**, 247.

Preliminary communication on studies of the digestive processes in pigs with intestinal cannulas. By R. BRAUDE, H. L. BUTTLE, F. HORSZCZARUK* and A. W. MYRES, *National Institute for Research in Dairying, Shinfield, Reading*

The use of intestinal cannulas has been found to facilitate the study of digestive processes in the living pig and to allow repeated observations on the same animal. One or two single cannulas, and one or two re-entrant cannulas were fitted into the small intestines of pigs weighing 60–100 kg. Preliminary observations were reported on the passage of the digesta, and particularly of nitrogen and polyethylene glycol (PEG) which was used as a marker. Movement of the digesta during the interval between two feeds showed two marked peaks, but very considerable variation between replicates from the same animal was observed. PEG was found unsatisfactory for tracing the passage of nitrogen, as it appears to leave the stomach faster than the solid material.

The influence of meat meal and other high-protein feeding-stuffs on the survival of chicks inoculated with *Salmonella gallinarum*. By R. HILL, *Royal Veterinary College, University of London*

The influence of level of dietary protein on infectious disease has been studied by a number of workers, often with conflicting results. Hill, Colburn & Schneider (1962) showed that certain substances other than protein, present in the diet, could modify the protein effect, and recently it became evident that the source of protein also influenced the result (Hill & Smith, 1969).

In the experiments described here the total protein contents of the diets were similar, about 20%, and near to the level recommended for chicks (National Research Council, 1966). For each diet one high-protein feeding-stuff was mixed with wheat-meal and an appropriate supplement of minerals, vitamins and amino acids to permit reasonably good growth in control chicks. From day-old to 12 days all chicks were

*Present address: Institute of Animal Physiology and Nutrition, Jablonna, Warsaw, Poland.