

p-Coumaric acid, a common dietary phenol, inhibits platelet activity *in vitro* and *in vivo*

Cristina Luceri^{1*}, Lucia Giannini¹, Maura Lodovici¹, Emilia Antonucci², Rosanna Abbate², Emanuela Masini¹ and Piero Dolaro¹

¹Department of Preclinical and Clinical Pharmacology, University of Florence, Azienda Ospedaliero-Universitaria Careggi, 50139 Florence, Italy

²Thrombosis Center, University of Florence, Azienda Ospedaliero-Universitaria Careggi, 50139 Florence, Italy

(Received 22 March 2006 – Revised 4 August 2006 – Accepted 5 September 2006)

p-Coumaric acid (3-(4-hydroxyphenyl)-2-propenoic acid; 4CA), is a ubiquitous plant metabolite with antioxidant and anti-inflammatory properties. The antiplatelet activity of this compound was analysed both *ex vivo* and *in vitro*. 4-CA, administered to rabbits for 2 weeks at the dose of 5 mg/kg, mixed with food, inhibited ADP-induced platelet aggregation without affecting blood coagulation. This effect was associated with a marked increase in plasma antioxidant activity, measured as ferric reducing ability of plasma, and with the reduction of thromboxane B₂ production. The antiplatelet effect was confirmed by *in vitro* experiments on human blood: 4CA (500 µM and 1 mM) reduced ADP-induced platelet aggregation (55.2 (SE 4.01) % and 35.6 (SE 2.35) % relative to basal level, respectively). 4CA was able to modify platelet function, measured with PFA-100™, a shear-inducing device that simulates primary haemostasis. 4CA interfered also with arachidonic acid cascade, reducing thromboxane B₂ production and lipopolysaccharide-induced prostaglandin E₂ generation (IC₅₀ 371 and 126 µM, respectively). The data show that 4CA is an antioxidant compound with good antiplatelet activity at doses that can be obtained with dietary intervention, suggesting possible applications for primary prevention of vascular disease.

Phenols: Platelet aggregation: Antioxidant activity

Epidemiological studies suggest that moderate red wine consumption may reduce the risk of CVD (de Gaetano & Cerletti, 2001) and this effect can be partially due to some naturally occurring constituents of this beverage. Wine contains a wide variety of polyphenols, including tannins, anthocyanins and phenolic flavonoids (including flavonols and flavones) (Rasmussen *et al.* 2005) that may contribute to the cardioprotective effects of grape products, as suggested by several studies (Duffy *et al.* 2001; da Luz & Coimbra, 2004; Vita, 2005).

One proposed mechanism of the apparent benefit of various sources of polyphenols is their favourable effect on platelet aggregation (Ruf, 2004; Cordova *et al.* 2005). Platelet aggregation plays a critical role in the pathogenesis of vascular disorders, and there is evidence that antiplatelet therapy reduces the risk of CVD (Youssef *et al.* 2005).

Polyphenols may inhibit platelet aggregation through a number of different mechanisms, including inhibition of cyclooxygenase (COX), lipoxygenase (Schubert *et al.* 1999; Hong *et al.* 2001) and phosphodiesterase activities (Dell'Agli *et al.* 2005). Inhibition of phosphodiesterase activity, increasing platelet levels of cAMP or cGMP, lowers platelet calcium levels and inhibits platelet activation both *in vitro* and *in vivo* (Demrow *et al.* 1995).

Other inhibitory effects of polyphenols on platelet aggregation include scavenging of reactive oxygen species such as superoxide anion (Frei & Higdon, 2003), decreasing phospholipase C (Pignatelli *et al.* 2000) and inhibition of lipid peroxidation (Aviram *et al.* 2002).

A relevant effect of polyphenols on the cardiovascular system could be due to their ability to interact with the L-arginine–nitric oxide pathway leading to the generation of NO from vascular endothelium (Andriambeloso *et al.* 1997), since this molecule has vasorelaxant and antiaggregation properties. Freedman *et al.* (2001) examined the effects of grape juice on platelet function, observing that the addition of grape juice to platelets *ex vivo* was associated with a decrease in the production of superoxide anion from platelets, a reduction in their aggregation and a significant increase in platelet NO production, effects related to reduced activation of protein kinase C. The findings were also reproduced *in vivo* in platelets isolated from healthy volunteers who had consumed grape juice (7 ml/kg per d) for 2 weeks (Freedman *et al.* 2001). Grape seed and grape skin, in combination, exhibited a high antiplatelet effect *in vitro* and *ex vivo*, in dog platelets at doses ranging between 2 and 20 mg/kg (Shanmuganayagam *et al.* 2002).

Abbreviations: AA, arachidonic acid; 4CA, *p*-coumaric acid; COX, cyclooxygenase; PGE₂, prostaglandin E₂; PRP, platelet-rich plasma; TXB₂, thromboxane B₂.

* **Corresponding author:** Dr Cristina Luceri, fax +39 055 4271 280, email cristina.luceri@unifi.it

Results of studies on the effect of red or white wine consumption on platelet aggregation are contrasting (Pignatelli *et al.* 2002). Some studies have focused on the effect of single wine constituents such as resveratrol, a phytoalexin present at low concentration in grapes and some wines, which is able to reduce several platelet functions (Olas *et al.* 2002), and the flavonoid quercetin, which inhibits collagen-induced platelet aggregation and adhesion (Pignatelli *et al.* 2000; Kikura *et al.* 2004).

p-Coumaric acid (3-(4-hydroxyphenyl)-2-propenoic acid; 4CA) is another wine constituent, widely distributed in cereals, fruits and vegetables (being a ubiquitous plant metabolite), intermediate in the synthesis of many phenols (Clifford, 2000). Studies performed in our laboratory demonstrate that 4CA has marked antioxidant and anti-inflammatory properties in rat colonic mucosa (Guglielmi *et al.* 2003; Luceri *et al.* 2004).

The purpose of the present study was to evaluate the effects of 4CA on systemic oxidant–antioxidant status and on platelet function.

Materials and methods

Venous blood was sampled from rabbits and healthy human volunteers using a 21-gauge needle (Pentaven[®]; Pentaferte, Campli, Teramo, Italy) and collected in evacuated polypropylene tubes containing 3.8% sodium citrate (9:1, v/v; Vacutest[®]; Greiner bio-one, Kremsmünster, Austria) or EDTA (Vacutest[®]; Vacutest Kima, Azzergande, Padova, Italy).

4CA and arachidonic acid (AA) were purchased from the Sigma Chemical Co. (St Louis, MO, USA) and ADP was purchased from Mascia Brunelli (Milan, Italy). Thromboxane B₂ (TXB₂) and prostaglandin E₂ (PGE₂) enzyme immunoassay kits were obtained from the Cayman Chemical Co. (Ann Arbor, MI, USA).

Ex vivo studies in rabbit blood

All the procedures adopted were carried out in agreement with the European Union Regulations on the Care and Use of Laboratory Animals (OJ of ECL 358/1, 12/18/1986) and the experiments were conducted according to Italian regulations on the protection of animals used for experimental and other scientific purposes (DM 116/1992). Blood samples from rabbits were obtained by puncture of an ear vein.

The first experiment was a dose–response study: twelve albino rabbits (2.5–3 kg) obtained from Charles River (Lecco, Italy) were randomly divided into three groups of four animals, each housed in a single cage: controls were fed a powdered standard lab chow purchased from Harlan (Milan, Italy), the 2030 Global Diet for rabbits, whereas the treated groups received the same diet enriched with 4CA; the proper amount of 4CA was mixed with the daily dietary mass (190 g) to a final dosage of 5 and 500 mg/kg, for 4 weeks.

The second experiment was a time-course study, performed using the lowest effective 4CA dose (5 mg/kg). Blood samples were obtained from five rabbits at time 0 and then after 2, 4, 6, 8 and 10 weeks of treatment.

Platelet aggregation. At the indicated time-points, blood samples were collected from rabbits into tubes containing

3.8% trisodium citrate solution (9:1, v/v). Platelet-rich plasma (PRP) was obtained by centrifuging the blood samples (80 g, 10 min, 20°C). After PRP removal, the remaining plasma was centrifuged (900 g, 10 min, 20°C) and the supernatant was collected to obtain platelet-poor plasma, used to dilute autologous PRP and to equilibrate the aggregometer. Platelet aggregation was determined by a four-channel aggregometer (model PAP-4; Biodata Corporation, Horsham, PA, USA) according to the method of Born (1962). Aggregation was induced by the addition of ADP or AA (final concentrations 3 and 2.86 mM, respectively). Aggregation was measured over 5 min and the extent of aggregation was estimated by measuring the maximal curve height above the baseline level.

Determination of anticoagulant activity. An aliquot of venous blood samples collected into citrate-containing tubes was used to determine prothrombin time and activated partial thromboplastin time, using a coagulometer (Dade Behring, Newark, DE, USA).

Whole blood cell counts. Blood cell counts, Hb concentration, haematocrit value and mean platelet volume were measured in whole blood collected in EDTA-containing tubes, using a haematology analyser (Coulter Electronics, Hialeah, FL, USA).

Antioxidant activity. Plasma total antioxidant power (ferric reducing ability of plasma) measures the ability of the antioxidants contained in a sample to reduce ferric ions (Benzie & Strain, 1996). The method is based on the reduction of a ferric-tripyridyltriazine [Fe(III)–tripyridyltriazine] complex to the ferrous [Fe(II)] form, which has an absorption maximum at 593 nm. Fe(III) is used in excess so that the colour formation is linearly related to the amount of reductant (antioxidant) present. Data are expressed as $\mu\text{mol/l}$.

Thromboxane B₂ release from activated platelets. TXB₂, the stable metabolite of thromboxane A₂ released during aggregation, was used as an index of COX-1 activity (Patrignani *et al.* 1994). Immediately after platelet aggregation induced by ADP or AA, the tubes containing PRP were transferred to an ice bath and later centrifuged in an Eppendorf microcentrifuge for 5 min. The supernatant was collected and frozen at –20°C for the determination of TXB₂ concentration by enzyme immunoassay kit, according to the manufacturer's instructions.

In vitro study in human blood

The study protocol followed our institutional guidelines and an informed consent was obtained from each subject. Ten healthy, non-smoking volunteers (six women, four men) aged 23–39 years participated in the study. None of the participants had reported previous CVD, metabolic or systemic disease, or had taken any drug in the 2 weeks before the study. Women were premenopausal and were not taking oestrogens.

Venous blood was carefully drawn from an antecubital vein by trauma-free venepuncture with rapid flow of blood. Within 10 min, whole blood was incubated for 60 min at room temperature with 4CA (dissolved in 2% 1 M-NaOH) or the vehicle alone. 4CA final concentrations were 10, 100, 500 and 1000 μM .

Platelet aggregation. After 60 min incubation with 10, 100, 500 and 1000 μM -4CA, blood samples were centrifuged (160g, 10 min, 20°C) and the supernatant separated as PRP. After PRP removal, the remaining plasma was centrifuged (2000g, 10 min, 20°C) and the supernatant was separated to obtain platelet-poor plasma. PRP was diluted to $2\text{--}3 \times 10^{11}$ platelets/l with autologous platelet-poor plasma. Platelet aggregation was performed at 37°C using a four-channel photometric aggregometer (model PAP-4; Biodata Corporation) based on the method of Born (1962). Aggregation was induced by ADP (final concentration 3 μM). This concentration was chosen to ensure a high probability of aggregation based on previous experience. The aggregation was allowed to proceed for 5 min and the extent was estimated quantitatively by measuring the maximal curve height above baseline level.

Whole blood cell counts. Whole blood cell counts were measured as described earlier for rabbit samples.

PFA-100™ closure time. Citrated whole blood samples incubated at room temperature for 60 min with 500 μM or 1 mM 4CA were analysed by PFA-100™ (Dade Behring), a high-shear-inducing device that simulates primary haemostasis. We determined closure times on duplicate samples of 800 μl using cartridges containing collagen–epinephrine or collagen–ADP membranes.

Prostaglandin E_2 production. Plasma PGE_2 levels were determined as an index of whole blood COX-2 activity. Aliquots of citrated whole blood were incubated with 4CA (dissolved in 2% 1M-NaOH) or the vehicle alone, in the presence of endotoxin (lipopolysaccharide) derived from *Escherichia coli* 026:B6 (Sigma Chemical Co.) at 37°C for 60 min. The contribution of platelet COX-1 was suppressed by the addition of 200 μM -aspirin (Patrignani *et al.* 1994). The final concentrations of 4CA were 10, 100, 500 μM and 1 mM. Aliquots of citrated whole blood were incubated with celecoxib (final concentration 1 μM) in the presence of 10 $\mu\text{g}/\text{ml}$ lipopolysaccharide, as positive control. Plasma was separated by centrifugation (1000g, 20 min, 20°C) and assayed for PGE_2 using an enzyme immunoassay kit according to the manufacturer's instructions.

Thromboxane B_2 release from human activated platelets. The determination of TXB_2 production was performed as described earlier for rabbit samples.

Statistical analysis

Experiments were performed at least in duplicate and the results were expressed as means and their standard errors. Statistical analysis was conducted using a two-way ANOVA.

Results

Ex vivo studies in rabbit blood

Inhibitory effects of 4CA on platelet aggregation. The dose–response study was performed using two extreme doses: 5 and 500 mg/kg 4CA, chosen considering that the estimated human daily intake of 4CA is about 15 mg/kg. At both doses 4CA was able to inhibit AA-induced platelet aggregation in a not statistically significant dose-dependent way (Fig. 1).

In the second experiment we observed that ADP-induced platelet aggregation was inhibited by 4CA (5 mg/kg) in a




Fig. 1. Inhibition of platelet aggregation in blood samples from rabbits fed a standard diet (controls) or a diet enriched with *p*-coumaric acid (4CA; 5 and 500 mg/kg) for 4 weeks. For details of procedures, see p. 459. Values are means with their standard errors depicted by vertical bars. Mean values were significantly different from those of the controls: * $P < 0.05$.

time-dependent manner: the effect started after 2 weeks, reaching the maximum inhibitory effect after 8 weeks of treatment (Fig. 2).

Blood cell counts and coagulation tests. Treatments with 4CA did not modify blood cell counts, Hb concentration, haematocrit value, mean platelet volume, platelet count, prothrombin time or activated partial thromboplastin time (data not shown).

Inhibition of thromboxane B_2 production. TXB_2 release was measured in both dose–response and time-course experiments. In rabbits treated with 4CA (5 or 500 mg/kg for 4 weeks), TXB_2 release, measured in PRP after AA-induced aggregation, was significantly reduced and the differences between the two doses were not statistically significant (Fig. 3(A)).

In rabbits treated with 4CA (5 mg/kg) TXB_2 release during ADP-induced aggregation was significantly reduced after 2 weeks of treatment (Fig. 3(B)).

Ferric reducing ability of plasma assay. The administration of 4CA (5 mg/kg) significantly increased the plasma antioxidant activity from 160 (SE 15) $\mu\text{mol}/\text{l}$ at time 0 to 310 (SE 30) $\mu\text{mol}/\text{l}$ after 1 week ($P < 0.05$) and to 380 (SE 50) $\mu\text{mol}/\text{l}$ after 3 weeks of feeding ($P < 0.05$).




Fig. 2. Inhibition of platelet aggregation in blood samples from rabbits fed a standard diet enriched with 5 mg/kg *p*-coumaric acid (4CA). For details of procedures, see p. 459. Values are means with their standard errors depicted by vertical bars. Mean values were significantly different from those at time 0: * $P < 0.05$.




Fig. 3. A, Thromboxane B₂ (TXB₂) production after arachidonic acid-induced platelet aggregation in blood samples from rabbits fed *p*-coumaric acid (4CA; 5 or 500 mg/kg) for 4 weeks. B, TXB₂ production after ADP-induced aggregation in blood samples from rabbits treated with 4CA (5 mg/kg). For details of procedures, see p. 459. Values are means with their standard errors depicted by vertical bars. Mean values were significantly different from those of the controls (A) or at time 0 (B): **P*<0.05.

In vitro studies in human blood

Inhibitory effects of 4CA on platelet function. All volunteers had normal platelet function and the vehicle alone (2% 1 M-NaOH) had no effect on platelet aggregation. At the doses of 500 μM and 1 mM, 4CA significantly reduced ADP-induced platelet aggregation (35.6 (SE 2.35) % and 55.18 (SE 4.01) % of the basal level, respectively, *P*<0.05), while at lower concentrations (10 and 100 μM) it did not induce a significant effect (data not shown).

Platelet function measured as PFA-100™ closure times. All volunteers had normal values and the vehicle alone (2% 1 M-NaOH) had no effect on closure time. Blood samples incubated for 60 min with 500 μM-4CA showed a prolonged closure time. When incubated with 1 mM-4CA the difference was statistically significant, using cartridges containing both collagen–epinephrine and collagen–ADP membranes (Table 1).

Inhibition of thromboxane B₂ production. At the doses of 500 μM and 1 mM, 4CA was able to reduce TXB₂ release after ADP-induced platelet aggregation in PRP from healthy volunteers, with an IC₅₀ of 371 μM.

Inhibition of prostaglandin E₂ production. Incubation of blood samples from healthy donors with lipopolysaccharide, in the presence of aspirin to block COX-1 activity, resulted

Table 1. Effect of blood incubation with different *p*-coumaric acid (4CA) doses on platelet function†

(Mean values with their standard errors)

	Collagen–epinephrine closure time (s)		Collagen–ADP closure time (s)	
	Mean	SE	Mean	SE
Control	127.3	6.8	96.2	5.4
500 μM-4CA	178.9	29.5	166.0	32.5
1 mM-4CA	216.7	31.4*	178.4	31.0*

Mean values were significantly different from those of the control group: **P*<0.05.

† For details of procedures, see p. 459.

in marked induction of PGE₂ formation. The vehicle alone (2% 1 M-NaOH) had no effect on PGE₂ production, whereas 4CA reduced lipopolysaccharide-induced PGE₂ generation with an IC₅₀ of 126 μM.

Discussion

Platelets play a pivotal role in health and disease, given their central involvement in haemostasis and thrombosis. Treatment with drugs that specifically inhibit platelet function has proven to be beneficial in the prevention of thrombosis.

The relationship between a diet high in vegetable products and a decreased risk of thrombosis is well established, although the explanation for this is far from clear (Liu & Williett, 2002). A number of studies have focused on the effects of dietary component, such as flavonoids on platelet function. Quercetin (40–100 μM) and catechin (100–420 μM) inhibit platelet aggregation *in vitro*, inhibiting the release of platelet hydrogen peroxide elicited by collagen, with almost complete suppression when used in combination (Pignatelli *et al.* 2000).

Resveratrol is another wine compound with several health-promoting properties; the most widely accepted mechanism of resveratrol-mediated cardioprotection is its ability to inhibit platelet aggregation. At the concentration of 3.56 mg/l, it lowers platelet aggregation by 50% in healthy volunteers (Bertelli *et al.* 1995). Moreover, resveratrol and quercetin inhibited thrombin-induced ADP and ATP secretion from platelets in a concentration-dependent manner (Kaneider *et al.* 2004).

In the present study we analysed the effect of another wine component, 4CA, which is also widely distributed in fruits, vegetables, cereals, tea and wine, mostly as conjugates. It is present as a free acid (about 22 mg/l) and as *p*-coumaroyltartaric acid (about 139 mg/l) in red wine (Ghiselli *et al.* 1998). Other abundant sources of 4CA are apples and berries (69–1700 mg/kg) and maize bran (2.9 g/kg) (Stich, 1991; Clifford, 2000; Zhao *et al.* 2005).

Administered *in vivo*, 4CA was able to reduce significantly platelet aggregation induced by AA and ADP after 2 weeks of treatment and at a dose comparable to the daily intake for cinnamates (Clifford, 2000). In addition, the TXB₂ production measurements were well correlated with platelet aggregation in both experiments.

In vitro, 4CA inhibited platelet aggregation without effects on prothrombin time, activated partial thromboplastin time, Hb concentration, haematocrit value, mean platelet volume and platelet count. In PFA-100™ analysis, which simulates

the process of platelet adhesion and aggregation induced by vascular injury, we also observed a significant effect of 4CA at high concentrations, using ADP as well as epinephrine cartridges. *In vivo* and *in vitro* studies suggested that 4CA modulates platelet aggregation through both COX-dependent and independent mechanisms.

There are many platelet pathways which could be influenced by 4CA: polyphenols may inhibit platelet aggregation because of their antioxidant activity (Salvemini & Botting, 1993; Neiva *et al.* 1999). 4CA is a potent inhibitor of LDL oxidation *in vitro* (Satué-Gracia *et al.* 1999; Morton *et al.* 2000). Owen *et al.* (2000) reported the antioxidant activity of 4CA against reactive oxygen species produced by hypoxanthine/xanthine oxidase and Lodovici *et al.* (2001) described its ability to reduce oxidative DNA damage induced *in vitro* by iron and cumene hydroperoxide. The present results show that the administration of 4CA (5 mg/kg) significantly increases plasma antioxidant activity after 1 week of treatment and more markedly after 3 weeks, confirming our previous data on the antioxidant activity of this molecule *in vivo*, on rat colon mucosa (Guglielmi *et al.* 2003).

It has also been suggested that the effect of polyphenols on platelet aggregation is related to the inhibition of the COX pathway, reducing the formation of TXB₂, a potent aggregating and vasoconstricting agent. The anti-inflammatory effects of 4CA are controversial since it has anti-inflammatory activity if administered locally, but demonstrates no effects *in vivo* on carrageenan-induced paw edema when administered at 200 mg/kg per d, 1 h before carrageenan injection (Fernandez *et al.* 1998). Recently we demonstrated that an acute intestinal inflammation induced by dextran sulphate sodium can be inhibited by pretreatment with 4CA (50 mg/kg) and that this effect was associated with the suppression of COX-2 expression and activity (Luceri *et al.* 2004). In the present study we demonstrate that the reduction of platelet aggregation induced by 4CA is correlated with decreased TXB₂ production, associated with the inhibition of COX-1 and COX-2 activities.

In summary, 4CA, a dietary phenolic compound, acts as a potent systemic antioxidant and shows good antiaggregant properties at concentrations that can be obtained through regular consumption of vegetables, cereals, fruits or a moderate quantity of red wine. It has been estimated that the daily dietary intake for total cinnamates and hydroxybenzoic acid derivatives should be 1 g/d, corresponding to about 15 mg/kg per d in man (Clifford, 2000; Tomás-Barberán & Clifford, 2000).

Healthy subjects with no dietary restrictions had a relatively low concentration of 4CA in faecal water, ranging from 0.3 to 1.44 µM (Jenner *et al.* 2005). However, in a recent paper by Garrat *et al.* (2006) the authors demonstrated that 4CA is absorbed by all the digestive tracts and, when orally administered, is mainly eliminated in the urine. Konishi *et al.* (2003), studying 4CA intestinal absorption, demonstrated that 4CA is actively transported by the colonic monocarboxylate transporter in Caco-2 cells; after an oral dose of 16.4 mg/kg, the serum C_{max} was 165.7 µM and the area under the curve was almost 3 mM in rats (Konishi *et al.* 2004).

Free phenolic acids are well absorbed but in foods they are mainly in bound form, esterified to cell wall polymers. It has

been reported that IR irradiation or fermentation can liberate covalently bound phenolic compounds with antioxidant activities and that the irradiation of sesame seeds significantly increases the amount of free 4CA (Lee *et al.* 2005); these data suggest that some industrial processes can enrich the amount of free 4CA in food.

In conclusion, the present data suggest that dietary intervention with foods rich in 4CA may be a possible strategy for primary prevention of vascular diseases. Further studies on patients at risk could help to determine the potential beneficial effects of 4CA as an antiplatelet and systemic antioxidant agent for prevention and treatment of thrombotic diseases.

Acknowledgements

This work was financially supported by the Ministero della Università e della Ricerca Scientifica e Tecnologica, by the University of Florence, Italy, by the EU program Polybind QLRT 1999-00 505 and by the Network of Excellence in Nutrigenomics, NuGO. We thank Mary Forrest for revision of the manuscript.

References

- Andriambelison E, Kleschyov AL, Muller B, Beretz A, Stoclet JC & Andriantsitohaina R (1997) Nitric oxide production and endothelium-dependent vasorelaxation induced by wine polyphenols in rat aorta. *Br J Pharmacol* **120**, 1053–1058.
- Aviram M, Dornfeld L, Kaplan M, *et al.* (2002) Pomegranate juice flavonoids inhibit low-density lipoprotein oxidation and cardiovascular diseases: studies in atherosclerotic mice and in humans. *Drugs Exp Clin Res* **28**, 49–62.
- Benzie IF & Strain JJ (1996) The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': the FRAP assay. *Anal Biochem* **239**, 70–76.
- Bertelli AA, Giovannini L, Giannessi D, Migliori M, Bernini W, Fregoni M & Bertelli A (1995) Antiplatelet activity of synthetic and natural resveratrol in red wine. *Int J Tissue React* **17**, 1–3.
- Born GV (1962) Aggregation of blood platelets by adenosine diphosphate and its reversal. *Nature* **194**, 927–929.
- Clifford MN (2000) Chlorogenic acids and other cinnamates - nature, occurrence, dietary burden, absorption and metabolism. *J Sci Food Agric* **80**, 1033–1043.
- Cordova AC, Jackson LS, Berke-Schlessel DW & Sumpio BE (2005) The cardiovascular protective effect of red wine. *J Am Coll Surg* **200**, 428–439.
- da Luz PL & Coimbra SR (2004) Wine, alcohol and atherosclerosis: clinical evidences and mechanisms. *Braz J Med Biol Res* **37**, 1275–1295.
- de Gaetano G & Cerletti C (2001) European project. FAIR CT 97 3261 Project participants. Wine and cardiovascular disease. *Nutr Metab Cardiovasc Dis* **11**, 47–50.
- Dell'Agli M, Galli GV, Vrhovsek U, Mattivi F & Bosisio E (2005) *In vitro* inhibition of human cGMP-specific phosphodiesterase-5 by polyphenols from red grapes. *J Agric Food Chem* **53**, 1960–1965.
- Demrow HS, Slane PR & Folts JD (1995) Administration of wine and grape juice inhibits *in vivo* platelet activity and thrombosis in stenosed canine coronary arteries. *Circulation* **91**, 1182–1188.
- Duffy SJ, Vita JA, Holbrook M, Swerdloff PL & Keaney JF Jr (2001) Effect of acute and chronic tea consumption on platelet aggregation in patients with coronary artery disease. *Arterioscler Thromb Vasc Biol* **21**, 1084–1089.

- Fernandez MA, Saenz MT & Garcia MD (1998) Anti-inflammatory activity in rats and mice of phenolic acids isolated from *Scrophularia frutescens*. *J Pharm Pharmacol* **50**, 1183–1186.
- Freedman JE, Parker C 3rd, Li L, Perlman JA, Frei B, Ivanov V, Deak LR, Iafrafi MD & Folts JD (2001) Select flavonoids and whole juice from purple grapes inhibit platelet function and enhance nitric oxide release. *Circulation* **103**, 2792–2798.
- Frei B & Higdon JV (2003) Antioxidant activity of tea polyphenols *in vivo*: evidence from animal studies. *J Nutr* **133**, 3275S–3284S.
- Garrait G, Jarrige JF, Blanquet S, Beyssac E, Cardot JM & Alric M (2006) Gastrointestinal absorption and urinary excretion of trans-cinnamic and p-coumaric acids in rats. *J Agric Food Chem* **54**, 2944–2950.
- Ghiselli A, Nardini M, Baldi A & Scaccini C (1998) Antioxidant activity of different phenolic fractions separated from an Italian red wine. *J Agric Food Chem* **46**, 361–367.
- Guglielmi F, Luceri C, Giovannelli L, Dolara P & Lodovici M (2003) Effect of 4-coumaric and 3,4-dihydroxybenzoic acid on oxidative DNA damage in rat colonic mucosa. *Br J Nutr* **89**, 581–587.
- Hong J, Smith TJ, Ho CT, August DA & Yang CS (2001) Effects of purified green and black tea polyphenols on cyclooxygenase- and lipoxygenase-dependent metabolism of arachidonic acid in human colon mucosa and colon tumor tissues. *Biochem Pharmacol* **62**, 1175–1183.
- Jenner AM, Raftar J & Halliwell B (2005) Human fecal water content of phenolics: the extent of colonic exposure to aromatic compounds. *Free Radic Biol Med* **38**, 763–772.
- Kaneider NC, Mosheimer B, Reinisch N, Patsch JR & Wiedermann CJ (2004) Inhibition of thrombin-induced signaling by resveratrol and quercetin: effects on adenosine nucleotide metabolism in endothelial cells and platelet-neutrophil interactions. *Thromb Res* **114**, 185–194.
- Kikura M, Levy JH, Safon RA, Lee MK & Szlam F (2004) The influence of red wine or white wine intake on platelet function and viscoelastic property of blood in volunteers. *Platelets* **15**, 37–41.
- Konishi Y, Hitomi Y & Yoshioka E (2004) Intestinal absorption of p-coumaric and gallic acids in rats after oral administration. *J Agric Food Chem* **52**, 2527–2532.
- Konishi Y, Kobayashi S & Shimizu M (2003) Transepithelial transport of p-coumaric acid and gallic acid in Caco-2 cell monolayers. *Biosci Biotechnol Biochem* **67**, 2317–2324.
- Lee SC, Jeong SM, Kim SY, Nam KC & Ahn DU (2005) Effect of far-infrared irradiation on the antioxidant activity of defatted sesame meal extracts. *J Agric Food Chem* **9**, 1495–1498.
- Liu S & Willett WC (2002) Dietary glycemic load and atherothrombotic risk. *Curr Atheroscler Rep* **4**, 454–461.
- Lodovici M, Guglielmi F, Meoni M & Dolara P (2001) Effect of natural phenolic acids on DNA oxidation *in vitro*. *Food Chem Toxicol* **39**, 1205–1210.
- Luceri C, Guglielmi F, Lodovici M, Giannini L, Messerini L & Dolara P (2004) Plant phenolic 4-coumaric acid protects against intestinal inflammation in rats. *Scand J Gastroenterol* **39**, 1128–1133.
- Morton LW, Croft KD, Puddey IB & Byrne L (2000) Phenolic acids protect low density lipoproteins from peroxynitrite-mediated modification *in vitro*. *Redox Rep* **5**, 124–125.
- Neiva TJ, Morais L, Polack M, Simoes CM & D'Amico EA (1999) Effects of catechins on human blood platelet aggregation and lipid peroxidation. *Phytother Res* **13**, 597–600.
- Olas B, Wachowicz B, Saluk-Juszczak J & Zielinski T (2002) Effect of resveratrol, a natural polyphenolic compound, on platelet activation induced by endotoxin or thrombin. *Thromb Res* **107**, 141–145.
- Owen RW, Mier W, Giacosa A, Hull WE, Spiegelhalter B & Bartsch H (2000) Phenolic compounds and squalene in olive oils: the concentration and antioxidant potential of total phenols, simple phenols, secoiridoids, lignans and squalene. *Food Chem Toxicol* **38**, 647–659.
- Patrignani P, Panara MR, Greco A, Fusco O, Natoli C, Iacobelli S, Cipollone F, Ganci A, Creminon C & Maclouf J (1994) Biochemical and pharmacological characterization of the cyclooxygenase activity of human blood prostaglandin endoperoxide synthases. *J Pharmacol Exp Ther* **271**, 1705–1712.
- Pignatelli P, Lenti L, Pulcinelli FM, Catasca R, Sacconi G, Germano G, Marcoccia A, Silvestri MA, Ghiselli A & Viola F (2002) Red and white wine differently affect collagen-induced platelet aggregation. *Pathophysiol Haemost Thromb* **32**, 356–358.
- Pignatelli P, Pulcinelli FM, Celestini A, Lenti L, Ghiselli A, Gazzaniga PP & Viola F (2000) The flavonoids quercetin and catechin synergistically inhibit platelet function by antagonizing the intracellular production of hydrogen peroxide. *Am J Clin Nutr* **72**, 1150–1155.
- Rasmussen SE, Frederiksen H, Struntze Krogholm K & Poulsen L (2005) Dietary proanthocyanidins: occurrence, dietary intake, bioavailability, and protection against cardiovascular disease. *Mol Nutr Food Res* **49**, 159–174.
- Ruf JC (2004) Alcohol, wine and platelet function. *Biol Res* **37**, 209–215.
- Salvemini D & Botting R (1993) Modulation of platelet function by free radicals and free-radical scavengers. *Trends Pharmacol Sci* **14**, 36–42.
- Satué-Gracia MT, Andres-Lacueva C, Lamuela-Raventos RM & Frankel EN (1999) Spanish sparkling wines (Cavas) as inhibitors of *in vitro* human low-density lipoprotein oxidation. *J Agric Food Chem* **47**, 2198–2202.
- Schubert SY, Lansky EP & Neeman I (1999) Antioxidant and eicosanoid enzyme inhibition properties of pomegranate seed oil and fermented juice flavonoids. *J Ethnopharmacol* **66**, 11–17.
- Shanmuganayagam D, Beahm MR, Osman HE, Krueger CG, Reed JD & Folts JD (2002) Grape seed and grape skin extracts elicit a greater antiplatelet effect when used in combination than when used individually in dogs and humans. *J Nutr* **132**, 3592–3598.
- Stich HF (1991) The beneficial and hazardous effects of simple phenolic compounds. *Mutat Res* **259**, 307–324.
- Tomàs-Barberan FA & Clifford MN (2000) Dietary hydroxybenzoic acid derivatives - nature, occurrence and dietary burden. *J Sci Food Agric* **80**, 1024–1032.
- Vita JA (2005) Polyphenols and cardiovascular disease: effects on endothelial and platelet function. *Am J Clin Nutr* **81**, 292S–297S.
- Youssef F, Gupta P, Mikhailidis DP & Hamilton G (2005) Risk modification in patients with peripheral arterial disease: a retrospective survey. *Angiology* **56**, 279–287.
- Zhao Z, Egashira Y & Sanada H (2005) Phenolic antioxidants richly contained in corn bran are slightly bioavailable in rats. *J Agric Food Chem* **53**, 5030–5035.