

- LECHIN, F., GOMEZ, F., VAN DER DIJS, B. & LECHIN, E. (1980a) Distal colon motility in schizophrenic patients. *Journal of Clinical Pharmacology*, **20**, 459–64.
- VAN DER DIJS, B., GOMEZ, F., VALLS, J. M., ACOSTA, E. & AROCHA, L. (1980b) Pharmacomanometric studies of colonic motility as a guide to the chemotherapy of schizophrenia. *The Journal of Clinical Pharmacology*, **20**, 664–71.
- (1979a) Effects of dopaminergic blocking agents on distal colon motility. *Journal of Clinical Pharmacology*, **19**, 617–25.
- (1979b) Dopamine and distal colon motility. *Digestive Diseases and Sciences*, **24**, 86–7.
- (1981a) Colon motility and psychological traits in irritable bowel syndrome. *Digestive Diseases and Sciences*, **26**, 474–5.
- (1981b) Clonidine therapy for psychosis and tardive dyskinesia. *American Journal of Psychiatry*, **138**, 390.
- (in press) Intestinal pharmacomanometry and glucose tolerance: evidences of two antagonistic dopaminergic mechanisms in the human. *Biological Psychiatry*.

DRUG SIDE-EFFECTS AND BRAIN DAMAGE

DEAR SIR,

Neuroleptic drugs induce extrapyramidal disturbances which mimic Parkinson's disease, but this is not taken to mean that they can cause Parkinson's disease: a patient whose Parkinsonism continues long after neuroleptic withdrawal is assumed to have been suffering from incipient Parkinson's disease. Involuntary movement disorders are another group of extrapyramidal disturbances and, *a priori*, it would hardly be surprising if neuroleptics induced conditions mimicking these. Such extrapyramidal disturbances would likewise be expected to be reversible on drug withdrawal; if the disorder persisted, the directly analogous inference to be drawn is that an underlying disease process was responsible.

For several years psychological (Johnstone *et al*, 1981; Owens and Johnstone, 1980) and radiological (Weinberger *et al*, 1979) evidence has been accumulating of organic brain damage in chronic schizophrenics unrelated to drug treatment, and further studies (Kleinman, 1981; Andreasen, 1981; Owens, 1981), described at the recent Annual Meeting of the Royal College of Psychiatrists, indicate an association between brain damage, irreversible movement disorder, and negative symptoms in chronic schizophrenic patients. It is thus becoming increasingly apparent that the involuntary movement disorders seen in neuroleptic-treated schizophrenic patients may be categorized as either drug-induced, if drug withdrawal

is followed by their disappearance, or as the result of structural brain disease with or without superimposed drug-induced effects, if the movements persist following drug withdrawal.

The term 'tardive dyskinesia' should be reserved for those cases in whom irreversible movement disorders occur late in the course of the schizophrenic or other brain disease processes.

A. A. SCHIFF

E. R. Squibb & Sons Ltd,
147–49 Staines Road,
Hounslow TW3 3JA

References

- ANDREASEN, N. (1981) Positive and Negative Symptoms in Relation to CT Scan Findings. Paper given at Annual Meeting of The Royal College of Psychiatrists.
- JOHNSTONE, E. C., CROW, T. J., FRITH, C. D., STEVENS, M., KREEL, L. & HUSBAND, J. (1978) The dementia of dementia praecox. *Acta Psychiatrica Scandinavica*, **57**, 305–24.
- KLEINMAN, J. (1981) CT Scan Studies at N.I.M.H. Paper given at Annual Meeting of The Royal College of Psychiatrists.
- OWENS, D. G. C. (1981) Do Neuroleptic Drugs Cause Tardive Dyskinesia? Paper given at Annual Meeting of The Royal College of Psychiatrists.
- & JOHNSTONE, E. C. (1980) The disabilities of chronic schizophrenia—their nature and the factors contributing to their development. *British Journal of Psychiatry*, **136**, 384–95.
- WEINBERGER, D. R., TORREY, E. F., NEOPHYTIDES, A. N. & WYATT, R. J. (1979) Lateral cerebral ventricular enlargement in chronic schizophrenia. *Archives of General Psychiatry*, **36**, 735–9.

SUICIDE IN FAMILIES: DRAW A LIFE-CHART

DEAR SIR,

The suicidal family described by Dr Khin-Maung-Zaw (*Journal*, July 1981, **139**, 68–69) is of considerable interest as an example of violent suicide and non-fatal deliberate self harm in several members of a family extending over more than one generation and including suicide in identical twins. Whether 'it is at least likely that there was a genetic predisposition to violent suicide in the family' does however still seem to be a matter for conjecture on the basis of the evidence presented in the paper, and the interactional effects of illness events may well have been more important than the author suggests.

To illustrate this, I have tabulated the data along the lines of a Meyer life chart which demonstrates how the suicides seem to have triggered the onset of