

ON THE CHARACTERISATION OF ALTERNATING GROUPS BY CODEGREES

MALLORY DOLORFINO , LUKE MARTIN , ZACHARY SLONIM ,
YUXUAN SUN  and YONG YANG  

(Received 16 October 2023; accepted 18 November 2023; first published online 26 January 2024)

Abstract

Let G be a finite group and $\text{Irr}(G)$ the set of all irreducible complex characters of G . Define the codegree of $\chi \in \text{Irr}(G)$ as $\text{cod}(\chi) := |G : \ker(\chi)|/\chi(1)$ and let $\text{cod}(G) := \{\text{cod}(\chi) \mid \chi \in \text{Irr}(G)\}$ be the codegree set of G . Let A_n be an alternating group of degree $n \geq 5$. We show that A_n is determined up to isomorphism by $\text{cod}(A_n)$.

2020 Mathematics subject classification: primary 20D06; secondary 20C15.

Keywords and phrases: codegree, alternating group.

1. Introduction

Let G be a finite group and $\text{Irr}(G)$ the set of all irreducible complex characters of G . For any $\chi \in \text{Irr}(G)$, define the codegree of χ as $\text{cod}(\chi) := |G : \ker(\chi)|/\chi(1)$ and the codegree set of G as $\text{cod}(G) := \{\text{cod}(\chi) \mid \chi \in \text{Irr}(G)\}$. We refer the reader to the authors' previous paper [8] for the current literature on codegrees.

The following conjecture appears in the *Kourovka Notebook of Unsolved Problems in Group Theory* [12, Question 20.79].

CODEGREE VERSION OF HUPPERT'S CONJECTURE. Let H be a finite nonabelian simple group and G a finite group such that $\text{cod}(H) = \text{cod}(G)$. Then $G \cong H$.

In [8], the authors verified the conjecture for all sporadic simple groups. In this paper, we provide a general proof verifying this conjecture for all alternating groups of degree greater than or equal to 5.

THEOREM 1.1. *Let A_n be an alternating group of degree $n \geq 5$ and G a finite group. If $\text{cod}(G) = \text{cod}(A_n)$, then $G \cong A_n$.*

This research was conducted under NSF-REU grant DMS-1757233, DMS-2150205 and NSA grant H98230-21-1-0333, H98230-22-1-0022 by Dolorfino, Martin, Slonim and Sun during the Summer of 2022 under the supervision of Yang. Yang was also partially supported by a grant from the Simons Foundation (#918096).

© The Author(s), 2024. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

Throughout the paper, we follow the notation used in Isaacs’ book [10] and the ATLAS of Finite Groups [6].

2. Proof of Theorem 1.1

First, we note that the cases $n = 5, 6$ and 7 have already been proven in [1, 2], so in the following, we always assume that $n > 7$. Now, let G be a minimal counterexample and N be a maximal normal subgroup of G . So $\text{cod}(G) = \text{cod}(A_n)$ and G/N is simple. By [8, Lemma 2.5], $\text{cod}(G/N) \subseteq \text{cod}(A_n)$. Then, by [9, Theorem B], $G/N \cong A_n$ so $N \neq 1$ since $G \not\cong A_n$.

Step 1: N is a minimal normal subgroup of G .

Suppose L is a nontrivial normal subgroup of G with $L < N$. Then by [8, Lemma 2.6], $\text{cod}(G/N) \subseteq \text{cod}(G/L) \subseteq \text{cod}(G)$. However, $\text{cod}(G/N) = \text{cod}(A_n) = \text{cod}(G)$, so equality must be attained in each inclusion. Thus, $\text{cod}(G/L) = \text{cod}(A_n)$ which implies that $G/L \cong A_n$ since G is a minimal counterexample. This is a contradiction since we also have $G/N \cong A_n$, but $L < N$.

Step 2: N is the only nontrivial, proper normal subgroup of G .

Otherwise, we assume M is another proper nontrivial normal subgroup of G . If N is included in M , then $M = N$ or $M = G$ since G/N is simple, which is a contradiction. Then $N \cap M = 1$ and $G = N \times M$. Since M is also a maximal normal subgroup of G , we have $N \cong M \cong A_n$. Choose $\psi_1 \in \text{Irr}(N)$ and $\psi_2 \in \text{Irr}(M)$ such that $\text{cod}(\psi_1) = \text{cod}(\psi_2) = \max(\text{cod}(A_n))$. Set $\chi = \psi_1 \cdot \psi_2 \in \text{Irr}(G)$. Then $\text{cod}(\chi) = (\max(\text{cod}(A_n)))^2 \notin \text{cod}(G)$, which is a contradiction.

Step 3: χ is faithful, for each nontrivial $\chi \in \text{Irr}(G|N) := \text{Irr}(G) - \text{Irr}(G/N)$.

From the proof of [8, Lemma 2.5],

$$\text{Irr}(G/N) = \{\hat{\chi}(gN) = \chi(g) \mid \chi \in \text{Irr}(G) \text{ and } N \leq \ker(\chi)\}.$$

By the definition of $\text{Irr}(G|N)$, it follows that if $\chi \in \text{Irr}(G|N)$, then $N \not\leq \ker(\chi)$. Thus, since N is the unique nontrivial, proper, normal subgroup of G , $\ker(\chi) = G$ or $\ker(\chi) = 1$. Therefore, $\ker(\chi) = 1$ for all nontrivial $\chi \in \text{Irr}(G|N)$.

Step 4: N is an elementary abelian group.

Suppose that N is not abelian. Since N is a minimal normal subgroup, by [7, Theorem 4.3A(iii)], $N = S^n$, where S is a nonabelian simple group and $n \in \mathbb{Z}^+$. By [14, Lemma 4.2] and [11, Theorem 4.3.34], there is a nontrivial character $\chi \in \text{Irr}(N)$ which extends to some $\psi \in \text{Irr}(G)$. Now, $\ker(\psi) = 1$ by Step 3, so $\text{cod}(\psi) = |G|/\psi(1) = |G/N| \cdot |N|/\chi(1)$. However, by assumption, $\text{cod}(G) = \text{cod}(A_n) = \text{cod}(G/N)$. Thus, $\text{cod}(\psi) \in \text{cod}(G) = \text{cod}(G/N)$, so $\text{cod}(\psi) = |G/N|/\phi(1)$ for some $\phi \in \text{Irr}(G/N)$. Hence, $|G/N|$ is divisible by $\text{cod}(\psi)$ which contradicts the fact that $\text{cod}(\psi) = |G/N| \cdot |N|/\chi(1)$, as $\chi(1) \neq |N|$. Thus, N must be abelian.

Now to show that N is elementary abelian, let a prime p divide $|N|$. Then N has a p -Sylow subgroup K , and K is the unique p -Sylow subgroup of N since N is abelian,

so K is characteristic in N . Thus, K is a normal subgroup of G , so $K = N$ as N is minimal. Thus, $|N| = p^n$. Now, take the subgroup $N^p = \{n^p \mid n \in N\}$ of N , which is proper by Cauchy's theorem. Since N^p is characteristic in N , it must be normal in G , so N^p is trivial by the uniqueness of N . Thus, every element of N has order p and N is elementary abelian.

Step 5: $C_G(N) = N$.

First note that since N is normal, $C_G(N) \leq G$. Additionally, since N is abelian by Step 4, $N \leq C_G(N)$. By the maximality of N , we must have $C_G(N) = N$ or $C_G(N) = G$. If $C_G(N) = N$, we are done.

If not, then $C_G(N) = G$, so N must be in the centre of G . Then since N is the unique minimal normal subgroup of G by Step 2, $|N|$ must be prime. If not, there always exists a proper nontrivial subgroup K of N , and K is normal since it is contained in $Z(G)$, contradicting the minimality of N . Hence, we have $N \leq Z(G)$ which implies that $Z(G) \cong N$. This is because N is a maximal normal subgroup of G so if not, we would have $Z(G) = G$, implying G is abelian which is a contradiction. Thus, N is isomorphic to a subgroup of the Schur multiplier of G/N by [10, Corollary 11.20].

Now, we note that it is well known that for $n > 7$, the Schur multiplier of A_n is \mathbb{Z}_2 , so $G \cong 2.A_n$ [17]. From [3, Theorem 4.3], $2.A_n$ always has a faithful irreducible character χ of degree $2^{\lfloor (n-2)/2 \rfloor}$. Recall that by Step 2, there is only one nontrivial proper normal subgroup of $G \cong 2.A_n$. In particular, $N \cong \mathbb{Z}_2$ is the only nontrivial proper normal subgroup of G . Thus, $|\ker(\chi)| = 1$ or 2 . Then $\text{cod}(\chi) = |2.A_n : \ker(\chi)|/\chi(1)$. If $|\ker(\chi)| = 1$, then $\text{cod}(\chi) = n!/2^{\lfloor (n-2)/2 \rfloor}$, and if $|\ker(\chi)| = 2$, then $\text{cod}(\chi) = (n!/2)/2^{\lfloor (n-2)/2 \rfloor} = n!/2^{\lfloor n/2 \rfloor}$. In either case, for any prime $p \neq 2$, $|\text{cod}(\chi)|_p = |n!|_p = |A_n|_p$. However, $\text{cod}(\chi) \in \text{cod}(A_n)$ since $\text{cod}(G) = \text{cod}(A_n)$. Therefore, there is a character degree of A_n which is a power of 2.

However, from [13], for $n > 7$, A_n only has a character degree equal to a power of 2 when $n = 2^d + 1$ for some positive integer d . In this case, $2^d = n - 1 \in \text{cd}(A_n)$ so we need $|A_n|/n - 1 = |2.A_n|/2^{\lfloor (n-2)/2 \rfloor}$ or $|2.A_n|/2^{\lfloor n/2 \rfloor}$. Hence,

$$\frac{1}{n - 1} = \frac{2}{2^{\lfloor (n-2)/2 \rfloor}} = \frac{1}{2^{\lfloor (n-2)/2 \rfloor - 1}} \quad \text{or} \quad \frac{1}{2^{\lfloor n/2 \rfloor - 1}}$$

so $n - 1 = 2^{\lfloor (n-2)/2 \rfloor - 1}$ or $2^{\lfloor n/2 \rfloor - 1}$. However, the only integer solution to either of these equations occurs when $n = 9$ and $9 - 1 = 8 = 2^3 = 2^{\lfloor 9/2 \rfloor - 1}$. In this case, we check the ATLAS [6] to find that the codegree sets of A_9 and $2.A_9$ do not have the same order. This is a contradiction, so $C_G(N) = N$.

Step 6. Let λ be a nontrivial character in $\text{Irr}(N)$ and $\vartheta \in \text{Irr}(I_G(\lambda)|\lambda)$, the set of irreducible constituents of $\lambda^{I_G(\lambda)}$, where $I_G(\lambda)$ is the inertia group of λ in G . Then $|I_G(\lambda)|/\vartheta(1) \in \text{cod}(G)$. Also, $\vartheta(1)$ divides $|I_G(\lambda)/N|$ and $|N|$ divides $|G/N|$. Lastly, $I_G(\lambda) < G$, that is, λ is not G -invariant.

Let λ be a nontrivial character in $\text{Irr}(N)$ and $\vartheta \in \text{Irr}(I_G(\lambda)|\lambda)$. Let χ be an irreducible constituent of ϑ^G . By [10, Corollary 5.4], $\chi \in \text{Irr}(G)$, and by [10, Definition 5.1], we have $\chi(1) = (|G|/|I_G(\lambda)|) \cdot \vartheta(1)$. Moreover, $\ker(\chi) = 1$ by Step 2, and thus

$\text{cod}(\chi) = |G|/\chi(1) = |I_G(\lambda)|/\vartheta(1)$, so $|I_G(\lambda)|/\vartheta(1) \in \text{cod}(G)$. Now, since N is abelian, $\lambda(1) = 1$, so we have $\vartheta(1) = \vartheta(1)/\lambda(1)$ which divides $|I_G(\lambda)|/|N|$, so $|N|$ divides $|I_G(\lambda)|/\vartheta(1)$. Moreover, $\text{cod}(G) = \text{cod}(G/N)$, and all elements in $\text{cod}(G/N)$ divide $|G/N|$, so $|N|$ divides $|G/N|$.

Next, we want to show $I_G(\lambda)$ is a proper subgroup of G . To reach a contradiction, assume $I_G(\lambda) = G$. Then $\ker(\lambda) \trianglelefteq G$. From Step 2, $\ker(\lambda) = 1$, and from Step 4, N is a cyclic group of prime order. Thus, by the Normaliser–Centraliser theorem, $G/N = \mathbf{N}_G(N)/\mathbf{C}_G(N) \leq \text{Aut}(N)$ so G/N is abelian, which is a contradiction.

Step 7: Final contradiction.

From Step 4, N is an elementary abelian group of order p^m for some prime p and integer $m \geq 1$. By the Normaliser–Centraliser theorem, $A_n \cong G/N = \mathbf{N}_G(N)/\mathbf{C}_G(N) \leq \text{Aut}(N)$ and $m > 1$. Note that in general, $\text{Aut}(N) \cong \text{GL}(m, p)$. By Step 6, $|N|$ divides $|G/N|$, so $|N| = p^m$ divides $|A_n|$ and $G/N \cong A_n \leq \text{GL}(m, p)$. We prove by contradiction that this cannot occur.

First, we claim that if p^m divides $|A_n|$ and $A_n \leq \text{GL}(m, p)$, then p must equal 2. To show this, we note that for $p > 2$, by [4], if p^m divides $|A_n|$, then $m < n/2$. However, by [16, Theorem 1.1], if $n > 6$, the minimal faithful degree of a modular representation of A_n over a field of characteristic p is at least $n - 2$. Since embedding A_n as a subgroup of $\text{GL}(m, p)$ is equivalent to giving a faithful representation of degree m over a field of characteristic p , we have $m \geq n - 2$. This is a contradiction since $n/2 > n - 2$ implies $n < 4$. Therefore, $p = 2$.

Now, let $p = 2$. As above, from [4], we obtain $|n!|_2 \leq 2^{n-1}$. Thus, if 2^m divides $|A_n|$, then $2^m \leq |A_n|_2 \leq 2^{n-2}$ so $m \leq n - 2$. We will deal first with $n > 8$ and then treat the case $n = 8$ later. For $n > 8$, [15, Theorem 1.1] shows that the minimal faithful degree of a modular representation of A_n over a field of characteristic 2 is at least $n - 2$. Therefore, we must have $m \geq n - 2$, so we have equality, $m = n - 2$.

Let $\lambda \in \text{Irr}(N)$, $\vartheta \in \text{Irr}(I_G(\lambda)|\lambda)$ and $T := I_G(\lambda)$. Then $1 < |G : T| < |N| = 2^{n-2}$ for $|G : T|$ is the number of all conjugates of λ . By Step 5, $|T|/\vartheta(1) \in \text{cod}(G)$ and moreover $|N|$ divides $|T|/\vartheta(1)$. Since $|N| = |N|_2 = |A_n|_2$ and $\text{cod}(G) = \text{cod}(A_n)$, it follows that $\|T|/\vartheta(1)\|_2 = |N|$. Thus, $\|T/N|/\vartheta(1)\|_2 = 1$ so the 2-parts of $|T/N|$ and $\vartheta(1)$ are equal. Thus, for every $\vartheta \in \text{Irr}(T | \lambda)$, we have $|\vartheta(1)|_2 = |T/N|_2$. However, $|T/N| = \sum_{\vartheta \in \text{Irr}(T|\lambda)} \vartheta(1)^2$. Hence, if $|\vartheta(1)|_2 = 2^k \geq 2$ for every $\vartheta \in \text{Irr}(T | \lambda)$, we would have $|T/N|_2 = 2^{2k}$, which contradicts the fact that $|\vartheta(1)|_2 = |T/N|_2$. Therefore, $|T/N|_2 = 1$. Thus, since $|G/N|_2 \geq |N| = 2^{n-2}$, we have $|G : T|_2 = |G/N : T/N|_2 \geq 2^{n-2}$, so $|G : T| \geq 2^{n-2} = |N|$, which is a contradiction.

Now we turn to the case $n = 8$. We have $p = 2$ and $m = 4, 5$ or 6 . In this case, $A_8 \cong \text{GL}(4, 2)$ and 2^6 divides $|A_8|$. We look at each possibility for m in turn. If $m = 6$, then $|N|_2 = |A_8|_2$. For this case, the same argument as above holds since $6 = 8 - 2$, and we reach a contradiction.

Second, let $m = 5$. As above, $|G : T| < |N| = 2^5$ and $|T|/\vartheta(1) \in \text{cod}(G)$ such that 2^5 divides $|T|/\vartheta(1)$. Further, $\|T/N|/\vartheta(1)\|_2 \leq 2$ so $|T/N|_2 \leq 4$ and $|G/N : T/N|_2 \geq 16$. Thus, 16 divides $|G/N : T/N|$ and $|G/N : T/N| < 32$. However, we may check the index

of all subgroups of $G/N \cong A_8$ using [6] and find that none of them satisfy these two properties.

Third, let $m = 4$. Then $G/N \cong A_8 \cong \text{GL}(4, 2)$ and $N = (\mathbb{Z}_2)^4$ so G is an extension of $\text{GL}(4, 2)$ by N . We may computationally calculate the codegree set for any such group using MAGMA [5]. There are only four such nonisomorphic extensions and we find that none of them have the same codegree set as A_8 . (The MAGMA code is available at <https://github.com/zachslonim/Characterizing-Alternating-Groups-by-Their-Codegrees>.) In every case, $|N| = p^m$ produces a contradiction, so $N = 1$ and $G \cong A_n$.

Acknowledgements

The authors gratefully acknowledge the financial support of NSF and NSA, and also thank Texas State University for providing a great working environment and support. The authors would also like to thank Professor Richard Stanley for his help.

References

- [1] N. Ahanjideh, ‘Nondivisibility among irreducible character co-degrees’, *Bull. Aust. Math. Soc.* **105** (2022), 68–74.
- [2] A. Bahri, Z. Akhlaghi and B. Khosravi, ‘An analogue of Huppert’s conjecture for character codegrees’, *Bull. Aust. Math. Soc.* **104**(2) (2021), 278–286.
- [3] C. Bessenrodt and J. B. Olsson, ‘Prime power degree representations of the double covers of the symmetric and alternating groups’, *J. London Math. Soc.* **66**(2) (2002), 313–324.
- [4] C. Bessenrodt, H. P. Tong-Viet and J. Zhang, ‘Huppert’s conjecture for alternating groups’, *J. Algebra* **470** (2017), 353–378.
- [5] J. C. W. Bosma and C. Playoust, ‘The Magma algebra system I: the user language’, *J. Symbolic Comput.* **24** (1997), 235–265.
- [6] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups* (Clarendon Press, Oxford, 1985).
- [7] J. D. Dixon and B. Mortimer, *Permutation Groups*, Graduate Texts in Mathematics, 163 (Springer, New York, 1996).
- [8] M. Dolorfino, L. Martin, Z. Slonim, Y. Sun and Y. Yang, ‘On the characterisation of sporadic simple groups by codegrees’, *Bull. Aust. Math. Soc.*, to appear. Published online (27 March 2023); doi:[10.1017/S0004972723000187](https://doi.org/10.1017/S0004972723000187).
- [9] N. N. Hung and A. Moreto, ‘The codegree isomorphism problem for finite simple groups’, Preprint, 2023, [arXiv:2301.00446](https://arxiv.org/abs/2301.00446).
- [10] I. M. Isaacs, *Character Theory of Finite Groups* (Academic Press, New York, 1976).
- [11] G. James and A. Kerber, *The Representation Theory of the Symmetric Group* (Addison-Wesley, Reading, MA, 1981).
- [12] E. I. Khukrho and V. D. Mazurov, *Unsolved Problems in Group Theory*, The Kourovka Notebook, 20 (Russian Academy of Sciences, Novosibirsk, 2022).
- [13] G. Malle and A. E. Zalesskii, ‘Prime power degree representations of quasi-simple groups’, *Arch. Math. (Basel)* **77** (2001), 461–468.
- [14] A. Moretó, ‘Complex group algebra of finite groups: Brauer’s problem 1’, *Adv. Math.* **208** (2007), 236–248.
- [15] A. Wagner, ‘The faithful linear representations of least degree of S_n and A_n over a field of characteristic 2’, *Math. Z.* **151** (1976), 127–138.
- [16] A. Wagner, ‘The faithful linear representations of least degree of S_n and A_n over a field of odd characteristic’, *Math. Z.* **154** (1977), 104–113.
- [17] R. A. Wilson, *The Finite Simple Groups* (Springer, London, 2009).

MALLORY DOLORFINO, Department of Mathematics,
Kalamazoo College, Kalamazoo, Michigan, USA
e-mail: mallory.dolorfino19@kzoo.edu

LUKE MARTIN, Department of Mathematics,
Gonzaga University, Spokane, Washington, USA
e-mail: lwmartin2019@gmail.com

ZACHARY SLONIM, Department of Mathematics,
University of California, Berkeley, Berkeley, California, USA
e-mail: zachslonim@berkeley.edu

YUXUAN SUN, Department of Mathematics and Statistics,
Haverford College, Haverford, Pennsylvania, USA
e-mail: ysun1@haverford.edu

YONG YANG, Department of Mathematics,
Texas State University, San Marcos, Texas, USA
e-mail: yang@txstate.edu