



## Predictive capacity and cut-off points of adiposity indices for body fat prediction according to adolescent periods

Núbia de Souza de Moraes<sup>1\*</sup>, Valter Paulo Neves Miranda<sup>2</sup>, Sarah Aparecida Vieira Ribeiro<sup>1</sup>, Patrícia Feliciano Pereira<sup>1</sup>, Eliana Carla Gomes de Souza<sup>1</sup>, Sylvia do Carmo Castro Franceschini<sup>1</sup> and Sílvia Eloiza Priore<sup>1</sup>

<sup>1</sup>Department of Nutrition and Health, Federal University of Viçosa, Avenida Peter Henry Rolfs, s/n. CEP: 36.570-000 Viçosa, Minas Gerais, Brazil

<sup>2</sup>Department of Physical Education, Federal University of Viçosa, Avenida Peter Henry Rolfs, s/n. CEP: 36.570-000 Viçosa, Minas Gerais, Brazil

(Submitted 14 July 2020 – Final revision received 20 January 2021 – Accepted 21 January 2021 – First published online 29 January 2021)

### Abstract

Overweight is increasing in the adolescent population and became a public health problem in the world. This study aimed to identify the body adiposity indices (BAI) with the best capacity to predict excess fat tissue and propose cut-off points for them, according to sex and adolescence period. This is a cross-sectional study. We calculated BMI, conicity index (CI), BAI, body roundness index (BRI), waist:height ratio (WtHR) and waist:hip ratio. Predictive capacity and cut-off points of adiposity indices were established by ROC (receiver operating characteristic) curves. We determined AUC-ROC and CI, stratified by sex and adolescence period. The best index to identify excess body fat in 10–13-year-old female adolescents was the WtHR (AUC = 0.92), like the BAI in girls aged from 14 to 16 years old (AUC = 0.87) and 17 to 19 years old (AUC = 0.80). In male adolescents aged from 10 to 13 years old and 14 to 16 years old, the best index was the WtHR (AUC = 0.93 and AUC = 0.8, respectively), like the BAI in boys aged from 17 to 19 years old (AUC = 0.95). The use of indices with specific cut-off points for each period of adolescence and according to sex is important for the reliable diagnosis of excess body fat. It is advisable to use indices together to obtain a more accurate assessment. Thus, the WtHR and BAI are reproducible and reliable, with high sensitivity and specificity values, and can be used together with the BMI.

**Key words:** Adolescent: Anthropometry: Body composition: Receiver operating characteristic curve

Overweight is increasing in the adolescent population and became a public health issue in Brazil and worldwide<sup>(1,2)</sup>. In Brazil, from 1974/1975 to 2008/2009, overweight increased from 3.7% to 21.7% among males from 10 to 19 years old, and from 7.6% to 19.4% among females in the same age range<sup>(3)</sup>. In 2015, among Brazilian adolescents aged between 13 and 17 years, 23.8% of the girls and 23.7% of the boys were overweight<sup>(4)</sup>. At the global scale, the prevalence of obesity among children and adolescents aged between 5 and 19 years increased from 0.7% and 0.9% in 1975, to 5.6% and 7.8% in 2016 in females and males, respectively<sup>(5)</sup>.

During adolescence, changes occur in body composition that vary according to sex, age, stage of sexual maturation and environmental factors of each individual<sup>(6)</sup>. The initial period is marked by growth spurt and appearance of secondary sexual characteristics,

as well as an increase in body fat accumulation, which occurs mainly in female sex, being an essential physiological gain for growth and necessary for the beginning and maintenance of menstrual cycles. In boys, the gain of adipose tissue is slower, and even loss may occur. In the middle and late periods, the completion of growth and morphological development occur<sup>(7)</sup>.

Due to the changes that occur in this period of life, adolescence becomes a vulnerable period for the occurrence of obesity and its associated complications<sup>(8,9)</sup>. Excess body fat is related to the development of diseases such as diabetes mellitus<sup>(10,11)</sup>, hypertension<sup>(12)</sup>, the metabolic syndrome<sup>(13)</sup> and CVD<sup>(12)</sup>, whose onset occurs in adolescence and progresses through adulthood<sup>(8,12)</sup>. In addition, obesity is a risk factor for psychosocial diseases such as body image disorders, which may evolve and contribute to the development of body image disorders<sup>(14)</sup>.

**Abbreviations:** BAI, body adiposity index; BF%, body fat percentage; BRI, body roundness index; CI, conicity index; DEXA, Dual-Energy X-ray absorptiometry; ROC, receiver operating characteristic; WC, waist circumference; WHR, waist:hip ratio; WtHR, waist:height ratio.

\* **Corresponding author:** N. S. Moraes, email [nubia.s.moraes25@gmail.com](mailto:nubia.s.moraes25@gmail.com)

This context highlights the importance and need of monitoring the health situation of adolescents. Accordingly, there is a clear need to carry out this assessment according to sex and age, as the several changes in body composition influence the reliability of the indices used in anthropometric evaluation<sup>(15)</sup>.

BMI is the most commonly used indicator to assess the nutritional status of individuals at any age. However, this is not the most reliable method for the assessment of excess body fat and the risk of associated diseases, such as cardiometabolic diseases<sup>(16–18)</sup>. A limitation that needs to be considered regarding this index is that it does not distinguish between lean and fat mass<sup>(17,19)</sup>. Particularly in adolescents, studies show that individuals classified as eutrophic based on their BMI have excess body fat, thus confirming that this limitation should be accounted for when assessing nutritional status<sup>(18,20)</sup>. Hence, in addition to the waist:height ratio (WtHR), other indices have been proposed to enable a more reliable assessment of excess body fat and, consequently, to predict the risk of developing cardiometabolic alterations.

The conicity index (CI)<sup>(21)</sup>, body adiposity index (BAI)<sup>(22)</sup> and the body roundness index (BRI)<sup>(23)</sup> were developed and have been used to assess body fat in adults. However, there is a limited number of studies that analyse the relevant capacity of these and other anthropometric indexes to evaluate the body fat in paediatric population. In addition, there are no reliable cut-off points that can verify with scientific authenticity the excess accumulation of adipose tissue in the adolescents, as well as the associated risk factors. Thus, this study aims to identify the BAI with the best capacity to predict excess fat and propose cut-off points for them, according to sex and period of adolescence.

## Methodology

### Study design and sample

This is a cross-sectional study which comprised the three periods of adolescence (early: from 10 to 13 years old; middle: from 14 to 16 years old; late: from 17 to 19 years old)<sup>(6)</sup> with 1188 adolescents: 804 female and 384 male, aged from 10 to 19 years old.

We conducted the research with data collected in cities located in the microregion of Viçosa, comprising twenty municipalities, which is part of Zona da Mata mesoregion, state of Minas Gerais, Brazil. The microregion has an estimated population of adolescents between 10 and 19 years of 39 270<sup>(10)</sup>; thus, the evaluated sample corresponds to 3% of the total population of the same age group.

The data used in this article were obtained from twelve cross-sectional studies developed by a research group of Graduate Program in Nutrition Sciences at the Federal University of Viçosa. Data collections were carried out with students from public schools, between 2002 and 2017.

In general, the eligibility criteria used in all databases were: adolescents aged between 10 and 19 years, girls who had already undergone menarche at least 1 year earlier, voluntary participants in good health conditions and whose participation was allowed by an authorisation signed by their parents or legal guardians (if under 18 years of age). As exclusion criteria, we verified whether the adolescents could normally engage in any kind of physical activity, whether they had any chronic or

communicable disease, some obesity-related factor or inflammatory diseases, whether they have been using some kind of controlled drug or medicine likely to influence metabolic function and whether they are currently participating in any other study addressing either body composition assessment or nutritional status control. Both criteria were observed throughout the selection and data collection processes, through contact with the students at the school, the Health Division and the research laboratory at the Federal University of Viçosa.

The databases of these studies were analysed separately and thoroughly until they were included in the study. After the analysis, all these databases were merged and we proceeded with the selection of the adolescents. We excluded duplicate data of adolescents evaluated more than once during the same period. Data from participants assessed more than once, but in different periods, were retained.

We calculated the study power by the OpenEpi® online programme ([www.OpenEpi.com](http://www.OpenEpi.com)), considering the outcome of the WtHR among the groups with high and adequate body fat percentage (BF%). Based on the means and standard deviations of the WtHR in the group of adolescents with excess body fat (0.48 (SD 0.069) cm) and eutrophic (0.41 (SD 0.032) cm), the result of the analysis showed power equal to 100%.

### Anthropometric and body composition assessment

All anthropometric measurements were performed by researchers, who were trained during the conduct of the pilot studies, before the start of data collection. The same protocols, instruments and devices were used for anthropometric and body composition measurements in all studies, so the information collected are homogeneous, which guarantees the relevance of the results found in this research.

The weight was measured on an electronic digital scale (Kratos®), with a maximum capacity of 150 kg and a sensitivity of 50 g. Weighing was performed according to the techniques proposed by WHO<sup>(24)</sup>. For greater measurement reliability, the calibration of the scale was verified with a standard weight of 1 kg.

We measured height using a portable stadiometer (Altuxata®), up to 220 cm in length. We performed the measurements following the techniques recommended by WHO<sup>(24)</sup>.

We used a flexible and inelastic tape measure (Cardiomed®) with a maximum length of 2 m to measure waist (WC) and hip circumferences, being careful not to compress soft tissues. We made the measurements in duplicate, assuming a maximum variation of 0.5 cm and in case of variation greater than this value, we performed a third measurement, using the mean between the two nearest measurements.

WC was measured at the midpoint between the lower margin of the last rib and the iliac crest, in the horizontal plane<sup>(25)</sup> and the hip circumference in the gluteal region, surrounding the largest horizontal portion between the waist and knees<sup>(26)</sup>.

We used the gold standard in the assessment of body composition, Dual-Energy X-ray absorptiometry (DEXA) equipment (Lunar Prodigy Advance DXA System – analysis version: 13.31, GE Healthcare), to estimate the BF%, in the morning, with all participants fasting for 12 h and following the assessment protocol. We used sex-specific cut-off points to classify excess body fat. Accordingly, we classified values  $\geq 25$  and 20% for females and males, respectively, as excess fat<sup>(27)</sup>.



### Body adiposity indices

The WtHR was calculated by the quotient of WC measurement (cm) by the height measurement (cm), waist:hip ratio (WHR) by the relationship between WC (cm) and hip circumference (cm). BMI was assessed according to age and sex (BMI/I) by WHO AnthroPlus software, using *z*-score values<sup>(28)</sup>.

The CI was calculated according to the formula proposed by Valdez<sup>(21)</sup>; the BAI according to Bergman *et al.*<sup>(22)</sup> and the BRI are based on the proposal of Thomas *et al.*<sup>(23)</sup>.

### Statistical analyses

The databases were elaborated in Excel software, and to guarantee the validity of the data, they were double-typed. We performed the statistical analysis with the aid of Statistical Package for Social Sciences (SPSS) software for Windows, version 20.0 and MedCalc version 9.3. We adopted as level of rejection of the null hypothesis  $\alpha = 5\%$ . We performed the Shapiro–Wilk test, in which was found that the variables did not present normal distribution.

We presented median and interquartile range values for the descriptive analysis of continuous variables. The Mann–Whitney *U* test was used to compare the sexes and the Kruskal–Wallis test to compare the periods of adolescence for each sex. After verifying statistical difference ( $P < 0.05$ ) by Kruskal–Wallis, we performed Bonferroni's correction for *P* value and again Mann–Whitney *U* test to identify which groups differed from each other.

We established the predictive capacity and cut-off points of BAI by building ROC (receiver operating characteristic) curves. We determined the AUC-ROC and the respective CI (95% CI), stratified by sex and period of adolescence. The sensitivity and 1-specificity values generated by the ROC curves were assessed to identify the best cut-off points, considering the best balance between these values. To classify the AUC values, we used:  $< 0.6$  bad; from 0.6 to 0.7 sufficient; from 0.7 to 0.8 good; from 0.8 to 0.9 very good and from 0.9 to 1.0 excellent<sup>(29)</sup>.

### Ethical aspects

All studies whose databases were included in this research were submitted to and approved by the Ethics Committee on Research with Human Beings of the Federal University of Viçosa. In addition, the present study was submitted to the Ethics Committee on Research with Human Beings of the Federal University of Viçosa and obtained a single approval number (2879661).

When the surveys were conducted, participants and their guardians were informed about their objectives and received the Informed Consent and Agreement Consent Terms for those under 18 years old. Only those who submitted duly signed terms were included.

The adolescents' participation was voluntary, so they could leave the research at any time without any harm and their data were kept confidential.

### Results

Most were female (67.7%), of the total 37.6% were in the middle period of adolescence and 35.2% in the late, with a median age of 16 years old (interquartile range: 13–17). According to BMI assessment, 22.1 and 21.5% of the male and female adolescents, respectively, were with overweight, and by DEXA assessment, 29.4 and 69.4%, respectively, had a high BF%.

Regarding nutritional status classified by BF% (DEXA), a lower proportion of excess body fat was observed in early adolescence (20.0%) compared with the middle (43.8%) and late (36.2%) ( $P < 0.001$ ) period. There was also a difference in the proportion between the middle and late period ( $P = 0.021$ ).

We observed that all measurements and indices showed differences ( $P < 0.05$ ) between periods in females and only WtHR and BRI did not differ ( $P > 0.05$ ) in any period in males (Table 1).

The analysis of the ROC curves showed AUC values considered very good or excellent ( $> 0.8$ ) for BMI, WtHR, BAI and BRI in adolescents in general (Table 2).

In the analysis by period of adolescence and sex, ROC curves indicated AUC values considered very good or excellent ( $> 0.8$ ) for BMI and BAI in females in all periods. In males, WHR had an AUC value below 0.8 in the early and late period of adolescence. The best combinations of sensitivity and specificity values to determine the cut-off points of these indices in predicting body fat in adolescents are shown in Tables 3 and 4.


In girls, from 10 to 13 years old, the cut-off points of 18.03 kg/m<sup>2</sup> for BMI, 0.46 for WtHR, 26.10 for BAI and 2.70 for BRI showed higher predictive capacity of excess body fat. Among those in the middle period, the values of 19.65 kg/m<sup>2</sup> for BMI, 0.42 for WtHR, 25.30 for BAI and 2.12 for BRI had the best combinations between sensitivity and specificity. In those aged from 17 to 19 years, the ideal cut-off values were 20.66 kg/m<sup>2</sup> for BMI, 0.44 for WtHR, 27.52 for BAI and 2.58 for BRI (Table 3).

In boys, in early adolescence, the ideal cut-off values for BMI were 18.18 kg/m<sup>2</sup>, 0.45 for WtHR, 24.41 for BAI and 2.43 for BRI. In the middle period, the cut-off point for BMI was 21.26 kg/m<sup>2</sup>, for WtHR 0.44, for BAI 22.56 and for BRI 2.44. In boys aged from 17 to 19 years, the cut-off points of 24.79 kg/m<sup>2</sup> for BMI, 0.46 for WtHR, 24.25 for BAI and 2.73 for BRI showed better predictive capacity of excess body fat (Table 4).

In female adolescents, BMI was the index with the highest predictive capacity for body fat in the three periods. In the early period, between BMI and WtHR, BRI and BAI, there was no difference ( $P > 0.05$ ). In the middle period, the BMI and BAI showed the same value of AUC ( $P > 0.05$ ). In the late, BMI was higher than all other indices ( $P < 0.05$ ), although BAI displayed good predictive power for body fat (Fig. 1).

For males in the early period, WtHR and BRI showed better predictive capacity for body fat, with equal AUC value ( $P > 0.05$ ). However, WtHR did not differ from BMI, BAI and BRI ( $P > 0.05$ ), as well as BRI showed no difference for BAI and BMI ( $P > 0.05$ ). BRI showed better capacity to predict body fat in the middle period, and it was similar to BMI, WtHR and BAI ( $P > 0.05$ ). In the late, BAI showed higher value of AUC, being similar to BMI, WtHR, CI and BRI ( $P > 0.05$ ) (Fig. 1).





**Figure 1.** Comparison between the receiver operating characteristic (ROC) curves of adiposity indices used to predict body fat, according to the sex and period of adolescence (*n* 1188). —, BMI; ·····, WtHR; - - -, WHR; - · - ·, CI; ·····, BAI; —, BRI.

**Table 1** Comparison of anthropometric and body composition variables between periods of adolescence according to the adolescents' sex, Minas Gerais (*n* 1188)†,‡

Variables	Sex/period of adolescence					
	Female			Male		
	Early ( <i>n</i> 155)	Middle ( <i>n</i> 360)	Late ( <i>n</i> 289)	Early ( <i>n</i> 168)	Middle ( <i>n</i> 87)	Late ( <i>n</i> 129)
Height (m)	1.49 <sup>a</sup>	1.62 <sup>bc</sup>	1.62 <sup>bc</sup>	1.47 <sup>a</sup>	1.70 <sup>b</sup>	1.74 <sup>c</sup>
Weight (kg)	41.50 <sup>a</sup>	54.15 <sup>bc</sup>	55.25 <sup>bc</sup>	38.45 <sup>a</sup>	59.00 <sup>b</sup>	65.45 <sup>c</sup>
BMI (kg/m <sup>2</sup> )	18.03 <sup>a</sup>	20.60 <sup>bc</sup>	21.26 <sup>bc</sup>	17.29 <sup>a</sup>	20.28 <sup>b</sup>	21.45 <sup>c</sup>
WP (cm)	66.50 <sup>a</sup>	71.00 <sup>bc</sup>	73.00 <sup>bc</sup>	63.50 <sup>a</sup>	72.00 <sup>b</sup>	74.60 <sup>c</sup>
WtHR	0.44 <sup>abc</sup>	0.44 <sup>ab</sup>	0.45 <sup>ac</sup>	0.43 <sup>*</sup>	0.42 <sup>*</sup>	0.43 <sup>*</sup>
HP (cm)	79.35 <sup>a</sup>	91.75 <sup>bc</sup>	93.50 <sup>bc</sup>	73.25 <sup>a</sup>	87.50 <sup>b</sup>	92.00 <sup>c</sup>
WHR	0.85 <sup>a</sup>	0.78 <sup>bc</sup>	0.79 <sup>bc</sup>	0.88 <sup>a</sup>	0.83 <sup>bc</sup>	0.83 <sup>bc</sup>
CI	1.18 <sup>a</sup>	1.14 <sup>b</sup>	1.14 <sup>c</sup>	1.17 <sup>a</sup>	1.14 <sup>bc</sup>	1.13 <sup>bc</sup>
BAI (%)	25.27 <sup>a</sup>	26.35 <sup>b</sup>	27.34 <sup>c</sup>	23.21 <sup>a</sup>	21.58 <sup>bc</sup>	22.16 <sup>bc</sup>
BRI	2.34 <sup>abc</sup>	2.25 <sup>ab</sup>	2.49 <sup>ac</sup>	2.17 <sup>*</sup>	1.97 <sup>*</sup>	2.17 <sup>*</sup>
%BF(DEXA)	23.85 <sup>a</sup>	29.54 <sup>bc</sup>	29.22 <sup>bc</sup>	17.47 <sup>a</sup>	13.98 <sup>bc</sup>	13.45 <sup>bc</sup>

WP, waist perimeter; WtHR, waist:height ratio; HP, hip perimeter; WHR, waist:hip ratio; CI, conicity index; BAI, body adiposity index; BRI, body roundness index; %BF, body fat percentage.

Medians in the same row followed by the same letter do not differ for each sex by the Kruskal–Wallis test, supplemented by the Mann–Whitney with Bonferroni correction (*P* < 0.016).

\* *P* > 0.05.

† Early: 10–13 years; Middle: 14–16 years; Late: 17–19 years.

‡ Reference Method: DEXA.



**Table 2** Cut-off points, sensitivity, specificity and AUC of the adiposity indices for body fat prediction in adolescents, Minas Gerais (*n* 1188)†,‡ (95 % confidence intervals)

Adiposity indices	Cut-off point	Sensitivity	95 % CI	Specificity	95 % CI	AUC	95 % CI
BMI	19.65	77.6	74.3, 80.7	71.7	67.7, 75.6	0.83*	0.80, 0.85
WtHR	0.45	62.5	58.8, 66.3	90.3	87.4, 92.7	0.82*	0.80, 0.84
WHR	0.78	36.8	33.2, 40.6	76.9	73.1, 80.5	0.53	0.50, 0.56
CI	1.20	32.7	29.2, 36.5	9.6	90.1, 94.7	0.63	0.60, 0.66
BAI	24.82	82.4	79.3, 85.2	81.4	77.8, 84.7	0.90*	0.88, 0.92
BRI	2.48	65.2	61.5, 68.9	88.7	85.7, 91.4	0.82*	0.80, 0.84

WtHR, waist:height ratio; WHR, waist:hip ratio; CI, conicity index; BAI, body adiposity index; BRI, body roundness index.

\* AUC value rated very good or excellent (> 0.8).

† Reference Method: DEXA.

‡ All AUC values were significant (*P* < 0.05).

**Table 3** Cut-off points, sensitivity, specificity and AUC of the adiposity indices for body fat prediction, according to the period of adolescence, in female adolescents, Minas Gerais (*n* 804)†,‡,§ (95 % confidence intervals)

Adiposity indices	Cut-off Point	Sensitivity	95 % CI	Specificity	95 % CI	AUC	95 % CI
<b>Early</b>							
BMI	18.03	92.7	83.9, 97.6	84.8	75.5, 91.7	0.94*	0.89, 0.97
WtHR	0.46	75.3	63.5, 84.9	95.3	88.5, 98.7	0.92*	0.87, 0.96
WHR	0.86	69.5	57.3, 80.1	79.0	69.0, 87.1	0.71	0.64, 0.78
CI	1.21	68.1	55.8, 78.8	94.1	86.9, 98.1	0.82*	0.75, 0.88
BAI	26.10	75.3	63.5, 84.9	93.0	85.4, 97.4	0.92*	0.87, 0.96
BRI	2.70	76.8	65.1, 86.1	95.3	88.5, 98.7	0.93*	0.87, 0.96
<b>Middle</b>							
BMI	19.65	77.5	72.1, 82.4	82.9	73.4, 90.1	0.87*	0.84, 0.91
WtHR	0.42	71.3	65.6, 76.6	81.8	72.2, 89.2	0.84*	0.80, 0.87
WHR	0.74	79.0	73.7, 83.7	32.9	23.3, 43.8	0.57	0.52, 0.62
CI	1.09	77.9	72.5, 82.7	42.0	31.6, 53.0	0.61	0.56, 0.66
BAI	25.30	76.8	71.4, 81.7	82.9	73.4, 90.1	0.87*	0.84, 0.91
BRI	2.12	70.5	64.8, 75.9	82.9	73.4, 90.1	0.83*	0.79, 0.87
<b>Late</b>							
BMI	20.66	71.8	65.5, 77.8	81.9	71.1, 90.0	0.84*	0.80, 0.88
WtHR	0.44	64.9	58.2, 71.3	84.7	74.3, 92.1	0.79	0.74, 0.83
WHR	0.81	35.9	29.6, 42.7	87.5	77.6, 94.1	0.59	0.53, 0.65
CI	1.14	56.6	49.8, 63.4	69.4	57.5, 79.8	0.60	0.54, 0.65
BAI	27.52	58.5	51.7, 65.2	90.2	81.0, 96.0	0.80*	0.75, 0.85
BRI	2.58	58.5	51.7, 65.2	93.0	84.5, 97.7	0.79	0.74, 0.84

WtHR, waist:height ratio; WHR, waist:hip ratio; CI, conicity index; BAI, body adiposity index; BRI, body roundness index.

\* AUC value rated very good or excellent (> 0.8).

† Early: 10–13 years; Middle: 14–16 years; Late: 17–19 years.

‡ Reference Method: DEXA.

§ All AUC values were significant (*P* < 0.05).

**Discussion**

Among all anthropometric indices assessed, WtHR, BMI, BAI and BRI showed good capacity to predict the excess body fat estimated by DEXA in both sexes and in all periods of adolescence.

Most of the measures and indices assessed in this study showed higher values in female adolescents. In both sexes, most measures and indices showed differences between the periods of adolescence, except for WtHR and BRI. In addition, except for WHR and CI in girls, and WHR, CI, BAI and BF% (DEXA) in boys, the other indices had lower values in the early period.

Other studies assessing the nutritional status of adolescents in relation to age have found that younger individuals had a higher proportion of overweight, according to the BMI classification<sup>(30,31)</sup>. Contrary to what was observed in this study, in which was found that there was no difference in nutritional status classified by BMI in relation to the period of adolescence. However, higher prevalence

of excess body fat was observed in adolescents aged 14 years or more, in both sexes.

In adolescence, body composition varies according to age, sexual maturation and physical growth, with physiological increase of adipose tissue, being higher in females, because this accumulation of body fat is essential for the beginning and maintenance of menstrual cycles and consequently for the reproductive process<sup>(2,31)</sup>. However, it should be considered that this is an age of many changes and therefore is a critical period for the development of obesity<sup>(32,33)</sup>. High-energy eating habits and sedentary lifestyles of adolescents in general may lead to excessive body fat accumulation and obesity<sup>(34,35)</sup>. This condition may lead to the development of cardiometabolic diseases, which may originate in adolescence<sup>(36,37)</sup>.

Changes in physical, cognitive and physiological development are influenced by genetics and environmental factors, so

**Table 4** Cut-off points, sensitivity, specificity and AUC of the adiposity indices for body fat prediction, according to the period of adolescence, in male adolescents, Minas Gerais (*n* 384)†,‡,§ (95 % confidence intervals)

Adiposity indices	Cut-off point	Sensitivity	95 % CI	Specificity	95 % CI	AUC	95 % CI
<b>Early</b>							
BMI	18.18	80.0	68.2, 88.9	87.3	79.4, 93.1	0.89*	0.84, 0.93
WtHR	0.45	75.3	63.1, 85.2	97.0	91.7, 99.4	0.93*	0.88, 0.96
WHR	0.89	61.5	48.6, 73.3	78.6	69.5, 86.1	0.75	0.68, 0.81
CI	1.17	78.4	8.6, 73.3	78.6	69.5, 86.1	0.84*	0.78, 0.89
BAI	24.41	78.4	66.5, 87.7	90.2	82.9, 95.2	0.91*	0.85, 0.94
BRI	2.43	81.5	66.5, 87.7	92.2	85.3, 96.6	0.93*	0.88, 0.96
<b>Middle</b>							
BMI	21.26	90.9	70.8, 98.6	86.1	75.3, 93.5	0.95*	0.89, 0.98
WtHR	0.44	95.4	77.1, 99.2	92.3	82.9, 97.4	0.98*	0.92, 0.99
WHR	0.87	72.7	49.8, 89.2	87.6	77.2, 94.5	0.83*	0.74, 0.90
CI	1.18	72.7	49.8, 89.2	93.8	85.0, 98.3	0.89*	0.81, 0.95
BAI	22.56	100.0	84.4, 100.0	89.2	79.1, 95.5	0.96*	0.90, 0.99
BRI	2.44	95.4	77.1, 99.2	95.3	87.1, 99.0	0.98*	0.92, 0.99
<b>Late</b>							
BMI	24.79	76.9	56.3, 91.0	93.2	86.5, 97.2	0.92*	0.86, 0.96
WtHR	0.46	92.3	74.8, 98.8	85.4	77.1, 91.6	0.94*	0.88, 0.97
WHR	0.83	84.6	65.1, 95.5	62.1	52.0, 71.5	0.74	0.66, 0.81
CI	1.14	92.3	74.8, 98.8	69.9	60.1, 78.5	0.86*	0.79, 0.92
BAI	24.25	88.4	69.8, 97.4	91.2	84.1, 95.9	0.95*	0.90, 0.98
BRI	2.73	92.3	74.8, 98.8	86.4	78.2, 92.4	0.94*	0.88, 0.97

WtHR, waist:height ratio; WHR, waist:hip ratio; CI, conicity index; BAI, body adiposity index; BRI, body roundness index.

\* AUC value rated very good or excellent (> 0.8).

† Early: 10–13 years; Middle: 14–16 years; Late: 17–19 years.

‡ Reference Method: DEXA.

§ All AUC values were significant ( $P < 0.05$ ).

physical characteristics differ among individuals, with bone mass and the ratio between muscle tissue and adipose tissue vary according to sex and age. These differences make it difficult to adopt specific cut-off points for anthropometric indices in adolescence. Therefore, the assessment of body composition is necessary, and the identification of more accurate adiposity indices for estimating body fat and the determination of cut-off points for these indices, in this age group, are important to expand the methods of assessment and monitoring of overweight and obesity. These strategies are essential for disease prevention and health promotion in the adolescent population<sup>(38,39)</sup>.

Therefore, BMI should be used with caution, especially in adolescents, since during the growth spurt, this index may not reliably reflect increases in body fat. In addition, the detection of cardiometabolic risk factors in this age group is complex, mainly due to quick and intense variations that occur in body composition<sup>(40)</sup>.

BMI is widely used as a surrogate measure of obesity. However, its calculation is not specific to assess body composition components, and its limitations should be considered<sup>(41)</sup>. A well-known limitation of BMI is that it does not distinguish between fat and lean mass<sup>(18)</sup>. Indeed, BMI has been shown to be a poor predictor of BF% in children and adolescents<sup>(16)</sup>.

Therefore, BMI should be used with caution, especially in adolescents, for during the growth spurt, this index does not reliably reflect increases in body fat<sup>(40)</sup>. Perhaps, this is because neither the BMI nor other indicator or anthropometric indices have taken into account the changes caused by sexual maturation in the body composition of adolescents of each sex throughout the periods of this life phase. Since this is not an effective index for

assessing excess body fat, it is also not effective for predicting cardiometabolic risk factors<sup>(18,19)</sup>.

A high prevalence of eutrophic adolescents with high BF% has been observed. In a meta-analysis that included thirty-seven studies comparing BMI with gold standard methods such as DEXA, the authors found that BMI had a sensitivity of 73 % to identify excess body fat in children and adolescents, concluding that more than a quarter of undiagnosed subjects with obesity due to BMI may have excess adiposity<sup>(42)</sup>. Another study showed that eutrophic adolescents with BMI and high BF% have high cardiometabolic risk factors similar to those with overweight by BMI and high BF%<sup>(43)</sup>. Therefore, other anthropometric indices have been used to assess the adult population, but do not have well-defined cut-off points for adolescents.

WtHR demonstrates good predictive capacity for body fat and is a good index to be used in the adolescents' assessment, as in the calculation height and waist are considered<sup>(44,45)</sup>. Pelegrini *et al.*<sup>(46)</sup> assessed 1197 Brazilian adolescents and found that BMI and WtHR had a greater capacity to estimate body fat in both sexes, unlike CI, which as in this study, had the lowest values of AUC.

WHR did not perform well in the estimation of body fat, as in other study that showed that BMI and WC correlated with BF% in Swedish adolescents of both sexes, while WHR did not correlate well<sup>(47)</sup>. Similar to Motlagh *et al.*<sup>(38)</sup> who assessed 2444 adolescents aged from 12 to 14 years and observed that WC and WtHR show higher AUC than WHR in predicting overweight and obesity.

In adolescence, WHR is not a good parameter for assessing body composition because it uses hip measurement and it changes quickly during the growth spurt<sup>(44)</sup>. Similar to previous results, in male children and adolescents from Spain, BMI and

WC had good capacity to estimate body fat, showing values of AUC above 0.80<sup>(48)</sup>.

As observed in this study, Weiss *et al.*<sup>(49)</sup> assessed Brazilian adolescents from 11 to 14 years old and found that the CI did not perform well to diagnose high body fat in female adolescents, with a value of AUC of 0.52. In boys, the value of the AUC was also low, 0.69, showing that CI was not a good predictor of high body fat.

Corroborating the results presented in this study, Frignani *et al.*<sup>(50)</sup> found that the BAI showed association and high agreement with the BF% in adolescents, in both sexes. Studies assessing the performance of BAI and BRI to estimate body fat are scarce in the literature, especially conducted specifically with adolescents. In this study, both showed good capacity to predict body fat in this population. According to Thomas *et al.*<sup>(23)</sup>, BRI has better performance to estimate the percentage of total and visceral body fat compared with BMI and WC.

Authors state that BRI can be considered a better predictor of cardiovascular risk when compared with BMI, since it estimates the location of excess body fat, better reflecting the estimation of visceral fat, since its calculation considers the WC<sup>(23,51)</sup>.

The analysis of the ROC curves allowed to determine cut-off points for the assessed indices, to estimate the excess of body fat, considering this population in general and according to sex and period of adolescence. The AUC for BMI, WtHR, BAI and BRI were considered very good or excellent in adolescents in general, but most of these values were lower compared with those analysed by sex and period of adolescence. Assessing the accuracy of anthropometric indices to estimate body fat is necessary, especially in this age group and due to changes in body composition that occur during adolescence<sup>(2)</sup>. We suggest the use of specific cut-off points for the various adiposity indices, according to the adolescent's sex and age, in order to obtain a more accurate and reliable diagnosis of excess body fat.

The ideal cut-off values for predicting excess body fat for the assessed indices varied according to adolescent's sexes and periods. Except for BMI, up to the present moment, no Brazilian study has proposed cut-off points for the other indices to be used in the adolescent population, considering sex and period of adolescence. Moreover, the known cut-off points for BMI are not specific for assessing excess body fat, as proposed and presented in this study for all indices.

The ideal BMI cut-off points increased with age in both sexes, as did BRI in males. In the middle period, the values for WtHR, BAI and BRI were lower in girls, as well as the WtHR and BAI in boys.

It is important to highlight that the ideal BMI cut-off values for the diagnosis of excess body fat in this population, in both sexes and all periods of adolescence, were below the cut-off point for overweight diagnosis according to WHO<sup>(28)</sup> standards. Similarly, ideal values of WtHR cut-offs in girls and boys of all ages in this study were below 0.50, a value proposed to assess abdominal obesity regardless of age and sex<sup>(45)</sup>. This suggests that the biological risk associated with increased body fat may be underestimated in adolescents.

The range of the sample can be considered a limitation of the study, since adolescents from only one city were assessed. Also, studies have shown that there may be distinctions in body

composition between different ethnicities<sup>(52,53)</sup>, although there is still no consensus on the subject<sup>(54)</sup>. However, the study has a significant sample number and comprised the whole age group of adolescents of different races and socio-economic status. In addition, the study used a gold standard method for assessing body composition (DEXA). Thus, we noticed that the results presented could be used to assess the accumulation of body fat in adolescents, as well as to prevent risk factors for health and development of this population.

In conclusion, besides the BMI, the adiposity index, WtHR, BAI and BRI have shown to perform well to identify excess body fat in these adolescents. Considering the characteristics of each index evaluated, we identified that the best index to predict excess body fat in female adolescents aged from 10 to 13 years was the WtHR, whereas the BAI was the best to those aged between 14 and 19 years. In male adolescents, aged from 10 to 13 years and 14 to 16 years, the best index was the WtHR and the BAI showed better predictive capacity in boys from 17 to 19 years old.

The results of this study reinforce the importance of using indices with specific cut-off points for each period of adolescence and according to sex for the reliable diagnosis of excess body fat. It is advisable to use indices together to obtain a more accurate assessment. Thus, the WtHR and BAI are a low-cost and non-invasive method that is reproducible and reliable, with high sensitivity and specificity values and can be used together with the BMI. This allows an assessment and monitoring of the nutritional and health status of adolescents, in order to prevent the development of diseases at this age and into adulthood. However, further studies are necessary to confirm the reproducibility of the results.

### Acknowledgements

The authors would like to thank the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

N. S. M. worked on the design of the article, analysis and interpretation of the results, on the writing of the manuscript and approved the final version to be published, S. A. V. R. worked on the design, analysis and interpretation of the results, on the revision of the writing of the manuscript and approved the final version to be published, P. F. P. worked on the design, analysis and interpretation of the results, on the revision of the writing of the manuscript and approved the final version to be published, V. P. N. M. worked on the analysis and interpretation of the results, on the revision of the writing of the manuscript and approved the final version to be published, E. C. G. S. worked on the analysis and interpretation of the results, on the revision of the writing of the manuscript and approved the final version to be published, S. C. C. F. worked on the design, analysis and interpretation of the results, on the revision of the writing of the manuscript and approved the final version to be published



and S. E. P. participated in the conception and design of the article, interpretation of results, revision of the writing of the manuscript and approved the final version to be published.

The authors declare that there are no conflicts of interest.

## References

1. Reis CEG, Vasconcelos IAL & Barros JFN (2011) Políticas públicas de nutrição para o controle da obesidade infantil. *Rev Paul Pediatr* **29**, 625–633.
2. Sumi A, Iwase M, Nakamura U, *et al.* (2018) Impact of age at menarche on obesity and glycemic control in Japanese patients with type 2 diabetes: Fukuoka Diabetes Registry. *J Diabetes Investig* **9**, 1216–1223.
3. Instituto Brasileiro de Geografia e Estatística - IBGE (2010) Pesquisa de Orçamentos Familiares 2008–2009. Antropometria e estado nutricional de crianças, adolescentes e adultos no Brasil. <https://biblioteca.ibge.gov.br/visualizacao/livros/liv45419.pdf> (accessed 15 February 2020).
4. Instituto Brasileiro de Geografia e Estatística - IBGE (2016) *Pesquisa de Nacional de Saúde do Escolar 2015. IBGE, Coordenação de População e Indicadores Sociais*. Rio de Janeiro: IBGE.
5. NCD Risk Factor Collaboration-NCD-RISC (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* **390**, 2627–2642.
6. World Health Organization (2005) *Nutrition in Adolescence: Issues and Challenges for the Health Sector: Issues in Adolescent Health and Development*. Geneva: WHO.
7. Priore SE, Faria FR & Franceschini SCC (2010) Adolescência. In: *PRIORE, Nutrição e Saúde na Adolescência*, Editora Rúbio, Ed. 1, Rio de Janeiro, P. 1–7.
8. Juonala M, Magnussen CG, Berenson GS, *et al.* (2011) Childhood adiposity, adult adiposity, and cardiovascular risk factors. *N Engl J Med* **365**, 1876–1885.
9. Wang H, Steffen LM, Vessby B, *et al.* (2011) Obesity modifies the relations between serum markers of dairy fats and inflammation and oxidative stress among adolescents. *Obesity* **19**, 2404–2410.
10. Bozza R, Campos W, Barbosa Filho VC, *et al.* (2016) Pressão Arterial Alterada em Adolescentes de Curitiba: Prevalência e Fatores Associados. *Arq Bras Cardiol* **106**, 411–418.
11. Lascar N, Brown J, Pattison H, *et al.* (2017) Type 2 diabetes in adolescents and young adults. *Lancet Diabetes Endocrinol* **6**, 69–80.
12. Buscot MJ, Thomson RJ, Juonala M, *et al.* (2018) Distinct child-to-adult body mass index trajectories are associated with different levels of adult cardiometabolic risk. *Eur Heart J* **39**, 2263–2270.
13. Giudici KV, Rolland-Cachera M-F, Gusto G, *et al.* (2017) Body mass index growth trajectories associated with the different parameters of the metabolic syndrome at adulthood. *Int J Obes* **41**, 1518–1525.
14. Miranda VPN, Amorim PRS, Bastos RR, *et al.* (2019) Body image disorders associated with lifestyle and body composition of female adolescents. *Public Health Nutr* 1–11.
15. Filgueiras MDS, Vieira SAV, Fonseca PCA, *et al.* (2018) Waist circumference, waist-to-height ratio and conicity index to evaluate android fat excess in Brazilian children. *Public Health Nutr* **22**, 140–146.
16. Vanderwall C, Randall Clark R, Eickhoff J, *et al.* (2017) BMI is a poor predictor of adiposity in young overweight and obese children. *BMC Pediatr* **17**, 135.
17. Filgueiras MDS, Cecon RS, Faria ER, *et al.* (2018) Agreement of body adiposity index (BAI) and paediatric body adiposity index (BAIp) in determining body fat in Brazilian children and adolescents. *Public Health Nutr* **22**, 132–139.
18. Ripka WL, Orsso CE, Haqq AM, *et al.* (2020) Validity, accuracy of body fat prediction equations using anthropometrics measurements in adolescents. *Eating Weight Disord-Stud Anorexia, Bulimia Obes* 1–8.
19. Woolcott OO & Bergman RN (2019) Relative Fat Mass as an estimator of whole-body fat percentage among children and adolescents: a cross-sectional study using NHANES. *Sci Rep* **9**, 15279.
20. Schröder H, Ribas L, Koebnick C, *et al.* (2014) Prevalence of abdominal obesity in Spanish children and adolescents. Do we need waist circumference measurements in pediatric practice? *PLoS One* **9**, e87549.
21. Valdez R (1991) A simple model-based index of abdominal adiposity. *J Clin Epidemiol* **44**, 955–956.
22. Bergman RN, Stefanovski D, Buchanan TA, *et al.* (2011) A better index of body adiposity. *Obesity* **19**, 1083–1089.
23. Thomas DM, Bredlau C, Bosy-Westphal A, *et al.* (2013) Relationships between body roundness with body fat and visceral adipose tissue emerging from a new geometrical model. *Obesity* **21**, 2264–2271.
24. World Health Organization (1995) *WHO Expert Committee on Physical Status. Physical status: the use and gnterpretation of anthropometry. WHO Technical Report Series no. 854*. Geneva: WHO.
25. World Health Organization (2008) *Waist Circumferences and Waist-Hip Ratio: Report of a WHO Expert Consultation*. Geneva: World Health Organization.
26. Heyward VH & Stolarczyk LM (2000) *Métodos de dobras cutâneas. Avaliação da composição corporal aplicada*. São Paulo: Manole.
27. Lohman TG (1992) *Assesing Fat Distribution. Advances in Body Composition Assessment: Current Issues in Exercise Science*. Champaign, IL: Human Kinetics.
28. De Onis M, Onyango AW, Borghi E, *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660–667.
29. Borges LSR (2016) Medidas de Acurácia Diagnóstica na Pesquisa Cardiovascular. *Int J Cardiovasc Sci* **29**, 218–222.
30. Gontijo CA, Faria ER, Oliveira RMS, *et al.* (2010) Síndrome Metabólica em Adolescentes Atendidos em Programa de Saúde de Viçosa – MG. *Rev Bras Cardiol* **23**, 324–333.
31. Costa MCD, Barreto ADC, Bleil RAT, *et al.* (2011) Estado nutricional de adolescentes atendidos em uma unidade de referência para adolescentes no Município de Cascavel, Estado do Paraná, Brasil. *Epidemiol Serv Saúde* **20**, 355–361.
32. Cumpian-Silva J, Rinaldi AEM, Mazzeti CMS, *et al.* (2018) Fenótipos corporais na adolescência e a maturação sexual. *Cad Saúde Pública* **34**, e00057217.
33. Guinhouya BC, Samouda H & Beaufort C (2013) Level of physical activity among children and adolescents in Europe: a review of physical activity assessed objectively by accelerometry. *Public Health* **127**, 301–311.
34. Longo-Silva G, Toloni MHA, Menezes RCE, *et al.* (2015) Introdução de refrigerantes e sucos industrializados na dieta de lactentes que frequentam creches públicas. *Rev Paul Pediatr* **33**, 34–41.
35. Miranda VPN, Amorim PRS, Bastos RR, *et al.* (2019) Evaluation of lifestyle of female adolescents through latent class analysis approach. *BMC Public Health* **19**, 184.
36. Kobayashi F, Oliveira FLC, Escrivão MAMS, *et al.* (2010) Obesidade e Fatores de Risco Cardiovascular em Adolescentes de escolas públicas. *Arq Bras Cardiol* 200–206.







37. Moraes MM & Veiga G (2014) Acurácia da gordura corporal e do perímetro da cintura para prever alterações metabólicas de risco cardiovascular em adolescentes. *2*, 341–351.
38. Motlagh M, Seiyed S, Zahra HR, *et al.* (2018) Assessment of overweight and obesity in Iranian adolescents: optimal cut-off values of anthropometric indices. *EMHJ* **24**, 975–987.
39. Ribeiro VB, Kogure GS, Lopes IP, *et al.* (2019) Association of measures of central fat accumulation indices with body fat distribution and metabolic, hormonal, and inflammatory parameters in women with polycystic ovary syndrome. *Arch Endocrinol Metab* **63**, 417–426.
40. Telford R, Cunningham RB & Abhayaratna WP (2013) Temporal divergence of percent body fat and body mass index in pre-teenage children: the LOOK longitudinal study. *Pediatr Obes* **9**, 448–454.
41. Swainson MG, Batterham AM, Tsakirides C, *et al.* (2017) Prediction of whole-body fat percentage and visceral adipose tissue mass from five anthropometric variables. *PLoS One* **12**, e0177175.
42. Javed A, Jumean M, Murad MH, *et al.* (2015) Diagnostic performance of body mass index to identify obesity as defined by body adiposity in children and adolescents: a systematic review and meta-analysis. *Pediatr Obes* **10**, 234–244.
43. Serrano HMS, Carvalho GQ, Pereira PF, *et al.* (2010) Composição Corpórea, Alterações Bioquímicas e Clínicas de Adolescentes com Excesso de Adiposidade. *Arq Bras Cardiol* **95**, 464–472.
44. Alvarez MM, Vieira ACR, Sichieri R, *et al.* (2008) Associação das medidas antropométricas de localização de gordura central com os componentes da síndrome metabólica em uma amostra probabilística de adolescentes de escolas públicas. *Endocrinol Metab* **52**, 649–657.
45. Ashwell M & Gibson S (2014) A proposal for a primary screening tool: “Keep your waist circumference to less than half your height”. *BMC Med* **12**, 207.
46. Pelegrini A, Silva DAS, Silva JMFL, *et al.* (2015) Indicadores antropométricos de obesidade na predição de gordura corporal elevada em adolescentes. *Rev Paul Pediatr* **33**, 56–62.
47. Neovius M, Linne Y & Rossner S (2005) BMI, waist-circumference and waist-hip-ratio as diagnostic tests for fatness in adolescents. *Int J Obes* **29**, 163–169.
48. Sarriá A, Moreno LA, García-Llop LA, *et al.* (2001) Body mass index, triceps skinfold and waist circumference in screening for adiposity in male children and adolescents. *Acta Paediatr* **90**, 387–392.
49. Weiss KM, Leal DB, Assis MAA, *et al.* (2015) Diagnostic accuracy of anthropometric indicators to predict excess body fat in adolescents aged 11–14 years. *Rev Bras Cineantropom Desempenhobum* **18**, 114–126.
50. Frignani RR, Passos MAS, Ferrari GLM, *et al.* (2015) Reference curves of the body fat index in adolescents and their association with anthropometric variables. *J Pediatr* **91**, 248–255.
51. Maessen MFH, Eijsvogels TM, Verheggen RJ, *et al.* (2014) Entering a new era of body indices: the feasibility of a body shape index and body roundness index to identify cardiovascular health status. *PLoS One* **9**, 1–8.
52. Wang D, Li Y, Lee SG, *et al.* Ethnic differences in body composition and obesity related risk factors: study in Chinese and white males living in China. *PLoS One* **6**, e19835.
53. Heymsfield SB, Peterson CM, Thomas DM, *et al.* (2016) Why are there race/ethnic differences in adult body mass index-adiposity relationships? A quantitative critical review. *Obes Rev* **17**, 262–275.
54. Alenaini W, Parkinson JRC, McCarthy JP, *et al.* (2020) Ethnic differences in body fat deposition and liver fat content in two UK-based cohorts. *Obesity* 1–11.