

Predictors of *Toxoplasma gondii* infection in Czech and Slovak populations: the possible role of cat-related injuries and risky sexual behavior in the parasite transmission

J. FLEGR^{1,2*}

¹ Division of Biology, Faculty of Science, Charles University in Prague, Prague, Klecany 250 67, Czech Republic

² National Institute of Mental Health, Klecany 250 67, Czech Republic

Received 26 August 2016; Final revision 27 December 2016; Accepted 10 January 2017;
first published online 10 February 2017

SUMMARY

The protozoan *Toxoplasma gondii* infects about one-third of the world's population. The consumption of raw meat, contact with cats, contact with soil, and ingestion of food or water contaminated with soil are considered to be the most important sources of infection. Still in most women who were infected during pregnancy, no definitive source of infection is found. In 2014–2016, independent sources of *T. gondii* infection were searched for by gathering epidemiological data from 1865 (519 infected) responders. Touching garden soil (odds ratio (OR) 3·14, 95% confidence interval (CI) 1·3–6·35), sustaining cat-related injuries (OR 2·16, 95% CI 1·25–3·74), and eating improperly washed root vegetables (OR 1·71, 95% CI 1·02–2·87), but not risky sexual behavior (OR 1·22, 95% CI 0·79–1·90), were the predictors of infection. The seroprevalence of *T. gondii* infection had been increasing up to ages 35–50 in men and ages 50–54 in women. Past those ages, seroprevalence of toxoplasmosis has been decreasing. This suggests that the natural decrease of anamnestic antibodies concentrations over time leads to positivity-to-negativity seroconversion in many subjects. If this is true, then the prevalence of *T. gondii* infection in a general population and its potential impacts on public health could be much larger than generally believed.

Key words: Cats, dogs, pets, risk factors, sexually transmitted diseases, STD, Toxoplasmosis, zoonosis.

INTRODUCTION

Toxoplasma gondii is probably the most common parasitic protozoon in humans in the developed world. Its prevalence is estimated to be more than 30% and varies between 5% and 100% in different countries depending on environmental conditions, hygienic standards and kitchen habits [1, 2]. *T. gondii*

is transmitted from an intermediate host, which can be any homeothermic animal, to the definitive host, which can be any species of small or large cats, by predation [1] and from definitive hosts to new intermediate hosts by ingestion of oocysts, the product of sexual reproduction of *T. gondii* in the cat's intestine, which are excreted by the millions in the feces of infected cats [3].

Despite the life cycle of *T. gondii* being known for a very long time, some aspects of its transmission to humans are still mysterious. For example, in more than two thirds of women that acquired infection during pregnancy, no contact with any known or

* Author for correspondence: J. Flegr, Division of Biology, Faculty of Science, Charles University in Prague, Prague, Viničná 7, 128 44, Czech Republic and National Institute of Mental Health, Klecany 250 67, Czech Republic.
(Email: flegr@cesnet.cz)

suspected source of infection was identified [4, 5]. Many studies showed that contact with cats, touching garden soil or eating raw meat were important epidemiological risk factors [6–8], while other studies did not demonstrate their role in transmitting the infection [9, 10]. A few studies detected some unusual predictors of being *T. gondii* infected, such as keeping rabbits for meat [9, 10], having a dog at home [11], having mice in a house [12], eating raw shellfish [13], being in a close contact with livestock [14], and having high rate of sexual activity [15, 16].

The aim of the present study was to search for independent predictors of *T. gondii* infection in a Czech and Slovak internet population of 1865 subjects by using a detailed electronic questionnaire for acquiring epidemiological data and a complex model for identifying the most probable risk factors for *T. gondii* infection. The specific aim of the study was to search for the possible roles of contact with cats and dogs in the transmission of *T. gondii* to human hosts.

METHODS

Participants and procedure

The participants were invited to participate in the study using a Facebook-based snowball method [17]. It was done by posting an invitation to participate in ‘an experiment searching for associations between keeping dogs and cats and health status and personality of a subject’ on the wall of the Facebook page ‘Guinea Pigs’ for people willing to take part in diverse evolutionary psychological experiments (www.facebook.com/pokusnikralici). This Facebook community consists of people of various ages, education levels, occupations, and locations. To keep the study blind and avoid a possible bias, no form of the term toxoplasmosis or *T. gondii* was mentioned during recruitment, not even in the informed consent form. This omission was approved by the Institutional Review Board (IRB). The participants were informed about the general aims of the study on the first page of the questionnaire. They were also provided with the following information: ‘The questionnaire is anonymous and obtained data will be used exclusively for scientific purposes. Your cooperation in the project is voluntary and you can terminate it at any time by closing this web page. Please share the link to this questionnaire with your friends, for example on Facebook’. The share button had been pressed by 967 participants, which resulted in data from 9125 responders

in total from 20 August 2014 to 5 November 2016. The entire study and method of acquiring electronic informed content were approved by the IRB of the Faculty of Science, Charles University in Prague (No. 2015/07).

Immunological tests for *T. gondii* infection

Most of the women and nearly all of men who know their *T. gondii*-infection status were tested for *T. gondii* infection during systematic research of behavioral effects of latent *T. gondii* infection which has been running at the Faculty of Science for 20 years. During this period of time, we tested more than 3000 students of biology. Most of them have been tested within past 5 years (approximately 300 per year). Within the past 3 years we have also tested about 800 other members of the Guinea Pigs community; we suppose that these subjects are overrepresented among participants of the current study. All testing was performed at the National Reference Laboratory for Toxoplasmosis, National Institute of Public Health, Prague. The complement-fixation test (CFT), which determines the overall levels of IgM and IgG antibodies of particular specificity, and Enzyme-Linked Immunosorbent Assays (ELISA) (IgG ELISA: SEVAC, Prague) were used to detect *T. gondii* infection status of the subjects. ELISA assay cut-point values were established using positive and negative standards according to the manufacturer’s instructions. In CFT, the titer of antibodies against *T. gondii* in sera was measured in dilutions between 1:8 and 1:1024. The subjects with CFT titers between 1:8 and 1:128 were considered *T. gondii* infected. Only subjects with clearly positive results of CFT or IgG ELISA tests were diagnosed as *T. gondii*-infected or *T. gondii*-free.

Electronic questionnaire

The questionnaire was distributed as a Qualtrics survey. The responders used the five-points Likert scale (1, never; 2, minimally (1–2 times in life); 3, rarely; 4, from time to time; 5, very often), to rate their past contacts with six known or suspected risk factors (eating or tasting raw meat, touching soil during gardening, eating root vegetables that were not washed properly, living in countries of lower hygienic standard, drinking water from suspicious sources such as creeks, being at risk of acquiring sexually transmitted diseases (by having sex without a condom with

various people). The responders were asked about the size of the settlement where they spent their childhood (1, less than 1000 inhabitants; 2, 1–5 thousands inhabitants; 3, 5–50 thousands inhabitants; 4, 50–100 thousands inhabitants; 5, 100–500 thousands inhabitants; 6, more than 500 thousands inhabitants). The anamnestic questionnaire also contained questions about the intensity and nature of contact with dogs and cats and about animal-related injuries. Participants were asked to rate the intensity of their life-long contact with dogs and cats using the seven-points scale: (1, never; 2, we kept a dog (cat) only in the past and only for a short time; 3, we kept a dog (cat) only in the past but for a long time; 4, we have one dog (cat); 5, we have two dogs (cats); 6, we have three dogs (cats); 7, we have more than three dogs (cats). Using seven-points scale the responders rated how often they stroke a dog (cat) (1, never; 2, maximally once per year; 3, maximally once per month; 4, maximally once per week; 5, maximally once per day; 6, maximally five times a day; 7, more often), whether he/she consider pleasant the smell of a clean dog (cat) fur (1, definitively no; 2, rather no; 3, neither pleasant, nor unpleasant; 4, rather yes; 5, definitively yes; 6, I love the smell of dog (cat) fur). By moving sliders on graphic scales 0–100, the responders expressed their agreement with following four statements: I like dogs (cats), I am not afraid of dogs (cats), I like to touch a dog (cat), A dog (cat) is a member of a family. In a different part of the questionnaire the responders were also asked to rate the intensity of sustained animal-related injuries using the following scale: 1, never; 2, just in play; 3, yes but just to warn me; 4, yes but not seriously; 5, yes with mediate seriousness; 6, yes seriously; 7, yes very seriously, I had to seek for medical help; 8, I was seriously bitten (scratched) by several dogs (cats). Responders were asked to check the variant describing the most serious injury they suffered. Before the analyses, the responses concerning animal-related injuries were recoded to five-points scale 1 (1 or 2), 2 (3), 3 (4), 5 (6 or 7) to get more regular distributions; however, the results of analyses that used the unreduced scale were approximately the same. The participants were also asked whether they are *T. gondii* infected. They were reminded that *T. gondii* is ‘a parasite of cats, dangerous especially to pregnant women’. Implicitly, the response ‘I do not know, I am not sure’ was selected. Responders could change this by selecting either ‘No, I was tested by a doctor and the result of my laboratory tests was negative’ or ‘Yes, I was tested by a doctor and I had

antibodies against *Toxoplasma*’. The responders of our questionnaires always had three options: they could complete any questionnaire absolutely anonymously, they could sign the finished questionnaire by a code obtained after anonymous registration, or they could sign the finished questionnaire by a code obtained after non-anonymous registration (see <http://pokusnikralici.cz>). Some questionnaires were ‘signed’ by <10% of non-anonymously registered subjects (e.g. the questionnaire about sexual behavior); however, the present questionnaire was ‘signed’ by 31% of the participants. When we checked the information about the *T. gondii* infection status provided in the questionnaire by 3827 participants of our past experiments with corresponding information in our records, we found nearly perfect (99.5%) agreement. The questionnaire also contained many questions concerning health status of responders, a standard Czech version of the Ten Item Personality Inventory (TIPI) [18], Beck Depression Inventory [19], Obsessive-Compulsive Inventory – Revised (OCI-R) [20], and Toxo1 questionnaire [21]; however, only the responses concerning epidemiological risk factors including the contacts with dogs and cats were used in the present epidemiological study. The questionnaire took participants 20–40 min to complete and about 72% of participants who started the study finished the whole questionnaire.

Statistics

Univariate tests and logistic regressions were performed by the statistical software Statistica v. 10.0, while parallel analysis and factor analysis were performed using R software. Significance was at a 5% level (two-sided). Before statistical analysis, I filtered out <1% of data, because it appeared suspicious (too high or too short body height, too low or too high body mass or age, etc.) The associations between ordinal predictors and *T. gondii* infection were studied by Spearman correlation. This non-parametric test is not sensitive to the presence of outliers in the data set. Spearman tests and Logistic regression tests (Quasi-Newton method, maximum-likelihood loss function and casewise deletion of missing data) were performed for all participants together and also separately for men and women. The optimal number of factors to be extracted (seven) was computed by parallel analysis before the factor analysis (principal axis factoring, direct oblimin normalized rotation, mean substitution of missing data) was performed. The false

discovery rate (preset to 0.1) was controlled with the Benjamini–Hochberg procedure [22]. In contrast to the simple Bonferroni's correction, this procedure also takes into account the distribution of *P* values of performed multiple tests. Therefore, the number of significant results after the correction could be higher than before the correction.

RESULTS

Descriptive statistics

Between 20 August 2014 and 5 November 2016, 3106 men (age: 35.8, standard deviation (s.d.) 12.91) and 6019 women (age 33.2, s.d. 12.73) completed the questionnaire. Four hundred and twenty-two (422) men (age: 34.4, s.d. 13.5) and 1443 women (age 33.1, s.d. 11.77), $t_{6179} = 1.82$, $P = 0.07$ were aware of their *T. gondii* infection status. *T. gondii*-infected participants were significantly older than *T. gondii*-free participants (36.4 vs. 32.2, $t_{1862} = -6.72$, $P < 0.00001$). Prevalence of *T. gondii* infection was higher in 1443 women (30.1%, $N = 434$), than in 422 men (20.1%, $N = 85$), $\chi^2 = 16.0$, $P < 0.0001$. For the age structure of the population, see Figure 1 and for frequency and intensity of contacts with potential risk factors see Table 1.

Identification of risk factors for acquiring *T. gondii* infection: univariate analyses

Table 2 shows the associations between toxoplasmosis and 24 ordinal variables describing the intensity of contact with particular risk factors. To decrease an impact of potential outliers, the associations with *T. gondii* infection were estimated with Spearman's correlation tests, however, the results obtained with parametric tests (logistic regression) were approximately the same (results not shown).

Reduction of dimensionality

A complicated network of relations between animal-related variables and between animal-related variables vs. *T. gondii* infection probably exists. To reduce the dimensionality of this data for the subsequent analyses, a factor analysis with 17 variables that characterize relations of particular participants with dogs and cats was performed. Seventeen variables were reduced to seven independent factors, which explained 62% of between-individual variability. After oblimin rotation, it became possible to identify the nature of

particular factors on the basis of factor loadings; see Table 3.

Association between *T. gondii* infection and various risk factors: multivariate analyses

The association between *T. gondii* infection and all analyzed risk factors – sex, age and all seven factors obtained with the factor analysis – was studied by logistic regression containing all independent variables in one model. The number of participants older than 50 was relatively low and prevalence of toxoplasmosis decreased after the age of 50 in men and of 54 in women (Fig. 1) suggesting that serological tests provide false negative results in subjects who were infected a rather long time ago. Therefore, only the participants younger than 50 were included into the final analyses; however, the analyses of the whole set gave very similar results. The results showed that independent risk factors for having anamnestic antibody titers to *T. gondii* were spending one's childhood in small communities, past direct contacts with garden soil, having consumed unwashed root vegetables, and having suffered a cat-related injury (Table 4). The results were similar when men and women were analyzed separately or when sex–age interaction was included into the model. Due to the lower number of men in our sample (422 vs. 1443), all effects except for the effect of age were non-significant in men. However, the strength of this association estimated on the basis of odds ratio (OR) was sometimes higher in men than in women (cat-related injury: OR = 3.88 vs. 1.93; eating unwashed vegetable OR = 3.64 vs. OR = 1.46). The seroprevalence of *T. gondii* infection increased with age in women up to age 50–54 and decreased after age of 54. In men, this seroprevalence increased only up to age of 39, stayed stable until age of 49, and decreased after age of 49; see Figure 1.

DISCUSSION

Analysis performed on the large sample of a population showed that having contact with garden soil, eating improperly washed root vegetables, and being injured by a cat were the most important predictors of the *T. gondii* infection in our population. The probability of being infected with *T. gondii* was higher in women than in men and in participants who spent their childhood in small communities. This probability increased with the age of the subject.




Fig. 1. Prevalence of *T. gondii* infection in men and women of different age. Numbers at the top of the columns show the prevalences (in %, vertical orientation) and a number of female/male participants.

Surprisingly, the prevalence, which was increasing up to the age 39 in men and up to age 54 in women, generally began showing a decrease when those ages were reached.

The important role of contact with garden soil and consuming unwashed root vegetables is in agreement with our knowledge of the life cycle of *T. gondii* and the ethology of its definitive host – a cat [23]. After primo-infection a cat may excrete millions of *T. gondii* oocysts. Domestic cats living around and in human communities defecate near houses, often in gardens. The oocysts survive and remain infectious in humid soil for years; therefore, the soil in garden patches near houses is often heavily contaminated with oocysts [24, 25].

The relationship between *T. gondii* infection and various aspects of pet dogs and cats was in the center of the interest through the present study. It was suggested earlier that a cat as well as a dog can be the proximal source of infection [11], that pets could be just an indication of a more rustic style of life [26] and that the animal-related injuries, especially cat bites, can be a source of *T. gondii* infection [27, 28]. It has been also published that, in a double blind experiment, *T. gondii*-infected male students consider a smell of highly diluted cat urine more pleasant than *T. gondii*-free control subjects [29]. Therefore, it can be speculated that infected persons are more enamored with cats and own cats more often than *T. gondii*-free persons. The results of the complex analysis showed that neither the number of cats nor dogs

in the house nor love of cats or dogs had a significant impact on the risk of being *T. gondii* infected; however, cat-related injuries were found to be predictive of such an infection.

It can be hypothesized that *T. gondii* parasites could be directly transmitted from the body of a cat to the body of a human by biting or scratching (nearly all participants bitten by a cat were also scratched by a cat). Another hypothetical way of acquiring infection is the contamination of scars with oocysts contained in soil on the cat's claws, paws and other parts of the pets' bodies; however, no non-oral route of infection by *T. gondii* oocysts has been described. If this latter possibility holds, then even non-infected cats could transmit *T. gondii* to a human host if they defecate or just rest at garden where other cats have defecated. Pregnant women are commonly urged not to change cat litter boxes. The present results suggest that a better recommendation is for pregnant women to entirely limit interaction with cats.

The present results confirmed that spending one's childhood in small communities increases the risk of *T. gondii* infection even when other associated variables like contact with cats and garden soil as well as eating unwashed root vegetables are controlled for. This means that other unknown risk factors associated with living in small communities probably also exist. A study performed on a population of 3250 soldiers showed that the strongest predictor of *T. gondii* seropositivity was keeping rabbits for meat (OR 1.45, 95% CI 1.22–1.77) [30]. In future studies, this

Table 2. Association between *T. gondii* infection and potential risk factors

	All <i>R</i>	All <i>P</i>	Men <i>R</i>	Men <i>P</i>	Women <i>R</i>	Women <i>P</i>
No. of dogs in house	0·01	0·77	0·02	0·77	0·00	1·00
Stroking dogs (frequency)	-0·01	0·72	0·04	0·38	-0·03	0·25
Bitten by a dog (intensity)	0·08	0·00	0·05	0·28	0·08	0·00
Dog's smell (pleasantness)	-0·05	0·07	-0·05	0·45	-0·06	0·06
Loving dogs	0·00	0·84	0·01	0·88	-0·02	0·38
No fear of dogs	-0·04	0·06	-0·07	0·16	-0·04	0·14
Touching dogs (pleasantness)	-0·02	0·29	-0·04	0·39	-0·04	0·16
Dog as a family member (agreement)	0·00	0·92	0·02	0·70	-0·03	0·30
Number of cats in house	0·08	0·00	0·18	0·00	0·05	0·08
Stroking cats (frequency)	0·04	0·11	0·13	0·01	0·00	0·89
Bitten by a cat (intensity)	0·06	0·01	0·17	0·00	0·04	0·15
Scratched by a cat (intensity)	0·07	0·00	0·13	0·01	0·06	0·02
Cat's smell (pleasantness)	-0·01	0·83	-0·04	0·47	-0·02	0·49
Loving cats	0·05	0·03	0·12	0·02	0·02	0·48
No fear of cats	-0·02	0·38	-0·07	0·15	-0·01	0·83
Touching cats (pleasantness)	0·02	0·31	0·06	0·26	0·00	0·89
Cat as a family member (agreement)	0·03	0·25	0·12	0·03	-0·02	0·53
Size of place of living in childhood	-0·10	0·00	-0·04	0·40	0·12	0·00
Eating raw meat	0·02	0·40	-0·05	0·37	0·04	0·15
Touching soil	0·16	0·00	0·10	0·06	0·17	0·00
Unwashed vegetables	0·11	0·00	0·15	0·01	0·10	0·00
Low hygienic standard country	0·01	0·67	0·02	0·68	0·02	0·43
Suspicious water	-0·01	0·70	0·00	0·98	0·00	1·00
Risky sexual behavior	0·07	0·01	0·00	0·96	0·08	0·00

Table shows results of Spearman's correlations. Positive (negative) Spearman *R* denotes positive (negative) association between particular variables. Significant associations (in two-side tests) are printed in bold. $P < 0·005$ is coded as 0·00. All significant results remained significant after the Benjamini-Hochberg procedure correction for multiple tests (with tolerated false discovery rate = 0·10).

factor should be monitored and included into statistical models.

On the basis of several independent lines of indirect evidence, the existence of a sexual route of transmission of *T. gondii* infection has been suggested [15, 16]. It has been shown that the prevalence of *T. gondii* infection in particular countries correlates with the incidence of sexually transmitted diseases [31]. Seemingly, the present results, namely the positive association between *T. gondii* infection and risky sexual behavior observed in univariate analyses, provided further support for this hypothesis. However, the results of logistic analysis of the complex model did not support the existence of this association. Because of high clinical importance of this potential route of infection (risk of congenital toxoplasmosis) the problem of the sexual transmission of *T. gondii* infection deserves more attention in future studies.

Neither eating raw meat nor drinking potentially contaminated water was significantly associated with being *T. gondii* infected. This contrasted with some earlier results and with our knowledge of the life

cycle of *T. gondii*. The most frequent source of raw meat in Czech cuisine is steak Tartar, the minced raw beef with egg and spice. However, the prevalence of *T. gondii* infection in cattle from large farms is low, the number and viability of tissue cysts in beef is very low and steak Tartar is prepared from frozen meat [23]. It is highly probable that all *T. gondii* bradyzoites are killed in frozen and possibly even cooled meat. The growing frequency of consumption of frozen and cooled meat can be, in fact, a reason for the systematic decrease of the prevalence of *T. gondii* infection in the developed world, including Czechia, during past 20 years [2].

A non-significant negative association between *T. gondii* infection and drinking suspicious water as well as visiting low hygienic standard countries (nearly significant in women) can be the side-effect of a changed personality profile of infected subjects, which is usually considered to be a product of manipulative activity of the parasite. *T. gondii* is known to increase the concentration of neurotransmitter dopamine in the brain tissue of infected animals [32, 33] (but see also

Table 3. *Nine independent variables describing relation of participants to dogs and cats*

	Loving dogs	Loving cats	No. of cats at house	No. of dogs at house	Cat-related injury	Pet as a family member	Likes animals' smell
No. of dogs in house	-0.06	0.01	0.03	0.97	-0.01	0.00	-0.03
Stroking dogs	0.17	0.02	-0.04	0.67	0.02	0.02	0.07
Bitten by a dog	-0.07	-0.01	-0.02	0.14	0.16	-0.01	0.03
Dog's smell	0.29	-0.14	-0.10	0.19	0.00	-0.01	0.49
Loving dogs	0.84	0.01	0.02	0.03	-0.01	0.06	-0.01
No fear of dogs	0.60	0.08	0.03	0.02	0.00	-0.08	-0.01
Touching dogs	0.85	0.06	-0.01	0.01	0.01	0.04	0.04
Dog as a family member	0.36	-0.09	0.01	0.08	0.02	0.62	0.04
No. of cats in house	0.02	-0.06	0.96	0.04	0.01	-0.02	-0.03
Stroking cats	0.00	0.16	0.68	-0.05	0.03	0.05	0.11
Bitten by a cat	-0.01	0.00	0.02	-0.02	0.75	0.01	0.02
Scratched by a cat	0.01	-0.01	-0.02	0.02	0.73	-0.02	-0.04
Cat's smell	-0.11	0.19	0.17	-0.08	0.03	0.03	0.58
Loving cat	0.02	0.88	0.06	-0.01	0.00	-0.01	0.00
No fear of cats	0.18	0.35	0.14	-0.02	-0.08	-0.10	0.00
Touching cats	0.06	0.90	-0.03	0.01	0.03	-0.01	0.04
Cat as a family member	-0.13	0.48	0.08	0.03	0.01	0.50	0.07
Explained variability (%)	13	14	10	9	7	5	5

Table shows factors loadings for particular factors after oblimin rotation. The variables with high positive (low negative) loadings have the highest positive (negative) correlation with a particular factor.

[34]), including humans, and increased level of this neurotransmitter is associated with decreased personality factor novelty seeking [35, 36]. The decreased level of novelty seeking monitored with Cloninger TCI questionnaire [37] was observed in both men and women infected with *T. gondii* [38, 39]. It is highly probable that subjects with low tendency to seek for new stimuli will consume exotic meals such as steak Tartar less often, and possibly also less often drink potentially contaminated water, e.g. water from creeks.

This study confirmed earlier results concerning non-monotonic change in the prevalence of *T. gondii* infection with age in the Czech male subjects [30]. In the current study, the prevalence of *T. gondii* infection in men achieved a maximum of 29.7% at ages 35–39, while a similar study performed on the population of 3250 male soldiers showed a maximum of 35% at ages 30–35. Because the first study was performed 10–15 years earlier than the current study, the observed decrease of prevalence of *T. gondii* infection after age 39 cannot be the effect of cohort (otherwise it should be shifted to age 45–50). On the basis of experience of reactivation of toxoplasmosis in

immunosuppressed patients, including AIDS patients, it is widely believed that dormant stages of the parasite survive in the body of infected subjects, especially in their brains, for their whole lives. However, results of longitudinal serological studies as well as case studies and anecdotal observations show that the concentration of anti-*T. gondii* antibodies irregularly decreases with age of infected subjects, and that some infected subjects could turn seronegative in older age [40]. On the basis of a permutation test for contaminated data, the fraction of seronegative subjects who are definitively *T. gondii*-infected was estimated 5%–10% among subjects age 20–40 [41, 42]. In the higher age strata, the fraction of such subjects is probably even higher. It can be speculated that the rate of acquiring new *T. gondii* infection slows down as subjects become older, and in the age ranges of 35–49 in men and 50–54 in women becomes equal with rate of positivity-to-negativity seroconversions caused by the natural decrease of concentration of anti-*T. gondii* antibodies. If this model is correct, then the prevalence of *T. gondii* infection is in fact much higher than the current serological surveys show. It must be reminded that except for the studies

Table 4. Association between *T. gondii* infection and potential risk factors

	<i>P</i> all	OR all	95% CI all	<i>P</i> men	OR men	<i>P</i> women	OR women
Age	0·00	6·00	3·62–9·94	0·02	4·23	0·00	7·32
Sex	0·01	1·50	1·11–2·03				
Size of place of living in childhood	0·00	0·53	0·38–0·75	0·64	0·81	0·00	0·50
Eating raw meat	0·79	0·95	0·64–1·41	0·12	0·43	0·74	1·08
Touching soil	0·00	3·14	1·55–6·35	0·47	1·84	0·00	3·44
Unwashed vegetables	0·04	1·71	1·02–2·87	0·05	3·64	0·19	1·46
Low-hygienic standard country	0·25	0·75	0·46–1·22	0·18	2·16	0·07	0·60
Suspicious water	0·47	0·83	0·51–1·37	0·19	0·44	0·96	1·02
Risky sexual behavior	0·38	1·22	0·79–1·90	0·75	0·83	0·20	1·38
Loving cats	0·38	0·72	0·35–1·49	0·10	0·27	0·85	0·92
Loving dogs	0·76	1·11	0·56–2·18	0·75	1·27	0·87	1·07
No. of cats at house	0·65	1·14	0·65–2·01	0·15	2·72	0·81	0·93
No. of dogs at house	0·62	0·85	0·45–1·61	0·94	0·94	0·56	0·81
Cat-related injury	0·01	2·16	1·25–3·74	0·06	3·88	0·03	1·93
Pet as a family member	0·70	1·16	0·55–2·42	0·06	4·50	0·58	0·79
Likes animals' smell	0·19	0·56	0·24–1·32	0·37	0·38	0·27	0·59

Associations were measured with Logistic regression. Men are coded as 1s, women 2s; therefore OR > 1 reflects the higher probability of being *Toxoplasma*-infected in women. Significant range Odd Ratios (OR) (in two-side tests) are printed in bold. *P* value 0·00 means *P* < 0·005.

performed on relatively young and age-homogenous populations of subjects, such as pregnant women and soldiers, the earlier epidemiological studies were performed mostly on mixed-gender populations with a rather low number of participants older than 50 years. The opposite trends (decrease *vs.* increase) of change in the prevalence of *T. gondii* infection in 35–54-year-old men and women, together with a usually rather low number of participants in higher age strata in earlier epidemiological surveys; see for example [43], might explain why this phenomenon – i.e., the decrease of prevalence of *T. gondii* infection in men older than 35 and women older than 54 – has escaped the attention of other researchers and why most of the previous studies report a stagnation of prevalence of *T. gondii* infection after the age of 30. The exception was a large study of Kodym *et al.* [44] (*N* = 3431). This study included nearly 500 participants in each 10-years strata and it showed the decrease of prevalence of *T. gondii* infection in men older than 59 and in women older than 49.

Limits of the present study

A major limit of the present study is the fact that its participants provided information about their *T. gondii* infection status themselves. At least some of them probably provided incorrect information and some of them (only the *T. gondii*-free) may have provided

obsolete information because they acquired the infection only after their test for anti-*T. gondii* antibodies had been done. It could be advocated that at least some of old individuals reported the historic status rather than the current status. However, even in such a case the prevalence in older age strata would be either same or higher than in younger age strata. Moreover, the decreased prevalence of toxoplasmosis in older age strata has been reported also in two previous studies in which the participants were directly serologically examined for the presence of specific antibodies [43, 44]. It must be also stressed that stochastic error could only cause false negative results, such as the failure to detect some weak risk factors, but not false positive results, such as identification of an actually non-existent risk factor [45]. There is no indication pointing to the existence of a systematic bias in the present data (for example, for higher probability of reporting positive toxoplasmosis status by cat keepers). Our independent analysis showed nearly perfect (99·5%) agreement between toxoplasmosis status reported by 3827 subjects during their electronic registration to Guinea Pigs community and toxoplasmosis status obtained in our serological tests. Similarly, we observed nearly perfect agreement (99·2%) in 393 responders who signed their questionnaire in the present study and also reported their toxoplasmosis status during registration to the Guinea Pigs community. Moreover, the only animal-related

variables positively associated with *Toxoplasma* infection were animal-related injuries. These variables are not considered to be toxoplasmosis risk factors by most parasitologists and laymen. In the context of our results on observed rapid decrease of seroprevalence of *T. gondii* infection in higher age strata and presumed frequent positivity-to-negativity seroconversion in older subjects, it can be argued that the epidemiological studies based on the reported results of passed serological tests could be actually more precise and sensitive than the cross-sectional studies, which rely on the serological examination of the current seropositivity/seronegativity status of participants.

Our systematic testing of subjects for *T. gondii* infection started about 25 years ago. Therefore, only 87 participants of the present study were older than 60 years. Therefore, the estimated prevalence of *T. gondii* infection in this age category could be relatively imprecise.

The current study involved mostly healthy people with the so-called latent toxoplasmosis, i.e. the form of the *T. gondii* infection characterized by the presence of non-sterile immunity against *T. gondii* and no clinical signs of the disease. It is possible that different sets of risk factors exist for this asymptomatic form of the infection and for the acute and chronic forms of toxoplasmosis. Theoretically, a more serious form of toxoplasmosis could be associated with the infection of humans with tissue cysts than with the oocysts because the tissue cysts are primarily destined for the infection of definitive host, a cat. Many predation transmitted-parasites harm their intermediate hosts, but not their longevous definitive host, which often serve them mainly as a meeting point for sexual partners and as reservoirs and vehicles of gene flow across space and time.

The present study was performed on a sample of internet users, very often previous students of biology at the Faculty of Science, Charles University. It is not clear which results of the study could be generalized to the standard Czech population.

CONCLUSIONS

The present study identified new risk factor for the acquisition of *T. gondii* infection for a modern population, namely cat-related injuries. The most important result of the present study is the observed decrease of seroprevalence of *T. gondii* infection, which starts around the age of 50 in men and age 55 in women. Experience with reactivation of

toxoplasmosis in the infected immunosuppressed patients shows that *T. gondii* survives in immunoprivileged organs of human hosts until end of life [46]. Therefore, the existence of such a decrease can be explained either by increased mortality rate of *T. gondii*-infected subjects [31, 47] or by frequent positivity-to-negativity seroconversions in infected subjects. If this is true, then either the prevalence of *T. gondii* infection or its potential impacts on public health could be much larger than generally believed.

ACKNOWLEDGEMENTS

The work was supported by project UNCE 204004 (Charles University in Prague) and the Czech Science Foundation (Grant No. P407/16/20958). I would like to thank to Raymon Gongora, Gary Hawkins, Charlie Lotterman and Lenka Příplatová for their help with final version of the paper.

DECLARATION OF INTERESTS

None.

ETHICAL STANDARDS

The author asserts that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

REFERENCES

1. Tenter AM, Heckeroth AR, Weiss LM. *Toxoplasma gondii*: from animals to humans. *International Journal for Parasitology* 2000; **30**: 1217–1258.
2. Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis. *International Journal for Parasitology* 2009; **39**: 1385–1394.
3. Dubey JP. History of the discovery of the life cycle of *Toxoplasma gondii*. *International Journal for Parasitology* 2009; **39**: 877–882.
4. Boyer KM, *et al.* Risk factors for *Toxoplasma gondii* infection in mothers of infants with congenital toxoplasmosis: implications for prenatal management and screening. *American Journal of Obstetrics and Gynecology* 2005; **192**: 564–571.
5. Petersen E, *et al.* What do we know about risk factors for infection in humans with *Toxoplasma gondii* and how can we prevent infections? *Zoonoses and Public Health* 2010; **57**: 8–17.

6. **Ferreira AIC, et al.** Risk factors for ocular toxoplasmosis in Brazil. *Epidemiology and Infection* 2014; **142**: 142–148.
7. **Abu EK, et al.** Infection risk factors associated with seropositivity for *Toxoplasma gondii* in a population-based study in the Central Region, Ghana. *Epidemiology and Infection* 2015; **143**: 1904–1912.
8. **Cong W, et al.** Seroprevalence and associated risk factors of *Toxoplasma gondii* infection in psychiatric patients: a case-control study in eastern China. *Epidemiology and Infection* 2015; **143**: 3103–3109.
9. **Minbaeva G, et al.** *Toxoplasma gondii* infection in Kyrgyzstan: seroprevalence, risk factor analysis, and estimate of congenital and AIDS-related toxoplasmosis. *PLoS Neglected Tropical Diseases* 2013; **7**.
10. **Sharbatkhori M, et al.** Seroprevalence of *Toxoplasma gondii* infections in pregnant women in Gorgan City, Golestan Province, Northern Iran-2012. *Iranian Journal of Parasitology* 2014; **9**: 181–187.
11. **Etheredge GD, et al.** The roles of cats and dogs in the transmission of *Toxoplasma* infection in Kuna and Embera children in eastern Panama. *Revista Panamericana de Salud Publica-Pan American Journal of Public Health* 2004; **16**: 176–186.
12. **de Moura FL, et al.** Prevalence and risk factors for *Toxoplasma gondii* infection among pregnant and postpartum women attended at public healthcare facilities in the City of Niteroi, State of Rio de Janeiro, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* 2013; **46**: 200–207.
13. **Chiang TY, et al.** Risk factors for acute *Toxoplasma gondii* diseases in Taiwan: a population-based case-control study. *PLoS ONE* 2014; **9**.
14. **Markovich MP, et al.** Seroepidemiology of *Toxoplasma gondii* infection in the Israeli population. *Epidemiology and Infection* 2014; **142**: 149–155.
15. **Flegr J, Klapilová K, Kaňková Š.** Toxoplasmosis can be a sexually transmitted infection with serious clinical consequences. Not all routes of infection are created equal. *Medical Hypotheses* 2014; **83**: 286–289.
16. **Alvarado-Esquivel C, et al.** High seroprevalence of *Toxoplasma gondii* infection in female sex workers: a case-control study. *European Journal of Microbiology and Immunology (Bp)* 2015; **5**: 285–292.
17. **Kankova S, Flegr J, Calda P.** The influence of latent toxoplasmosis on women's reproductive function: four cross-sectional studies. *Folia Parasitologica* 2015; **62**.
18. **Gosling SD, Rentfrow PJ, Swann WB.** A very brief measure of the Big-Five personality domains. *Journal of Research in Personality* 2003; **37**: 504–528.
19. **Beck AT, Steer RA, Brown GK.** *BDI-II: Depression Inventory Manual*, 2nd edn. Sant Antonio: Psychological Corporation, 1996.
20. **Foa EB, et al.** The obsessive-compulsive inventory: development and validation of a short version. *Psychological Assessment* 2002; **14**: 485–496.
21. **Flegr J.** Influence of latent toxoplasmosis on the phenotype of intermediate hosts. *Folia Parasitologica* 2010; **57**: 81–87.
22. **Benjamini Y, Hochberg Y.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B – Methodological* 1995; **57**: 289–300.
23. **Dubey JP.** Transmission of *Toxoplasma gondii* – From land to sea, a personal perspective. In: Janovy J and Esch GW, eds. *A Century of Parasitology: Discoveries, Ideas and Lessons Learned by Scientists who Published in the Journal of Parasitology, 1914–2014, First Edition*. New York: John Wiley & Sons, 2016, pp. 148–164.
24. **Gotteland C, et al.** Spatial distribution of *Toxoplasma gondii* oocysts in soil in a rural area: influence of cats and land use. *Veterinary Parasitology* 2014; **205**: 629–637.
25. **Torrey EF, Yolken RH.** *Toxoplasma* oocysts as a public health problem. *Trends in Parasitology* 2013; **29**: 380–384.
26. **Flegr J, Hrdá Š, Tachezy J.** The role of psychological factors in questionnaire-based studies on routes of human toxoplasmosis transmission. *Central European Journal of Public Health* 1998; **6**: 45–50.
27. **Hanauer DA, Ramakrishnan N, Seyfried LS.** Describing the relationship between cat bites and human depression using data from an electronic health record. *PLoS ONE* 2013; **8**: e70585.
28. **Flegr J, Hodny Z.** Cat scratches, not bites, are associated with unipolar depression – cross-sectional study. *Parasites & Vectors* 2016; **9**.
29. **Flegr J, et al.** Fatal attraction phenomenon in humans: cat odour attractiveness increased for *Toxoplasma*-infected men while decreased for infected women. *PLoS Neglected Tropical Diseases* 2011; **5**: e1389.
30. **Kolbeková P, et al.** New and old risk-factors for *Toxoplasma gondii* infection: prospective cross-sectional study among military personnel in the Czech Republic. *Clinical Microbiology and Infection* 2007; **13**: 1012–1017.
31. **Flegr J, et al.** Toxoplasmosis - A global threat. Correlation of latent toxoplasmosis with specific disease burden in a set of 88 countries. *PLoS ONE* 2014; **9**.
32. **Gatkowska J, et al.** Sex-dependent neurotransmitter level changes in brains of *Toxoplasma gondii* infected mice. *Experimental Parasitology* 2013; **133**: 1–7.
33. **Prandovszky E, et al.** The neurotropic parasite *Toxoplasma gondii* increases dopamine metabolism. *PLoS ONE* 2011; **6**: e23866.
34. **Wang ZT, et al.** Reassessment of the role of aromatic amino acid hydroxylases and the effect of infection by *Toxoplasma gondii* on host dopamine. *Infection and Immunity* 2015; **83**: 1039–1047.
35. **Cloninger CR.** The genetics and psychobiology of the seven-factor model of personality. In: Silk KR, ed. *Biology of Personality Disorders*. Washington, DC: American Psychiatric Press, Inc., 1998, pp. 63–92.
36. **Wiesbeck GA, et al.** Neuroendocrine support for a relationship between “novelty seeking” and dopaminergic function in alcohol-dependent men. *Psychoneuroendocrinology* 1995; **20**: 755–761.

37. **Cloninger CR, et al.** *The Temperament and Character Inventory (TCI): A Guide to its Development and Use*. St. Louis, Missouri: Center for Psychobiology of Personality, Washington University Press, 1994.
38. **Flegr J, et al.** Decreased level of psychobiological factor novelty seeking and lower intelligence in men latently infected with the protozoan parasite *Toxoplasma gondii*. Dopamine, a missing link between schizophrenia and toxoplasmosis? *Biological Psychology* 2003; **63**: 253–268.
39. **Skallová A, et al.** Decreased level of novelty seeking in blood donors infected with *Toxoplasma*. *Neuroendocrinology Letters* 2005; **26**: 480–486.
40. **Kodym P, et al.** Evaluation of a commercial IgE ELISA in comparison with IgA and IgM ELISAs, IgG avidity assay and complement fixation for the diagnosis of acute toxoplasmosis. *Clinical Microbiology and Infection* 2007; **13**: 40–47.
41. **Flegr J, Havlíček J.** Changes in the personality profile of young women with latent toxoplasmosis. *Folia Parasitologica* 1999; **46**: 22–28.
42. **Flegr J, Hrdá Š, Kodym P.** Influence of latent ‘asymptomatic’ toxoplasmosis on body weight of pregnant women. *Folia Parasitologica* 2005; **52**: 199–204.
43. **Jones JL, Kruszon-Moran D, Wilson M.** *Toxoplasma gondii* infection in the United States, 1999–2000. *Emerging Infectious Diseases* 2003; **9**: 1371–1374.
44. **Kodym P, et al.** *Toxoplasma* in the Czech Republic 1923–1999: first case to widespread outbreak. *International Journal for Parasitology* 2000; **30**: 11–18.
45. **Flegr J.** Could contamination of data with misclassified individuals increase the probability of false positive results of statistical tests? *Fishare* 2016; dx.doi.org/10.6084/m9.figshare.3806553.v1.
46. **Flegr J.** How and why *Toxoplasma* makes us crazy. *Trends in Parasitology* 2013; **29**: 156–163.
47. **Flegr J, Escudero D.** Impaired health status and increased incidence of diseases in *Toxoplasma*-seropositive subjects – An explorative cross-sectional study. *Parasitology* 2016; **143**: 1974–1989.