

Research Article

Cite this article: Yang L, Wei Z, Su Y, Wang X, Du Y, Farooq MZ, Li J, Xu Q (2024) An organic zinc-chelating-peptide GQ-Zn supplementation ameliorates diarrhea through regulate gut microbiota and metabolites in pre-weaning Holstein calves. *Animal Nutriomics*, e17, 1–12. <https://doi.org/10.1017/anr.2024.11>

Received: 27 February 2024

Revised: 9 May 2024

Accepted: 11 June 2024


Keywords:

calf diarrhea; GQ-Zn; inflammation; gut microbiota; metabolomics

Corresponding author: Qingbiao Xu;

Email: qbxu@mail.hzau.edu.cn

An organic zinc-chelating-peptide GQ-Zn supplementation ameliorates diarrhea through regulate gut microbiota and metabolites in pre-weaning Holstein calves

Linhai Yang^{1,2}, Zihai Wei¹, Yanjing Su¹, Xianghuang Wang² , Yufeng Du², Muhammad Zahid Farooq^{2,3}, Jianxiong Li⁴ and Qingbiao Xu²

¹Ministry of Agriculture and Rural Affairs Key Laboratory of Dairy Cattles Genetic Improvement in Southern China, Bright Farming Co., Ltd., Shanghai, P.R. China; ²College of Animal Sciences and Technology, Huazhong Agricultural University, Wuhan, P.R. China; ³Department of Animal Science, University of Veterinary and Animal Science, Lahore, Pakistan and ⁴Wuhan Jason Biotech Co., Ltd., Wuhan, P.R. China

Abstract

Newborn calf diarrhea has led to widespread overuse of antibiotics. Therefore, it is crucial to find effective solutions for calf diarrhea. In this study, we aimed to evaluate the impact of the synthetic organic zinc-chelating-peptide glycine-glutamine-Zn (GQ-Zn) on the microbiota and metabolites in the gut of calves with diarrhea. The results showed that GQ-Zn alleviated diarrhea in calves. Additionally, 16S rDNA sequencing and metabolomic analysis revealed that GQ-Zn improved antioxidant capacity, relieved inflammation, altered the gut microbiota by decreasing the number of harmful bacteria *Prevotella denticola*, *Fusobacterium necrophorum* and influenced metabolomic profiles via the linoleic acid metabolic pathway in calves. In conclusion, GQ-Zn supplementation alleviated diarrhea through regulating the gut microbiota and metabolites in pre-weaning Holstein calves.

Introduction

Ranch managers have long struggled with problems related to animal industrialization, such as diarrhea (Klein-Jobstl et al. 2014). In livestock, diarrhea is typically classified as either infectious or noninfectious. Infectious agents include a variety of bacteria, viruses and parasites (Cho et al. 2013). Causes of noninfectious diarrhea include poor management of breeding, changes in diet and malnutrition (Murray et al. 2016). In a pasture, an outbreak of neonatal calf diarrhea in 4-week-old calves causing 80% morbidity rate and 20% mortality rate (Dall Agnol et al. 2021). In the Netherlands, the incidence of calf diarrhea is 24% in the first 3 weeks, while in Canada, the incidence of calf diarrhea is 20% during weaning (Bartels et al. 2010). Diarrhea is a major threat to cow production. Therefore, it is crucial to search for new feed additives to reduce the incidence of calf diarrhea, which could increase the number of Holstein calves that reach maturity and in turn improve the profitability of related enterprises.

Intestinal microbiota play an important role in the occurrence and treatment of calf diarrhea. Kim found that fecal microbial transplantation could improve fecal score and alleviate diarrhea of pre-weaning calves, resulting in the gradual maturation of intestinal microbiota of calves, indicating the importance of microbiota in alleviating diarrhea of calves (Kim et al. 2021). Certain probiotics, such as *Lactobacillus*, have demonstrated a significant reduction in the incidence of diarrhea in cattle (Schwaiger et al. 2023). Certain pathogenic microbiota such as *Escherichia coli* K99 and *Salmonella* can lead to diarrhea in calves (Du et al. 2023b). The fecal samples of calves were analyzed for microbiota, and there was a significant correlation between pathogenic microbiota such as *Escherichia coli* K99, *Salmonella*, *Clostridium* and diarrhea of calves (Cho et al. 2013). *Escherichia coli* K99 and *Salmonella* are the main microbiota of antibiotic resistance (Du et al. 2023a).

Antibiotics was useful for promoting growth and curing disorders caused by microbial infections such as diarrhea but accompanied with the reduction of absorption rate in livestock (Heinrichs et al. 2003). Considerable evidence showed that misuse of these antibiotics is a major contributor to current health crises (Allahverdiyev et al. 2011). Antibiotics are no longer as effective as they once were because drug-resistant microbiota proliferate (Mobarki et al. 2019). Varieties of antibiotics are accumulated by humans because of biomagnification (Chee-Sanford et al. 2009). Calf diarrhea treated with medicines often relapses after a few days (Shin et al. 2015). It is possible that antibiotics are bactericidal against a wide range of microbiota, which leads to

© The Author(s), 2024. Published by Cambridge University Press on behalf of Zhejiang University and Zhejiang University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

an imbalance of gut microbiome. Furthermore, even if calves recover from diarrhea, antibiotics are not conducive to their future growth potential because they interfere with the development and colonization of rumen microbiota (Okada *et al.* 2023). Overall, the excessive application of antibiotics in livestock should be restricted, and additives substitutes should be developed.

Zinc is an essential mineral for metabolism and has been widely used in animal husbandry, which was proven to promote animal growth and alleviate diarrhea (Hill and Shannon 2019). Zinc supplements are divided into two main types: inorganic zinc and organic zinc. Inorganic zinc (e.g., ZnO, ZnSO₄) is widely used during the weaning period to alleviate diarrhea in piglets by strengthening intestinal barrier function (Zhu *et al.* 2017). Although ZnO is beneficial to young animals, it often results in the loss of significant amounts of zinc, which causes environmental pollution (Ogiyama *et al.* 2005). Organic zinc has higher bioavailability, allowing animals to digest, absorb and utilize it better, especially when zinc is chelated with amino acids or small peptides (Case and Carlson 2002). The bioavailability of ZnO is lower than Zn-methionine and Zn-lysine in high concentrations of Zn in weanling pigs (Schell and Kornegay 1996). Previous study found that glycine-zinc significantly improved the weekly weight gain, feed intake, feed conversion ratio of control and inorganic zinc in broilers (Gul and Alhidary 2024). In our study, zinc was chelated with glycyl-glutamine dipeptides (GQ-Zn) to form new organic zinc.

The amount of zinc added to calf milk supplements in China cannot exceed 180 mg/kg. Previous studies suggested that different doses of zinc have different effects on animals (Wright and Spears 2004). Due to the negative effects of high doses of zinc, China completely banned the addition of excessive zinc in 2017 (Liu *et al.* 2020). To determine the optimal level of supplementation, a two-pronged approach was used in this study. The aim of this research was to study a novel organic zinc GQ-Zn on the growth performance, health status and microbiome of the cattle.

Materials and methods

Animals and management

The trial was conducted on the Modern Farming Saibei Phase III Pasture (Guyuan County, Zhangjiakou City, Hebei, China). The experiment lasted for a total of 14 days. Thirty Holstein female calves with similar birth dates, weights and genetic backgrounds were randomly assigned to three groups ($n = 10$). Every morning during the experiment, 50 mL of purified water was given to the control group (Con), 50 mL of glycine-glutamine-Zn (GQ-Zn) solution (80 mg/d) was given to the low-zinc group (Low) and 50 mL of GQ-Zn solution (160 mg/d) was given to the high-zinc group (High). During the experiment, the calves were kept on the calf island and fed 4 kg of pasteurized milk and 250 g of starter every day, with free access to food and water. If the temperature drops, a calf warmer garment will come in handy. Calves that are severely unwell or have diarrhea should be treated or euthanized.

Data and sample collection

A group of calves were fed organic zinc to further examine its impact on calves with diarrhea. The severity of diarrhea in the calves was recorded daily, and the rectal contents were collected at 0, 7 and 14 days. The weights of the calves were measured with

a weighbridge on Days 0, 7, 14, 60 and 365. On the mornings of Days 7 and 14, 10 mL of blood was collected from the jugular vein by using a blood collection vessel without anticoagulants. To isolate the serum, the blood was left at room temperature for 1 hour and centrifuged at 4000 rpm for 20 min. The upper layer of serum was extracted with a pipette gun and released into centrifuge tubes, which were stored at -20°C until analysis of the serum biochemical indices. On Days 0, 7 and 14, 5 g of feces was collected from the calf rectum and packed into 5 mL sterile cryogenic tubes, which were subsequently stored in a -80°C refrigerator.

Determination of diarrhea incidence

The criteria for determining the fecal score were established according to previous methods (Magalhaes *et al.* 2008). Fecal scored as 1 when firm, 2 when soft or of moderate consistency, 3 when runny or mild diarrhea and 4 when watery and profuse diarrhea. Calf feces were observed and scored daily during the experiment. Score ≥ 3 was considered to calf diarrhea, and the diarrhea rate was calculated based on the score. In the control group, one calf died due to severe diarrhea on Day 9.

Analysis of biochemical indicators

Serum biochemical indicators, including the antioxidant indicators CAT, MDA, SOD, GSH-Px and T-AOC and the immune indicators IgA, IgG, SIgA, IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-17, IL-22, IFN- γ , TGF- β and TNF- α , were analyzed using an enzyme-linked immunosorbent assay kit (Meimian, Jiangsu, China) according to the manufacturer's instructions.

Determination of short-chain fatty acids (SCFAs)

Gas chromatography was used to determine the concentration of SCFAs in calf feces. A 1 g sample was placed into a 2 mL tube, and 1 mL of methanol was added to the tube, which was centrifuged at $12000 \times g$ for 10 min. One milliliter of supernatant and 0.2 mL of 25% metaphosphate solution were mixed and centrifuged at $12000 \times g$ for 10 min. Finally, the sample was filtered through a $0.45 \mu\text{m}$ membrane, and the sample was injected into a gas chromatograph (Yan *et al.* 2019). One microliter of sample was injected into a gas chromatograph (Trace 1300, Thermo Fisher Scientific, MA, USA). A flame ionization detector was used with an oven temperature of 75°C , an inlet and detector temperature of 230°C and a column flow and purge flow of 3 mL/min and 5 mL/min for determination of acetate, propionate, butyrate, isobutyrate, valerate and isovalerate concentrations.

Determination of the fecal microbiome

Total genomic DNA was extracted from feces samples using the TGuide S96 Magnetic Soil /Stool DNA Kit (Tiangen Biotech [Beijing] Co., Ltd.) according to manufacturer's instructions. The quality and quantity of the extracted DNA were examined using electrophoresis on a 1.8% agarose gel, and DNA concentration and purity were determined with NanoDrop 2000 UV-Vis spectrophotometer (Thermo Scientific, Wilmington, USA). The full-length 16S rRNA gene were amplified with primer pairs 27F: AGRGTTTGATYNTGGCTCAG and 1492R: TASGGHTACCTTGTASGACTT. Both the forward and reverse 16S primers were tailed with sample-specific PacBio barcode

sequences to allow for multiplexed sequencing. We chose to use barcoded primers because this reduces chimera formation as compared to the alternative protocol in which primers are added in a second polymerase chain reaction (PCR) reaction. The KOD One PCR Master Mix (TOYOBO Life Science) was used to perform 25 cycles of PCR amplification, with initial denaturation at 95°C for 2 min, followed by 25 cycles of denaturation at 98°C for 10 s, annealing at 55°C for 30 s, extension at 72°C for 1 min 30 s and a final step at 72°C for 2 min. The total of PCR amplicons were purified with VAHTSTM DNA Clean Beads (Vazyme, Nanjing, China) and quantified using the Qubit dsDNA HS Assay Kit and Qubit 3.0 Fluorometer (Invitrogen, Thermo Fisher Scientific, Oregon, USA). After the individual quantification step, amplicons were pooled in equal amounts. SMRTbell libraries were prepared from the amplified DNA by SMRTbell Express Template Prep Kit 2.0 according to the manufacturer's instructions (Pacific Biosciences). Purified SMRTbell libraries from the pooled and barcoded samples were sequenced on a PacBio Sequel II platform (Beijing Biomarker Technologies Co., Ltd., Beijing, China) using Sequel II binding kit 2.0.

The bioinformatics analysis of this study was performed with the aid of the BMKCloud (<http://www.biocloud.net/>). The raw reads generated from sequencing were filtered and demultiplexed using the SMRT Link software (version 8.0) with the $\text{minPasses} \geq 5$ and $\text{minPredictedAccuracy} \geq 0.9$, in order to obtain the circular consensus sequencing (CCS) reads. Subsequently, the lima (version 1.7.0) was employed to assign the CCS sequences to the corresponding samples based on their barcodes. CCS reads containing no primers and those reads beyond the length range (1,200–1,650 bp) were discarded through the recognition of forward and reverse primers and quality filtering using the Cutadapt (Bolger Anthony et al. 2014) (version 2.7) quality control process. The UCHIME algorithm (v8.1) (Kechin et al. 2017) was used in detecting and removing chimera sequences to obtain the clean reads. Sequences with similarity $>97\%$ were clustered into the same operational taxonomic unit (OTU) by USEARCH (Edgar Robert 2013) (v10.0), and the OTUs counts less than 2 in all samples were filtered.

Clean reads then were conducted on feature classification to output an ASV (amplicon sequence variant) by DADA2 (Abreu et al. 2020), and the ASVs counts less than 2 in all samples were filtered. Taxonomy annotation of the OTUs/ASVs was performed based on the Naive Bayes classifier in QIIME2 (Evan et al. 2019) using the SILVA database (Christian et al. 2013) (release 138.1) with a confidence threshold of 70%. The alpha diversity was calculated and displayed by the QIIME2 and R software, respectively. Beta diversity was determined to evaluate the degree of similarity of microbial communities from different samples using QIIME. Principal coordinate analysis (PCoA), heatmaps, unweighted pair-group method with arithmetic means (UPGMA) and nonmetric multidimensional scaling were used to analyze the beta diversity. To explore the dissimilarities of the microbiome among different factors, a redundancy analysis was performed in R using the package *vegan*.

The qualified sequences with more than 97% similarity thresholds were allocated to one OTU using USEARCH (version 10.0). Taxonomy annotation of the OTUs/ASVs was performed based on the Naive Bayes classifier in QIIME2 (Evan et al. 2019) using the SILVA database (Christian et al. 2013) (release 138.1) with a confidence threshold of 70%. Alpha was performed to identify the complexity of species diversity of each sample utilizing QIIME2 software. Beta diversity calculations were analyzed by PCoA to

assess the diversity in samples for species complexity. One-way analysis of variance (ANOVA) was used to compare bacterial abundance and diversity. The online platform BMKCloud (<https://www.biocloud.net>) was used to analyze the sequencing data.

Determination of fecal metabolites

After processing the sample, LC–MS was used for detection. The raw data were first converted to mzXML format by MSConvert in the ProteoWizard software package (v3.0.8789) and processed using XCMS for feature detection, retention time correction and alignment. The metabolites were identified by accurate mass (<30 ppm) and MS/MS data, which were matched with HMDB, MassBank, LipidMaps, mzCloud and KEGG data. The robust locally Estimated Scatterplot Smoothing (LOESS) signal correction (Gagnebin et al. 2017) was applied for data normalization to correct for any systematic bias. After normalization, only ion peaks with relative standard deviations less than 30% in the quality control (QC) were retained to ensure proper metabolite identification.

Ropls software was used for all multivariate data analyses and modeling. The data were mean-centered using scaling. Models were built via principal component analysis, orthogonal partial least-square discriminant analysis (OPLS-DA) and partial least-square discriminant analysis. The *P* value, variable importance projection produced by OPLS-DA and fold change were applied to discover the contributable variables for classification. The differentially abundant metabolites were subjected to pathway analysis via MetaboAnalyst, which combines the results from powerful pathway enrichment analysis with pathway topology analysis. The metabolites identified via metabolomics were subsequently mapped to the KEGG pathway for biological interpretation of higher-level systemic functions. The metabolites and corresponding pathways were visualized using the KEGG Mapper tool.

Statistical analysis

The data were analyzed using GraphPad Prism software with *t*-tests and one-way ANOVA. All the figures were created with GraphPad Prism 9.0 (GraphPad Software, Inc., La Jolla, CA, USA). In all analyses, *P* values < 0.05 were considered to indicate statistical significance ($*P < 0.05$).

Results

GQ-Zn ameliorated diarrhea in pre-weaning calves

The results showed that GQ-Zn reduced the incidence of diarrhea in calves (Fig. 1A–C). Evaluation of stool fluidity using the stool score method showed that the stool score in the high-GQ-Zn group was significantly lower than that in the control group on Days 7, 9, 10 and 12 (Fig. 1D). The calves were weighed on Days 0, 7, 14, 60 and 365 and the growth ability of the calves was evaluated. The results indicated that GQ-Zn had no significant effect on the growth rate of the calves (Table 1).

GQ-Zn enhanced the antioxidant capacity of calves and reduced inflammatory reactions

Several serum biomarkers have been shown to be significantly different between different treatment groups. As expected, these indicators changed after processing (Fig. 2). The antioxidant capacity of the calves was assessed. GQ-Zn significantly increased




Figure 1. The effect of GQ-Zn on the incidence of diarrhea of pre-weaning Holstein calves. Images of the fecal samples collected from the recta of control (A), low (B) and high concentration of GQ-Zn (C). The incidence of diarrhea was represented by the score of fecal samples (D). The data were presented as the mean \pm SEM and evaluated using one-way ANOVA ($n = 10$).

Table 1. Body weight and ADG of low and high concentration of supplementation with GQ-Zn in pre-weaned calves

Item	Con ^a	Low	High	SEM	<i>P</i> -value
BW (kg)					
d0 ^b	38.15	36.60	37.00	0.61	0.5794
d7	50.80	49.95	50.05	0.68	0.8650
d14	57.17	55.45	55.80	0.72	0.6151
d60	110.20	104.20	106.40	1.63	0.3213
d365	335.10	332.50	349.40	3.53	0.0935
ADG ^c (kg/d)					
d0–7	1.81	1.91	1.86	0.10	0.9186
d8–14	0.80	0.79	0.82	0.06	0.9709
d15–60	1.04	0.96	0.99	0.03	0.3935
d60–365	0.78	0.74	0.84	0.02	0.1362

Notes: ^aCon = supplementation with 50 mL purified water, Low = supplementation with 50 mL 80 mg/d GQ-Zn solution, High = supplementation with 50 mL 160 mg/d GQ-Zn solution.

^bDays after supplementation with GQ-Zn.

^cADG = average daily gain.

the level of CAT and decreased the serum MDA concentration (Fig. 2A and B). In terms of immunity, GQ-Zn significantly

increased the level of sIgA and decreased the serum levels of IL-2 and IFN- γ (Fig. 3D, E and M). However, no significant differences in the other indicators were observed between the groups (Fig. 3).

SCFAs produced by the gut microbiota

We analyzed the SCFAs in rectal contents of calves in different treatment groups by GC-MS (Fig. 4). We observed that after GQ-Zn supplementation, different concentrations of zinc had different effects on calves, and high concentrations of GQ-Zn tended to reduce the concentration of propionate on Day 14 (Fig. 4B), while low concentrations of GQ-Zn increased the concentration of valerate on Day 7 (Fig. 4E).

GQ-Zn alters the gut microbiota composition of calves

We further investigated the gut microbiota of the three different experimental groups by 16S rDNA sequencing. A Venn diagram was constructed to display the number of OTUs annotated from the different experimental groups: control, low concentration of GQ-Zn and high concentration of GQ-Zn had 69, 13 and 9 unique OTUs, respectively (Fig. 5A). We used alpha diversity and beta diversity to assess the diversity of the gut microbiota. Alpha-diversity-based analysis revealed little microbial difference between different groups (Fig. 5C–F). According to the beta diversity determined via PCoA, there was




Figure 2. Effects of GQ-Zn on the level of antioxidant indicators in serum in pre-weaning Holstein calves. The concentration of catalase (CAT) (A), malondialdehyde (MDA) (B), superoxide dismutase (SOD) (C), glutathione peroxidase (GSH-Px) (D), total antioxidant capacity (T-AOC) (E) in serum. The data were presented as the mean \pm SEM and evaluated using one-way ANOVA ($n = 10$). * $P < 0.05$.

little difference in the microbial structure between the different groups (Fig. 5B). After conducting a histogram analysis of the annotated OTUs, an obvious difference in the taxonomic composition was found between the three experimental groups at the phylum (Fig. 5G) and genus (Fig. 5H) levels. Metastats analysis was used to evaluate the differences in microbiota composition between group Con, Low and High (Fig. 6) in species level. Five highest differential relative abundance bacteria were presented. Low concentrations of GQ-Zn decreased *Cenchrus americanus*, *Enterococcus faecium*, *Lentibacillus massiliensis*, *Prevotella denticola* and *unclassified Alphaproteobacteria*. High concentrations of GQ-Zn decreased *Agathobaculum butyriciproducens*, *Cenchrus americanus*, *Fusobacterium necrophorum*, *Prevotella denticola* and *Roseburia inulinivorans*.

GQ-Zn alters the fecal metabolomic profile of diarrheic calves

To clarify the application potential of GQ-Zn, it is important to understand how the products of microorganisms affect the incidence of calf diarrhea. We analyzed the fecal metabolome of

calves in different experimental groups. After feeding, the calves were treated with different doses of zinc. PCoA revealed that the data points from different experimental groups became clustered, and there was significant separation between groups (Fig. 7A). Moreover, the amount of fecal metabolites also changed, showing an overall upward trend (Fig. 7B). The relative abundances of various metabolites were identified and annotated in rectal content samples. Enrichment of these metabolites leads to upregulation of multiple metabolic pathways, and the most significant difference is in the digestion and absorption of vitamins. Hierarchical clustering analysis was used to cluster the metabolites in calf feces from different experimental groups. Heatmap analysis revealed that the fecal metabolite profile of calves fed GQ-Zn was different from that of the other groups (Fig. 7C). Most of the metabolites tended to increase, but a small portion of the metabolites were down-regulated (Fig. 7D). Based on these findings, it seems likely that GQ-Zn has a major impact on fecal metabolites, which may be the cause of the reversal of diarrhea in calves. The majority of these metabolites play important roles in the metabolism of linoleic acid (Fig. 7E-L).




Figure 3. Effects of GQ-Zn on the level of immune indicators in serum in pre-weaning Holstein calves. The concentration of TNF- α (A), IL-1 β (B), TGF- β (C), IFN- γ (D), IL-2 (E), IL-4 (F), IL-6 (G), IL-8 (H), IL-10 (I), IL-17 (J), IL-22 (K), IgA (L), IgG (N) in serum and the concentration of sIgA (M) in feces. The data were presented as the mean \pm SEM and evaluated using one-way ANOVA ($n = 10$). * $P < 0.05$, ** $P < 0.01$.

Discussion

GQ-Zn formed via chelation of Gly-Gln and zinc. The results of the present study indicated that GQ-Zn supplementation had no effect on ADG during days 1–365 of the experiment. Although some experiments have shown that organic zinc can promote growth, there is still controversy over whether zinc improves growth performance (Jenkins and Hidirolou 1991). The reason for these conflicting results is still unclear. It may be due to the different environments in which the animals are located, or it has an impact on long-term growth. In a preliminary laboratory study, zinc was found to play a role in promoting growth rather than alleviating diarrhea (Jiang *et al.* 2009). Our result showed that GQ-Zn had no significant effect on calf performance.

In the present study, GQ-Zn significantly reduced the stool score in days 7–14 of the experiment. A high concentration of GQ-Zn significantly reduced the incidence of diarrhea, indicating that

GQ-Zn helped to reduce the incidence and severity of diarrhea. Several studies have shown that zinc is a guardian of immune function. Zinc alleviates diarrhea and is associated with the immune system (Chasapis *et al.* 2020). Animal diarrhea is accompanied by intestinal inflammation (Hodges and Gill 2010).

Inflammation occurs when the intestine is subjected to a series of complex and harmful stimuli (e.g., pathogens), and excessive inflammation is harmful to the body (Reuter *et al.* 2010). Zinc induces regulatory T cells to inhibit the production of IFN- γ (Maywald and Rink 2017). IFN- γ is the main inflammatory cytokine secreted by Th1 cells and promotes the upregulation of multiple proinflammatory factors to activate macrophages (Askenasy 2015; Lees 2015). IFN- γ inhibits the replication of *Listeria*, *Salmonella* and *Mycobacterium tuberculosis* in cells by promoting the iron ion output of pathogen macrophages (Abreu *et al.* 2020). GQ-Zn decreases IFN- γ in serum possibly through the




Figure 4. Effects of GQ-Zn on short-chain fatty acids in pre-weaning Holstein calves. Concentrations of acetate (A), propionate (B), butyrate (C), isobutyrate (D), pentanoate (E), isovalerate (F) and total short-chain fatty acids (SCFAs) (G) in the gut of calves. The data are presented as the mean \pm SEM ($n = 10$) and evaluated using one-way ANOVA. * $P < 0.05$.

action of zinc in vivo. IL-2 has a variety of biological activities and is highly important for immune regulation. IL-2 binds to specific receptors on the surface of T cells, B cells and monocytes, causes the activation and proliferation of T cells and enhances native killer cell activity; moreover, it is an inflammatory marker that helps to determine the occurrence of inflammation (Brandhorst et al. 2015; Nelson 2004). In addition, sIgA can interfere with early pathogen infections by blocking intestinal epithelial cell receptors (Mantis et al. 2006). sIgA acts as a competitive inhibitor to interrupt the binding of pathogens to the intestinal epithelium (Mestecky and Russell 2009). Therefore, the level of sIgA in the gut is essential for assessing gut mucosal immunity (Lin et al. 2014). After 14 days treatment of GQ-Zn, the decrease of IL-2 and the increase of sIgA reflected a reduction in intestinal inflammation.

Oxidative stress is a normal phenomenon and has dual effects. Free radicals can prevent microbial invasion, but they can also cause cell damage during inflammation (Liu et al. 2019; Rahal et al. 2014). Under normal circumstances, free radicals in the body are maintained at a low level; however, the concentration of free radicals abnormally increases when diseases occur. Oxidative stress has been observed in many animal infectious diseases (Lykkesfeldt and Svendsen 2007). CAT is a ubiquitous enzyme in various organisms (Kirkman and Gaetani 2007),

and it can eliminate excess ROS produced by organisms when exposed to external stimuli (Zamocky et al. 2008). Intestinal epithelial cells are shielded by CAT, hence preventing the onset of intestinal inflammation (Bhattacharyya et al. 2014). MDA is a marker of oxidative stress, and its excessive production has certain genotoxicity (Niedernhofer et al. 2003). Infection with *Escherichia coli* and rotavirus increases the levels of MDA (Aydin et al. 2022). Several studies have shown that feeding calves different concentrations of ZnO results in a linear decrease in the serum MDA concentration (Wei et al. 2019). These findings suggest that inflammation and oxidative stress should be controlled within a certain range during the health of calves. The concentration of MDA decreased and CAT increased after GQ-Zn treatment reflecting a reduction of oxidative stress in calves.

The concentration of SCFAs in the rectal contents also changed. SCFAs are generally produced by gut microbiota and are nutrients for intestinal epithelial cells (Park et al. 2016). SCFAs are one of the mediators through which the gut microbiota can control inflammation (Vinolo et al. 2011). In a previous study, general attention was given to the functions of acetic acid, propionic acid and butyric acid; however, a recent study suggested that valeric acid plays a certain role in immunity (Luu et al. 2019). The decrease in the propionic acid concentration may be due to the antibacterial effect of




Figure 5. Effects of GQ-Zn on rectal contents microbiota in pre-weaning calves. Venn analysis based on OTUs (A). Beta-diversity visualized as principal coordinate analysis (PCoA) plots for group Con, Low and High (B). Effect of GQ-Zn supplementation on bacterial α -diversity indices: ACE (C), Chao1 (D), Shannon (E) and Simpson index (F). Histogram analysis of microbial taxonomic composition at phylum (G) and genus (H) levels. The data are presented as the mean \pm SEM and evaluated using Metastats ($n = 10$).

GQ-Zn, which reduced the relative abundance of some bacteria. It was reported that zinc reduces the concentration of propionate in the intestines of piglets (Starke et al. 2014). In the present study, the reduction in propionate concentration may have been due to changes in intestinal ecology and microbial metabolism.

After birth, various of microbiota colonize the gastrointestinal tract in mammals. This complex microbiota plays an important role in the health of newborns (Yizhao et al. 2024). GQ-Zn supplementation ameliorates diarrhea may through regulate gut microbiota in pre-weaning Holstein calves. In our study, some pathogenic bacteria, such as *Enterococcus faecium*, *Prevotella denticola* and *Fusobacterium necrophorum*, were decreased after treatment. *Enterococcus faecium* is considered as a pathogenic bacteria causing adult cat diarrhea (Hélie and Higgins 1999), but research also found that *Enterococcus faecium* decreased inflammatory responses in goat (Dong et al. 2023). *Prevotella denticola* infection caused necrotizing fasciitis and enhances caries-associated virulence of plaque biofilms in some research (Jeremiah and Takashi 2022; Niu et al. 2023). *Fusobacterium necrophorum* is

considered as pathogenic bacteria causing empyema, mastoiditis and septic shock (Arias Miranda et al. 2005; Farinas Salto et al. 2007; Lechiche et al. 2001).

Microbiota changes can be link to metabolites in pre-weaning calves (Yizhao et al. 2024). We further hypothesized GQ-Zn changed the microbiota, then altered the metabolites leading to the alleviation of calf diarrhea. The levels of biotin, niacinamide, riboflavin, retinol, α -tocopherol, 9-oxoODE, 13S-HODE, etc. were found to increase in after treating with GQ-Zn, indicating that changes in some microbiota led to the alter of metabolites. After retinol supplementation in diarrheal mice, the expression of tight junction proteins was upregulated, indicating that the intestinal mechanical barrier was improved. Retinol can decrease the mortality and incidence rate of infectious gastrointestinal diseases (Thornton et al. 2014). α -tocopherol is a kind of vitamin E. Vitamin E has antioxidant functions and can improve intestinal bleeding and diarrhea (Kennedy et al. 2001). Calves infected with viruses will experience a decrease in vitamin E levels (Nonnecke et al. 2014). Supplement chicks with vitamin E can




Figure 6. Metastats analysis was used to analyze the differences in microbiota abundance in feces in day 14 after treatment ($n = 10$), different species richness between Con and Low group (A), Con and High group (B). The red box indicates the top 5 relative abundances in Con and Low group (A) or Con and High group (B).

protect them from *E. coli* infection (Heinzerling et al. 1974). 9-OxoODE is a metabolite of conjugated linoleic acid that can inhibit the secretion of inflammatory factors by macrophages through

PPAR γ (Wysoczanski et al. 2019). We hypothesized that GQ-Zn alleviates calf diarrhea by altering the structure of microbiota and thereby altering metabolites. The impact of metabolites on diarrhea




Figure 7. Metabolomics analysis of rectal contents. OPLS-DA score plot of metabolites (A). Statistic histograms of differently expressed metabolite (B). Metabolite sets enrichment overview (C). The enrichment analysis of KEGG pathways by differential metabolites of rectal contents (D). Metabolites that formed significant different metabolic pathways (E–L). The data are presented as the mean \pm SEM and evaluated using one-way ANOVA; $n = 10$. * $P < 0.05$, ** $P < 0.01$.

and the validation of selected metabolites should be further studied.

Conclusion

In conclusion, supplementation with GQ-Zn alleviated diarrhea in calves, improved fecal scores and reduced oxidative stress,

intestinal inflammation and the relative abundance of microbiome in pre-weaning Holstein calves. GQ-Zn exhibited beneficial effects on the health of calves.

Author contributions. Linhai Yang, Zihai Wei, Yanjing Su, and Xianghuang Wang have contributed equally to this work.

Financial support. This work was funded by the grants from the Shanghai Agriculture Applied Technology Development Program (No. T20220201), National Key R&D Program of China (2022YFD1300705 and 2022YFD1301004) and National Natural Science Foundation of China (31601962).

Conflicts of interest. The authors declare that they have no conflicts of interest.

Ethical standards. The animal study was approved by Animal Welfare and Ethics Committee of Huazhong Agricultural University.

References

- Abreu R, Essler L, Giri P *et al.* (2020) Interferon-gamma promotes iron export in human macrophages to limit intracellular bacterial replication. *PLoS One* **15**(12), e0240949.
- Allahverdiyev AM, Abamor ES, Bagirova M *et al.* (2011) Antimicrobial effects of TiO₂ and Ag₂O nanoparticles against drug-resistant bacteria and leishmania parasites. *Future Microbiology* **6**(8), 933–940.
- Arias Miranda IM, Nuno Mateo FJ, Fonseca Aizpuru EM *et al.* (2005) Empyema caused by *Fusobacterium necrophorum*. *Anales de Medicina Interna (Madrid, Spain)* **22**(7), 352–353.
- Askenasy N (2015) Interferon and tumor necrosis factor as humoral mechanisms coupling hematopoietic activity to inflammation and injury. *Blood Reviews* **29**(1), 11–15.
- Aydin O, Ulas N, Genc A *et al.* (2022) Investigation of hemogram, oxidative stress, and some inflammatory marker levels in neonatal calves with *Escherichia coli* and coronavirus diarrhea. *Microbial Pathogenesis* **173**, 105802.
- Bartels CJ, Holzhauer M, Jorritsma R *et al.* (2010) Prevalence, prediction and risk factors of enteropathogens in normal and non-normal faeces of young Dutch dairy calves. *Preventive Veterinary Medicine* **93**(2–3), 162–169.
- Bhattacharyya A, Chattopadhyay R, Mitra S *et al.* (2014) Oxidative stress: An essential factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiological Reviews* **94**(2), 329–354.
- Bolger Anthony M, Marc L and Bjoern U (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**(15), 2114–2120.
- Brandhorst G, Petrova DT, Weigand S *et al.* (2015) Lack of correlation between Treg quantification assays in inflammatory bowel disease patients. *World Journal of Gastroenterology* **21**(11), 3325–3329.
- Case CL and Carlson MS (2002) Effect of feeding organic and inorganic sources of additional zinc on growth performance and zinc balance in nursery pigs. *Journal of Animal Science* **80**(7), 1917–1924.
- Chasapis CT, Ntoupa PSA, Spiliopoulou CA *et al.* (2020) Recent aspects of the effects of zinc on human health. *Archives of Toxicology* **94**(5), 1443–1460.
- Chee-Sanford JC, Mackie RI, Koike S *et al.* (2009) Fate and transport of antibiotic residues and antibiotic resistance genes following land application of manure waste. *Journal of Environmental Quality* **38**(3), 1086–1108.
- Cho YI, Han JI, Wang C *et al.* (2013) Case-control study of microbiological etiology associated with calf diarrhea. *Veterinary Microbiology* **166**(3–4), 375–385.
- Christian Q, Elmar P, Pelin Y *et al.* (2013) The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research* **41**(Database issue), D590–596.
- Dall Agnol AM, Lorenzetti E, Leme RA *et al.* (2021) Severe outbreak of bovine neonatal diarrhea in a dairy calf rearing unit with multifactorial etiology. *Brazilian Journal of Microbiology* **52**(4), 2547–2553.
- Dong J, Jiang Y, Li Z *et al.* (2023) *Enterococcus faecium* supplementation prevents enteritis caused by *Escherichia coli* in goats. *Beneficial Microbes* **14**(5), 477–491.
- Du WJ, Wang XH, Hu MY *et al.* (2023a) Modulating gastrointestinal microbiota to alleviate diarrhea in calves. *Frontiers in Microbiology* **14**, 1181545.
- Du YF, Gao Y, Hu M Y *et al.* (2023b) Colonization and development of the gut microbiome in calves. *Journal of Animal Science and Biotechnology* **14**(1), 46.
- Edgar Robert C (2013) UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* **10**(10), 996–+.
- Evan B, Jai Ram R, Dillon Matthew R *et al.* (2019) Reproducible, interactive, scalable and extensible microbiome data science using qiime 2 (vol 37, pg 852, 2019). *Nature Biotechnology* **37**(9), 1091–1091.
- Farinas Salto M, Santos Sebastian MM, Gutierrez Triguero M *et al.* (2007) Mastoiditis due to *Fusobacterium necrophorum*. *Anales de Pediatría (Barcelona, Spain)* **66**(2), 193–194.
- Gagnebin Y, Tonoli D, Lescuyer P *et al.* (2017) Metabolomic analysis of urine samples by UHPLC-QTOF-MS: Impact of normalization strategies. *Analytica Chimica Acta* **955**(27), 27–35.
- Gul S and Alhidary IA (2024) Effect of inclusion of zinc-glycine chelate and zinc sulphate on live performance, immunity and lipid peroxidation in broilers. *Journal of Applied Animal Research* **52**(1), 2317889.
- Heinrichs AJ, Jones CM and Heinrichs BS (2003) Effects of mannan oligosaccharide or antibiotics in neonatal diets on health and growth of dairy calves¹. *Journal of Dairy Science* **86**(12), 4064–4069.
- Heinzerling RH, Nockels CF, Quarles CL *et al.* (1974) Protection of chicks against *Escherichia coli* infection by dietary supplementation with vitamin-E. *Proceedings of the Society for Experimental Biology and Medicine* **146**(1), 279–283.
- Hélie P and Higgins R (1999) Diarrhea associated with *Enterococcus faecium* in an adult cat. *Journal of Veterinary Diagnostic Investigation* **11**(5), 457–458.
- Hill GM and Shannon MC (2019) Copper and zinc nutritional issues for agricultural animal production. *Biological Trace Element Research* **188**(1), 148–159.
- Hodges K and Gill R (2010) Infectious diarrhea cellular and molecular mechanisms. *Gut Microbes* **1**(1), 4–21.
- Jenkins KJ and Hidirolou M (1991) Tolerance of the preruminant calf for excess manganese or zinc in milk replacer¹. *Journal of Dairy Science* **74**(3), 1047–1053.
- Jeremiah L and Takashi H (2022) Necrotizing fasciitis due to *Prevotella denticola* infection in an intravenous drug user. *Cureus* **14**(1), e20901–e20901.
- Jiang ZY, Sun LH, Lin YC *et al.* (2009) Effects of dietary glycyl-glutamine on growth performance, small intestinal integrity, and immune responses of weaning piglets challenged with lipopolysaccharide. *Journal of Animal Science* **87**(12), 4050–4056.
- Kechin A, Boyarskikh U, Kel A *et al.* (2017) cutPrimers: A new tool for accurate cutting of primers from reads of targeted next generation sequencing. *Journal of Computational Biology* **24**(11), 1138–1143.
- Kennedy M, Bruninga K, Mutlu EA *et al.* (2001) Successful and sustained treatment of chronic radiation proctitis with antioxidant vitamins E and C. *American Journal of Gastroenterology* **96**(4), 1080–1084.
- Kim HS, Whon TW, Sung H *et al.* (2021) Longitudinal evaluation of fecal microbiota transplantation for ameliorating calf diarrhea and improving growth performance. *Nature Communications* **12**(1), 161.
- Kirkman HN and Gaetani GF (2007) Mammalian catalase: A venerable enzyme with new mysteries. *Trends in Biochemical Sciences* **32**(1), 44–50.
- Klein-Jobstl D, Iwersen M and Drillich M (2014) Farm characteristics and calf management practices on dairy farms with and without diarrhea: A case-control study to investigate risk factors for calf diarrhea. *Journal of Dairy Science* **97**(8), 5110–5119.
- Lechiche C, Corne P, Rivière S *et al.* (2001) *Fusobacterium necrophorum* septic shock from colonic tract with multiple abscesses. *Revue De Medecine Interne* **22**(2), 198–199.
- Lees JR (2015) Interferon gamma in autoimmunity: A complicated player on a complex stage. *Cytokine* **74**(1), 18–26.
- Lin M, L. D, Brandtzaeg P *et al.* (2014) IgA subclass switch recombination in human mucosal and systemic immune compartments. *Mucosal Immunology* **7**(3), 511–520.
- Liu JL, Wang YH, Ma JZ *et al.* (2019) A review on bidirectional analogies between the photocatalysis and antibacterial properties of ZnO. *Journal of Alloys and Compounds* **783**, 898–918.
- Liu WR, Zeng D, She L *et al.* (2020) Comparisons of pollution characteristics, emission situations, and mass loads for heavy metals in the manures of different livestock and poultry in China. *Science of the Total Environment* **734**, 139023.

- Luu M, Pautz S, Kohl V *et al.* (2019) The short-chain fatty acid pentanoate suppresses autoimmunity by modulating the metabolic-epigenetic crosstalk in lymphocytes. *Nature Communications* **10**(1), 760.
- Lykkesfeldt J and Svendsen O (2007) Oxidants and antioxidants in disease: Oxidative stress in farm animals. *Veterinary Journal* **173**(3), 502–511.
- Magalhaes VJA, Susca F, Lima FS *et al.* (2008) Effect of feeding yeast culture on performance, health, and immunocompetence of dairy calves. *Journal of Dairy Science* **91**(4), 1497–1509.
- Mantis NJ, Mcguinness CR, Sonuyi O *et al.* (2006) Immunoglobulin A antibodies against ricin A and B subunits protect epithelial cells from ricin intoxication. *Infection and Immunity* **74**(6), 3455–3462.
- Maywald M and Rink L (2017) Zinc supplementation induces CD4(+)CD25(+)Foxp3(+) antigen-specific regulatory T cells and suppresses IFN-gamma production by upregulation of Foxp3 and KLF-10 and downregulation of IRF-1. *European Journal of Nutrition* **56**(5), 1859–1869.
- Mestecky J and Russell MW (2009) Specific antibody activity, glycan heterogeneity and polyreactivity contribute to the protective activity of S-IgA at mucosal surfaces. *Immunology Letters* **124**(2), 57–62.
- Mobarki N, Almerabi B and Hattan A (2019) Antibiotic resistance crisis. *International Journal of Medicine In Developing Countries* **3**(6), 561–564.
- Murray CF, Fick LJ, Pajor EA *et al.* (2016) Calf management practices and associations with herd-level morbidity and mortality on beef cow-calf operations. *Animal* **10**(3), 468–477.
- Nelson BH (2004) IL-2, regulatory T cells, and tolerance. *Journal of Immunology* **172**(7), 3983–3988.
- Niedernhofer LJ, Daniels JS, Rouzer CA *et al.* (2003) Malondialdehyde, a product of lipid peroxidation, is mutagenic in human cells. *Journal of Biological Chemistry* **278**(33), 31426–31433.
- Niu YF, Zhang CY, Sun YF *et al.* (2023) Symbiotic relationship between *Prevotella denticola* and *Streptococcus mutans* enhances virulence of plaque biofilms. *Archives of Oral Biology* **151**, 105714.
- Nonnecke BJ, McGill JL, Ridpath JF *et al.* (2014) Acute phase response elicited by experimental bovine diarrhea virus (BVDV) infection is associated with decreased vitamin D and E status of vitamin-replete preruminant calves. *Journal of Dairy Science* **97**(9), 5566–5579.
- Ogiyama S, Sakamoto K, Suzuki H *et al.* (2005) Accumulation of zinc and copper in an arable field after animal manure application. *Soil Science and Plant Nutrition* **51**(6), 801–808.
- Okada S, Inabu Y, Miyamoto H *et al.* (2023) Estimation of silent phenotypes of calf antibiotic dysbiosis. *Scientific Reports* **13**(1), 6359.
- Park JH, Kotani T, Konno T *et al.* (2016) Promotion of intestinal epithelial cell turnover by commensal bacteria: Role of short-chain fatty acids. *PLoS One* **11**(5), e0156334.
- Rahal A, Kumar A, Singh V *et al.* (2014) Oxidative stress, prooxidants, and antioxidants: The interplay. *Biomed Research International* **2014**, 761264.
- Reuter S, Gupta SC, Chaturvedi MM *et al.* (2010) Oxidative stress, inflammation, and cancer: How are they linked? *Free Radical Biology and Medicine* **49**(11), 1603–1616.
- Schell TC and Kornegay ET (1996) Zinc concentration in tissues and performance of weanling pigs fed pharmacological levels of zinc from ZnO, Zn-methionine, Zn-lysine, or ZnSO₄. *Journal of Animal Science* **74**(7), 1584–1593.
- Schwaiger K, Storch J, Bauer C *et al.* (2023) *Lactobacillus* (*Limosilactobacillus*) *reuteri*: A probiotic candidate to reduce neonatal diarrhea in calves. *Frontiers in Microbiology* **14**, 1266905.
- Shin NR, Whon TW and Bae JW (2015) Proteobacteria: Microbial signature of dysbiosis in gut microbiota. *Trends in Biotechnology* **33**(9), 496–503.
- Starke IC, Pieper R, Neumann K *et al.* (2014) The impact of high dietary zinc oxide on the development of the intestinal microbiota in weaned piglets. *FEMS Microbiology Ecology* **87**(2), 416–427.
- Thornton KA, Mora-Plazas M, Marin C *et al.* (2014) Vitamin A deficiency is associated with gastrointestinal and respiratory morbidity in school-age children. *Journal of Nutrition* **144**(4), 496–503.
- Vinolo MAR, Rodrigues HG, Nachbar RT *et al.* (2011) Regulation of inflammation by short chain fatty acids. *Nutrients* **3**(10), 858–876.
- Wei JY, Ma FT, Hao LY *et al.* (2019) Effect of differing amounts of zinc oxide supplementation on the antioxidant status and zinc metabolism in newborn dairy calves. *Livestock Science* **230**, 103819.
- Wright CL and Spears JW (2004) Effect of zinc source and dietary level on zinc metabolism in Holstein calves. *Journal of Dairy Science* **87**(4), 1085–1091.
- Wysoczanski R, Kendall AC, Motwani M *et al.* (2019) Ulcerative colitis is characterized by amplified acute inflammation with delayed resolution. *bioRxiv*.
- Yan Y, B. X, Yin B *et al.* (2019) Modulation of gut microbial community and metabolism by dietary glycyl-glutamine supplementation may favor weaning transition in piglets. *Frontiers in Microbiology* **10**, 3125.
- Yizhao S, Yan L, Tingting W *et al.* (2024) Early microbial intervention reshapes phenotypes of newborn *Bos taurus* through metabolic regulations. *GigaScience* **13**, giad118.
- Zamocky M, Furtmuller PG and Obinger C (2008) Evolution of catalases from bacteria to humans. *Antioxidants & Redox Signaling* **10**(9), 1527–1548.
- Zhu C, H. LV, Chen Z *et al.* (2017) Dietary zinc oxide modulates antioxidant capacity, small intestine development, and jejunal gene expression in weaned piglets. *Biological Trace Element Research* **175**(2), 331–338.