

Research Article

\*Current address: Department of Entomology, Texas A&M University, College Station, TX 77843, USA.

A similar manuscript has been uploaded to biorxiv – <https://www.biorxiv.org/content/10.1101/2023.06.26.546593v1>



**Cite this article:** Herb H, González J, Ferreira FC, Fonseca DM (2023). Multiple piroplasm parasites (Apicomplexa: Piroplasmida) in northeastern populations of the invasive Asian longhorned tick, *Haemaphysalis longicornis* Neumann (Ixodida: Ixodidae), in the United States. *Parasitology* **150**, 1063–1069. <https://doi.org/10.1017/S0031182023000914>

Received: 29 July 2023  
 Revised: 20 September 2023  
 Accepted: 21 September 2023  
 First published online: 4 October 2023

**Keywords:**  
 agriculture; invasive spillover; livestock; One Health; public health; wildlife

**Corresponding author:**  
 Dina M. Fonseca;  
 Email: [dina.fonseca@rutgers.edu](mailto:dina.fonseca@rutgers.edu)

# Multiple piroplasm parasites (Apicomplexa: Piroplasmida) in northeastern populations of the invasive Asian longhorned tick, *Haemaphysalis longicornis* Neumann (Ixodida: Ixodidae), in the United States

Heidi Herb<sup>1,2</sup>, Julia González<sup>1,\*</sup>, Francisco C. Ferreira<sup>1,\*</sup>   
 and Dina M. Fonseca<sup>1,2,3</sup> 

<sup>1</sup>Center for Vector Biology, Rutgers University, New Brunswick, NJ 08901, USA; <sup>2</sup>Department of Ecology and Evolution, Rutgers University, New Brunswick, NJ 08901, USA and <sup>3</sup>Department of Entomology, Rutgers University, New Brunswick, NJ 08901, USA

## Abstract

Piroplasms, which include the agents of cattle fever and human and dog babesiosis, are a diverse group of blood parasites of significant veterinary and medical importance. The invasive Asian longhorned tick, *Haemaphysalis longicornis*, is a known vector of piroplasms in its native range in East Asia and invasive range in Australasia. In the USA, *H. longicornis* has been associated with *Theileria orientalis* Ikeda outbreaks that caused cattle mortality. To survey invasive populations of *H. longicornis* for a broad range of piroplasms, 667 questing *H. longicornis* collected in 2021 from 3 sites in New Jersey, USA, were tested with generalist piroplasm primers targeting the 18S small subunit rRNA (395–515 bp, depending on species) and the cytochrome b oxidase loci (1009 bp). Sequences matching *Theileria cervi* type F (1 adult, 5 nymphs), an unidentified *Theileria* species (in 1 nymph), an undescribed *Babesia* sensu stricto (‘true’ *Babesia*, 2 adults, 2 nymphs), a *Babesia* sp. Coco (also a ‘true *Babesia*’, 1 adult, 1 nymph), as well as *Babesia microti* S837 (1 adult, 4 nymphs) were recovered. *Babesia microti* S837 is closely related to the human pathogen *B. microti* US-type. Additionally, a 132 bp sequence matching the cytochrome b locus of deer, *Odocoileus virginianus*, was obtained from 2 partially engorged *H. longicornis*. The diverse assemblage of piroplasms now associated with *H. longicornis* in the USA spans 3 clades in the piroplasm phylogeny and raises concerns of transmission amplification of veterinary pathogens as well as spillover of pathogens from wildlife to humans.

## Introduction

Invasive ticks, mosquitoes and other blood-feeding arthropods may introduce and transmit (i.e. vector) exotic pathogens for which local populations have little or no immunity. Resulting disease can range from mild to severe to fatal and can have a significant impact on human health (e.g. Zika fever, West Nile virus encephalitis), animal health (e.g. redwater fever, blue tongue virus), hasten extinction (e.g. bird malaria) and cause economic damage to agriculture, tourism and other industries (Athni *et al.*, 2021). Invasive vectors can also potentially spread existing wildlife pathogens by creating new transmission pathways, which can have significant ecological and public health implications, particularly in the context of One Health (*sensu* Lerner and Berg, 2015).

The phylum Apicomplexa includes well-known blood-borne protozoa such as *Plasmodium falciparum* and *P. vivax*, the primary agents of human malaria (Votýpka *et al.*, 2016). The Apicomplexa class Piroplasmida includes *Babesia*, *Theileria* and *Cytauxzoon* that are primarily transmitted by hard ticks (Ixodida: Ixodidae) and can affect a wide range of hosts (Onyiche *et al.*, 2021; Almazán *et al.*, 2022). While piroplasms were once classified based on morphology and host associations alone, the advent of molecular methods has greatly advanced the overall understanding of the diversity of Piroplasmida (Garrett *et al.*, 2019). A recent analysis by Jalovecka *et al.* (2019) indicates that there are at least 10 distinct clades within Piroplasmida, with both *Babesia* spp. and *Theileria* spp. comprising polyphyletic groups in need of taxonomic revision.

*Babesia* spp. are broadly divided into *Babesia* sensu stricto and *Babesia* sensu lato, with the former representing a monophyletic group considered as ‘true *Babesia*’, distinguishable from other piroplasms by their ability to infect the reproductive organs of the tick and to be transmitted to the eggs (transovarial transmission) (Schreeg *et al.*, 2016; Jalovecka *et al.*, 2019; Schnittger *et al.*, 2022). Within *Babesia* sensu lato, one of the best-characterized clades is the *Babesia microti* group, which contains the piroplasms responsible for most human babesiosis cases worldwide, especially in the northern United States (Renard and Mamoun, 2021). Importantly, *B. microti* includes at least 2 different genetic lineages pathogenic to humans

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

and several only known from reservoir hosts such as mice, voles and skunks (Goethert, 2021). Moreover, although still relatively rare, human disease caused by other *Babesia* species such as *B. divergens* in Europe, *B. venatorum* in Asia and Europe and *B. duncani* in North America has been increasingly reported (Scott and Scott, 2018; Hong *et al.*, 2019; Kumar *et al.*, 2021; Scott *et al.*, 2021).

Rates of human babesiosis have been increasing in the USA, particularly in the northeastern states (Almazán *et al.*, 2022; Swanson *et al.*, 2023). While those infected with *Babesia* may experience fever, chills, headache, muscle aches, fatigue and red or brown urine, some may not have any symptoms at all, especially if their immune systems are not compromised (Almazán *et al.*, 2022). People may remain infected for years, and even if asymptomatic, can transmit piroplasms through blood transfusions or organ transplants (Bloch *et al.*, 2019), and transmission from infected mothers to developing fetuses has been demonstrated (Horowitz and Freeman, 2020). Piroplasmid infections are typically treated with various combinations of atovaquone, azithromycin, clindamycin and quinine; however, concerns regarding side-effects, drug resistance and drug efficacy indicate the need for development of novel treatment options (Renard and Mamoun, 2021).

Bovine babesiosis caused by *Babesia bigemina* and *Babesia bovis* has long been a concern to American cattle ranchers and several *Theileria* species can sicken horses, cervids and bovids (Almazán *et al.*, 2022; Osbrink *et al.*, 2022). Both babesiosis and theileriosis cause significant economic losses annually to North American and Australasian agricultural industries due to reduced production, death, abortions, restrictions on animal movement and costs associated with preventive measures and treatments (Dinkel *et al.*, 2021; Almazán *et al.*, 2022; Osbrink *et al.*, 2022; Schnittger *et al.*, 2022). Companion animals are also at risk, with infections of *Babesia vulpes*, *Babesia conradae*, *Babesia vogeli*, *Babesia gibsoni* and *Babesia* sp. Coco capable of causing mild to severe disease in dogs in the United States (Dear and Birkenheuer, 2022).

Since the initial discovery of the invasive Asian longhorned tick (*Haemaphysalis longicornis*) in the United States in 2017 (Rainey *et al.*, 2018), there have been concerns regarding the potential threats this ectoparasite may pose. In North America, as in Australasia where it expanded to in the early 20th century, *H. longicornis* reproduces asexually (clonally) by parthenogenesis (Schappach *et al.*, 2020), which underlies the ability of this species to develop large populations very quickly. In its Australasian range, *H. longicornis* represents a major threat to domestic livestock, heavily parasitizing large ruminants, and impeding production (Heath, 2016).

Globally, *H. longicornis* is a known vector of piroplasms that infect humans, livestock and companion animals, including *B. ovata*, *B. gibsoni*, *B. microti*, *T. uilenbergi* and *T. orientalis* (Li *et al.*, 2009; Wu *et al.*, 2017; Gray *et al.*, 2019; Dinkel *et al.*, 2021; Dear and Birkenheuer, 2022) and may also be a vector of *Babesia caballi*, the agent of equine babesiosis (Bautista *et al.*, 2001). As in Australia (Marendy *et al.*, 2020), in Virginia, USA, *H. longicornis* has already been implicated in the transmission of the virulent *Theileria orientalis* Ikeda genotype that resulted in multiple cattle deaths (Thompson *et al.*, 2020; Dinkel *et al.*, 2021).

Although humans are not favoured hosts of *H. longicornis*, opportunistic feeding is well documented both in the native and invasive ranges (Bickerton and Toledo, 2020; Wormser *et al.*, 2020). In East Asia, *H. longicornis* vectors severe fever with thrombocytopenia syndrome virus (SFTSV), an emerging human tick-borne disease recently reclassified as Dabie bandavirus (Liu *et al.*, 2015; Luo *et al.*, 2015; Li *et al.*, 2021). Under laboratory conditions, US lineages of *H. longicornis* can vector

the closely related Heartland virus (Rainey *et al.*, 2022a) as well as Powassan virus (Rainey *et al.*, 2022b), 2 native pathogenic viruses emergent in parts of the USA. They can also vector *Rickettsia rickettsii*, the causative agent of Rocky Mountain spotted fever (Stanley *et al.*, 2020). In addition, Bourbon virus has been detected from a larval pool, 2 nymphs and 1 adult field collected *H. longicornis* in Virginia, USA (Cumbie *et al.*, 2022). While *H. longicornis* is currently not perceived as a major public health threat in the USA, this status may change given the enormous densities it can reach in favourable habitats (Bickerton and Toledo, 2020; Schappach *et al.*, 2020; González *et al.*, 2023; Rochlin *et al.*, 2023).

The objective of this study was to assess the potential role of *H. longicornis* as a vector of piroplasms in New Jersey (NJ), the most urbanized US state that, maybe surprisingly to many, also boasts the highest density of horses (Rankins and Malinowski, 2020).

## Materials and methods

### Study areas

This study was conducted in 3 sites approximately 1.2 km from each other within the Rutgers University Cook Campus in New Brunswick, NJ (please refer to Ferreira *et al.*, 2023 for a map). Surveys for *H. longicornis* were initiated at these sites in 2018 when the species was first detected on a grassy area next to a goat pen (Egizi *et al.*, 2019b), a site that became known as the 'Goat Farm' (40.47444° N, 74.43683° W). The 'Rutgers Gardens' site (40.47455° N, 74.42030° W) is inside a 180-acre botanical garden, consisting of designed gardens, plant collections and natural habitats. Finally, the 'University Inn' site (40.48413° N, 74.43051° W) is a meadow and forested park behind the Rutgers University Inn & Conference Center. At all sites, local forest is dominated by oak and maple trees and huckleberry and blueberry shrubs (Breden *et al.*, 2001), with grassy ecotones.

### Tick surveillance

From June through September 2021, concomitant with surveys for ticks on mammals at the same sites, questing ticks were sampled from 50–75 m<sup>2</sup> of the vegetation at each site (Ferreira *et al.*, 2023). Tick sampling was performed using a white crib flannel sweep measuring 50 × 100 cm with a PVC pipe handle (Egizi *et al.*, 2019b). The sweep cloth was checked in 1–2 m intervals since *H. longicornis* does not attach firmly to the flannel and often drops off over longer intervals (Bickerton *et al.*, 2021). Ticks were collected from both sides of the sweep and identified morphologically in the laboratory to the species level using a stereomicroscope (Leica S8 APO, Leica Microsystems, Deerfield, IL, USA) following appropriate taxonomical keys (Keirans and Litwak, 1989; Egizi *et al.*, 2019a). The larvae of *H. longicornis* were not stored during these surveys and were not available for pathogen testing. A few questing *H. longicornis* that were found partially engorged (*sensu* Price *et al.*, 2022) were processed separately (see section below on 'Bloodmeal analysis of partially engorged specimens').

### DNA extraction and pathogen detection

Each tick was placed in 180 µl of Qiagen buffer ATL with 20 µl Qiagen Proteinase K (10 mg mL<sup>-1</sup>) in microfuge tubes and homogenized with a 5 mm sterile glass bead (Fisher Scientific, Waltham, MA, USA) in a TissueLyser (Qiagen Inc., Valencia, CA, USA). DNA from individual ticks was extracted using Dneasy Blood and Tissue 96-well plate kits (Qiagen Inc., Valencia, CA, USA) following the manufacturer's instructions.

DNA was eluted from each column twice with 50  $\mu$ l of Qiagen's elution buffer AE into separate labelled microtubes.

After reviewing the literature, primers were chosen targeting the multi copy 18S rRNA locus (Table 1) to match a broad range of piroplasm species (Casati *et al.*, 2006) and all ticks were tested individually. To further characterize an undescribed new *Babesia* sp. detected, specimens positive for that *Babesia* were also tested using primers targeting the mitochondrial cytochrome oxidase b (*cytb*) locus (Table 1) shown to work across multiple *Babesia* species (Rajapakshage *et al.*, 2012).

The targeted loci were amplified in 20  $\mu$ l reactions with Amplitaq Gold Master Mix (ThermoFisher Scientific, Waltham, MA, USA) following the manufacturer's protocol. After visualizing the amplification in a 1% agarose gel, polymerase chain reactions (PCR) were cleaned with ExoSAP-IT (ThermoFisher Scientific) and Sanger sequenced separately with both primers at Azenta Genewiz (South Plainfield, NJ, USA). The sequences were trimmed and aligned with Geneious Prime 2023.0.1 (Biomatters Inc., San Diego, CA, USA) and the consensus was used as a query in NCBI's Basic Local Alignment Search Tool, BLASTn (Altschup *et al.*, 1990).

### Phylogenetic analysis

Consensus sequences were aligned to available NCBI Genbank sequences (Benson *et al.*, 2012) representative of the major piroplasm clades (Jalovecka *et al.*, 2019) and the alignments were trimmed to the same size (557 bp). Maximum likelihood phylogenetic trees based on the 18S rRNA and *cytb* loci were constructed using IQ-TREE with 1000 ultrafast bootstrap replicates (Nguyen *et al.*, 2015; Hoang *et al.*, 2018). ModelFinder was used to choose the best-fitting substitution model based on Bayesian information criterion (Kalyaanamoorthy *et al.*, 2017). *Cardiosporidium cionae* (GenBank accession number EU052685) and *B. microti* (GenBank accession number NC034637) were used as outgroups for the 18S rRNA and *cytb* phylogenetic trees, respectively.

### Statistical analysis

To determine whether there are statistically significant differences in infection rates in nymphal vs female *H. longicornis*, a binomial generalized linear model using the glm function in R (R Core Team, 2022) was used. To determine the correlation between sample size and infection rates, the lm function in R was used.

### Bloodmeal analysis of partially engorged specimens

DNA from 2 partially engorged *H. longicornis* collected on 15 July 2021 (1 nymph and 1 adult) was isolated using DNeasy Blood and Tissue columns (Qiagen, Valencia, CA, USA). An extraction control was included, and all work was performed in a dedicated clean lab inside a laminar flow hood (Mystaire, Creedmor, NC, USA). Primers CytbVertR1 (5'-GGACGAGGACTATACTAC GG-3' from Egizi *et al.*, 2013) and BMF1 (5'-AAACTGC AGCCCCTCAGAATGATATTTGTCCTCA-3'), originally called H15149 (Kocher *et al.*, 1989), were used to amplify a 132-nucleotide fragment in the cytochrome oxidase b locus using an annealing temperature of 55°C. A PCR product obtained from the adult *H. longicornis* was purified (ExoSAP-IT, Affymetrix, Santa Clara, CA, USA) then sequenced at Azenta Genewiz (South Plainfield). The sequences were trimmed and aligned with Geneious Prime 2023.0.1 (Biomatters Inc.) and the consensus was used as a query in NCBI's Basic Local Alignment Search Tool, BLASTn (Altschup *et al.*, 1990).

## Results

Overall, 667 *H. longicornis* nymphs and adults collected from the environment were screened and evidence of piroplasm parasites was found in 18 ticks (2.7%, Table 2). Adult and nymph infection rates did not differ (infection rates of 2.4 and 2.8%, respectively,  $P$  value = 0.60). Infection rates among sampling sites reflected sample size ( $r^2 = 0.92$ ,  $P$  value < 0.01; using adult and nymph data separately to increase the statistical power).

There was no evidence of piroplasm coinfections (such as double chromatogram peaks) in the positive ticks. The primers targeting the 18S rRNA gene amplified fragments ranging in size from 395 to 515 base pairs (bp) spanning the V4 hypervariable region (Cauvin *et al.*, 2019).

The 18S rRNA fragment from a nymph collected at University Inn had a 100% pairwise identity to 5 *Theileria* sp. sequences (GenBank accession numbers MW008536, MW008531, MK262962, MK262963 and MK262959) that are considered 'Type X' or 'divergent' (Cauvin *et al.*, 2019; Olafson *et al.*, 2020). This sequence differed by 20 bp (17 mismatches and 3 deletions) from 6 other sequences obtained from *H. longicornis* also collected from the University Inn site. The closest match for these 6 sequences (99.8–100% pairwise identity) was a *Theileria cervi* type F sequence (GenBank U97054). Three of the 6 sequences, 2 from nymphs and 1 from an adult, were identical to U97054 while the remaining 3, all from nymphs, differed by 1 bp (GenBank accession numbers OR612075, OR612078, OR612080). All sequences clustered within a clade containing *T. cervi* and *T. orientalis* (Fig. 1a).

Two 18S rRNA sequences, one from a Goat Farm adult and the other from a University Inn nymph, were identical to a *Babesia* sp. isolate NYT-435 (GenBank accession number MW665118) recently sequenced from a pool of lone star ticks (*Amblyomma americanum*) from Staten Island, NYC (Jain *et al.*, 2021). The sequence of the *Babesia* sp. isolate NYT-435 was also the closest match for 2 other sequences, 1 from another Goat Farm adult and a second University Inn nymph (99.7 and 99.5% pairwise identities with a 1 and 2 bp difference, respectively; GenBank accession numbers OR612068, OR612072, respectively). The next best match for these 4 sequences (97.9–98.2% pairwise identity) was a *Babesia* sp. from a white-tailed deer in Texas (GenBank accession number HQ264120). These 4 sequences clustered within the *Babesia* sensu stricto clade with *B. bovis* and *Babesia ovis* (Fig. 1a). From one of the specimens that were positive for the unknown *Babesia*, a 1009 bp DNA fragment was amplified with the *cytb* primers (GenBank accession number OR610156). The closest match to this sequence was *Babesia motasi* isolate Lintan (GenBank acc. num. MN605889) with 90.8% pairwise identity. The sequence from this *H. longicornis* clustered with *B. motasi* in the *Babesia* sensu stricto clade (Fig. 1b).

The 18S rRNA from 2 sequences, one from a University Inn nymph and another from a Goat Farm adult, matched *Babesia* sp. Coco. The sequence from the Goat Farm adult was 100% pairwise identical to GenBank EU109716, whereas the sequence from the University Inn nymph differed by 1 bp and had a 99.8% pairwise identity (GenBank accession number OR612082).

Finally, 5 18S rRNA sequences, from 2 nymphs and 1 adult from the Goat Farm and 2 nymphs from the University Inn, had a 100% pairwise identity to *B. microti* isolate S837 (GenBank accession number AY144698). In the phylogenetic tree, these sequences clustered with *B. vulpes* within the *B. microti*-like group (Fig. 1a; Jalovecka *et al.*, 2019).

A 132 bp DNA fragment was amplified and sequenced from a partially engorged adult *H. longicornis* and was 100% identical to a *cytb* fragment from white-tailed deer, *Odocoileus virginianus* (GenBank accession number AF535863).



**Table 1.** Primers used for piroplasm detection in *H. longicornis*

Target	Locus	Primer	Primer sequence (5' to 3')	T <sub>a</sub>
<i>Piroplasmida</i> spp.	18S rRNA	BJ1	GTCTTGAATTGGAATGATGG	55°C
		BN2	TAGTTTATGGTTAGGACTACG	
<i>Babesia</i> spp.	mtDNA cytochrome b	cytbF1	ATGTTGTCCTATTGGTTCC	*54–50°C
		cytbR1	ATATGCAAACCTCCCGGCTA	

T<sub>a</sub> = annealing temperature. The expected 18S rRNA amplicon size varies depending on the species of piroplasm from 395 to 515 bp; the expected cytochrome b amplicon size is 1009 bp. \*Instead of a specific T<sub>a</sub>, a ‘touch-down’ approach was used starting with T<sub>a</sub> = 54°C and decreasing the T<sub>a</sub> by 1 degree for 4 additional cycles. This was followed by 40 additional cycles at T<sub>a</sub> = 52°C.

## Discussion

DNA sequences belonging to 3 piroplasm clades were detected in 2.7% of field-collected *H. longicornis* nymphs and adults in NJ, United States. Infection rates for *Theileria*, *Babesia* sensu stricto and *B. microti*, were 1, 0.9 and 0.8%, respectively. While these may be low piroplasm infection rates compared to studies that detected *T. orientalis* in 12.7% of *H. longicornis* in Virginia (Thompson et al., 2020, 2022), those were collected from the cattle farm where the first US outbreak of *T. orientalis* Ikeda occurred, which would have increased the likelihood that local vectors were infected. Overall, there have been few exploratory examinations of piroplasms in *H. longicornis* in the United States.

This is the first report of *Theileria* species besides *T. orientalis* Ikeda in *H. longicornis* in the United States. Most sequences matched *T. cervi* type F, a piroplasm that commonly infects deer and other cervids (Cauvin et al., 2019; Olafson et al., 2020). However, 1 sequence differed in at least 20 bp from *T. cervi* type F and instead matched a strain denoted ‘divergent’ or ‘type X’ found in wild and farmed deer in Florida and in *Anocenter nitens*, the tropical horse tick, parasitizing white-tailed deer in Texas (Cauvin et al., 2019; Olafson et al., 2020). US populations of *H. longicornis* have often been reported feeding on white-tailed deer (Tufts et al., 2021), and finding deer DNA in a partially engorged tick supports this. These findings indicate that *H. longicornis* may be involved in the transmission cycle of *Theileria* in NJ.

This is also the first report of *Babesia* in questing un-engorged *H. longicornis* in the United States. Specifically, the phylogenetic analyses using both 18S and cytochrome b loci indicate the unknown *Babesia* sequences fall within the *Babesia* sensu stricto clade. *Babesia* sensu stricto (also referred to as ‘true *Babesia*’) are pathogens of both veterinary and medical importance capable of transovarial transmission in their tick vectors, allowing the parasite to propagate in the absence of vertebrate reservoirs (Jalovecka et al., 2019).

*Babesia* sp. Coco, another ‘true *Babesia*’ was also detected in 2 *H. longicornis*. *Babesia* sp. Coco can be pathogenic to dogs but usually only if they are immunocompromised (Birkenheuer et al., 2004; Holman et al., 2009; Sikorski et al., 2010; Dear and Birkenheuer, 2022), so it is unclear whether dogs are incidental hosts or are an important reservoir species that manifest clinical disease when immunocompromised. Dogs are considered an important blood host for *H. longicornis* in the United States (Trout Fryxell et al., 2021; Thompson et al., 2022).

Finally, this is the first report of *B. microti* genotype S837 in *H. longicornis*. This genotype is found in skunks *Mephitis mephitis* (Goethert, 2021), which are an important host for this tick species in NJ (Ferreira et al., 2023) and, critically, *H. longicornis* has been shown to be a competent vector of *B. microti* under experimental conditions (Wu et al., 2017). Of note, there is an overall lack of knowledge of the biology and epidemiology of wildlife piroplasms in the northeastern USA. In areas endemic for human babesiosis, molecular studies of wildlife piroplasms rarely employ sequencing

**Table 2.** Numbers of *H. longicornis* nymphs and adults collected from the environment at 3 sites at Rutgers Cook campus and tested for piroplasm DNA (# Pos represents the number of ticks that were positive). Collections were started on 24 June 2021 and proceeded approximately bi-weekly until 10 September 2021 (refer to Ferreira et al., 2023 for details).

	June		July		August		September		Total	
	N	# Pos	N	# Pos	N	# Pos	N	# Pos	N	# Pos
<b>Goat Farm</b>	<b>35</b>	<b>2</b>	<b>100</b>	<b>4</b>	<b>14</b>	<b>0</b>	<b>4</b>	<b>0</b>	<b>153</b>	<b>6</b>
Nymph	20	2 <sup>1</sup>	38	0	7	0	1	0	66	2
Adult	15	0	62	4 <sup>123</sup>	7	0	3	0	87	4
<b>Rutgers Gardens</b>	<b>26</b>	<b>0</b>	<b>30</b>	<b>0</b>	<b>8</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>64</b>	<b>0</b>
Nymph	22	0	16	0	4	0	0	0	42	0
Adult	4	0	14	0	4	0	0	0	22	0
<b>University Inn</b>	<b>78</b>	<b>1</b>	<b>308</b>	<b>10</b>	<b>56</b>	<b>1</b>	<b>8</b>	<b>0</b>	<b>450</b>	<b>12</b>
Nymph	64	1 <sup>3</sup>	250	9 <sup>12345</sup>	33	1 <sup>1</sup>	7	0	354	11
Adult	14	0	58	1 <sup>4</sup>	23	0	1	0	96	1
<b>Grand Total</b>	<b>139</b>	<b>3</b>	<b>438</b>	<b>14</b>	<b>78</b>	<b>1</b>	<b>12</b>	<b>0</b>	<b>667</b>	<b>18</b>


<sup>1</sup>*Babesia microti* S837 (5 positive ticks)

<sup>2</sup>*Babesia* sp. Coco (2 positive ticks)

<sup>3</sup>*Babesia* sp. (4 positive ticks)

<sup>4</sup>*Theileria cervi* (a total of 6 positive ticks)

<sup>5</sup>*Theileria* sp. (1 positive ticks)



**Figure 1.** Phylogenetic trees with piroplasm sequences obtained from *Haemaphysalis longicornis* (in red). (a) Tree based on 18S rRNA locus. Constructed with TIM2 + F + I + G4 substitution model. (b) Tree based on cytochrome b locus. Constructed with K3Pu + F + I + G4 substitution model.

to confirm parasite identity, although piroplasm parasites thought to only infect wildlife and/or domestic animals have recently been reported infecting humans in North America (Scott *et al.*, 2021) and elsewhere (Hong *et al.*, 2019).

Although the health risk of *H. longicornis* to US livestock was established by the outbreak of *T. orientalis* Ikeda in cattle in the

state of Virginia (Thompson *et al.*, 2020; Dinkel *et al.*, 2021), there is limited research regarding the veterinary and medical significance of *H. longicornis* in the USA, which has focused primarily on testing the ability of US specimens to transmit pathogens of known public health concern (Breuner *et al.*, 2020; Stanley *et al.*, 2020; Raney *et al.*, 2022a). The discovery of various piroplasm

parasites in questing *H. longicornis* specimens in NJ brings forth new concerns. As a result, it becomes crucial to prioritize studies that delve into the role of *H. longicornis* as an actual vector for these pathogens.

**Data availability statement.** Nucleotide sequence data reported in this paper are available in GenBank™, EMBL and DDBJ databases under the accession numbers OR610156 and OR612065-OR612082.

**Acknowledgements.** We thank Nicole Wagner MS from the Rutgers Center for Vector Biology and Julia Brennan, now developing her PhD at Virginia Tech, for their technical support. We also thank Dr Glen Scoles, USDA-ARS Beltsville, for sharing his piroplasm expertise.

**Author contributions.** Heidi E. Herb: conceptualization, methodology, formal analysis, investigation, data curation, writing – original draft, writing – review and editing, visualization. Julia González and Francisco C. Ferreira: investigation, data curation, writing – review & editing. Dina M. Fonseca: conceptualization, methodology, resources, writing – review & editing, supervision, project administration, funding acquisition.

**Financial support.** This work was supported in part by a United States Department of Agriculture National Institute of Food and Agriculture Multistate Grant, NE1943, to D. M. F., a State of New Jersey FY22 Tick Research and Control-Special Purpose Funding to CVB/NJAES and by Cooperative Agreement Number IU01CK000509 between the Centers for Disease Control and Prevention (CDC) and the Northeast Center of Excellence in Vector Borne Diseases (PI at Rutgers, D. M. F.). The contents of this work are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.

**Competing interests.** None.

**Ethical standards.** Not applicable.

## References

- Almazán C, Scimeca RC, Reichard MV and Mosqueda J (2022) Babesiosis and theileriosis in North America. *Pathogens* **11**, 168.
- Altschup SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Athni TS, Shocket MS, Couper LI, Nova N, Caldwell IR, Caldwell JM, Childress JN, Childs ML, De Leo GA, Kirk DG, MacDonald AJ, Olivarius K, Pickel PG, Roberts SO, Winokur OC, Young HS, Cheng J, Grant EA, Kurzner DM, Kyaw S, Lin BJ, Lopez RC, Massihpour DS, Olsen EC, Roache M, Ruiz A, Schultz EA, Shafat M, Spencer RL, Bharti N and Mordecai EA (2021) The influence of vector-borne disease on human history: socio-ecological mechanisms. *Ecology Letters* **24**, 829–846.
- Bautista JLR, Ikadai H, You M, Battsetseg B, Igarashi I, Nagasawa H and Fujisaki K (2001) Molecular evidence of *Babesia caballi* (Nuttall and Strickland, 1910) parasite transmission from experimentally-infected SCID mice to the ixodid tick, *Haemaphysalis longicornis* (Neuman, 1901). *Veterinary Parasitology* **102**, 185–191.
- Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J and Sayers EW (2012) GenBank. *Nucleic Acids Research* **41**, D36–D42.
- Bickerton M and Toledo A (2020) Multiple pruritic tick bites by Asian longhorned tick larvae (*Haemaphysalis longicornis*). *International Journal of Acarology* **46**, 373–376.
- Bickerton M, McSorley K and Toledo A (2021) A life stage-targeted acaricide application approach for the control of *Haemaphysalis longicornis*. *Ticks and Tick-borne Diseases* **12**, 101581.
- Birkenheuer AJ, Neel J, Ruslander D, Levy MG and Breitschwerdt EB (2004) Detection and molecular characterization of a novel large *Babesia* species in a dog. *Veterinary Parasitology* **124**, 151–160.
- Bloch EM, Kumar S and Krause PJ (2019) Persistence of *Babesia microti* infection in humans. *Pathogens* **8**, 102.
- Breden TF, Alger Y, Strakosch Walz K and Windisch A (2001) Classification of vegetation communities of New Jersey: second iteration. Office of Natural Lands Management, Division of Parks and Forestry, NJDEP.
- Breuner NE, Ford SL, Hojgaard A, Osikowicz LM, Parise CM, Rosales Rizzo MF, Bai Y, Levin ML, Eisen RJ and Eisen L (2020) Failure of the Asian longhorned tick, *Haemaphysalis longicornis*, to serve as an experimental vector of the Lyme disease spirochete, *Borrelia burgdorferi* sensu stricto. *Ticks and Tick-borne Diseases* **11**, 101311.
- Casati S, Sager H, Gern L and Piffaretti J-C (2006) Presence of potentially pathogenic *Babesia* sp. for human in *Ixodes ricinus* in Switzerland. *Annals of Agricultural and Environmental Medicine: AAEM* **13**, 65–70.
- Cauvin A, Hood K, Shuman R, Orange J, Blackburn JK, Saylor KA and Wisely SM (2019) The impact of vector control on the prevalence of *Theileria cervi* in farmed Florida white-tailed deer, *Odocoileus virginianus*. *Parasites & Vectors* **12**, 100.
- Cumby AN, Trimble RN and Eastwood G (2022) Pathogen spillover to an invasive tick species: first detection of bourbon virus in *Haemaphysalis longicornis* in the United States. *Pathogens* **11**, 454.
- Dear JD and Birkenheuer A (2022) *Babesia* in North America. *Veterinary Clinics of North America: Small Animal Practice* **52**, 1193–1209.
- Dinkel KD, Herndon DR, Noh SM, Lahmers KK, Todd SM, Ueti MW, Scoles GA, Mason KL and Fry LM (2021) A U.S. isolate of *Theileria orientalis*, Ikeda genotype, is transmitted to cattle by the invasive Asian longhorned tick, *Haemaphysalis longicornis*. *Parasites & Vectors* **14**, 157.
- Egizi A, Healy SP and Fonseca DM (2013) Rapid blood meal scoring in anthropophilic *Aedes albopictus* and application of PCR blocking to avoid pseudogenes. *Infection, Genetics and Evolution* **16**, 122–128.
- Egizi AM, Robbins RG, Beati L, Nava S, Evans CR, Occi JL and Fonseca DM (2019a) A pictorial key to differentiate the recently detected exotic *Haemaphysalis longicornis* Neumann, 1901 (Acari, Ixodidae) from native congeners in North America. *ZooKeys* **818**, 117–128.
- Egizi AM, Occi JL, Price DC and Fonseca DM (2019b) Leveraging the expertise of the New Jersey mosquito control community to jump start standardized tick surveillance. *Insects* **10**, 219.
- Ferreira FC, González J, Milholland MT, Tung GA and Fonseca DM (2023) Ticks (Acari: Ixodida) on synanthropic small and medium-sized mammals in areas of the northeastern United States infested with the Asian longhorned tick, *Haemaphysalis longicornis*. *International Journal for Parasitology*. doi: 10.1016/j.ijpara.2023.06.003. Online ahead of print.
- Garrett KB, Hernandez SM, Balsamo G, Barron H, Beasley JC, Brown JD, Cloherty E, Farid H, Gabriel M, Groves B, Hamer S, Hill J, Lewis M, McManners K, Nemeth N, Oesterle P, Ortiz S, Peshock L, Schnellbacher R, Schott R, Straif-Bourgeois S and Yabsley MJ (2019) Prevalence, distribution, and diversity of cryptic piroplasm infections in raccoons from selected areas of the United States and Canada. *International Journal for Parasitology: Parasites and Wildlife* **9**, 224–233.
- Goethert HK (2021) What *Babesia microti* is now. *Pathogens* **10**, 1168.
- González J, Fonseca DM and Toledo A (2023) Seasonal dynamics of tick species in the ecotone of parks and recreational areas in Middlesex County (New Jersey, USA). *Insects* **14**, 258.
- Gray JS, Estrada-Peña A and Zintl A (2019) Vectors of babesiosis. *Annual Review of Entomology* **64**, 149–165.
- Heath A (2016) Biology, ecology and distribution of the tick, *Haemaphysalis longicornis* Neumann (Acari: Ixodidae) in New Zealand. *New Zealand Veterinary Journal* **64**, 10–20.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ and Vinh LS (2018) UFBBoot2: improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**, 518–522.
- Holman PJ, Backlund BB, Wilcox AL, Stone R, Stricklin AL and Bardin KE (2009) Detection of a large unnamed *Babesia* piroplasm originally identified in dogs in North Carolina in a dog with no history of travel to that state. *Journal of the American Veterinary Medical Association* **235**, 851–854.
- Hong S-H, Kim S-Y, Song BG, Roh JY, Cho CR, Kim C-N, Um T-H, Kwak YG, Cho S-H and Lee S-E (2019) Detection and characterization of an emerging type of *Babesia* sp. similar to *Babesia motasi* for the first case of human babesiosis and ticks in Korea. *Emerging Microbes & Infections* **8**, 869–878.
- Horowitz R and Freeman PR (2020) Archives of medical case reports case report healthy fetal outcomes using a novel treatment for maternal Lyme disease and babesiosis during consecutive pregnancies: a case study and literature review. *Archives of Medical Case Reports* **2**, 1–19.
- Jain K, Tagliafierro T, Marques A, Sanchez-Vicente S, Gokden A, Fallon B, Mishra N, Briese T, Kapoor V, Sameroff S, Guo C, Marcos LA, Hu L, Lipkin WI and Tokarz R (2021) Development of a capture sequencing assay for enhanced detection and genotyping of tick-borne pathogens. *Scientific Reports* **11**, 12384.



- Jalovecka M, Sojka D, Ascencio M and Schnittger L (2019) Babesia life cycle – when phylogeny meets biology. *Trends in Parasitology* **35**, 356–368.
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A and Jerminin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587–589.
- Keirans JE and Litwak TR (1989) Pictorial key to the adults of hard ticks, family ixodidae (Ixodida: Ixodoidea), East of the Mississippi River. *Journal of Medical Entomology* **26**, 435–448.
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Paabo S, Villablancatt FX and Wilson AC (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proceedings of the National Academy of Sciences* **86**, 6196–6200.
- Kumar A, O'Bryan J and Krause PJ (2021) The global emergence of human babesiosis. *Pathogens* **10**. doi: 10.3390/pathogens10111447
- Lerner H and Berg C (2015) The concept of health in One Health and some practical implications for research and education: what is One Health? *Infection Ecology & Epidemiology* **5**, 25300.
- Li Y, Luo J, Guan G, Ma M, Liu A, Liu J, Ren Q, Niu Q, Lu B, Gao J, Liu Z, Dang Z, Tian Z, Zhang B, He Z, Bai Q and Yin H (2009) Experimental transmission of *Theileria uilenbergi* infective for small ruminants by *Haemaphysalis longicornis* and *Haemaphysalis qinghaiensis*. *Parasitology Research* **104**, 1227–1231.
- Li A, Liu L, Wu W, Liu Y, Huang X, Li C, Liu D, Li J, Wang S, Li D and Liang M (2021) Molecular evolution and genetic diversity analysis of SFTS virus based on next-generation sequencing. *Biosafety and Health* **3**, 105–115.
- Liu K, Zhou H, Sun R-X, Yao H-W, Li Y, Wang L-P, Mu D, Li X-L, Yang Y, Gray GC, Cui N, Yin W-W, Fang L-Q, Yu H-J and Cao W-C (2015) A national assessment of the epidemiology of severe fever with thrombocytopenia syndrome, China. *Scientific Reports* **5**, 9679.
- Luo L-M, Zhao L, Wen H-L, Zhang Z-T, Liu J-W, Fang L-Z, Xue Z-F, Ma D-Q, Zhang X-S, Ding S-J, Lei X-Y and Yu X (2015) *Haemaphysalis longicornis* ticks as reservoir and vector of severe fever with thrombocytopenia syndrome virus in China. *Emerging Infectious Diseases* **21**, 1770–1776.
- Marendy D, Baker K, Emery D, Rolls P and Stutchbury R (2020) *Haemaphysalis longicornis*: the life-cycle on dogs and cattle, with confirmation of its vector status for *Theileria orientalis* in Australia. *Veterinary Parasitology* **277**, 100022.
- Nguyen L-T, Schmidt HA, von Haeseler A and Minh BQ (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**, 268–274.
- Olafson PU, Buckmeier BG, May MA and Thomas DB (2020) Molecular screening for rickettsial bacteria and piroplasms in ixodid ticks surveyed from white-tailed deer (*Odocoileus virginianus*) and nilgai antelope (*Boselaphus tragocamelus*) in southern Texas. *International Journal for Parasitology: Parasites and Wildlife* **13**, 252–260.
- Onyiche TE, Răileanu C, Fischer S and Silaghi C (2021) Global distribution of *Babesia* species in questing ticks: a systematic review and meta-analysis based on published literature. *Pathogens* **10**, 230.
- Osbrink WLA, Thomas DB, Lohmeyer KH and Temeyer KB (2022) Climate change and alternative hosts complicate the eradication of cattle fever ticks (Acari: Ixodidae) in the Southern United States, a review. *Annals of the Entomological Society of America* **115**, 39–55.
- Price KJ, Witmier BJ, Eckert RA and Boyer CN (2022) Recovery of partially engorged *Haemaphysalis longicornis* (Acari: Ixodidae) ticks from active surveillance. *Journal of Medical Entomology* **59**, 1842–1846.
- Rainey T, Occi JL, Robbins RG and Egizi A (2018) Discovery of *Haemaphysalis longicornis* (Ixodida: Ixodidae) parasitizing a sheep in New Jersey, United States. *Journal of Medical Entomology* **55**, 757–759.
- Rajapakshage BK, Yamasaki M, Hwang S-J, Sasaki N, Murakami M, Tamura Y, Lim SY, Nakamura K, Ohta H and Takiguchi M (2012) Involvement of mitochondrial genes of *Babesia gibsoni* in resistance to diminazene aceturate. *Journal of Veterinary Medical Science* **74**, 1139–1148.
- Raney WR, Perry JB and Hermance ME (2022a) Transovarial transmission of heartland virus by invasive Asian longhorned ticks under laboratory conditions. *Emerging Infectious Diseases* **28**, 726–729.
- Raney WR, Herslebs EJ, Langohr IM, Stone MC and Hermance ME (2022b) Horizontal and vertical transmission of Powassan virus by the invasive Asian longhorned tick, *Haemaphysalis longicornis*, under laboratory conditions. *Frontiers in Cellular and Infection Microbiology* **12**. doi: 10.3389/fcimb.2022.923914
- Rankins EM and Malinowski K (2020) Horse racing and veterinary practices in New Jersey. *Journal of Equine Veterinary Science* **85**, 102879.
- R Core Team (2022) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>
- Renard I and Mamoun CB (2021) Treatment of human babesiosis: then and now. *Pathogens* **10**, 1120.
- Rochlin I, Egizi A, Narvaez Z, Bonilla DL, Gallagher M, Williams GM, Rainey T, Price DC and Fonseca DM (2023) Microhabitat modeling of the invasive Asian longhorned tick (*Haemaphysalis longicornis*) in New Jersey, USA. *Ticks and Tick-borne Diseases* **14**, 102126.
- Schappach BL, Krell RK, Hornbostel VL and Connally NP (2020) Exotic *Haemaphysalis longicornis* (Acari: Ixodidae) in the United States: biology, ecology, and strategies for management. *Journal of Integrated Pest Management* **11**. doi: 10.1093/jipm/pmaa019
- Schnittger L, Ganzinelli S, Bhoora R, Omondi D, Nijhof AM and Florin-Christensen M (2022) The Piroplasmida *Babesia*, *Cytauxzoon*, and *Theileria* in farm and companion animals: species compilation, molecular phylogeny, and evolutionary insights. *Parasitology Research* **121**, 1207–1245.
- Schreeg ME, Marr HS, Tarigo JL, Cohn LA, Bird DM, Scholl EH, Levy MG, Wiegmann BM and Birkenheuer AJ (2016) Mitochondrial genome sequences and structures aid in the resolution of piroplasmida phylogeny. *PLoS ONE* **11**, e0165702.
- Scott JD and Scott CM (2018) Human babesiosis caused by *Babesia duncani* has widespread distribution across Canada. *Healthcare* **6**. doi: 10.3390/healthcare6020049
- Scott JD, Sajid MS, Pascoe EL and Foley JE (2021) Detection of *Babesia odocoilei* in humans with babesiosis symptoms. *Diagnostics* **11**, 947.
- Sikorski LE, Birkenheuer AJ, Holowaychuk MK, McCleary-Wheeler AL, Davis JM and Littman MP (2010) Babesiosis caused by a large *Babesia* species in 7 immunocompromised dogs. *Journal of Veterinary Internal Medicine* **24**, 127–131.
- Stanley HM, Ford SL, Snellgrove AN, Hartzler K, Smith EB, Krapianaya I and Levin ML (2020) The ability of the invasive Asian longhorned tick *Haemaphysalis longicornis* (Acari: Ixodidae) to acquire and transmit *Rickettsia rickettsii* (Rickettsiales: Rickettsiaceae), the agent of Rocky Mountain spotted fever, under laboratory conditions. *Journal of Medical Entomology* **57**, 1635–1639.
- Swanson M, Pickrel A, Williamson J and Montgomery S (2023) Trends in reported babesiosis cases – United States, 2011–2019. *Morbidity and Mortality Weekly Report* **72**, 273–277.
- Thompson AT, White S, Shaw D, Egizi A, Lahmers K, Ruder MG and Yabsley MJ (2020) *Theileria orientalis* Ikeda in host-seeking *Haemaphysalis longicornis* in Virginia, U.S.A. *Ticks and Tick-borne Diseases* **11**, 101450.
- Thompson AT, White SA, Doub EE, Sharma P, Frierson K, Dominguez K, Shaw D, Weaver D, Vigil SL, Bonilla DL, Ruder MG and Yabsley MJ (2022) The wild life of ticks: using passive surveillance to determine the distribution and wildlife host range of ticks and the exotic *Haemaphysalis longicornis*, 2010–2021. *Parasites & Vectors* **15**, 331.
- Trout Fryxell RT, Vann DN, Butler RA, Paulsen DJ, Chandler JG, Willis MP, Wyrosdick HM, Schaefer JJ, Gerhold RW, Grove DM, Ivey JZ, Thompson KW, Applegate RD, Sweaney J, Daniels S, Beaty S, Balthaser D, Freye JD, Mertins JW, Bonilla DL and Lahmers K (2021) Rapid discovery and detection of *Haemaphysalis longicornis* through the use of passive surveillance and collaboration: building a state tick-surveillance network. *International Journal of Environmental Research and Public Health* **18**, 7980.
- Tufts DM, Goodman LB, Benedict MC, Davis AD, VanAcker MC and Diuk-Wasser M (2021) Association of the invasive *Haemaphysalis longicornis* tick with vertebrate hosts, other native tick vectors, and tick-borne pathogens in New York City, USA. *International Journal for Parasitology* **51**, 149–157.
- Votýpka J, Modrý D, Oborník M, Šlapeta J and Lukeš J (2016) Apicomplexa. In Archibald JM, Simpson AGB, Slamovits CH, Margulis L, Melkonian M, Chapman DJ and Corliss JO (eds), *Handbook of the Protists*. Cham: Springer, pp 58. [https://doi.org/10.1007/978-3-319-32669-6\\_20-1](https://doi.org/10.1007/978-3-319-32669-6_20-1).
- Wormser GP, McKenna D, Piedmonte N, Vinci V, Egizi AM, Backenson B and Falco RC (2020) First recognized human bite in the United States by the Asian longhorned tick, *Haemaphysalis longicornis*. *Clinical Infectious Diseases* **70**, 314–316.
- Wu J, Cao J, Zhou Y, Zhang H, Gong H and Zhou J (2017) Evaluation on infectivity of *Babesia microti* to domestic animals and ticks outside the *Ixodes* genus. *Frontiers in Microbiology* **8**. doi: 10.3389/fmicb.2017.01915.