

Is *Glyphis cicatricosa* an indicator for ‘global warming’ or an ‘urban heat island’ effect in Japan?

During the course of floristic studies of Japanese lichens, *Glyphis cicatricosa* Ach. (*Graphidaceae*, Ascomycota), a pantropical species (Galloway 2007), was found on the bark of *Phellodendron amurense* Rupr. in a suburban area of Tsukuba-city (36°06'03"N, 140°06'43"E). This is on the Pacific side of eastern Honshu, with a warm-temperate climate and is the northernmost locality of *G. cicatricosa* in Japan. The sapling tree was planted in the Tsukuba Botanical Garden (c. 14 ha, 20 m elev.) in 1984 when the mean annual temperature was 12 °C, but this has gradually increased since then to almost 15 °C in 2015 when *G. cicatricosa* was discovered (Fig. 1). Does an increase in annual temperature at this site plausibly explain the unexpected phytogeographical distribution of this lichen in terms of ‘global warming’, or is it the result of an ‘urban heat island’ effect, or indeed is it related to one or more other factors?

A voucher specimen of *G. cicatricosa* collected during a field survey in 2015, now housed in the National Museum of Nature and Science (TNS), was compared with other specimens of the species in TNS listed below. Morphological observations were made using a dissecting microscope (Olympus SZX16) and a differential interference contrast microscope (Olympus BX51). Anatomical examination was undertaken using hand-cut sections mounted in GAW (glycerine: ethanol: water, 1:1:1).

Meteorological and air pollution data were obtained from the Japan Meteorological Agency and the National Institute for Environmental Studies respectively; it should be noted that the distances between the Botanical Garden and the meteorological and air pollution monitoring stations are 5.5 and 8.1 km, respectively.

Glyphis cicatricosa is characterized by its greyish white or brownish thallus with crowded lirellae formed in rounded to elliptical pseudostromata; lirellae with open discs that are covered with brownish pruina; clear hymenium; 8-spored asci with hyaline spores

which are 6–12-locular with lenticular cells, 30–55 × 7–12 µm; and absence of specific chemical substances (Nakanishi 1966; Archer 2009). The specimen collected from Tsukuba-city agrees with the morphological and chemical features mentioned above, with spore sizes ((min.–) mean ± 1SD (–max.)) as follows: (31.7–)35.1 ± 3.8(–45.1) × (6.7–)7.8 ± 0.7(–9.2) µm ($n = 10$).

Glyphis cicatricosa is widely distributed in tropical to warm-temperate regions such as south-eastern USA, Central and South America, South, South-East and East Asia, Australia, New Zealand and the Pacific (e.g. Archer 2009; Moon *et al.* 2015). In Japan, it is reported from warm-temperate and subtropical regions in the south and the Bonin Islands (Nakanishi 1966; Kurokawa 1969). The record from Tsukuba-city is the most northern locality within Japan (Fig. 2). Here the tree on which it occurs is isolated in a relatively open setting (Fig. 3). The tree on which *G. cicatricosa* is found was originally transplanted from a montane area of central Japan (Yamanashi Prefecture) where this lichen is absent and has almost certainly become established after the tree was collected. On the tree trunk c. 10 thalli (each 1–3 cm in diam.) grow, 30 to 150 cm above the ground, mainly in association with *Graphis handelii* Zahlbr. and scattered thalli of *Lecanora pulverulenta* Müll. Arg. and *Pertusaria pertusa* (L.) Tuck.

The distinctive phytogeographical distribution pattern of pantropical species in Japan is delimited south of the 15 °C annual isothermal line (with a mean annual minimum of –3.5 °C), known as the ‘Crinum Line’; this is based on the distribution of *Crinum asiaticum* L. var. *japonicum* Baker, a vascular plant belonging to the *Amaryllidaceae* (Koshimizu 1938). Many




FIG. 1. Mean annual temperature in Tsukuba-city, 1984 to 2015, at the monitoring station located 5.5 km S of the Botanical Garden (Japan Meteorological Agency; <http://www.jma.go.jp/>). Dashed line represents the threshold 15 °C isotherm, the ‘Crinum Line’, which correlates approximately with the distribution of *Glyphis cicatricosa*.

organisms have similar distribution patterns in Japan including pantropical lichens which, according to Sato (1959), are generally distributed in the warmer areas to the south of the 15 °C annual isothermal line. *Glyphis cicatricosa* in Japan (Fig. 2) is clearly pantropical, with a mainly tropical and subtropical distribution and only sporadic records from adjacent warm-temperate regions (see Kurokawa 2006). Other pantropical lichen species in Japan are represented by, for example, *Dictyonema sericeum* (Sw.) Berk., *D. moorei* (Nyl.) Henssen, *Lecanora leprosa* Feé, *Sticta gracilis* (Müll. Arg.) Zahlbr. and *S. duplolumbata* (Hue) Vain. (Sato 1959; Kurokawa 2006). Of these pantropical species, only *L. leprosa* and *G. cicatricosa* have been found in Tsukuba-city (Ohmura 2016).

The higher air temperature created in urban areas, the ‘urban heat island effect’, is well known (e.g. McCarthy *et al.* 2010; Kusaka *et al.* 2012) but it has not been widely recognized as contributing directly to lichen distribution (Coppins 1973), mainly due to the overriding impact of air pollution. In arid or sub-arid climates, both air pollution and drought can affect lichen distribution, as demonstrated by Nimis (1985). There have

been many studies which have suggested that an increase in mean temperature, for example through ‘global warming’, will influence lichen distribution (e.g. van Herk *et al.* 2002; Seaward 2004; Ellis *et al.* 2007, 2015; Aptroot 2009; Hauck 2009; Bengtsson & Paltto 2013; Davydov *et al.* 2013; Semenova *et al.* 2015). Northward shifts in distribution in Japan, as seen in *G. cicatricosa*, have already been reported in various plants and animals (Kitahara *et al.* 2001; Murakami *et al.* 2007; Japan Weather Association 2013). Therefore, the relatively recent occurrence of *G. cicatricosa*, as well as *L. leprosa*, in Tsukuba-city could be explained in terms of ‘global warming’, exacerbated by an ‘urban heat island’ effect. However, the documented distribution shifts for lichens that can be linked to global warming effects are still rather scanty, and corroborative reports for a few species, as above, have some severe limitations in the degree to which they can be generalized. Due to dramatic reductions in air pollution in many cities worldwide, as exemplified by Tsukuba-city (Fig. 4) and other cities in Japan (Ohmura *et al.* 2008, 2012, 2014), lichen distribution patterns no longer demonstrate a clear relationship with sulphur dioxide. This leads to a multivariate process involving urban




FIG. 2. Distribution map of *Glyphis cicatricosa* in Japan, with records from Tsukuba-city (★) and other localities (●); mean annual temperature from 1981 to 2010 delineated (Japan Meteorological Agency; <http://www.jma.go.jp/>).

heat islands which potentially represent higher latitude refugia for the colonization of species that are shifting their distributions as a consequence of global warming. Investigating this phenomenon is facilitated by declining levels of pollution in urban areas where lichen recolonization is of particular interest to biogeographers.

Representative specimens examined. **Japan:** Honshu, Prov. Hitachi (Pref. Ibaraki): Tsukuba Botanical Garden,

Amakubo, Tsukuba-city (36°06'03"N, 140°06'43"E), on bark of *Phellodendron amurense*, 20 m elev., 2015, Y. Ohmura 11001 (TNS). Prov. Izu (Pref. Shizuoka): Mishima, 1928, Y. Asahina 196 (TNS). Prov. Tohomi (Pref. Shizuoka): Saruwatari, Aigusa-mura, Ogasa-gun, 7 vii 1932, G. Hashimoto s. n. (TNS). Prov. Izumi (Pref. Osaka): Kami-Kyoshi, Kyoshi-mura, Semnan-gun, 1954, M. Togashi 549056 (TNS). Kyushu, Prov. Buzen (Pref. Fukuoka): Nakatsumiya, Ohshima, Munakata-gun, on tree bark, 20 m elev., 8 viii 1960, W. Yoshitake s. n. (TNS). Prov. Bungo (Pref. Ōita): Nishi-Arita-mura, Hita-gun, 21 ii 1922, Nakayama s. n. (TNS). Prov. Hyuga (Pref. Miyazaki): Nakaura, Nishi-Benbun,



FIG. 3. *Glyphis cicatricosa*, Tsukuba Botanical Garden. A, habitat; B, location on *Phellodendron amurense* indicated by arrow, associated mainly with *Graphis handelii*; C, thallus. Scale: C = 2 mm.

Nichinan, 1955, *S. Kurokawa* 550087 (TNS); Uto Shrine, Minami-Naka-gun, 1955, *S. Kurokawa* 550117 (TNS). *Izu Islands* (Tokyo Metropolis): Tairo-pond, Miyake Island, 1976, *H. Kashiwadani* 13710 (TNS). *Bonin Islands* (Tokyo Metropolis): Minami-Fukurozawa, Chichi-jima Island, 1977, *H. Kashiwadani* 14138, 14144 (TNS); Sekimon, Haha-jima Island, on twig of *Morus australis*, 250 m elev., 2013, *Y. Ohmura* 9794 (TNS). *Ryukyu Islands* (Pref. Okinawa): along the river, Nakama-gawa, Iriomote Island, Yaeyama Islands, 0–50 m elev., 1973, *I. Yoshimura* 248

(TNS); Sakamine, Kikai-jima, on bark of *Casuarina equisetifolia*, 1988, *H. Kashiwadani* 35650 (TNS).

Specimens noted in references but not examined. Japan: *Prov. Hizen* (Pref. Nagasaki): Arikawa, Arikawa-cho, 120 m elev., [undated], *M. Nakanishi* 9644 (HIRO, Nakanishi 1966). *Prov. Ohsumi* (Pref. Kagoshima): M. Nagata, Yakushima-seihokubu, [undated], *M. Oshio* 9154 (HIRO, Nakanishi 1966).

The authors are most grateful to Dr N. Tanaka and Ms T. Tanimura of the Tsukuba Botanical Garden for




FIG. 4. Annual mean concentration of sulphur dioxide, 1984 to 2013 measured at an air pollution monitoring station located 8.1 km SE of the Botanical Garden (National Institute for Environmental Studies; <http://tenbou.nies.go.jp/gis/>).

habitat information, to Dr M. Nakanishi for providing an additional specimen and valuable comments, and to anonymous referees for their helpful criticism of an earlier draft of this paper. This study is partly supported by JSPS KAKENHI Grant no. 24300314.

REFERENCES

- Aptroot, A. (2009) Lichens as an indicator of climate and global change. In *Climate Change: Observed Impacts on Planet Earth* (T. M. Letcher, ed.): 401–408. Oxford: Elsevier.
- Archer, A. W. (2009) *Graphidaceae*. In *Flora of Australia, Volume 57. Lichens 5* (P. M. McCarthy, ed.): 84–194. Canberra: ABRIS and CSIRO Publishing.
- Bengtsson, O. & Paltto, H. (2013) Changes in the epiphytic flora over 17 years in Härryda, southwest Sweden. *Svensk Botanisk Tidskrift* **107**: 100–105.
- Coppins, B. J. (1973) The ‘drought hypothesis’. In *Air Pollution and Lichens* (B. W. Ferry, M. S. Baddeley & D. L. Hawksworth, eds): 124–142. London: Athlone Press.
- Davydov, E. A., Insarov, G. E. & Sundetpaev, A. K. (2013) Lichen monitoring in Katon-Karagai National Park, eastern Kazakhstan, in context of climate change. *Problems of Ecological Monitoring and Ecosystem Modelling* **15**: 428–441.
- Ellis, C. J., Coppins, B. J., Dawson, T. P. & Seaward, M. R. D. (2007) Response of British lichens to climate change scenarios: trends and uncertainties in the projected impact for contrasting biogeographic groups. *Biological Conservation* **140**: 217–235.
- Ellis, C. J., Eaton, S., Theodoropoulos, M., Coppins, B. J., Seaward, M. R. D. & Simkin, J. (2015) *Lichen Epiphyte Scenarios. A Toolkit of Climate and Woodland Change for the 21st Century*. Edinburgh: Royal Botanic Garden.
- Galloway, D. J. (2007) *Flora of New Zealand Lichens. Volume 1*. 2nd ed. Lincoln, New Zealand: Manaaki Whenua Press.
- Hauck, M. (2009) Global warming and alternative causes of decline in arctic-alpine and boreal-montane lichens in north-western Central Europe. *Global Change Biology* **15**: 2653–2661.
- Japan Weather Association (2013) *Climate Change and its Impacts in Japan (FY2012)*. Available at: https://www.env.go.jp/en/earth/cc/impacts_FY2012.pdf
- Kitahara, M., Iriki, M. & Shimizu, G. (2001) On the relationship between the northward distributional expansion of the great mormon butterfly, *Papilio memnon* Linnaeus, and climatic warming in Japan. *Transactions of the Lepidopterological Society of Japan* **52**: 253–264.
- Koshimizu, T. (1938) On the ‘‘Criminum Line’’ in the flora of Japan. *Botanical Magazine, Tokyo* **52**: 135–139.
- Kurokawa, S. (1969) Lichens of Chichijima Island of the Bonin Islands collected by Dr. H. Inoue. *Bulletin of the National Science Museum, Tokyo* **12**: 685–692.
- Kurokawa, S. (2006) Phytogeographical elements of the lichen flora of Japan. *Journal of the Hattori Botanical Laboratory* **100**: 721–738.
- Kusaka, H., Hara, M. & Takane, Y. (2012) Urban climate projection by the WRF model at 3-km horizontal grid increment: dynamical downscaling and predicting heat stress in the 2070’s August for Tokyo, Osaka, and Nagoya metropolises. *Journal of the Meteorological Society of Japan* **90B**: 47–63.
- McCarthy, M. P., Best, M. J. & Betts, R. A. (2010) Climate change in cities due to global warming and urban effects. *Geophysical Research Letters* **37**: L09705. doi:10.1029/2010GL042845.

- Moon, K. H., Nakanishi, M., Futagami, Y. & Kashiwadani, H. (2015) Studies on Cambodian species of *Graphidaceae* (*Ostropales*, Ascomycota) (II). *Journal of Japanese Botany* **90**: 98–102.
- Murakami, K., Matsui, R. & Morimoto, Y. (2007) Northward invasion and range expansion of the invasive fern *Thelypteris dentata* (Forssk.) St. John into the urban matrix of three prefectures in Kinki District, Japan. *American Fern Journal* **97**: 186–198.
- Nakanishi, M. (1966) Taxonomical studies on the family *Graphidaceae* of Japan. *Journal of Science of Hiroshima University, Series B, Division 2 (Botany)* **11**: 51–126.
- Nimis, P. L. (1985) Urban lichen studies in Italy. I. The town of Trieste. *Studia Geobotanica* **5**: 49–74.
- Ohmura, Y. (2016) *The Handbook of Urban Lichens in Japan*. Tokyo: Bun-ichi Sogo Shuppan. (In Japanese)
- Ohmura, Y., Kawachi, M., Ohtara, K. & Sugiyama, K. (2008) Long-term monitoring of *Parmotrema tinctorum* and qualitative changes of air pollution in Shimizu Ward, Shizuoka City, Japan. *Journal of Japan Society for Atmospheric Environment* **43**: 47–54. (In Japanese with English summary)
- Ohmura, Y., Kashiwadani, H. & Moon, K. H. (2012) Recovery of macrolichen flora in the Imperial Palace Grounds, Tokyo, Japan. *Journal of Japanese Botany* **87**: 55–61.
- Ohmura, Y., Thor, G., Frisch, A., Kashiwadani, H. & Moon, K. H. (2014) Increase of lichen diversity in the Imperial Palace Grounds, Tokyo, Japan. *Memoirs of the National Museum of Nature and Science* **49**: 193–217.
- Sato, M. (1959) Range of the Japanese lichens (V). *Bulletin of the Faculty of Arts and Sciences, Ibaraki University, Natural Science* **10**: 77–87.
- Seaward, M. R. D. (2004) The use of lichens for environmental impact assessment. *Symbiosis* **37**: 293–305.
- Semenova, T. A., Morgado, L. N., Welker, J. M., Walker, M. D., Smets, E. & Geml, J. (2015) Long-term experimental warming alters community composition of ascomycetes in Alaskan moist and dry arctic tundra. *Molecular Ecology* **24**: 424–437.
- van Herk, C. M., Aptroot, A. & van Dobben, H. F. (2002) Long-term monitoring in the Netherlands suggests that lichens respond to global warming. *Lichenologist* **34**: 141–154.

**Yoshihito Ohmura and
Mark R. D. Seaward**

Y. Ohmura: Department of Botany, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, 305-0005, Japan. Email: ohmura-y@kahaku.go.jp.
M. R. D. Seaward: School of Archaeological Sciences, University of Bradford, Bradford BD7 1DP, UK.