


Erratum

Cite this article: Smits ML *et al* (2023). Day hospital versus intensive outpatient mentalization-based treatment: 3-year follow-up of patients treated for borderline personality disorder in a multicentre randomized clinical trial - ERRATUM. *Psychological Medicine* **53**, 605–608. <https://doi.org/10.1017/S0033291720004018>

First published online: 7 December 2020

Day hospital versus intensive outpatient mentalization-based treatment: 3-year follow-up of patients treated for borderline personality disorder in a multicentre randomized clinical trial - ERRATUM

Maaïke L. Smits¹ , Dine J. Feenstra¹, Dawn L. Bales², Matthijs Blankers^{3,4,5}, Jack J. M. Dekker^{3,6}, Zwaan Lucas⁷, Jan H. Kamphuis^{1,8}, Jan J. V. Busschbach^{1,9}, Roel Verheul¹⁰ and Patrick Luyten^{1,11,12}

¹Viersprong Institute for Studies on Personality Disorders, Halsteren, The Netherlands; ²Expertcentre MBT-Nederland, Bergen op Zoom, The Netherlands; ³Department of Research, Arkin Mental Health Care, Amsterdam, The Netherlands; ⁴Trimbos Institute, The Netherlands Institute of Mental Health and Addiction, Utrecht, The Netherlands; ⁵Department of Psychiatry, University of Amsterdam, Amsterdam, The Netherlands; ⁶Department of Clinical Psychology, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; ⁷Lentis, Groningen, The Netherlands; ⁸Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands; ⁹Department of Psychiatry, Section Medical Psychology and Psychotherapy, Erasmus MC, Rotterdam, The Netherlands; ¹⁰CEO De Viersprong, Halsteren, The Netherlands; ¹¹Faculty of Psychology and Educational Sciences, University of Leuven, Leuven, Belgium and ¹²Research Department of Clinical, Educational and Health Psychology, University College London, London, UK

DOI: <https://doi.org/10.1017/S0033291720002123>. Published online by Cambridge University Press, 30 June 2020

In the above published article there was an editing error in [Table 1](#). The corrected table is shown below.

The publishers apologise for this error.

Table 1. Predicted means, results from multilevel models and effect sizes 36 months after the start of treatment on primary and secondary outcome measures for patients randomly assigned to intensive outpatient mentalization-based treatment (MBT-IOP) ($n = 44$) or day hospital mentalization-based treatment (MBT-DH) ($n = 70$)

	Symptom severity (GSI)				Identity integration (SIPP)				Self-control (SIPP)			
	MBT-IOP		MBT-DH		MBT-IOP		MBT-DH		MBT-IOP		MBT-DH	
	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI
Baseline	1.97	1.84–2.11	1.88	1.78–1.97	1.88	1.76–1.99	1.86	1.79–1.94	2.05	1.95–2.15	2.16	2.08–2.23
18 months	1.45	1.25–1.64	1.30	1.16–1.44	2.34	2.13–2.54	2.64	2.78–2.50	2.57	2.40–2.74	2.86	2.75–2.98
36 months	1.36	1.11–1.61	1.16	0.96–1.36	2.71	2.43–2.99	2.59	2.36–2.81	2.88	2.64–3.13	3.00	2.82–3.19
Model: Wald χ^2	69.26 (df = 4)				96.82 (df = 6)				104.31 (df = 5)			
Linear change	–0.04	–0.05 to –0.14	$p = 0.352$		0.25*	0.05–0.47	$p = 0.016$		0.09*	0.01–0.18	$p = 0.032$	
Quadratic change	–0.02***	–0.09 to 0.05	$p < 0.001$		–0.07	–0.14 to 0.01	$p = 0.069$		–	–	–	
Cubic change	–	–	–		0.01*	0.00–0.02	$p = 0.037$		0.00	–0.00 to 0.00	$p = 0.138$	
Δ Linear change	–0.02	–0.09 to 0.05	$p = 0.610$		–0.27**	–0.44 to –0.10	$p = 0.002$		–	–	–	
Δ Quadratic change	–	–	–		0.04**	0.02–0.07	$p = 0.001$		–	–	–	
Δ Cubic change	–	–	–		–	–	–		0.00*	0.00–0.00	$p = 0.048$	
Δ Group 36 months	–0.20	–0.62 to 0.22	$P = 0.350$		–0.13	–0.60 to 0.35	$p = 0.606$		0.12	–0.30 to 0.54	$p = 0.576$	
Within-group ES (baseline–18/18–36 months)	0.86/0.12		1.04/0.21		0.80/0.49		1.58/–0.08		1.07/0.49		1.61/0.24	
(Baseline–36 months)	1.00		1.12		1.45		1.15		1.71		1.54	
Between-group ES (18/36 months)	0.24/0.26				0.50/0.15				0.57/0.17			
	Social concordance (SIPP)				Responsibility (SIPP)				Relational capacities (SIPP)			
	MBT-IOP		MBT-DH		MBT-IOP		MBT-DH		MBT-IOP		MBT-DH	
	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI
Baseline	2.63	2.53–2.73	2.75	2.68–2.83	2.49	2.37–2.61	2.67	2.60–2.75	2.29	2.15–2.42	2.16	2.08–2.24
18 months	2.81	2.65–2.97	3.00	2.89–3.11	2.74	2.55–2.92	2.86	2.74–2.97	2.41	2.21–2.61	2.58	2.46–2.70
36 months	2.99	2.78–3.20	3.25	3.11–3.39	2.98	2.76–3.21	3.04	2.88–3.20	2.54	2.29–2.79	2.58	2.41–2.75
Model: Wald χ^2	41.06 (df = 3)				43.27 (df = 3)				37.30 (df = 5)			
Linear change	0.06**	0.02–0.10	$p = 0.001$		0.08***	0.05–0.12	$p < 0.001$		0.08	–0.02 to 0.19	$p = 0.135$	
Quadratic change	–	–	–		–	–	–		–0.00	–0.02 to 0.01	$p = 0.570$	
Δ Linear change	0.02	–0.03 to 0.07	$p = 0.362$		–0.02	–0.07 to 0.02	$p = 0.337$		–0.15*	0.29 to –0.01	$p = 0.033$	
Δ Quadratic change									0.03**	0.01–0.05	$p = 0.005$	
Δ Group 36 months	0.26	–0.05 to 0.56	$p = 0.101$		0.06	–0.26 to 0.37	$p = 0.728$		–0.04	–0.18 to 0.86	$p = 0.859$	
Within-group ES (baseline–18/18–36 months)	0.38/0.32		0.59/0.53		0.45/0.39		0.15/0.62		0.21/0.29		0.91/0.00	
(Baseline–36 months)	0.77		1.09		0.90		0.75		0.55		0.79	
Between-group ES (18/36 months)	0.40/0.45				0.00/0.09				0.29/0.05			

	Borderline symptomatology (PAI-BOR)				Quality of life (EQ-5D-3L)				Interpersonal problems (IIP)			
	MBT-IOP		MBT-DH		MBT-IOP		MBT-DH		MBT-IOP		MBT-DH	
	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI	<i>M</i>	95% CI
Baseline	49.61	48.14–51.09	47.52	46.32–48.73	0.44	0.41–0.47	0.48	0.45–0.51	108.44	103.74–113.15	108.95	105.48–112.42
18 months	40.07	38.15–43.24	33.91	33.35–37.23	0.60	0.55–0.64	0.66	0.63–0.69	105.29	96.27–114.31	89.75	83.07–96.42
36 months	35.42	31.77–39.07	32.95	29.92–35.97	0.61	0.56–0.66	0.69	0.65–0.73	90.94	77.58–104.30	87.74	77.05–98.43
Model: Wald χ^2	116.44 (df = 5)				44.68 (df = 4)				19.44 (df = 5)			
Linear change	–1.56*	–3.01 to –0.10	$p = 0.036$		–0.00	–0.03 to 0.03	$p = 0.816$		–6.65	–14.47 to 1.17	$p = 0.096$	
Quadratic change	–	–	–		–	–	–		0.62	–0.51 to 1.76	$p = 0.283$	
Cubic change	–0.02	–0.05 to –0.01	$p = 0.160$		0.00**	0.00–0.00	$p = 0.007$					
Δ Linear change	–1.32	–0.55 to 3.20	$p = 0.165$		0.01	–0.02 to 0.03	$p = 0.606$		–8.85	–1.31 to 19.00	$p = 0.088$	
Δ Quadratic change	–	–	–		–	–	–		–1.58*	–3.05 to 0.10	$p = 0.036$	
Δ Cubic change	–0.04	–0.08 to 0.00	$p = 0.061$		–	–	–		–	–	–	
Δ Group 36 months	–2.47	–9.26 to 4.31	$p = 0.475$		0.08	–0.04 to 0.20	$p = 0.199$		–3.20	–28.28 to 21.87	$p = 0.802$	
Within-group ES (baseline–18/18–36 months)	1.23/0.56		1.68/0.24		1.10/0.09		1.34/0.24		0.12/0.40		0.81/0.00	
(Baseline–36 months)	1.95		1.63		1.20		1.57		0.67		0.70	
Between-group ES	0.66/0.23				0.43/0.56				0.53/0.08			
	Frequency of self-harm (SSHI)				Frequency of suicide attempts (SSHI)							
	MBT-IOP		MBT-DH		MBT-IOP		MBT-DH					
	Log <i>M</i>	95% CI	Log <i>M</i>	95% CI	Log <i>M</i>	95% CI	Log <i>M</i>	95% CI				
Baseline	1.63	1.38–1.87	0.89	0.70–1.08	0.17	0.09–0.24	0.22	0.11–0.20				
18 months	0.33	0.09–0.56	0.42	0.24–0.59	0.19	0.02–0.15	0.13	0.05–0.12				
36 months	0.27	0.11–0.42	0.41	0.290–0.53	0.11	0.00–0.07	0.04	0.00–0.02				
Model: Wald χ^2	83.04 (df = 7)				14.06 (df = 3)							
Linear change	–0.23	–0.69 to 0.22	$p = 0.315$		–0.03**	–0.05 to 0.01	$p = 0.012$					
Quadratic change	0.14	–0.04 to 0.32	$p = 0.127$		–	–	–					
Cubic change	–0.02*	–0.04 to –0.00	$p = 0.018$		–	–	–					
Δ Linear change	0.47	–0.12 to 1.05	$p = 0.119$		0.00	–0.02 to 0.03	$p = 0.830$					
Δ Quadratic change	–0.25*	–0.48 to –0.01	$p = 0.039$		–	–	–					
Δ Cubic change	0.03*	0.01 to 0.06	$p = 0.012$		–	–	–					
Δ Group 36 months	0.14	–0.33 to 0.62	$p = 0.558$		0.00	–0.09 to 0.09	$p = 0.919$					
Within-group ES (baseline–18/18–36 months)	1.45/0.10		0.56/0.01		0.31/0.57		0.39/0.72					
(Baseline–36 months)	1.51		0.64		0.62		0.90					
Between-group ES (18/36 months)	0.13/0.31				0.02/0.09							

MBT-IOP, intensive outpatient mentalization-based treatment; MBT-DH, day hospital mentalization-based treatment; GSI, Global Severity Index of the Brief Symptom Inventory; SIPP, Severity Indices of Personality Problems; PAI-BOR, Personality Assessment Inventory – Borderline Personality Disorder section; EQ-5D-3L, EuroQol-5D-3L; IIP, Inventory of Interpersonal Problems; SSHI, Suicide and Self-Harm Inventory. Linear/quadratic/cubic change represent main effects of (polynomials of) time and indicate changes over time independent of group; Δ linear/quadratic/cubic change, interaction term of time (polynomial) with group indicating differential change of MBT-DH compared to MBT-IOP over time; Δ group 36 months, difference of MBT-DH compared to MBT-IOP at 36 months; ES, effect size by means of Cohen's *d*.

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$.

Reference

- Smits, M., Feenstra, D., Bales, D., Blankers, M., Dekker, J., Lucas, Z., . . . Luyten, P. (2020). Day hospital versus intensive outpatient mentalization-based treatment: 3-year follow-up of patients treated for borderline personality disorder in a multicentre randomized clinical trial. *Psychological Medicine*, 1–11. doi:10.1017/S0033291720002123