

healthcare appointments; going to public places; returning to office-based working.

Questionnaires routinely completed at assessment and end of treatment: Montgomery–Åsberg Depression Rating Scale (MADRS); Beck's Anxiety Inventory (BAI); Beck's Depression Inventory (BDI). Adapted treatment with CBT included an extended assessment which helped differentiate anxiety symptoms from ASD. Main CBT adaptations included development of skills for the patient to identify and express emotional experiences and thoughts with the focus on physical sensations and behaviour. Graded exposure items were linked to concrete aims or interests and structured to fit around the patient's routine daily activities. Clinical data was analysed and compared outcomes from the initial standard and subsequent adapted treatment.

Results. The patient's response to the initial course of standard CBT showed a 14% increase in anxiety and 14% increase in symptoms of depression on self-rated measures. The subsequent adapted CBT showed a 31% improvement in anxiety and a 16% improvement in symptoms of depression on self-rated measures.

Conclusion. This case report supports literature describing the need to adapt standard assessment and treatment to differentiate experiences related to ASD from discrete anxiety disorders, although there may be some overlap. The promising results support using adapted CBT to ensure appropriate treatment of anxiety disorders in autistic people.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Huntington's Disease and Criminal Behaviour: An Exploration of Psychiatric Risk and Management in a High Secure Forensic Unit

Dr Thomas Gupta-Jessop^{1*} and Dr Ellice Caldwell-Dunn²

¹Forensicare, Melbourne, Australia and ²Central and North West London NHS Trust, London, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.665

Aims. Huntington's disease (HD) is an autosomal dominant neurodegenerative disorder characterised by a pathologically prolonged CAG nucleotide sequence in the huntingtin gene (HTT). Neuropsychiatric symptoms such as aggression, depression, impulsivity and psychosis are non-motor signs of HD. The association between HD and criminal behaviour is debated, and evidence lacking. This is particularly relevant in forensic psychiatry, which focusses on the risk assessment of mentally disordered offenders. This manuscript examines the antecedents of offending behaviour in a male diagnosed with HD during admission to a high secure unit, and the evolution of his risk profile from childhood to post-diagnosis. Additionally, through exploration of psychopharmacological management of psychiatric symptoms in HD, this study aims to further our understanding as to how we can best support people with HD in a forensic mental health setting.

Methods. Following review of relevant literature on criminal behaviour in the context of HD, we report the case of a 41-year-old man with a background of dissocial personality traits admitted to a high security unit with symptoms of a delusional disorder; manifesting as paranoia, delusional beliefs and aggression. These were believed to be organically induced within the context of HD, a diagnosis confirmed through genetic testing six months following admission. The patient's symptoms were

only partially responsive to first-line antipsychotics; however, good symptomatic control was achieved with clozapine and sodium valproate, enabling step-down to medium secure specialist services.

Results. In HD patients, there may be a challenge of discerning whether offending behaviour relates to prodromal presentation or whether there are pre-existing antisocial attitudes or behaviour; an uncertainty which was present in this case and within the literature. The age of HD onset is inversely correlated with CAG repeat length, and a longer repeat length has been associated with criminal behaviour. This has the potential for use as a marker to determine the time point in which presenting features are attributable to HD. In this case study it was possible to determine through analysis of the CAG repeat length that the delusional disorder was likely linked to the onset of HD; however, dissocial personality traits were not.

Conclusion. A patient's background relating to the life-course persistence of violence, suicidality and psychiatric symptoms in patients with HD informs the process of formulating their risk profile. Changes to the risk profile also reflect the progressing stages of HD. This highlights the need for awareness of how HD may contribute or predispose to criminal behaviour and how interventions could be targeted during critical periods where they benefit most.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Ghosts From the Past: A Juvenile Onset Huntington's Disease Case From Bahrain

Dr Hanan Husain*, Dr Shaima Bucheeri and Dr Feras Alsaif

Governmental Hospitals, Manama, Bahrain

*Presenting author.

doi: 10.1192/bjo.2024.666

Aims. Huntington's disease (HD) is a rare inherited disease in an autosomal dominant pattern, that is most prevalent among Caucasians.

Juvenile onset Huntington disease (JHD) is a rare subtype of the disease, defined by presence of the disease by the age of 20 or younger.

We report a case of a 28-year-old woman with JHD and discuss the challenges we faced in her diagnosis and management.

Methods. A now 28-year-old Arab woman, presented to the psychiatric hospital when she was an 18-year-old, complaining of restlessness and low mood. She was diagnosed to be having social phobia and panic attacks, and was given escitalopram. About 6 months after her first presentation, the patient's mother showed up reporting that the patient is doing well without the medication and that she is not going to take them anymore. However, the patient started developing anxiety symptoms two years later and started taking the same medication. Moreover, three years after her first presentation, the patient started developing movement symptoms and mentioned that her father passed away by Huntington's disease. The patient was immediately referred to a genetic lab and a Huntington disease diagnosis was given along with tetrabenazine and risperidone. Moreover, the patient attempted suicide multiple times after worsening of symptoms over the years. A brain magnetic resonance imaging of the patient showed bilateral caudate nuclei atrophy with similar changes affecting the putamen as well but to a lesser extent, changes that are associated with Huntington's disease.