

## HARTOGS TYPE THEOREMS FOR $CR$ $L^2$ FUNCTIONS ON COVERINGS OF STRONGLY PSEUDOCONVEX MANIFOLDS

ALEXANDER BRUDNYI

**Abstract.** We prove an analog of the classical Hartogs extension theorem for  $CR$   $L^2$  functions defined on boundaries of certain (possibly unbounded) domains on coverings of strongly pseudoconvex manifolds. Our result is related to a question formulated in the paper of Gromov, Henkin and Shubin [GHS] on holomorphic  $L^2$  functions on coverings of pseudoconvex manifolds.

### §1. Introduction

**1.1.** In this paper, following our previous work [Br4], we continue to study holomorphic  $L^2$  functions on coverings of strongly pseudoconvex manifolds. The subject was originally motivated by the paper [GHS] of Gromov, Henkin and Shubin. In [GHS] the von Neumann dimension was used to measure the space of holomorphic  $L^2$  functions on *regular* (i.e., *Galois*) coverings of a strongly pseudoconvex manifold  $M$ . In particular, it was shown that the space of such functions is infinite-dimensional. It was also asked whether the regularity of the covering is relevant for the existence of many holomorphic  $L^2$  functions on  $M'$  or it is just an artifact of the chosen method of the proof which requires a use of von Neumann algebras.

In an earlier paper [Br4] we proved that actually the regularity of  $M'$  is irrelevant for the existence of many holomorphic  $L^2$  functions on  $M'$ . Moreover, we obtained an extension of some of the main results of [GHS]. The method of the proof used in [Br4] is completely different and (probably) easier than that used in [GHS] and is based on  $L^2$  cohomology techniques, as well as, on the geometric properties of  $M$ . Also, in [Br1]–[Br3] the case of coverings of pseudoconvex domains in Stein manifolds was considered. Using the methods of the theory of coherent Banach sheaves together with

---

Received January 5, 2006.

Revised October 24, 2006.

2000 Mathematics Subject Classification: Primary 32V25; Secondary 32A40.

Research supported in part by Max-Planck-Institut für Mathematik.

Cartan's vanishing cohomology theorems, we proved some results on holomorphic  $L^p$  functions,  $1 \leq p \leq \infty$ , defined on such coverings.

**1.2.** The present paper is related to one of the open problems posed in [GHS], a Hartogs type theorem for coverings of strongly pseudoconvex manifolds. Let us recall that for a bounded open set  $D \subset \mathbb{C}^n$  ( $n > 1$ ) with a connected smooth boundary  $bD$  the classical Hartogs theorem states that any holomorphic function in some neighbourhood of  $bD$  can be extended to a holomorphic function on a neighbourhood of the closure  $\overline{D}$ . In [Bo] Bochner proved a similar extension result for functions defined on the  $bD$  only. In modern language his result says that for a smooth function defined on the  $bD$  and satisfying the tangential Cauchy-Riemann equations there is an extension to a holomorphic function in  $D$  which is smooth on  $\overline{D}$ . In fact, this statement follows from Bochner's proof (under some smoothness conditions). However at that time there was not yet the notion of a  $CR$ -function. Over the years significant contributions to the area of Hartogs theorem were made by many prominent mathematicians, see the history and the references in the paper of Harvey and Lawson [HL, Section 5]. A general Hartogs-Bochner type theorem for bounded domains  $D$  in Stein manifolds was proved by Harvey and Lawson [HL, Theorem 5.1]. The proof relies heavily upon the fact that for  $n \geq 2$  any  $\bar{\partial}$ -equation with compact support on an  $n$ -dimensional Stein manifold has a compactly supported solution. In [Br2] and [Br3] we proved some extensions of the theorem of Harvey and Lawson for certain (possibly unbounded) domains on coverings of Stein manifolds. In the present paper we prove an analogous result for  $CR$   $L^2$  functions defined on boundaries of certain domains on coverings of strongly pseudoconvex manifolds. More general Hartogs type theorems for  $CR$ -functions of slow growth on boundaries of such domains will be presented in a forthcoming paper.

**1.3.** Let  $M \subset\subset N$  be a domain with smooth boundary  $bM$  in an  $n$ -dimensional complex manifold  $N$ , specifically,

$$(1.1) \quad M = \{z \in N : \rho(z) < 0\}$$

where  $\rho$  is a real-valued function of class  $C^2(\Omega)$  in a neighbourhood  $\Omega$  of the compact set  $\overline{M} := M \cup bM$  such that

$$(1.2) \quad d\rho(z) \neq 0 \quad \text{for all } z \in bM.$$

Let  $z_1, \dots, z_n$  be complex local coordinates in  $N$  near  $z \in bM$ . Then the tangent space  $T_z N$  at  $z$  is identified with  $\mathbb{C}^n$ . By  $T_z^c(bM) \subset T_z N$  we denote the complex tangent space to  $bM$  at  $z$ , i.e.,

$$(1.3) \quad T_z^c(bM) = \left\{ w = (w_1, \dots, w_n) \in T_z(N) : \sum_{j=1}^n \frac{\partial \rho}{\partial z_j}(z) w_j = 0 \right\}.$$

The *Levi form* of  $\rho$  at  $z \in bM$  is a hermitian form on  $T_z^c(bM)$  defined in local coordinates by the formula

$$(1.4) \quad L_z(w, \bar{w}) = \sum_{j,k=1}^n \frac{\partial^2 \rho}{\partial z_j \partial \bar{z}_k}(z) w_j \bar{w}_k.$$

The manifold  $M$  is called *pseudoconvex* if  $L_z(w, \bar{w}) \geq 0$  for all  $z \in bM$  and  $w \in T_z^c(bM)$ . It is called *strongly pseudoconvex* if  $L_z(w, \bar{w}) > 0$  for all  $z \in bM$  and all  $w \neq 0, w \in T_z^c(bM)$ .

Equivalently, strongly pseudoconvex manifolds can be described as the ones which locally, in a neighbourhood of any boundary point, can be presented as strictly convex domains in  $\mathbb{C}^n$ . It is also known (see [C], [R]) that any strongly pseudoconvex manifold admits a proper holomorphic map with connected fibres onto a normal Stein space. In particular, if  $M$  is a strongly pseudoconvex non-Stein manifold of complex dimension  $n \geq 2$ , then the union  $C_M$  of all compact complex subvarieties of  $M$  of complex dimension  $\geq 1$  is a compact complex subvariety of  $M$ .

Let  $r : M' \rightarrow M$  be an unbranched covering of  $M$ . Assume that  $N$  is equipped with a Riemannian metric  $g_N$ . By  $d$  we denote the path metric on  $M'$  induced by the pullback of  $g_N$ . Consider a domain  $\tilde{D} \subset\subset M$  with a connected  $C^1$  smooth boundary  $b\tilde{D}$  such that

$$(1.5) \quad b\tilde{D} \cap C_M = \emptyset.$$

Let  $D$  be a connected component of  $r^{-1}(\tilde{D})$ . By  $bD$  we denote the boundary of  $D$  and by  $\bar{D} \subset M'$  the closure of  $D$ . Also, by  $\mathcal{O}(D)$  we denote the space of holomorphic functions on  $D$ . Now, recall that a continuous function  $f$  on  $bD$  is called *CR* if for every smooth  $(n, n - 2)$ -form  $\omega$  on  $M'$  with compact support one has

$$\int_{bD} f \cdot \bar{\partial} \omega = 0.$$

If  $f$  is smooth this is equivalent to  $f$  being a solution of the tangential *CR*-equations:  $\bar{\partial}_b f = 0$  (see, e.g., [KR]).

Let  $dV_{M'}$  and  $dV_{bD}$  be the Riemannian volume forms on  $M'$  and  $bD$  obtained by the pullback of the Riemannian metric  $g_N$  on  $N$ . By  $H^2(D)$  we denote the Hilbert space of holomorphic functions  $g$  on  $D$  with norm

$$\left( \int_{z \in D} |g(z)|^2 dV_{M'}(z) \right)^{1/2}.$$

Also,  $L^2(bD)$  stands for the Hilbert space of functions  $g$  on  $bD$  with norm

$$\left( \int_{z \in bD} |g(z)|^2 dV_{bD}(z) \right)^{1/2}.$$

The following question was asked in [GHS, Section 4]:

*Suppose that  $D$  is a regular covering of a strongly pseudoconvex manifold  $\tilde{D} \subset\subset M$ . Is it true that for every CR-function  $f \in L^2(bD) \cap C(\bar{D})$  there exists  $F \in H^2(D) \cap C(\bar{D})$  such that  $F|_{bD} = f$ ?*

In the present paper we give a particular answer to this question. To formulate our results we require the following definitions.

For every  $x$  from the closure of  $\tilde{D}$  we introduce the Hilbert space  $l_{2,x}(D)$  of functions  $g$  on  $r^{-1}(x) \cap \bar{D}$  with norm

$$(1.6) \quad |g|_x := \left( \sum_{y \in r^{-1}(x) \cap \bar{D}} |g(y)|^2 \right)^{1/2}.$$

Next, we introduce the Banach space  $\mathcal{H}_2(D)$  of holomorphic on  $D$  functions  $f$  with norm

$$|f|_D := \sup_{x \in \tilde{D}} |f|_x.$$

Similarly, we introduce the Banach space  $\mathcal{L}_2(bD)$  of continuous on  $bD$  functions  $g$  with norm

$$|g|_{bD} := \sup_{x \in b\tilde{D}} |g|_x.$$

Let  $\mathcal{U} = (U_i)_{i \in I}$  be a finite open cover of  $b\tilde{D}$  by open simply connected sets  $U_i \subset\subset M$ . Then  $r^{-1}(U_i) \cap bD$  is homeomorphic to  $(U_i \cap b\tilde{D}) \times Q$  where  $Q$  is the fibre of the covering  $r : D \rightarrow \tilde{D}$ . In what follows we identify  $r^{-1}(U_i) \cap bD$  with  $(U_i \cap b\tilde{D}) \times Q$ .

Suppose that  $f \in C(bD)$  is a CR-function satisfying the following conditions

- (1)  $f \in \mathcal{L}_2(bD)$ ;
- (2) for any  $i \in I$  and any  $z_1, z_2 \in b\tilde{D} \cap U_i$  there is a constant  $L_i$  such that

$$\left( \sum_{q \in Q} \left| \frac{f(z_1, q) - f(z_2, q)}{d((z_1, q), (z_2, q))} \right|^2 \right)^{1/2} \leq L_i.$$

(It is easy to show that condition (2) is independent of the choice of the cover.)

**THEOREM 1.1.** *For any CR-function  $f$  on  $bD$  satisfying conditions (1) and (2) there exists  $\hat{f} \in \mathcal{H}_2(D) \cap C(\bar{D})$  such that*

$$\hat{f}|_{bD} = f \quad \text{and} \quad |\hat{f}|_D = |f|_{bD}.$$

*Remark 1.2.* (A) If, in addition,  $bD$  is smooth of class  $C^k$ ,  $1 \leq k \leq \infty$ , and  $f \in C^s(bD)$ ,  $1 \leq s \leq k$ , then the extension  $\hat{f}$  belongs to  $\mathcal{O}(D) \cap C^s(\bar{D})$ . This follows from [HL, Theorem 5.1].

(B) From the Cauchy integral formula it follows that the hypotheses of the theorem are true if  $f$  is the restriction to  $bD$  of a holomorphic function from  $\mathcal{H}_2(W)$  where  $\tilde{W} := r(W) \subset M$  is a neighbourhood of  $b\tilde{D}$  and  $W$  is a connected component of  $r^{-1}(\tilde{W})$  containing  $bD$  (see [Br1, Proposition 2.4] for similar arguments).

(C) It was shown in [Br4, Theorem 1.1] that holomorphic functions from  $\mathcal{H}_2(M')$  separate points on  $M' \setminus C'_M$  where  $C'_M := r^{-1}(C_M)$ . Thus there are sufficiently many CR-functions  $f$  on  $bD$  satisfying conditions (1) and (2).

As before by  $\mathcal{L}_2(M')$  we denote the Banach space of continuous functions  $f$  on  $M'$  with norm

$$|f|_{M'} := \sup_{x \in M} |f|_x.$$

where  $|\cdot|_x$ ,  $x \in M$ , is defined as in (1.6) with  $M'$  substituted for  $\bar{D}$ . Also, for a measurable locally bounded  $(0, 1)$ -differential form  $\eta$  on  $M'$  by  $|\eta|_z$ ,  $z \in M'$ , we denote the norm of  $\eta$  at  $z$  defined by the natural hermitian metric on the fibres of the cotangent bundle  $T^*M'$  on  $M'$ . We say that such  $\eta$  belongs to the space  $\mathcal{E}_2(M')$  if

$$(1.7) \quad |\eta|_{M'} := \sup_{x \in M} \left( \sum_{z \in r^{-1}(x)} |\eta|_z^2 \right)^{1/2} < \infty.$$

(Note that this definition does not depend on the choice of the Riemannian metric on  $N$ , and that the expression in the brackets is correctly defined for almost all  $x \in M$ .)

By  $\text{supp } \eta$  we denote support of  $\eta$ , i.e., the minimal closed set  $K \subset M'$  such that  $\eta$  equals zero almost everywhere on  $M' \setminus K$ .

As mentioned above, the proof of the classical Hartogs theorem is based on the fact that for  $n \geq 2$  any  $\bar{\partial}$ -equation with compact support on an  $n$ -dimensional Stein manifold has a compactly supported solution. Similarly our proof of Theorem 1.1 is based on the following result.

**THEOREM 1.3.** *Let  $O \subset\subset M \setminus C_M$ . Assume that a  $(0, 1)$ -form  $\eta$  on  $M'$  belongs to  $\mathcal{E}_2(M')$ , is  $\bar{\partial}$ -closed (in the distributional sense) and*

$$r(\text{supp } \eta) \subset O.$$

*Then there are a function  $F \in \mathcal{L}_2(M')$  and a neighborhood  $U \subset M$  of  $bM$  such that  $\bar{\partial}F = \eta$  (in the distributional sense) and  $F|_{r^{-1}(U)} = 0$ .*

(Since  $M'$  can be thought of as a subset of a covering  $L'$  of a neighbourhood  $L$  of  $\bar{M}$ , the boundary  $bM'$  of  $M'$  is correctly defined.)

*Remark 1.4.* (A) Condition (2) in the formulation of Theorem 1.1 means that  $f$  is a Lipschitz section of a Hilbert vector bundle on  $b\tilde{D}$  with fibre  $l_2(Q)$  associated with the natural action of the fundamental group  $\pi_1(b\tilde{D})$  of  $b\tilde{D}$  on  $l_2(Q)$  (see [Br1, Example 2.2(b)] for a similar construction). This condition is required by the method of the proof. It would be interesting to know to what extent it is necessary.

(B) Another interesting question is whether a general extension theorem for  $CR$ -functions on  $bD$  without growth condition might hold.

**Acknowledgment.** This work was written during my stay at the Max-Planck-Institut für Mathematik in Bonn. I am deeply grateful to MPIM for hospitality and financial support.

## §2. Proof of Theorem 1.1

In this section we prove Theorem 1.1 modulo Theorem 1.3. Then in the next section we prove Theorem 1.3.

Since  $b\tilde{D}$  is a compact  $C^1$  smooth manifold, there are a neighbourhood  $O \subset\subset M \setminus C_M$  of  $b\tilde{D}$  and a  $C^1$  retraction  $p : O \rightarrow b\tilde{D}$ . (As such  $O$  one can

take, e.g., a neighbourhood of the zero section of the normal vector bundle on  $b\tilde{D}$ .) Without loss of generality we may assume also that fundamental groups  $\pi_1(O)$  and  $\pi_1(b\tilde{D})$  are isomorphic. Let  $O'$  be a connected component of  $r^{-1}(O) \subset M'$  containing  $bD$ . Then by the covering homotopy theorem there is a  $C^1$  retraction  $p' : O' \rightarrow bD$  such that  $r \circ p' = p \circ r$ .

Let  $\rho, 0 \leq \rho \leq 1$ , be a  $C^\infty$  function on  $M$  equals 1 in a neighbourhood of  $b\tilde{D}$  with  $\text{supp } \rho \subset\subset O$ . Consider the  $C^\infty$  function  $\rho' := \rho \circ r$  on  $M'$ .

Let  $\mathcal{V} = (V_j)_{j \in J}$  be a finite open cover of  $\tilde{D} \cup b\tilde{D}$  by simply connected coordinate charts  $V_j \subset\subset M$ . We naturally identify  $r^{-1}(V_j)$  with  $V_j \times S$  where  $S$  is the fibre of  $r : M' \rightarrow M$ . Then in these local coordinates on  $M'$  we have

$$(2.1) \quad p'(z, s) = (p(z), s), \quad \rho'(z, s) = \rho(z), \quad (z, s) \in O' \cap r^{-1}(V_j), \quad j \in J.$$

Next, for a CR-function  $f$  satisfying the assumptions of the theorem we define

$$(2.2) \quad f_1(z) := \rho'(z) \cdot f(p'(z)), \quad z \in \bar{D}.$$

LEMMA 2.1. *In the above local coordinates on  $M'$  one has*

$$\left( \sum_{s \in S} \left| \frac{f_1(z_1, s) - f_1(z_2, s)}{d((z_1, s), (z_2, s))} \right|^2 \right)^{1/2} \leq C_j, \quad (z_1, s), (z_2, s) \in \bar{D} \cap r^{-1}(V_j), \quad j \in J,$$

for some numerical constants  $C_j$ .

*Proof.* By  $d_N$  we denote the path metric on  $N$  determined by the Riemannian metric  $g_N$ . Since the path metric  $d$  on  $M'$  is obtained by the pullback of  $g_N$ , we have  $d((z_1, s), (z_2, s)) = d_N(z_1, z_2)$ . Also, by the definition of  $p'$  and  $\rho'$  we clearly have for some  $C > 0$ ,

$$\begin{aligned} d(p'(z_1, s), p'(z_2, s)) &\leq C d_N(z_1, z_2) \quad \text{for all } z_1, z_2 \in \text{supp } \rho, \text{ and} \\ |\rho'(z_1, s) - \rho'(z_2, s)| &\leq C d_N(z_1, z_2) \quad \text{for all } z_1, z_2 \in M. \end{aligned}$$

Using these inequalities, condition (2) of the theorem and the triangle inequality for  $l_2$  norms we obtain that there is  $A > 0$  such that for  $z_1, z_2 \in$

$\text{supp } \rho$

$$\begin{aligned}
& \left( \sum_{s \in S} \left| \frac{f_1(z_1, s) - f_1(z_2, s)}{d_N(z_1, z_2)} \right|^2 \right)^{1/2} \\
& \leq \left( \sum_{s \in S} \left\{ \left| \frac{\rho(z_1) - \rho(z_2)}{d_N(z_1, z_2)} \right| \cdot |f(p(z_1), s)| \right. \right. \\
& \quad \left. \left. + |\rho(z_2)| \cdot \left| \frac{f(p(z_1), s) - f(p(z_2), s)}{d_N(z_1, z_2)} \right| \right\}^2 \right)^{1/2} \\
& \leq C \left\{ \left( \sum_{s \in S} |f(p(z_1), s)|^2 \right)^{1/2} + \left( \sum_{s \in S} \left| \frac{f(p(z_1), s) - f(p(z_2), s)}{d((p(z_1), s), (p(z_2), s))} \right|^2 \right)^{1/2} \right\} \\
& \leq A.
\end{aligned}$$

Suppose now that, e.g.,  $z_1 \in \text{supp } \rho$  and  $z_2 \notin \text{supp } \rho$ . Then the term with  $|\rho(z_2)|$  in the second line of the above inequalities disappears and again we get the required estimate. Finally, the case  $z_1, z_2 \notin \text{supp } \rho$  is obvious.  $\square$

This lemma in particular implies that  $f_1$  is a bounded Lipschitz function on  $\overline{D}$ . Now, using the McShane extension theorem [M] we extend  $f_1$  to a Lipschitz function  $\tilde{f}$  on  $M'$ .

Further, since locally the metric  $d$  is equivalent to the Euclidean metric and since  $\tilde{f}$  is Lipschitz on  $M'$ , by the Rademacher theorem, see, e.g., [Fe, Section 3.1.6],  $\tilde{f}$  is differentiable almost everywhere. In particular,  $\overline{\partial} \tilde{f}$  is a  $(0, 1)$ -form on  $M'$  whose coefficients in its local coordinate representations are  $L^\infty$ -functions. Let  $\chi_D$  be the characteristic function of  $D$ . Consider the  $(0, 1)$ -form on  $M'$  defined by

$$\omega := \chi_D \cdot \overline{\partial} \tilde{f}.$$

Then repeating word-for-word the arguments of [Br3, Lemma 3.3] we get

LEMMA 2.2.  $\omega$  is  $\overline{\partial}$ -closed in the distributional sense.  $\square$

Also, the inequality of Lemma 2.1 implies immediately that  $\omega \in \mathcal{E}_2(M')$ , see (1.7). Moreover, by our construction  $r(\text{supp } \omega) \subset\subset M \setminus C_M$ . Thus according to Theorem 1.3 there is a continuous function  $F \in \mathcal{L}_2(M')$  such that  $\overline{\partial} F = \omega$  and  $F|_{r^{-1}(U)} = 0$  for a neighbourhood  $U \subset M$  of  $bM$ . Since



$D \subset M'$  is a domain with a connected boundary, and  $F$  is holomorphic outside  $\bar{D}$  (by the definition of  $\omega$ ), the latter implies that  $F|_{bD} = 0$ .

We set

$$\hat{f}(z) := f_1(z) - F(z), \quad z \in \bar{D}.$$

Using the above properties of  $f_1$  and  $F$  one obtains easily that

$$\hat{f} \in \mathcal{O}(D) \cap C(\bar{D}) \quad \text{and} \quad \hat{f}|_{bD} = f.$$

Since  $f_1$  and  $F|_{\bar{D}}$  belong to  $\mathcal{L}_2(\bar{D})$ ,  $\hat{f} \in \mathcal{H}_2(D)$ . Now, the identity  $|\hat{f}|_D = |f|_{bD}$  follows from the fact that the function  $z \mapsto |f|_z$ ,  $z \in \tilde{D} \cup b\tilde{D}$ , see (1.6), is continuous and plurisubharmonic on  $\tilde{D}$ .

This completes the proof of the theorem. □

### §3. Proof of Theorem 1.3

**3.1.** In Sections 3.1–3.6 we collect some auxiliary results required in the proof. Then in Section 3.7 we prove the theorem.

Let  $X$  be a complete Kähler manifold of dimension  $n$  with a Kähler form  $\omega$  and  $E$  be a hermitian holomorphic vector bundle on  $X$  with curvature  $\Theta$ . Let  $L_2^{p,q}(X, E)$  be the space of  $L^2$   $E$ -valued  $(p, q)$ -forms on  $X$  with the  $L^2$  norm, and let  $W_2^{p,q}(X, E)$  be the subspace of forms such that  $\bar{\partial}\eta$  is  $L^2$ . (The forms  $\eta$  may be taken to be either smooth or just measurable, in which case  $\bar{\partial}\eta$  is understood in the distributional sense.) The cohomology of the resulting  $L^2$  Dolbeault complex  $(W_2^{\cdot, \cdot}, \bar{\partial})$  is the  $L^2$  cohomology

$$H_{(2)}^{p,q}(X, E) = Z_2^{p,q}(X, E) / B_2^{p,q}(X, E),$$

where  $Z_2^{p,q}(X, E)$  and  $B_2^{p,q}(X, E)$  are the spaces of  $\bar{\partial}$ -closed and  $\bar{\partial}$ -exact forms in  $L_2^{p,q}(X, E)$ , respectively.

If  $\Theta \geq \epsilon\omega$  for some  $\epsilon > 0$  in the sense of Nakano, then the  $L^2$  Kodaira-Nakano vanishing theorem, see [D], [O], states that

$$(3.1) \quad H_{(2)}^{n,r}(X, E) = 0 \quad \text{for } r > 0.$$

Assume now that  $\Theta \leq -\epsilon\omega$  for some  $\epsilon > 0$  in the sense of Nakano. Then using a duality argument and the Kodaira-Nakano vanishing theorem (3.1) one obtains, see [L, Corollary 2.4],

$$(3.2) \quad H_{(2)}^{0,r}(X, E) = 0 \quad \text{for } r < n.$$

**3.2.** Let  $M \subset\subset N$  be a strongly pseudoconvex manifold. Without loss of generality we will assume that  $\pi_1(M) = \pi_1(N)$  and  $N$  is strongly pseudoconvex, as well. Then there exist a normal Stein space  $X_N$ , a proper holomorphic surjective map  $p : N \rightarrow X_N$  with connected fibres and points  $x_1, \dots, x_l \in X_N$  such that

$$p : N \setminus \bigcup_{1 \leq i \leq l} p^{-1}(x_i) \longrightarrow X_N \setminus \bigcup_{1 \leq i \leq l} \{x_i\}$$

is biholomorphic, see [C], [R]. By definition, the domain  $X_M := p(M) \subset X_N$  is strongly pseudoconvex, and so it is Stein. Without loss of generality we may assume that  $x_1, \dots, x_l \in X_M$ . Thus  $\bigcup_{1 \leq i \leq l} p^{-1}(x_i) = C_M$ .

Next, we introduce a complete Kähler metric on the complex manifold  $M \setminus C_M$  as follows.

First, according to [N] there is a proper one-to-one map  $i : X_M \hookrightarrow \mathbb{C}^{2n+1}$ ,  $n = \dim_{\mathbb{C}} X_M$ , which is an embedding in regular points of  $X_M$ . Then  $i(X_M)$  is a complex subvariety of  $\mathbb{C}^{2n+1}$ . By  $\omega_e$  we denote the (1,1)-form on  $M$  obtained as the pullback by  $i \circ p$  of the Euclidean Kähler form on  $\mathbb{C}^{2n+1}$ . Clearly,  $\omega_e$  is  $d$ -closed and positive outside  $C_M$ .

Similarly we can embed  $X_N$  into  $\mathbb{C}^{2n+1}$  as a closed complex subvariety. Let  $j : X_N \hookrightarrow \mathbb{C}^{2n+1}$  be an embedding such that  $j(X_M)$  belongs to the open Euclidean ball  $B$  of radius  $1/4$  centered at  $0 \in \mathbb{C}^{2n+1}$ . Set  $z_i := j(x_i)$ ,  $1 \leq i \leq l$ . By  $\omega_i$  we denote the restriction to  $M \setminus C_M$  of the pullback with respect to  $j \circ p$  of the form  $-\sqrt{-1} \cdot \partial \bar{\partial} \log(\log \|z - z_i\|^2)^2$  on  $\mathbb{C}^{2n+1} \setminus \{z_i\}$ . (Here  $\|\cdot\|$  stands for the Euclidean norm on  $\mathbb{C}^{2n+1}$ .) Since  $j(X_M) \subset B$ , the form  $\omega_i$  is Kähler. Its positivity follows from the fact that the function  $-\log(\log \|z\|^2)^2$  is strictly plurisubharmonic for  $\|z\| < 1$ . Also,  $\omega_i$  is extended to a smooth form on  $M \setminus p^{-1}(x_i)$ . Now, let us introduce a Kähler form  $\omega_M$  on  $M \setminus C_M$  by the formula

$$(3.3) \quad \omega_M := \omega_e + \sum_{1 \leq i \leq l} \omega_i.$$

**PROPOSITION 3.1.** *The path metric  $d$  on  $M \setminus C_M$  induced by  $\omega_M$  is complete.*

*Proof.* Assume, on the contrary, that there is a sequence  $\{w_j\}$  convergent either to  $C_M$  or to the boundary  $bM$  of  $M$  such that the sequence  $\{d(o, w_j)\}$  is bounded (for a fixed point  $o \in M \setminus C_M$ ). Then, since  $\omega_L \geq \omega_e$ ,

the sequence  $\{i(p(w_j))\} \subset \mathbb{C}^{2n+1}$  is bounded. This implies that  $\{w_j\}$  converges to  $C_M$ . But since  $\omega_L \geq \sum \omega_i$ , the latter is impossible. One can check this using single blow-ups of  $\mathbb{C}^{2n+1}$  at points  $z_i$  and rewriting the pullbacks to the resulting manifold of (1, 1)-forms  $-\sqrt{-1} \cdot \partial\bar{\partial} \log(\log \|z - z_i\|^2)^2$  in local coordinates near exceptional divisors, see, e.g., [GM] for similar arguments. □

Similarly one obtains complete Kähler metrics on unbranched coverings of  $M \setminus C_M$  induced by pullbacks to these coverings of the Kähler form  $\omega_M$  on  $M \setminus C_M$ .

**3.3.** We retain the notation of the previous section.

Let  $r : N' \rightarrow N$  be an unbranched covering. Consider the corresponding covering  $(M \setminus C_M)' := r^{-1}(M \setminus C_M)$  of  $M \setminus C_M$ . We equip  $(M \setminus C_M)'$  with the complete Kähler metric induced by the form  $\omega'_M := r^*\omega_M$ . Next we consider the function  $f := \sum_{0 \leq s \leq l} f_s$  on  $(M \setminus C_M)'$  such that  $f_0$  is the pullback by  $i \circ p \circ r$  of the function  $\|z\|^2$  on  $\mathbb{C}^{2n+1}$  and  $f_s$  is the pullback by  $j \circ p \circ r$  of the function  $-\log(\log \|z - z_s\|^2)^2$  on  $\mathbb{C}^{2n+1} \setminus \{z_s\}$ ,  $1 \leq s \leq l$ . Clearly we have

$$(3.4) \quad \omega'_M := \sqrt{-1} \cdot \partial\bar{\partial} f.$$

Let  $E := (M \setminus C_M)' \times \mathbb{C}$  be the trivial holomorphic line bundle on  $(M \setminus C_M)'$ . Let  $g$  be the pullback to  $(M \setminus C_M)'$  of a smooth plurisubharmonic function on  $M$ . We equip  $E$  with the hermitian metric  $e^{f+g}$  (i.e., for  $z \times v \in E$  the square of its norm in this metric equals  $e^{f(z)+g(z)}|v|^2$  where  $|v|$  is the modulus of  $v \in \mathbb{C}$ ). Then the curvature  $\Theta_E$  of  $E$  satisfies

$$(3.5) \quad \Theta_E := -\sqrt{-1} \cdot \partial\bar{\partial} \log(e^{f+g}) = -\omega'_M - \sqrt{-1} \cdot \partial\bar{\partial} g \leq -\omega'_M.$$

From here by (3.2) we obtain

$$(3.6) \quad H_{(2)}^{0,r}((M \setminus C_M)', E) = 0 \quad \text{for } r < n.$$

**3.4.** In the proof we also use the following result.

**LEMMA 3.2.** *Let  $h$  be a nonnegative piecewise continuous function on  $M$  equals 0 in some neighbourhood of  $C_M$  and bounded on every compact subset of  $M \setminus C_M$ . Then there exists a smooth plurisubharmonic function  $\hat{g}$  on  $M$  such that*

$$\hat{g}(z) \geq h(z) \quad \text{for all } z \in M.$$

*Proof.* Without loss of generality we identify  $M \setminus C_M$  with  $X_M \setminus \bigcup_{1 \leq j \leq l} \{x_j\}$ . Also, we identify  $X_M$  with a closed subvariety of  $\mathbb{C}^{2n+1}$  as in Section 3.2. Let  $U$  be a neighbourhood of  $\bigcup_{1 \leq j \leq l} \{x_j\}$  such that  $h|_U \equiv 0$ . By  $\Delta_r \subset \mathbb{C}^{2n+1}$  we denote the open polydisk of radius  $r$  centered at  $0 \in \mathbb{C}^{2n+1}$ . Assume without loss of generality that  $0 \in X_M \setminus U$ . Consider the monotonically increasing function

$$(3.7) \quad v(r) := \sup_{\Delta_r \cap X_M} h, \quad r \geq 0.$$

By  $v_1$  we denote a smooth monotonically increasing function satisfying  $v_1 \geq v$  (such  $v_1$  can be easily constructed by  $v$ ). Let us determine

$$v_2(r) := \int_0^{r+1} 2v_1(2t) dt, \quad r \geq 0.$$

By the definition  $v_2$  is smooth, convex and monotonically increasing. Moreover,

$$v_2(r) \geq \int_{\frac{r+1}{2}}^{r+1} 2v_1(2t) dt \geq (r+1)v(r+1).$$

Next we define a smooth plurisubharmonic function  $v_3$  on  $\mathbb{C}^{2n+1}$  by the formula

$$v_3(z_1, \dots, z_{2n+1}) := \sum_{j=1}^{2n+1} v_2(|z_j|).$$

Then the pullback of  $v_3$  to  $M$  is a smooth plurisubharmonic function on  $M$ . This is the required function  $\hat{g}$ . Indeed, under the identification described at the beginning of the proof for  $|z|_\infty := \max_{1 \leq i \leq 2n+1} |z_i|$  we have

$$\begin{aligned} \hat{g}(z) = v_3(z) &\geq (|z|_\infty + 1)v(|z|_\infty + 1) \\ &\geq \sup_{\Delta_{|z|_\infty + 1} \cap X_M} h \geq h(z) \quad \text{for all } z \in M. \quad \square \end{aligned}$$

**3.5.** In the proof of Theorem 1.3 we will assume without loss of generality that  $C_M$  is a *divisor with normal crossings*. Indeed, according to the Hironaka theorem, there is a *modification*  $m : N_H \rightarrow N$  of  $N$  from Section 1.3 such that  $m^{-1}(C_M)$  is a divisor with normal crossings and  $m : N_H \setminus m^{-1}(C_M) \rightarrow N \setminus C_M$  is biholomorphic. By the definition  $M_H := m^{-1}(M) \subset N_H$  is strongly pseudoconvex. Further, since  $M$  is a complex manifold,  $m$  induces an isomorphism of fundamental groups

$m_* : \pi_1(M_H) \rightarrow \pi_1(M)$ . Thus for an unbranched covering  $r : M' \rightarrow M$  of  $M$  there are a covering  $r_H : M'_H \rightarrow M_H$  and a modification  $m' : M'_H \rightarrow M'$  such that  $r \circ m' = m \circ r_H$  and  $m'$  induces an isomorphism of the corresponding fundamental groups.

Assume now that a  $(0, 1)$ -form  $\eta \in \mathcal{E}_2(M')$  satisfies the hypotheses of Theorem 1.3. Consider its pullback  $\tilde{\eta} := (m')^*\eta$  on  $M'_H$ . Clearly,  $\tilde{\eta}$  also satisfies the hypotheses of Theorem 1.3 with  $M$  replaced by  $M_H$ . Now, suppose that Theorem 1.3 is valid for  $M'_H$ , i.e., there is a continuous function  $\tilde{f} \in \mathcal{L}_2(M'_H)$  such that  $\bar{\partial}\tilde{f} = \tilde{\eta}$  and  $\tilde{f}$  vanishes in a neighbourhood of  $bM'_H$ . Since by the definition of  $\eta$  the function  $\tilde{f}$  is holomorphic in a neighbourhood of  $(r \circ m')^{-1}(C_M) \subset M'_H$  and  $m' : M'_H \rightarrow M'$  is a modification of  $M'$ , there is a function  $f \in \mathcal{L}_2(M')$  such that  $\tilde{f} = (m')^*f$ . Obviously,  $f$  satisfies the required statements of the theorem.

**3.6.** Let  $U_q \subset\subset M$  be a simply connected coordinate chart of  $q \in C_M$  with complex coordinates  $z = (z_1, \dots, z_n)$ ,  $n = \dim_{\mathbb{C}} M$ , such that  $z_1(q) = \dots = z_n(q) = 0$  and

$$(3.8) \quad C_M \cap U_q = \{f_q(z) = 0\}, \quad f_q(z) := z_1 \cdots z_k.$$

(Such coordinates exist by the definition of a divisor with normal crossings.)

Let  $\hat{f}$  be a function on  $M \setminus C_M$  such that  $r^*\hat{f} = f$ , see Section 3.3. From the definition of  $f$  we obtain

LEMMA 3.3.  *$e^{\hat{f}}$  extended by 0 to  $C_M$  is a continuous function on  $M$  such that  $e^{\hat{f}}/|f_q|$  is unbounded on  $U_q \setminus C_M$ . □*

Let  $\omega$  be the associated  $(1, 1)$ -form of a hermitian metric  $g_N$  on  $N$ . Since by the definition  $\omega_M \geq \omega_e$  and the latter form vanishes on  $C_M$ , we have locally near  $C_M \cap U_q$

$$(3.9) \quad \omega_M^n \geq c'|f_q|^{2m'}\omega^n$$

for some  $c' > 0$ ,  $m' \in \mathbb{N}$ . This and Lemma 3.3 imply that locally near  $C_M \cap U_q$

$$(3.10) \quad e^{\hat{f}}\omega_M^n \geq c|f_q|^{2m}\omega^n$$

for some  $c > 0$ ,  $m \in \mathbb{N}$ .

By  $E_n(M)$  we denote a holomorphic line vector bundle on  $M$  determined by the divisor  $nC_M$ ,  $n \in \mathbb{N}$ . Let  $s_1$  be a holomorphic section of  $E_1(M)$

defined in local coordinates on  $U_q$  by functions  $f_q$  from (3.8). Then  $(r^*s_1)^n$  is a holomorphic section of the bundle  $E'_n(M) := r^*E_n(M)$  on  $M'$ .

Since the hermitian bundle  $E$  from Section 3.3 is holomorphically trivial, we naturally identify sections of  $E$  with functions on  $(M \setminus C_M)'$ . Here and below we set  $X' := r^{-1}(X)$  for  $X \subset M$ . Also, the Banach space  $\mathcal{L}_2(X')$  of continuous functions on  $X'$  is defined similarly to  $\mathcal{L}_2(M')$ , see Section 1.3.

Let  $(U_i)_{i \in I}$  be a finite open cover of a neighbourhood  $\overline{M}$  ( $\subset\subset N$ ) by simply connected coordinate charts  $U_i \subset\subset N$ .

**PROPOSITION 3.4.** *Suppose  $h \in L_2((M \setminus C_M)', E)$  is such that for any  $U'_i$  there is a continuous function  $h_i \in \mathcal{L}_2(U'_i)$  so that  $c_i := h - h_i \in \mathcal{O}((U_i \setminus C_M)')$ . Then there is an integer  $n \in \mathbb{N}$  independent of  $h$  such that  $h \cdot (r^*s_1)^n$  admits an extension  $\hat{h} \in C(M', E'_n(M))$ . Moreover,  $h|_{O'} \in \mathcal{L}_2(O')$  for every  $O \subset\subset M \setminus C_M$ .*

*Proof.* Let  $U_q$  be a simply connected coordinate chart of  $q \in C_M$  with the local coordinates satisfying (3.8). We naturally identify  $U'_q$  with  $U_q \times S$  where  $S$  is the fibre of  $r$ . Then the hypotheses of the proposition imply that

$$(3.11) \quad \int_{z \in U_q \setminus C_M} \left( \sum_{s \in S} |h(z, s)|^2 \right) e^{\hat{f}(z) + \hat{g}(z)} \omega_M^n(z) < \infty$$

where  $\hat{g}$  is a smooth plurisubharmonic function on  $M$  such that  $r^*\hat{g} = g$ . Diminishing if necessary  $U_q$  assume that  $\hat{f}, \omega_M^n$  satisfy (3.10) there. Also, on  $U_q$  we clearly have  $\hat{g} \sim 1$ . From here and (3.11) we obtain on  $U_q$

$$(3.12) \quad \int_{z \in U_q \setminus C_M} \left( \sum_{s \in S} |h(z, s)|^2 \right) |f_q(z)|^{2m} \omega^n(z) < \infty$$

where  $f_q$  is defined by (3.8).

Further, according to the hypothesis of the proposition, there is a continuous function  $h_q \in \mathcal{L}_2(U'_q)$  such that  $c_q := h - h_q \in \mathcal{O}((U_q \setminus C_M)')$ . This and (3.12) imply that every  $f_q^m \cdot c_q(\cdot, s), s \in S$ , is  $L^2$  integrable with respect to the volume form  $(\sqrt{-1})^n \wedge_{i=1}^n dz_i \wedge d\bar{z}_i$ . Using these facts and the Cauchy integral formulas for coefficients of the Laurent expansion of  $f_q^m c_q(\cdot, s)$ , one obtains easily that every  $f_q^m c_q(\cdot, s)$  can be extended holomorphically to  $U_q$ . In turn, this gives a continuous extension  $\hat{h}$  of  $h \cdot (r^*f_q)^m$  to  $U'_q$ .

Let  $V_q \subset\subset U_q$  be another connected neighbourhood of  $q$ . By the Bergman inequality for holomorphic functions, see, e.g., [GR, Chapter 6, Theorem 1.3], we have

$$(3.13) \quad |h(y, s)f_q^m(y)|^2 \leq A \int_{z \in U_q} |h(z, s)f_q^m(z)|^2 \omega^n(z) \quad \text{for all } (y, s) \in W'_q$$

with some constant  $A$  depending on  $U_q, W_q$  and  $\omega$  only. Therefore from (3.12) and (3.13) we obtain

$$\sup_{z \in V_q} \left( \sum_{s \in S} |\hat{h}(z, s)|^2 \right)^{1/2} < \infty.$$

Equivalently,  $\hat{h}|_{V'_q} \in \mathcal{L}_2(V'_q)$ .

Next assume that  $U_q \subset (U_i)_{i \in I}$  is a simply connected coordinate neighbourhood of a point  $q \in M \setminus C_M$ . Without loss of generality we may assume that all such  $U_q$  are relatively compact in  $M \setminus C_M$ . Identifying  $U'_q$  with  $U_q \times S$  we have anew inequality of type (3.11) for  $h|_{U'_q}$ . Since  $U_q \subset\subset M \setminus C_M$  and  $\hat{f}, \hat{g}$  and  $\omega_M^n$  are smooth on  $M \setminus C_M$  by their definitions, we obviously have on  $U_q$

$$e^{\hat{f} + \hat{g}} \cdot \omega_M^n \sim \omega^n.$$

Similarly to (3.12)–(3.13) (with  $f_q = 1$ ) this implies that  $h|_{V'_q} \in \mathcal{L}_2(V'_q)$  for any connected neighbourhood  $V_q \subset\subset U_q$  of  $q$ . Choose the above neighbourhoods  $V_q$  so that they form a finite cover of a set  $O \subset\subset M \setminus C_M$ . Then from the implications  $h|_{V'_q} \in \mathcal{L}_2(V'_q)$  we obtain that  $h|_{O'} \in \mathcal{L}_2(O')$ . Now, choosing the neighbourhoods  $V_q, q \in C_M$ , so that they form a finite cover of  $C_M$  and taking as the  $n$  the maximum of the numbers  $m$  in the powers of  $f_q$ , see (3.10), we obtain that  $h \cdot (r^*s_1)^n$  admits an extension  $\hat{h} \in C(M', E'_n(M))$ . By our construction  $n$  is independent of  $h$ . □

**3.7. Proof of Theorem 1.3**

Assume that a  $(0, 1)$ -form  $\eta$  belongs to  $\mathcal{E}_2(M')$ , is  $\bar{\partial}$ -closed and  $r(\text{supp } \eta) \subset O \subset\subset M \setminus C_M$ .

Let us define the function  $g$  in the definition of the bundle  $E$  from Section 3.3 by Lemma 3.2. Namely, fix a neighbourhood  $U \subset\subset M$  of  $C_M$  and consider the function  $h$  on  $M$  defined by the formula

$$(3.14) \quad h(z) := \frac{\chi_{U^c}(z)}{\text{dist}(z, bM)}$$

where  $\chi_{U^c}$  is the characteristic function of  $U^c := M \setminus U$  and the distance to the boundary is defined by the path metric  $d_N$  on  $N$  induced by the Riemannian metric  $g_N$ . Further, according to Lemma 3.2 we can find a  $C^\infty$  plurisubharmonic function  $\hat{g}$  on  $M$  such that  $\hat{g}(z) \geq h(z)$  for all  $z \in M$ . Then in the definition of the metric on  $E$  we choose  $g := r^*\hat{g}$ .

LEMMA 3.5. *The form  $\eta$  belongs to  $L_2^{0,1}((M \setminus C_M)', E)$ .*

*Proof.* We retain the notation of Proposition 3.4. Consider the set  $U'_q \cong U_q \times S$  on  $M'$  for some  $q \in M$  such that  $U_q \subset\subset M \setminus C_M$ . Since  $\eta \in \mathcal{E}_2(M')$ ,  $r(\text{supp } \eta) \subset O \subset\subset M \setminus C_M$  and  $\hat{g}$ ,  $\hat{f}$  and  $\omega_M^n$  are bounded on  $O$ , for every such  $U_q$  we have

$$(3.15) \quad \int_{z \in U_q \setminus C_M} \left( \sum_{s \in S} |\eta|_{(z,s)}^2 \right) e^{\hat{f}(z) + \hat{g}(z)} \omega_M^n(z) < \infty.$$

(Recall that  $|\eta|_{(z,s)}^2$  stands for the norm of  $\eta$  at  $(z, s) \in M'$  defined by the natural hermitian metric on the fibres of the cotangent bundle  $T^*M'$  on  $M'$ .) Taking a finite open cover of  $O$  by such sets  $U_q$  we get the required statement.  $\square$

From Lemma 3.5 and the fact that  $\bar{\partial}\eta = 0$  we obtain by (3.6) that there exists a function  $F' \in L_2((M \setminus C_M)', E)$  such that  $\bar{\partial}F' = \eta$ . Moreover, by the definition of  $\eta$ , this function is holomorphic on  $(M \setminus C_M)' \setminus r^{-1}(\bar{O})$ . Also, since  $\eta \in \mathcal{E}_2(M')$  the equation  $\bar{\partial}G = \eta$  has local (continuous) solutions  $f_U \in \mathcal{L}_2(U')$  for every  $U \subset\subset M$  biholomorphic to an open Euclidean ball of  $\mathbb{C}^n$ . (In fact, since  $U' \cong U \times S$ , we can rewrite the equation  $\bar{\partial}G = \eta$  on  $U'$  as a  $\bar{\partial}$ -equation on  $U$  with a measurable Hilbert valued  $(0, 1)$ -form on the right-hand side. Then we apply the formula presented in the proof of Lemma 3.4 of [Br3] (see also [H, Section 4.2]) to solve this equation and to get a solution from  $\mathcal{L}_2(U')$ , for similar arguments see [Br1, Appendix A].)

Let us prove now

LEMMA 3.6. *There is a neighbourhood  $U \subset M$  of  $bM$  such that  $F'|_{r^{-1}(U)} = 0$ .*

*Proof.* Let  $q \in bM$  and  $U_q \subset\subset N \setminus C_M$  be a simply connected coordinate chart of  $q$ . Since  $\pi_1(M) = \pi_1(N)$  by our assumption, the covering  $M'$  of  $M$  is contained in the corresponding covering  $r : N' \rightarrow N$  of  $N$ . Thus



$r^{-1}(U_q) \subset N'$  can be naturally identified with  $U_q \times S$  where  $S$  is the fibre of  $r$ . Further, without loss of generality we may identify  $U_q$  with an open Euclidean ball  $B$  in  $\mathbb{C}^n$ . In this identification, on each component  $U_q \times \{s\}$ ,  $s \in S$ , the path metric  $d$  on  $N'$  is equivalent to the Euclidean metric on  $B$  with the constants of equivalence independent of  $s$ .

Next, for some  $s \in S$  let us consider the restriction  $F'_s$  of  $F'$  to  $U_q \times \{s\} = B$ . We set  $M'_s := M' \cap (U_q \times \{s\})$  and  $bM'_s := bM' \cap (U_q \times \{s\})$ . Diminishing if necessary  $U_q$ , without loss of generality we may assume that these sets are connected. Also by  $dv$  we denote the Euclidean volume form on  $\mathbb{C}^n$ . By the constructions of  $\omega_{M'}$ , see Section 3.2, and  $f$ , see Section 3.3, we clearly have

$$(3.16) \quad f|_{U_q \times \{s\}} \geq c \quad \text{and} \quad \omega_{M'}^n|_{U_q \times \{s\}} \geq c \, dv$$

for some  $c > 0$  independent of  $s \in S$ .

Further, by the definition  $F'_s \in L_2(M'_s, E)$ . So by the choice of  $g$  in the definition of the hermitian metric on  $E$  using (3.16) we obtain

$$(3.17) \quad \int_{z \in M'_s} |F'_s(z)|^2 e^{\frac{1}{\text{dist}(z, bM'_s)}} \, dv(z) < \infty.$$

Without loss of generality we may assume that  $U_q \cap r(\text{supp } \eta) = \emptyset$ . Thus  $F'_s$  is holomorphic on  $M'_s$  for each  $s \in S$ . Now, from (3.17) using the mean-value property for the plurisubharmonic function  $|F'_s|^2$  defined on  $M'_s$  we easily obtain that for any  $y \in bM'_s$

$$(3.18) \quad \lim_{z \rightarrow y} F'_s(z) = 0.$$

Indeed, for a point  $z$  sufficiently close to  $y \in bM'_s$  consider a Euclidean ball  $B_z$  centered at  $z$  of radius  $r_z := \text{dist}(z, bM'_s)/2$ . Choosing  $z$  closer to  $y$  we may assume that  $B_z \subset \subset M'_s$ . Then by the triangle inequality for the metric  $d_N$  we have

$$\text{dist}(w, bM'_s) \leq 3r_z/2 \quad \text{for all } w \in B_z.$$

Now from (3.17) by the mean-value property we get for some  $c_n > 0$  depending on  $n$  only:

$$\begin{aligned} c_n r_z^{2n} e^{2/(3r_z)} |F'_s(z)|^2 &\leq e^{2/(3r_z)} \int_{w \in B_z} |F'_s(w)|^2 \, dv(w) \\ &\leq \int_{w \in B_z} |F'_s(w)|^2 e^{\frac{1}{\text{dist}(w, bM'_s)}} \, dv(w) \leq A < \infty. \end{aligned}$$

Hence,

$$\lim_{z \rightarrow y} |F'_s(z)|^2 \leq \lim_{z \rightarrow y} \frac{Ae^{-2/(3r_z)}}{c_n r_z^{2n}} = 0.$$

Thus (3.18) is true for any  $y \in bM'_s$ .

Next, since  $M'_s$  is connected, (3.18) implies that  $F'_s \equiv 0$  on  $M'_s$  for each  $s \in S$ . Actually, let  $z \in bM'_s$ . Consider a complex line  $l_z$  passing through  $z$  and containing the normal to  $bM'_s$  at  $z$  (recall that  $bM'_s$  is smooth). Then  $l_z$  intersects  $bM'_s$  transversely in a neighbourhood of  $z$  in  $bM'_s$ . This implies that there is a simply connected domain  $W_z \subset l_z \cap M'_s$  whose boundary  $bW$  contains  $z$  such that  $F'_s|_{\overline{W}_z} \in C(\overline{W}_z)$  and it equals 0 on an open subset of  $bW_z$ . Thus by the uniqueness property for univariate holomorphic functions we have  $F'_s = 0$  on  $W_z$ . Observe that if  $z$  varies along  $bM'_s$  the union of the connected components of  $l_z \cap M'_s$  containing  $W_z$  contains an open subset of  $M'_s$ . This implies that  $F'_s \equiv 0$  on  $M'_s$ .

Finally, taking a finite open cover of  $bM$  by the above sets  $U_q$  and using similar arguments we obtain the required neighbourhood  $U$  of  $bM$  (as the union of such  $U_q$  intersected with  $M$ ). This completes the proof of the lemma. □

Let us finish the proof of the theorem. As established above, the function  $F'$  satisfies conditions of Proposition 3.4. According to this proposition there is a number  $n \in \mathbb{N}$  independent of  $F'$  such that  $F' \cdot (r^*s_1)^n$  is extended to a continuous section of  $E'_n(M)$  equals 0 on  $r^{-1}(U)$ . Moreover,  $F'|_O \in \mathcal{L}_2(O')$  for any  $O \subset\subset M \setminus C_M$ .

We set

$$\tilde{F} := e^{F'} - 1 \quad \text{and} \quad \tilde{\eta} := \bar{\partial}\tilde{F} = \tilde{F}\eta.$$

By the definitions of  $\eta$  and  $F'$  we have  $\text{supp } \tilde{\eta} = \text{supp } \eta$  and  $\tilde{F}$  is bounded on  $\text{supp } \eta$ . In particular,  $\tilde{\eta}$  is  $\bar{\partial}$ -closed and belongs to  $L_2^{0,1}((M \setminus C_M)', E)$ , as well. Then by (3.6) there is a function  $\tilde{F}' \in L_2((M \setminus C_M)', E)$  such that  $\bar{\partial}\tilde{F}' = \tilde{\eta}$ . Applying to  $\tilde{F}'$  the same arguments as to  $F'$  we conclude that  $\tilde{F}'|_{r^{-1}(U)} \equiv 0$  for some neighbourhood  $U \subset M$  of  $bM$ . Since by the definition  $\tilde{F} - \tilde{F}'$  is holomorphic on  $(M \setminus C_M)'$  and equals zero on a neighbourhood of  $bM'$ , from the connectedness of  $M'$  we get  $\tilde{F} = \tilde{F}'$ . Also, as in the case of  $F'$ ,  $\tilde{F}' \cdot (r^*s_1)^n$  is extended to a continuous section of  $E_n(M')$ .

Let  $q$  be a regular point of  $C'_M := r^{-1}(C_M)$ . The above properties of  $F'$  and  $\tilde{F}$  imply that for suitable complex coordinates  $z = (z_1, \dots, z_n)$  in a

neighbourhood  $U_q$  of  $q$  we have  $C'_M \cap U_q = \{z_1 = 0\}$  and

$$e^{F'(z)} = z_1^{-n}A(z), \quad F'(z) = z_1^{-n}B(z), \quad z \in U_q \setminus C_M,$$

where  $A, B \in \mathcal{O}(U_q)$ . Suppose that  $A(z) = z_1^l A'(z)$  for some  $0 \leq l < n$  with  $A' \in \mathcal{O}(U_q)$  not identically 0 on  $C'_M \cap U_q$ . Then there is a point  $p \in C'_M \cap U_q$  and its neighbourhood  $W \subset U_q$  so that  $A'(z) \neq 0$  for all  $z \in \overline{W}$ . Thus we can introduce complex coordinates  $y = (y_1, \dots, y_n)$  on  $W$  by  $y_1 := z_1(A'(z))^{1/(n-l)}$ ,  $y_2 = z_2, \dots, y_n = z_n$ . In these coordinates we have  $e^{F'(y)} = y_1^{l-n}$ ,  $y \in W \setminus C_M$ . Since  $F' \in \mathcal{O}(W \setminus C_M)$ , the latter is impossible. This contradiction shows that  $l \geq n$  and so  $e^{F'}$  admits a holomorphic extension to  $U_q$ . From here we obtain easily that  $F'$  admits a holomorphic extension to  $U_q$ , as well.

Taking an open cover of regular points of  $C'_M$  by such neighbourhoods  $U_q$ , from the above arguments we obtain that  $F'$  is extended holomorphically to  $C'_M$  (it is extended to nonregular points of  $C'_M$  by the Hartogs theorem because the complex codimension of the set of such points in  $M'$  is  $\geq 2$ ).

Finally, the extended function  $F$  (i.e., the extension of  $F'$ ) belongs to  $\mathcal{L}_2(M')$ . Indeed, by the definition  $F|_{O'} \in \mathcal{L}_2(O')$  for every  $O \subset\subset N \setminus C_M$ . Assume now that  $q \in C_M$ . Let  $U$  be a simply connected coordinate chart of  $q$  with complex coordinates  $z = (z_1, \dots, z_n)$  such that  $z_1(q) = \dots = z_n(q) = 0$ ,  $C_M \cap M = \{z_1 \cdots z_n = 0\}$  and  $\overline{U} = \{z \in M : \max_{1 \leq k \leq n} |z_k| \leq 1\}$ . We identify  $\overline{U}$  with the unit polydisk in  $\mathbb{C}^n$  and by  $\mathbb{T}^n$  we denote its boundary torus. Also, we naturally identify  $(\overline{U})' \subset M'$  with  $\overline{U} \times S$  where  $S$  is the fibre of  $r : M' \rightarrow M$ . Diminishing, if necessary,  $U$  we will assume that  $F$  is holomorphic in a neighbourhood  $O' := r^{-1}(O)$  of  $(\overline{U})'$  where  $O$  is a neighbourhood of  $\overline{U}$ .

Let  $\{S_l\}_{l \in \mathbb{N}} \subset S$  be an increasing sequence of finite subsets of  $S$  such that  $\bigcup_l S_l = S$ . Then from the Cauchy integral formula we obtain

$$\begin{aligned} & \lim_{l \rightarrow \infty} \left( \sum_{s \in S_l} |F(y, s)|^2 \right) \\ & \leq \left( \frac{1}{2\pi} \right)^n \int_{x \in \mathbb{T}^n} \sum_{s \in S} \frac{|F(x, s)|^2}{(1 - |z_1(y)|) \cdots (1 - |z_n(y)|)} dx, \quad y \in U, \end{aligned}$$

where  $dx$  is the volume form on  $\mathbb{T}^n$ . Since  $\mathbb{T}^n \subset\subset M \setminus C_M$ ,  $F|_{\mathbb{T}^n} \in \mathcal{L}_2(\mathbb{T}^n \times S)$ . This implies that  $F|_{r^{-1}(y)} \in l_2(S)$  for all  $y \in V_q := \{z \in U : \max_{1 \leq k \leq n} |z_k| \leq 1/2\}$  and the  $l_2$  norms  $|\cdot|_y$  of these functions are uniformly bounded. Choosing a finite cover of  $C_M$  by such  $V_q$  and taking into

account that  $F|_{O'} \in \mathcal{L}_2(O')$  for every  $O \subset\subset N \setminus C_M$ , from the above we obtain that  $F \in \mathcal{L}_2(M')$ . Also,  $\bar{\partial}F = \eta$ .

This completes the proof of the theorem.  $\square$

#### REFERENCES

- [Bo] S. Bochner, *Analytic and meromorphic continuation by means of Green's formula*, Ann. of Math., **44** (1943), 652–673.
- [Br1] A. Brudnyi, *Representation of holomorphic functions on coverings of pseudoconvex domains in Stein manifolds via integral formulas on these domains*, J. Funct. Anal., **231** (2006), 418–437.
- [Br2] A. Brudnyi, *Holomorphic functions of slow growth on coverings of pseudoconvex domains in Stein manifolds*, Compositio Math., **142** (2006), 1018–1038.
- [Br3] A. Brudnyi, *Hartogs type theorems on coverings of Stein manifolds*, Internat. J. Math., **17** (2006), no. 3, 339–349.
- [Br4] A. Brudnyi, *On holomorphic  $L^2$ -functions on coverings of strongly pseudoconvex manifolds*, Publications of RIMS, Kyoto University, **43** (2007), no. 4, 963–976.
- [C] H. Cartan, *Sur les fonctions de plusieurs variables complexes. Les espaces analytiques*, Proc. Intern. Congress Mathematicians Edinburgh 1958, Cambridge Univ. Press, 1960, pp. 33–52.
- [D] J.-P. Demailly, *Estimations  $L^2$  pour l'opérateur  $\bar{\partial}$  d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété kahlérienne complète*, Ann. Sci. Ecole Norm. Sup. (4), **15** (3) (1982), 457–511.
- [Fe] H. Federer, *Geometric measure theory*, Springer-Verlag, New York, 1969.
- [GHS] M. Gromov, G. Henkin and M. Shubin, *Holomorphic  $L^2$  functions on coverings of pseudoconvex manifolds*, GAFA, Vol. **8** (1998), 552–585.
- [GM] C. Grant and P. Milman, *Metrics for singular analytic spaces*, Pacific J. Math., **168** (1995), no. 1, 61–156.
- [GR] H. Grauert and R. Remmert, *Theorie der Steinschen Räume*, Springer-Verlag, Berlin, 1977.
- [HL] R. Harvey and H. B. Lawson, *On boundaries of complex analytic varieties, I*, Ann. of Math. (2), **102** (1975), no. 2, 223–290.
- [H] G. Henkin, *The method of integral representations in complex analysis*, Several complex variables, I, Introduction to complex analysis, A translation of *Sovremennyye problemy matematiki. Fundamental'nye napravleniya, Tom 7*, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekn. Inform., Moscow 1985. Encyclopaedia of Mathematical Sciences, 7, Springer-Verlag, Berlin, 1990.
- [KR] J. J. Kohn and H. Rossi, *On the extension of holomorphic functions from the boundary of a complex manifold*, Ann. of Math. (2), **81** (1965), 451–472.
- [L] F. Lárusson, *An extension theorem for holomorphic functions of slow growth on covering spaces of projective manifolds*, J. Geom. Anal., **5** (1995), no. 2, 281–291.
- [M] E. McShane, *Extension of range functions*, Bull. Amer. Math. Soc., **40** (1934), no. 12, 837–842.

- [N] R. Narasimhan, *Imbedding of holomorphically complete complex spaces*, Amer. J. Math., **82** (1960), no. 4, 917–934.
- [O] T. Ohsawa, *Complete Kähler manifolds and function theory of several complex variables*, Sugaku Expositions, **1** (1) (1988), 75–93.
- [R] R. Remmert, *Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes*, C. R. Acad. Sci. Paris, **243** (1956), 118–121.

*Department of Mathematics and Statistics  
University of Calgary  
Calgary  
Canada*