

## MILLIMETER-WAVE OBSERVATIONS OF PLANETARY NEBULAE

Hugh M. Johnson  
Lockheed Missiles and Space Co.  
Palo Alto, California

Observations of the integrated continuum flux density of several stellar planetary nebulae have been made at 90 GHz. These are reported, with some remarks on their significance for objects with positive radio-frequency spectral index, and to display graphically their relation to published microwave and near-infrared data. The 90-GHz data have been obtained with the 11-m dish of the National Radio Astronomy Observatory\* on Kitt Peak, under the Lockheed Independent Research Program. Three of the objects have been included among a dozen stellar planetary nebulae observed with the Fabry-Perrot interferometer at the Cerro Tololo Inter-American Observatory† so as to provide kinematical information with the microwave spectral data, since they are related in the theories of the positive spectral index that appeared in 1975. The Fabry-Perrot observations are reported in Astrophys. J. 1977 September 15. No other journal article is yet in press.

\*The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under contract with the National Science Foundation.

†Visiting Astronomer, Cerro Tololo Inter-American Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation.

## THE "BUTTERFLY" NEBULA M2-9: ITS POSSIBLE RELATION TO B[e] STARS AND PROTO-PLANETARIES

Y. Andrillat and J.P. Swings  
Observatoire de Haute Provence, France and Institut d'Astro-  
Physique, Université de Liège, Belgium, respectively.

M2-9 is not a bona fide planetary, nor simply a star: it most probably is a planetary nebula in the formative stages. High densities, the presence of [OI], FeII and [FeII] emission lines, and prominent infrared continua are characteristic of such objects. There is a striking progression of their spectra from B[e] stars such as HD45677 to