

# The role of L-carnitine in the control of oxidative stress and lipid $\beta$ -oxidation during *in vitro* follicle growth, oocyte maturation, embryonic development and cryopreservation: a review

## Review Article

**Cite this article:** Leitão AMF *et al.* (2024) The role of L-carnitine in the control of oxidative stress and lipid  $\beta$ -oxidation during *in vitro* follicle growth, oocyte maturation, embryonic development and cryopreservation: a review. *Zygote*, **32**: 335–340. doi: [10.1017/S096719942400039X](https://doi.org/10.1017/S096719942400039X)




Received: 15 March 2024  
Revised: 19 September 2024  
Accepted: 3 October 2024  
First published online: 7 November 2024

### Keywords:

cryopreservation; embryo; *in vitro* culture; L-carnitine; oocyte

### Corresponding author:

Jose Roberto V. Silva; Email: [jrsilva@ufc.br](mailto:jrsilva@ufc.br)

Allana Maria Freire Leitão<sup>1</sup> , Bianca Regia Silva<sup>1</sup>, Efigênia C. Barbalho<sup>1</sup>,  
Lais R.M. Paulino<sup>1</sup>, Francisco das Chagas Costa<sup>1</sup> , Fabricio Sousa Martins<sup>2</sup> and  
Jose Roberto V. Silva<sup>1</sup> 

<sup>1</sup>Laboratory of Biotechnology and Physiology of Reproduction (LABIREP), Federal University of Ceara, Av. Comandante Maurocêlio Rocha Ponte 100, postal code 62041-040, Sobral, CE, Brazil and <sup>2</sup>State University of Acaraú Valley, Center of Agricultural and Biological Sciences, postal code 62040370, Sobral, CE, Brazil

### Abstract

L-carnitine has an important role in the control of oxidative stress and lipid  $\beta$ -oxidation during *in vitro* culture and cryopreservation of ovarian follicles, oocytes and embryos. This substance balances the acetyl-CoA/CoA ratio, maintains glucose metabolism and increases energy production in mitochondria. It also plays a key role in reducing endoplasmic reticulum stress, by transferring palmitate to mitochondria or eliminating it to avoid toxicity. By eliminating reactive oxygen species, L-carnitine increases the percentages of mature oocytes with uniform mitochondrial distribution and improves embryo post-thaw cryotolerance. Therefore, L-carnitine controls lipid  $\beta$ -oxidation and oxidative stress during *in vitro* culture of ovarian follicles, oocyte maturation, embryonic development and cryopreservation.


## Introduction

The improvements in *in vitro* culture systems for ovarian follicles, oocytes and embryos are of great relevance to increase efficiency of assisted reproduction techniques in mammalian species. However, lipid peroxidation and imbalance in the production and elimination of reactive oxygen species (ROS) represent the main barriers to having healthy oocytes and embryos after *in vitro* culture and cryopreservation (Soto-Heras and Paramio, 2020). In this sense, the addition of natural substances to the culture media has been an alternative to control the damages caused by excessive ROS (Paulino *et al.*, 2022). L-carnitine is a water-soluble vitamin-like compound that is naturally produced and synthesized primarily from lysine and methionine in the liver to improve lipid breakdown and generate metabolic (Modak *et al.*, 2022). According to Carrillo-González *et al.* (2023), lipids are the most abundant reservoir of energy in bovine embryos, and triacylglycerol-containing lipid droplets represent the main stocks of fatty acids in oocytes. These authors showed that L-carnitine mobilizes fatty acids from oocyte cytoplasm to mitochondria, which results in  $\beta$ -oxidation and generation of energy. Additionally, acetyl-L-carnitine exhibits antioxidant effects and has beneficial effects on reproductive functions (Liu *et al.*, 2004; Cheng and Chen, 2008; Aliabadi *et al.*, 2012; Agarwal *et al.*, 2018). When administered exogenously, acetyl-L-carnitine has higher bioavailability than L-carnitine and regulates even the production of reproduction-associated hormones (Agarwal *et al.*, 2018).

This review aims to show the role of L-carnitine on hypothalamus-pituitary-gonad-axis and to discuss its influence on lipid  $\beta$ -oxidation and oxidative stress during *in vitro* culture of ovarian follicles, oocyte maturation, embryo development and cryopreservation.

## Oxidative stress

Free radicals are chemical specimens that have at least one unpaired electron in their outer orbitals, being highly reactive (Prevedello and Comachio, 2021). This characteristic enables the transfer of electrons between neighbouring molecules, causing changes in the molecular environment (Ferreira *et al.*, 2020; Martelli and Nunes, 2014). The ROS are naturally produced by cellular metabolism and play an important physiological role, being involved in several processes, such as energy production, phagocytosis, intercellular signalling, regulation of cell growth, immunity, cell defence and synthesis of biological substances (Prevedello and Comachio, 2021). However, when ROS production exceeds its degradation, it causes oxidative stress, being responsible for various damages to DNA, proteins and phospholipids in different cell types (Simas *et al.*, 2019). Controlling the production and neutralization of ROS is crucial for



**Figure 1.** Effects of L-carnitine on hypothalamus-pituitary gonad axis.


maintaining cellular integrity. *In vivo*, this control is performed through enzymatic and non-enzymatic antioxidant systems. Various endogenous enzymes, like catalase (CAT), peroxiredoxins (PRDX), superoxide dismutase (SOD) and glutathione reductase/peroxidase (GPX) constitute the endogenous antioxidant system (Souza *et al.*, 2020), which are capable of inactivating the harmful effects of free radicals. The non-enzymatic system includes low molecular weight compounds such as L-carnitine, ascorbic acid, tocopherol, selenium, zinc, taurine, hypotaurine, carotene, lipoic acid and other thiol compounds such as cystine, cysteine, cysteamine and beta-mercaptoethanol (Crocomo *et al.*, 2012).

During *in vitro* culture of different types of cells, the reduction of endogenous antioxidant protection linked to other factors, such as exposure to light and high concentrations of oxygen, favours a significant increase in ROS production (Alves *et al.*, 2019; Sadeesh *et al.*, 2014) and oxidative stress, which has been reported as one of the main limitations of *in vitro* culture of various types of cells (Del Collado *et al.*, 2017; Soto-Heras and Paramio, 2020). In excess, oxidative stress in granulosa cells results in follicular atresia (Saeed-Zidane *et al.*, 2017) and has been reported as one of the main factors associated with poor quality of cultured ovarian follicles (Sá *et al.*, 2018; Paulino *et al.*, 2022). Due to damages caused by oxidative stress during *in vitro* culture, several studies have sought to develop protocols to minimize it (Cordeiro *et al.*, 2023; Nascimento *et al.*, 2022).

#### Effects of L-carnitine on hypothalamus-pituitary-gonadal axis

The L-carnitine influences the hypothalamus-pituitary-gonad axis and upregulates gonadotropin-releasing hormone (GnRH) secretion from the hypothalamus and, consequently, induce depolarization of hypothalamic neuronal cells to increase secretory activity (Agarwal *et al.*, 2018; Krsmanovic, *et al.*, 1994). Also, L-carnitine has been reported to increase the levels of other hormones, like luteinizing hormone (LH), progesterone and oestradiol, while it decreases prolactin secretion in mammalian species (Agarwal *et al.*, 2018; Genazzani *et al.*, 2011; Krsmanovic *et al.*, 1992). Figure 1 illustrates the effects of L-carnitine on the reproductive system of mammalian females.

The L-carnitine and its primary ester have direct effects against oxidative stress, minimizing cell death by apoptosis and maintaining cellular energy (Agarwal *et al.*, 2018; Abdelrazik *et al.*, 2009; Infante *et al.*, 2002; Vanella *et al.*, 2000). To minimize oxidative stress, L-carnitine can also be used in combination with other antioxidant commonly known for quenching free radicals such as vitamins (C, E and  $\beta$ -carotene), and some metalloenzymes, including GPx, CAT and SOD (Nimse and Pal, 2015). Thus, due to its energy generation property combined with its antioxidant property, L-carnitine has been studied for use in reproductive technologies, including *in vitro* culture of ovarian



**Figure 2.** Direct effects of L-carnitine on oocytes of mammals.

follicles, *in vitro* maturation, *in vitro* embryo production and cryopreservation.

#### Effects of L-carnitine on *in vitro* follicle development and oocyte maturation

The role of L-carnitine in ovarian follicles *in vitro* is still little explored. Dunning and Robker (2012) reported that L-carnitine did not alter survival, growth or differentiation of mouse secondary follicles *in vitro*. However, it significantly increased  $\beta$ -oxidation and markedly improved fertilization rate and blastocyst development. Recently, Modak *et al.* (2022) reported that L-carnitine increased the rate of oocytes in metaphase II (MII) stage from early antral follicles cultured *in vitro*. Furthermore, the presence of L-carnitine decreased the rate of degeneration and even promoted the formation of structures similar to antrum after the *in vitro* culture of buffalo oocyte granulosa complexes.

In mouse oocytes, L-carnitine acts through the electrogenic force of voltage-gated  $\text{Na}^+$  channels and it is transported by  $\text{Na}^+$ /organic cationic transporter-2 (OCTN-2) to oocytes (Infante *et al.*, 2002; Dunning and Robker 2012). In the oocyte, L-carnitine is converted to acetyl-L-carnitine by carnitine palmitoyltransferase-I (CPT-I) in the mitochondria and can act on the endoplasmic reticulum, mitochondria and even in ooplasm (Mingorance *et al.*, 2011) (Figure 2). Various studies have shown that L-carnitine optimizes glucose metabolism by transferring fatty acids to the mitochondria and facilitating  $\beta$ -oxidation since lipid metabolism is one of the primary regulators of oocyte maturation (Stojkovic *et al.*, 2001; Dunning *et al.*, 2010; Agarwal *et al.*, 2018). Within the oocyte, L-carnitine is converted to acetyl-L-carnitine and keeps glucose metabolism through the citric acid cycle and, consequently, increases energy production (Infante *et al.*, 2002; Agarwal *et al.*, 2018) (Figure 2). The L-carnitine reduces pyruvate entry into the citric acid cycle and transports palmitate and other long-chain fatty acids to facilitate their utilization through  $\beta$ -oxidation

(Dunning *et al.*, 2010) (Figure 2). The L-carnitine reduces the levels of palmitate in the endoplasmic reticulum by transferring it to mitochondria or by eliminating it from where it can cause oocyte lipotoxicity by oxidative stress (Agarwal *et al.*, 2018). L-carnitine increases the proportion of mature oocytes with uniform mitochondrial distribution and supports *in vitro* oocyte maturation and embryonic development in mice and pigs (Zare *et al.*, 2015; Somfai *et al.*, 2011). Chankitisakul *et al.* (2013) showed that L-carnitine increases the rate of bovine embryo production after *in vitro* maturation and subsequent vitrification of oocytes. Marin *et al.* (2020) also reported that L-carnitine increased oocyte competence during buffalo oocyte maturation in the absence of foetal bovine serum.

*In vivo*, L-carnitine can stabilize the mitochondrial membrane and protect DNA against ROS-induced damage in oocytes of women with polycystic ovary syndrome (PCOS) (Mohd Shukri *et al.*, 2022; Ismail *et al.*, 2014; Fenkci, *et al.*, 2008). Using a mouse model of PCOS, oral administration of acetyl-L-carnitine alleviated ovarian dysfunction associated with that syndrome through its antioxidant/glycation activity and mitochondria potentiation (Di Emidio *et al.*, 2020).

#### Effects of L-carnitine on *in vitro* embryo development

The ROS may originate in the embryo itself or from exogenous sources. During oocyte *in vitro* fertilization (IVF), strategies to reduce ROS production, such as addition of free radical scavengers and lowering the oxygen tension are important for improving the fertility potential in assisted reproductive technologies (Agarwal *et al.*, 2014). The ROS are involved in defective embryo development and retardation of embryo growth and induce cell membrane damage, DNA damage and apoptosis (Volpe *et al.*, 2018). Apoptosis results in fragmented embryos, which have limited potential to implant and therefore, result in poor fertility rates (Agarwal *et al.*, 2014).

The antioxidant capacity of L-carnitine might account for its preferential use to improve *in vitro* oocyte and embryo development. The treatment of porcine embryos with antioxidants improved blastocyst production (Castillo-Martín, *et al.*, 2014). Moreover, the presence of L-carnitine in the culture medium was associated with increased cleavage and blastocyst rates in porcine species (Lowe *et al.*, 2017). Finally, the supplementation of L-carnitine to bovine embryo culture medium has been shown to scavenge ROS within two-cell stage embryos (Takahashi *et al.*, 2013).

#### Effects of L-carnitine on cryopreservation of ovarian follicles, oocytes and embryos

The cryopreservation technique allows the sub-zero storage of tissues or cells by dramatically reducing natural cellular biochemical processes for extended periods of time (Vining *et al.*, 2021). Currently, cryopreservation is a modern and safe method which assists in the preservation of genetic material from follicles, oocytes and embryos in human and animals (Sekhon *et al.*, 2018), but the cells can still be seriously damaged during cryopreservation (Truong, *et al.*, 2022; Spijkers *et al.*, 2017; Barsky *et al.*, 2016; Beyer and Griesinger, 2016). Unfortunately, frozen follicles, oocytes and embryos are still reported to contain a higher proportion of apoptotic cells compared to their non-frozen counterparts, with freezing procedures generally associated with triggering apoptotic cell death (Vining, *et al.*, 2021). Exposure to high concentrations of cryoprotectants, osmolarity and rapid

temperature changes during cryopreservation have been shown to affect gamete and embryo physiology (Somoskoi *et al.*, 2015; Dalcin *et al.*, 2013), as well as their gene expression (Sahraei *et al.*, 2018; Monzo *et al.*, 2012). These deleterious effects are strongly associated with the occurrence of oxidative stress during cryopreservation.

Production of ROS during the vitrification of gametes may be a crucial mediator of damage to proteins and DNA (Costa *et al.*, 2022; Zhang *et al.*, 2020). Disturbances in the oxidative metabolism and damage in cell membranes are other important stress factors related to vitrification. Together, these effects decrease glutathione (GSH) levels, alter expression of regulatory genes and are associated with decreasing maturation rate and developmental competence of follicles, oocytes and embryos after cryopreservation (Berteli *et al.*, 2022; Zare *et al.*, 2022; Costa *et al.*, 2022; Wu *et al.*, 2019; Pan *et al.*, 2018). Furthermore, cryopreservation of oocytes or embryos has been reported to cause mitochondrial dysfunction, such as changes in membrane potential and reduced adenosine triphosphate (ATP) production (Gualtieri *et al.*, 2021; Iwata, 2021). However, the detrimental effects of cumulative stress have been shown to be partly improved by adding antioxidants in vitrification media, such as L-carnitine. Some reports have already demonstrated that L-carnitine plays an important role in attenuating the deleterious effects of oxidative stress on cryopreserved follicles. For instance, Zhang *et al.* (2015) observed lower rates of apoptosis and malondialdehyde, as well as higher levels of oestradiol in mice ovarian follicles cryopreserved *in situ*. These results were translated into increased follicular survival. Zolini *et al.* (2019) demonstrated that the addition of L-carnitine in embryo culture medium improved post-thaw cryotolerance but had no effect on pregnancy and implantation rate after transfer of cryopreserved bovine embryos.

The L-carnitine is well known for its role in  $\beta$ -oxidation, ATP production and decreasing the lipid content during embryo development, providing improved cryo-survivability (Truong *et al.*, 2016). In buffaloes, the addition of L-carnitine to the medium significantly benefits embryonic developmental competence after vitrification, as evidenced by the high cleavage rate and the formation of morulae and blastocysts. Improving the cryotolerance of buffalo embryos directly after thawing may be through increased lipid metabolism (El-Sokary *et al.*, 2021). Furthermore, L-carnitine acts as an antioxidant blocking degenerative changes arising from oxidative stress during embryonic development (Bhakty *et al.*, 2021). Lowe *et al.* (2017) reported that the antioxidant capacity of L-carnitine was associated with the increased cleavage rate and the improved cryotolerance of resultant porcine blastocysts. Supplementation of L-carnitine to bovine embryo culture medium has been shown to scavenge ROS within two-cell stage embryos, antagonize the cryodamage and enhance the cryotolerance of blastocysts (Takahashi *et al.*, 2013). It is plausible to suggest that L-carnitine may be used to improve the freezing survival of oocytes or embryos (Li *et al.*, 2023). Table 1 shows some effects of L-carnitine during *in vitro* culture of follicles, oocytes and embryos in different species.

#### Final considerations

Many factors inherent to the oocyte itself and *in vitro* culture environment determine the chance of having a complete follicular development with success in the acquisition of oocyte competence *in vitro*. A culture system, with different combinations of hormones, growth factors and mainly antioxidant factors, at each



**Table 1.** Effects of L-carnitine during *in vitro* culture of follicles, oocytes and embryos in different species

Effects	Species	Reference
<b>Ovarian follicles</b>		
Inhibits apoptosis, alleviates oxidative damage and increases the survival and function of follicles cryopreserved <i>in situ</i> .	Murine	Zhang et al., 2015.
<b>Oocytes</b>		
Exposure of oocyte to L-carnitine prior to insemination increase cleavage and improve cryotolerance of the resulting blastocysts.	Porcine	Lowe et al., 2017.
Increases the percentage of oocytes reaching metaphase II.	Ovine	Bhakty et al., 2021
Supports <i>in vitro</i> oocyte maturation and embryonic development.	Porcine	Hashimoto, 2008
Increases the proportion of mature oocytes with uniform mitochondrial distribution during <i>in vitro</i> maturation.	Murine	Zare et al., 2015; Somfai et al., 2011
L-carnitine during <i>in vitro</i> maturation and subsequent oocyte vitrification improves the <i>in vitro</i> production rate of embryos.	Bovine	Chankitisakul et al., 2013
Increases the rate of nuclear maturation up to metaphase II (MII) stage, decreases the rate of degeneration and promotes the antrum formation.	Buffalo	Modak et al., 2022
Increases oocyte competence during <i>in vitro</i> maturation and in the absence of FBS.	Buffalo	Marin et al., 2020
<b>Embryos</b>		
Increases lipid metabolism, improves development and cryo- tolerance.	Bovine	Takahashi et al., 2013.
Improves post-thawing cryotolerance in cryopreserved embryos.	Bovine	Zolini et al., 2019.
In combination with N-acetylcysteine and $\alpha$ -lipoic acid, L-carnitine increases the number of cells in blastocyst.	Murine	Truong et al., 2016.
Increases the cleavage rate, as well as the formation of morulae and blastocysts.	Buffalo	El-Sokary et al., 2021.

stage of growth, is necessary to allow the follicles to present an adequate size in a long-term culture period. This review shows that L-carnitine can be used to regulate oxidative stress and lipid  $\beta$ -oxidation during *in vitro* culture of ovarian follicles, oocyte maturation, embryo production and cryopreservation, especially due to stimulation energy generation combined with its antioxidant properties.

**Competing interests.** The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

**Funding.** This work was supported by the National Council for Scientific and Technological Development (CNPq, Brazil, Grant No. 407992/2021-9) and Coordination for the Improvement of Higher Education Personnel (CAPES).

## References

- Abdelrazik, H., Sharma, R., Mahfouz, R. and Agarwal, A. (2009) L-carnitine decreases DNA damage and improves the *in vitro* blastocyst development rate in mouse embryos. *Fertility and Sterility* **91**(2), 589–596. doi: [10.1016/j.fertnstert.2007.11.067](https://doi.org/10.1016/j.fertnstert.2007.11.067)
- Agarwal, A., Durairajanayagam, D. and du Plessis, S. S. (2014) Utility of antioxidants during assisted reproductive techniques: an evidence based review. *Reproductive Biology and Endocrinology*, **12**, 112. doi: [10.1186/1477-7827-12-112](https://doi.org/10.1186/1477-7827-12-112).
- Agarwal, A., Sengupta, P. and Durairajanayagam, D. (2018) Role of L-carnitine in female infertility. *Reproductive Biology and Endocrinology* **16**(1), 1–18. doi: [10.1186/s12958-018-0323-4](https://doi.org/10.1186/s12958-018-0323-4)
- Aliabadi, E., Soleimani Mehranjani, M., Borzoei, Z., Talaei-Khozani, T., Mirkhani, H. and Tabesh, H. (2012) Effects of L-carnitine and L-acetylcarnitine on testicular sperm motility and chromatin quality. *Iranian Journal of Reproductive Medicine* **10**(2), 77–82.
- Alves, C. S., Furtado, R. A., Nascetes, G. A. N., Rumpf, R. and Tavares, D. C. (2019) Evaluation of melatonin effects on the production of bovine embryos

- obtained by *in vitro* fertilization and somatic cell nuclear transfer. *Revista Brasileira de Reprodução Animal* **43**(4), 815–823. doi: [10.5555/20203153822](https://doi.org/10.5555/20203153822)
- Barsky, M., St Marie, P., Rahil, T., Markenson, G. R. and Sites, C. K. (2016) Are perinatal outcomes affected by blastocyst vitrification and warming? *American Journal of Obstetrics and Gynecology* **215**(5), 603. doi: [10.1016/j.ajog.2016.06.002](https://doi.org/10.1016/j.ajog.2016.06.002)
- Berteli, T. S., Vireque, A. A., Da Luz, C. M., Borges, E. D., Ferreira, C. R. and Navarro, P. A. (2022) Equilibration solution composition and extended exposure to equilibration phase affect embryo development and lipid profile of mouse oocytes. *Reproductive Biomedicine Online* **44**(6), 961–975. doi: [10.1016/j.rbmo.2022.01.006](https://doi.org/10.1016/j.rbmo.2022.01.006)
- Beyer, D. A. and Griesinger, G. (2016) Vitrified-warmed embryo transfer is associated with mean higher singleton birth weight compared to fresh embryo transfer. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* **203**, 104–107. doi: [10.1016/j.ejogrb.2016.05.041](https://doi.org/10.1016/j.ejogrb.2016.05.041)
- Bhakty, Z. W., Kaiine, M., Karjan, W. K. and Setiadim, A. (2021) L-carnitine supplementation enhances nuclear and cytoplasmic maturation rates of sheep oocytes. *In vitro Tropical Animal Science Journal* **44**(2), 131–137. doi: [10.5398/tasj.2021.44.2.131](https://doi.org/10.5398/tasj.2021.44.2.131)
- Carrillo-González, D. F., Hernández-Herrera, D. Y. and Maldonado-Estrada, J. G. (2023) The role of L-carnitine in bovine embryo metabolism. A review of the effect of supplementation with a metabolic modulator on *in vitro* embryo production. *Animal Biotechnology* **34**(2), 413–423. doi: [10.1080/10495398.2021.1938593](https://doi.org/10.1080/10495398.2021.1938593)
- Castillo-Martín, M., Bonet, S., Morató, R. and Yeste, M. (2014) Comparative effects of adding  $\beta$ -mercaptoethanol or L-ascorbic acid to culture or vitrification-warming media on IVF porcine embryos. *Reproduction, Fertility, and Development* **26**(6), 875–882. doi: [10.1071/RD13116](https://doi.org/10.1071/RD13116)
- Chankitisakul, V., Somfai, T., Inaba, Y., Techakumphu, M. and Nagai, T. (2013) Supplementation of maturation medium with L-carnitine improves cryo-tolerance of bovine *in vitro* matured oocytes. *Theriogenology* **79**(4), 590–598. doi: [10.1016/j.theriogenology.2012.11.011](https://doi.org/10.1016/j.theriogenology.2012.11.011)
- Cheng, H. J. and Chen, T. (2008) Clinical efficacy of combined L-carnitine and acetyl-L-carnitine in idiopathic asthenospermia. *Zhonghua nan ke xue= National Journal of Andrology* **14**(2), 149–151.

- Cordeiro, E., Silva, B., Paulino, L., Barroso, P., Barrozo, L., de Lima Neto, M. and Silva, J. (2023) Effects of N-acetylcysteine on growth, viability and reactive oxygen species levels in small antral follicles cultured *in vitro*. *Asian Pacific Journal of Reproduction* 12(1), 42–48. doi: [10.4103/2305-0500.365231](https://doi.org/10.4103/2305-0500.365231)
- Costa, F. C., Vasconcelos, E. M., Azevedo, V. A. N., de Assis, E. I. T., Paulino, L. R. F. M., Silva, A. W. B., SILVA, J. R. V. and Batista, A. L. P. S. (2022) Aloe vera increases collagen fibres in extracellular matrix and mRNA expression of peroxiredoxin-6 in bovine ovarian cortical tissues cultured *in vitro*. *Zygote*, 30(3), 365–372. doi: [10.1017/S0967199421000824](https://doi.org/10.1017/S0967199421000824)
- Crocorno, L. F., Marques Filho, W. C., Landim-Alvarenga, F. D. C. and Bicudo, S. D. (2012) *In vitro* embryo production: oxidative stress and antioxidants. *Veterinária e Zootecnia* 19(4), 470–479. doi: [10.5555/20133263819](https://doi.org/10.5555/20133263819)
- Dalcin, L., Silva, R. C., Paulini, F., Silva, B. D., Neves, J. P. and Lucci, C. M. (2013) Cytoskeleton structure, pattern of mitochondrial activity and ultrastructure of frozen or vitrified sheep embryos. *Cryobiology* 67(2), 137–145. doi: [10.1016/j.cryobiol.2013.05.012](https://doi.org/10.1016/j.cryobiol.2013.05.012)
- Del Collado, M., da Silveira, J. C., Oliveira, M. L., Alves, B. M., Simas, R. C., Godoy, A. T., Coelho, M. B., Marques, L. A., Carriero, M. M., Nogueira, M. F. G., Eberlin, M. N., Silva, L. A., Meireles, F. V. and Perecin, F. (2017) *In vitro* maturation impacts cumulus–oocyte complex metabolism and stress in cattle. *Reproduction* 154(6), 881–893. doi: [10.1530/REP-17-0134](https://doi.org/10.1530/REP-17-0134)
- Di Emídio, G., Rea, F., Placidi, M., Rossi, G., Cocciolone, D., Virmani, A., Macchiarelli, G., Palmerini, M. G., D'Alessandro, M. A., Artini P. G. and Tatone, C. (2020) Regulatory functions of L-Carnitine, acetyl, and propionyl L-Carnitine in a PCOS mouse model: focus on antioxidant/antiglycative molecular pathways in the ovarian microenvironment. *Antioxidants* 9(9), 867. doi: [10.3390/antiox9090867](https://doi.org/10.3390/antiox9090867)
- Dunning, K. R., Cashman, K., Russell, D. L., Thompson, J. G., Norman, R. J. and Robker, R. L. (2010) Beta-oxidation is essential for mouse oocyte developmental competence and earl embryo development. *Biology of Reproduction* 83(6), 909–918. doi: [10.1095/biolreprod.110.084145](https://doi.org/10.1095/biolreprod.110.084145)
- Dunning, K. R. and Robker, R. L. (2012) Promoting lipid utilization with l-carnitine to improve oocyte quality. *Animal Reproduction Science* 134(1–2), 69–75. doi: [10.1016/j.anireprosci.2012.08.013](https://doi.org/10.1016/j.anireprosci.2012.08.013)
- El-Sokary, M. M. M., El-Naby, A. A. H., Hameed, A. R. A. E., Mahmoud, K. G. M. and Scholkamy, T. H. (2021) Impact of L-carnitine supplementation on the *in vitro* developmental competence and cryotolerance of buffalo embryos. *Veterinary World* 14(12), 3164–3169. doi: [10.14202/vetworld.2021.3164-3169](https://doi.org/10.14202/vetworld.2021.3164-3169)
- Fenkci, S. M., Fenkci, V., Oztekin, O., Rota, S. and Karagenc, N. (2008) Serum total L- carnitine levels in non-obese women with polycystic ovary syndrome. *Human Reproduction (Oxford, England)* 23(7), 1602–1606. doi: [10.1093/humrep/den109](https://doi.org/10.1093/humrep/den109)
- Ferreira, J. G. S., da Silva Ferreira, V. V., de Almeida Costa, F., de Lima Santos, I. L. V. and da Silva, C. R. C. (2020) Envelhecimento e a influência degenerativa dos radicais livres nesse processo. Campina Grande.
- Genazzani AD., Lanzoni C., Ricchieri F., Santagni S., Rattighieri E., Chierchia E., Monteleone P. and Jassoni VM (2011) Acetyl-L-carnitine (ALC) administration positively affects reproductive axis in hypogonadotropic women with functional hypothalamic amenorrhea. *Journal of Endocrinological Investigation* 34, 287–91. doi: [10.1007/BF03347087](https://doi.org/10.1007/BF03347087)
- Gualtieri, R., Kalthur, G., Barbato, V., Di Nardo, M., Adiga, S. K. and Talevi, R. (2021) Mitochondrial Dysfunction and Oxidative Stress Caused by Cryopreservation in Reproductive Cells. *Antioxidants (Basel, Switzerland)* 10(3), 337. doi: [10.3390/antiox10030337](https://doi.org/10.3390/antiox10030337)
- Hashimoto, S. (2008) L-Carnitine decreased the apoptosis of granulosa cells and improved the meiotic competence of porcine growing oocytes. *Reproduction Domestic Animal* 43, 190–191. doi: [10.3390%2Fani12151957](https://doi.org/10.3390%2Fani12151957)
- Infante, J. P., Tschanz, C. L., Shaw, N., Michaud, A. L., Lawrence, P. and Brenna, J. T. (2002) Straight-chain acyl-CoA oxidase knockout mouse accumulates extremely long chain fatty acids from alpha-linolenic acid: evidence for runaway carousel-type enzyme kinetics in peroxisomal beta-oxidation diseases. *Molecular Genetics and Metabolism* 75(2), 108–119. doi: [10.1006/mgme.2001.3279](https://doi.org/10.1006/mgme.2001.3279)
- Ismail, A. M., Hamed, A. H., Saso, S. and Thabet, H. H. (2014) Adding L-carnitine to clomiphene resistant PCOS women improves the quality of ovulation and the pregnancy rate. A randomized clinical trial. *European Journal of Obstetrics, Gynecology, and Reproductive Biology* 180, 148–152. doi: [10.1016/j.ejogrb.2014.06.008](https://doi.org/10.1016/j.ejogrb.2014.06.008)
- Iwata H. (2021) Resveratrol enhanced mitochondrial recovery from cryopreservation-induced damages in oocytes and embryos. *Reproductive Medicine and Biology* 20(4), 419–426. doi: [10.1002/rmb2.12401](https://doi.org/10.1002/rmb2.12401)
- Krsmanović, L. Z., Virmani, M. A., Stojilković, S. S. and Catt, K. J. (1992) Actions of acetyl- L-carnitine on the hypothalamo-pituitary-gonadal system in female rats. *The Journal of Steroid Biochemistry and Molecular Biology* 43(4), 351–358. doi: [10.1016/0960-0760\(92\)90170-n](https://doi.org/10.1016/0960-0760(92)90170-n)
- Krsmanovic, L. Z., Virmani, M. A., Stojilkovic, S. S. and Catt, K. J. (1994) Stimulation of gonadotropin-releasing hormone secretion by acetyl-L-carnitine in hypothalamic neurons and GT1 neuronal cells. *Neuroscience Letters* 165(1–2), 33–36. doi: [10.1016/0304-3940\(94\)90702-1](https://doi.org/10.1016/0304-3940(94)90702-1)
- Li, X., Wu, X., Ma, T., Zhang, Y., Sun, P., Qi, D. and Ma, H. (2023) Protective effect of L-carnitine against oxidative stress injury in human ovarian granulosa cells. *Experimental and Therapeutic Medicine* 25(4), 1–11. doi: [10.3892/etm.2023.11860](https://doi.org/10.3892/etm.2023.11860)
- Liu, J., Head, E., Kuratsune, H., Cotman, C. W. and Ames, B. N. (2004) Comparison of the effects of L-carnitine and acetyl-L-carnitine on carnitine levels, ambulatory activity, and oxidative stress biomarkers in the brain of old rats. *Annals of the New York Academy of Sciences*, 1033(1), 117–131. doi: [10.1196/annals.1320.011](https://doi.org/10.1196/annals.1320.011)
- Lowe, J. L., Bartolac, L. K., Bathgate, R. and Grupen, C. G. (2017) Cryotolerance of porcine blastocysts is improved by treating *in vitro* matured oocytes with L-carnitine prior to fertilization. *The Journal of Reproduction and Development* 63(3), 263–270. doi: [10.1262/jrd.2016-141](https://doi.org/10.1262/jrd.2016-141)
- Marin, D. F. D., Nogueira da Costa, N., di Paula Bessa Santana, P., Baia de Souza, E., Rolim Filho, S. T., da Silva Cordeiro, M. and Ohashi, O. M. (2020) Influence of l-carnitine on lipid metabolism of buffalo cumulus-oocyte complexes matured in either fetal bovine serum or fatty acid-free bovine serum albumin. *Theriogenology* 158, 384–390. doi: [10.1016/j.theriogenology.2020.09.030](https://doi.org/10.1016/j.theriogenology.2020.09.030)
- Martelli, F. and Nunes, F. M. F. (2014) Radicais livres: em busca do equilíbrio. *Ciência e cultura* 66(3), 54–57. doi: [10.21800/S0009-67252014000300017](https://doi.org/10.21800/S0009-67252014000300017)
- Mingorance, C., Rodriguez-Rodriguez, R., Justo, M. L., Herrera, M. D. and de Sotomayor, M. A. (2011) Pharmacological effects and clinical applications of propionyl-L-carnitine. *Nutrition Reviews* 69(5), 279–290. doi: [10.1111/j.1753-4887.2011.00387.x](https://doi.org/10.1111/j.1753-4887.2011.00387.x)
- Modak, A. K., Alam, M. H., Islam, M. N., Paul, N., Akter, I., Hashem, M. A., Kabir, A. A. and Moniruzzaman, M. (2022) L-Carnitine Supports the *In Vitro* growth of buffalo oocytes. *Animals: an Open Access Journal From MDPI* 12(15), 1957. doi: [10.3390/ani12151957](https://doi.org/10.3390/ani12151957)
- Mohd Shukri, M. F., Norhayati, M. N., Badrin, S. and Abdul Kadir, A. (2022) Effects of L- carnitine supplementation for women with polycystic ovary syndrome: a systematic review and meta-analysis. *PeerJ*, 10, 13992. doi: [10.7717/peerj.13992](https://doi.org/10.7717/peerj.13992)
- Monzo, C., Haouzi, D., Roman, K., Assou, S., Dechaud, H. and Hamamah, S. (2012) Slow freezing and vitrification differentially modify the gene expression profile of human metaphase II oocytes. *Human Reproduction (Oxford, England)* 27(7), 2160–2168. doi: [10.1093/humrep/des153](https://doi.org/10.1093/humrep/des153)
- Nascimento, D. R., Azevedo, V. A. N., Barroso, P. A. A., Barrozo, L. G., Silva, B. R., Silva, A. W. B., Donato, M. A. M., Peixoto, C. A. and Silva, J. R. V. (2022) Effects of N-acetylcysteine on growth, viability, and ultrastructure of *in vitro* cultured bovine secondary follicles. *Animals: an Open Access Journal From MDPI*, 12(22), 3190. doi: [10.3390/ani12223190](https://doi.org/10.3390/ani12223190)
- Nimse, S. B. and Pal, D. (2015) Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Advances* 5(35), 27986–28006. doi: [10.1039/C4RA13315C](https://doi.org/10.1039/C4RA13315C)
- Pan, B., Yang, H., Wu, Z., Qazi, I. H., Liu, G., Han, H., Meng, Q. and Zhou, G. (2018) Melatonin improves parthenogenetic development of vitrified-warmed mouse oocytes potentially by promoting G1/S cell cycle progression. *International Journal of Molecular Sciences* 19(12), 4029. doi: [10.3390/ijms19124029](https://doi.org/10.3390/ijms19124029)
- Paulino, L. R. F. M., Barroso, P. A. A., Silva, B. R., Barrozo, L. G., Barbalho, E. C., Bezerra, F. T. G., Souza, A. L. P., Monte, A. P. O., Silva, A. W. B., Matos, M. H. T. and Silva, J. R. V. (2022) Immunolocalization of melatonin receptors in bovine ovarian follicles and *in vitro* effects of melatonin on

- growth, viability and gene expression in secondary follicles. *Domestic Animal Endocrinology* **81**, 106750. doi: [10.1016/j.domaniend.2022.106750](https://doi.org/10.1016/j.domaniend.2022.106750)
- Prevedello, M. T. and Comachio, G.** (2021) Antioxidants and their relationship with free radicals, and chronic non communicable diseases: a literature review. *Brazilian Journal of Development* **7**(6), 55244–55285. doi: [10.34117/bjdv7n6-096](https://doi.org/10.34117/bjdv7n6-096)
- Sá, N. A. R., Bruno, J. B., Guerreiro, D. D., Cadenas, J., Alves, B. G., Cibin, F. W. S., Leal-Cardoso, J. H., Gastal, E. L. and Figueiredo, J. R.** (2018) Anethole reduces oxidative stress and improves *in vitro* survival and activation of primordial follicles. *Brazilian Journal of Medical and Biological Research* **51**(8), e7129. doi: [10.1590/1414-431x20187129](https://doi.org/10.1590/1414-431x20187129)
- Sadeesh, E. M., Shah, F., Balhara, A. K., Thirumaran, S. M. K., Yadav, S. and Yadav, P. S.** (2014) Effect of growth factor and antioxidant on *in vitro* maturation of oocytes and cleavage rates of *in vitro* produced Indian buffalo (*Bubalus bubalis*) embryos. *Veterinarski Arhiv* **84** (5), 459–474.
- Saeed-Zidane, M., Linden, L., Salilew-Wondim, D., Held, E., Neuhoﬀ, C., Tholen, E., Hoelker, M., Schellander, K. and Tesfaye, D.** (2017) Cellular and exosome mediated molecular defense mechanism in bovine granulosa cells exposed to oxidative stress. *PLoS One* **12**(11), 0187569. doi: [10.1371/journal.pone.0187569](https://doi.org/10.1371/journal.pone.0187569)
- Sahraei, S. S., Shahhoseini, M. and Movaghar, B.** (2018) Vitrification has an effect like culture on gene expression and histone modification in mouse embryos. *CryoLetters* **39**(2), 102–112
- Sekhon, L., Lee, J. A., Flisser, E., Copperman, A. B. and Stein, D.** (2018) Blastocyst vitrification, cryostorage and warming does not affect live birth rate, infant birth weight or timing of delivery. *Reproductive Biomedicine Online* **37**(1), 33–42. doi: [10.1016/j.rbmo.2018.03.023](https://doi.org/10.1016/j.rbmo.2018.03.023)
- Simas, L. A. W., Granzoti, R. O. and Porsch, L.** (2019) Oxidative stress and its impact on aging: a literature review. *Brazilian Journal of Natural Sciences* **2**(2), 80. doi: [10.31415/bjns.v2i2.53](https://doi.org/10.31415/bjns.v2i2.53)
- Somfai, T., Kaneda, M., Akagi, S., Watanabe, S., Haraguchi, S., Mizutani, E., Dang-Nguyen, T. Q., Geshi, M., Kikuchi, K. and Nagai, T.** (2011) Enhancement of lipid metabolism with L-carnitine during *in vitro* maturation improves nuclear maturation and cleavage ability of follicular porcine oocytes. *Reproduction, Fertility, and Development* **23**(7), 912–920. doi: [10.1071/RD10339](https://doi.org/10.1071/RD10339)
- Somoskoi, B., Martino, N. A., Cardone, R. A., Lacalandra, G. M., Dell'Aquila, M. E. and Cseh, S.** (2015) Different chromatin and energy/redox responses of mouse morulae and blastocysts to slow freezing and vitrification. *Reproductive Biology and Endocrinology: RB&E* **13**, 22. doi: [10.1186/s12958-015-0018-z](https://doi.org/10.1186/s12958-015-0018-z)
- Soto-Heras, S. and Paramio, M. T.** (2020) Impact of oxidative stress on oocyte competence for *in vitro* embryo production programs. *Research in Veterinary Science* **132**, 342–350. doi: [10.1016/j.rvsc.2020.07.013](https://doi.org/10.1016/j.rvsc.2020.07.013)
- Souza, L. M. V., Costa, R. de A., Santos, J. D. M. dos, Santos, J. L. dos, Costa, L. S., Oliveira, J. U. de, Silva, R. J. dos S. and Estevam, C. dos S.** (2020) High-intensity interval training and oxidative stress: a brief presentation. *Research, Society and Development* **9**(8), e741986478. doi: [10.33448/rsd-v9i8.6478](https://doi.org/10.33448/rsd-v9i8.6478)
- Spijkers, S., Lens, J. W., Schats, R. and Lambalk, C. B.** (2017) Fresh and frozen-thawed embryo transfer compared to natural conception: differences in perinatal outcome. *Gynecologic and Obstetric Investigation* **82**(6), 538–546. doi: [10.1159/000468935](https://doi.org/10.1159/000468935)
- Stojkovic, M., Machado, S. A., Stojkovic, P., Zakhartchenko, V., Hutzler, P., Goncalves, P. B. and Wolf, E.** (2001) Mitochondrial distribution and adenosine triphosphate content of bovine oocytes before and after *in vitro* maturation: correlation with morphological criteria and developmental capacity after *in vitro* fertilization and culture. *Biology of Reproduction* **64**(3), 904–909. doi: [10.1095/biolreprod64.3.904](https://doi.org/10.1095/biolreprod64.3.904)
- Takahashi, T., Inaba, Y., Somfai, T., Kaneda, M., Geshi, M., Nagai, T. and Manabe, N.** (2013) Supplementation of culture medium with L-carnitine improves development and cryotolerance of bovine embryos produced *in vitro*. *Reproduction, Fertility and Development* **25**(4), 589–599. doi: [10.1071/rd11262](https://doi.org/10.1071/rd11262)
- Truong, T., Harvey, A. J. and Gardner, D. K.** (2022) Antioxidant supplementation of mouse embryo culture or vitrification media support more in-vivo-like gene expression post-transfer. *Reproductive Biomedicine Online* **44**(3), 393–410. doi: [10.1016/j.rbmo.2021.11.013](https://doi.org/10.1016/j.rbmo.2021.11.013)
- Truong, T. T., Soh, Y. M. and Gardner, D. K.** (2016) Antioxidants improve mouse preimplantation embryo development and viability. *Human Reproduction (Oxford, England)* **31**(7), 1445–1454. doi: [10.1093/humrep/dew098](https://doi.org/10.1093/humrep/dew098)
- Vanella, A., Russo, A., Acquaviva, R., Campisi, A., Di Giacomo, C., Sorrenti, V. and Barcellona, M. L.** (2000) L-propionyl-carnitine as superoxide scavenger, antioxidant, and DNA cleavage protector. *Cell Biology and Toxicology* **16**(2), 99–104. doi: [10.1023/a:1007638025856](https://doi.org/10.1023/a:1007638025856)
- Vining, L. M., Zak, L. J., Harvey, S. C. and Harvey, K. E.** (2021) The role of apoptosis in cryopreserved animal oocytes and embryos. *Theriogenology* **173**, 93–101. doi: [10.1016/j.theriogenology.2021.07.017](https://doi.org/10.1016/j.theriogenology.2021.07.017)
- Volpe, C. M. O., Villar-Delfino, P. H., Dos Anjos, P. M. F. and Nogueira-Machado, J. A.** (2018) Cellular death, reactive oxygen species (ROS) and diabetic complications. *Cell Death & Disease* **9**(2), 119. doi: [10.1038/s41419-017-0135-z](https://doi.org/10.1038/s41419-017-0135-z)
- Wu, Z., Pan, B., Qazi, I. H., Yang, H., Guo, S., Yang, J., Zhang, Y., Zeng, C., Zhang, M., Han, H., Meng, Q. and Zhou, G.** (2019) Melatonin improves *in vitro* development of vitrified-warmed mouse germinal vesicle oocytes potentially via modulation of spindle assembly checkpoint-related genes. *Cells* **8**(9), 1009. doi: [10.3390/cells8091009](https://doi.org/10.3390/cells8091009)
- Zare, Z., Masteri Farahani, R., Salehi, M., Piryaei, A., Ghaffari Novin, M., Fadaei Fathabadi, F., Mohammadi, M. and Dehghani-Mohammadabadi, M.** (2015) Effect of L-carnitine supplementation on maturation and early embryo development of immature mouse oocytes selected by brilliant cresyl blue staining. *Journal of Assisted Reproduction and Genetics* **32**(4), 635–643. doi: [10.1007/s10815-015-0430-5](https://doi.org/10.1007/s10815-015-0430-5)
- Zare, Z., Rezaei, N. and Mohammadi, M.** (2022) Treatment of mouse cumulus-oocyte complexes with L-carnitine during vitrification and *in vitro* maturation affects maturation and embryonic developmental rate after parthenogenetic activation. *Anatomia, Histologia, Embryologia* **51**(1), 44–50. doi: [10.1111/ahel.12750](https://doi.org/10.1111/ahel.12750)
- Zhang, Q., Wang, S. M., Yao, P. B., Zhang, L., Zhang, Y. J., Chen, R. X., Fu, Y. and Zhang, J. M.** (2015) Effects of L-carnitine on follicular survival and graft function following autotransplantation of cryopreserved-thawed ovarian tissues. *Cryobiology* **71**(1), 135–140. doi: [10.1016/j.cryobiol.2015.04.008](https://doi.org/10.1016/j.cryobiol.2015.04.008)
- Zhang, S., Yao, H., Liu, Y., Ren, L., Xiang, D., and Wang, Y.** (2020) Hypothermic machine perfusion after static cold storage improves ovarian function in rat ovarian tissue transplantation. *Journal of Assisted Reproduction and Genetics* **37**, 1745–1753. doi: [10.1007/s10815-020-01797-4](https://doi.org/10.1007/s10815-020-01797-4)
- Zolini, A. M., Carrascal-Triana, E., Ruiz de King, A., Hansen, P. J., Alves Torres, C. A. and Block, J.** (2019) Effect of addition of l-carnitine to media for oocyte maturation and embryo culture on development and cryotolerance of bovine embryos produced *in vitro*. *Theriogenology* **133**, 135–143. doi: [10.1016/j.theriogenology.2019.05.005](https://doi.org/10.1016/j.theriogenology.2019.05.005)