

Triaxial orbit-based model of NGC 4365

Remco C. E. van den Bosch,¹ Glenn van de Ven,²
Michele Cappellari¹ and P. Tim de Zeeuw¹

¹Sterrewacht Leiden, Universiteit Leiden, Postbus 9513, 2300 RA Leiden, The Netherlands

²Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA

email: Bosch@strw.leidenuniv.nl

Abstract. We have developed an orbit-based method for constructing triaxial models of elliptical galaxies, which fit their observed surface brightness and kinematics (van den Bosch *et al.*). We have tested this extended Schwarzschild method (1979) against analytical models with general distribution functions (DF) and find that we can recover the DF (van de Ven *et al.*). Here, we present a model of NGC 4365.

Keywords. galaxies: elliptical and lenticular, cD – galaxies: kinematics and dynamics

1. NGC 4365

NGC 4365 is an E3 galaxy with a kinematically de-coupled core (KDC). Its rotation axis is misaligned with respect to that of the main body of the galaxy (Surma & Bender 1994; Davies *et al.* 2001; Statler *et al.* 2004). The bulk of the galaxy appears to rotate around the major axis, but the stars in the central 6'' rotate around the minor axis. These properties can be explained by an intrinsically triaxial shape.




Figure 1. Stellar kinematics of NGC 4365 on the top row. From left to right: the mean velocity, velocity dispersion and the higher order moments h_3 and h_4 (measuring deviations from Gaussian). The contours are representative isophotes. The best-fitting model is shown on the bottom row and it is a good fit, reproducing the observed features in the symmetrized kinematics (middle row) with great accuracy ($\chi^2/\text{degrees-of-freedom} \sim 1$).

We measured the stellar kinematics of NGC 4365 (Fig. 1) with the SAURON integral-field spectrograph (Bacon *et al.* 2001). We constructed triaxial models using both the integral-field stellar kinematics and HST and ground-based imaging and including a central black hole. Our best-fit model (Fig. 1) is close to oblate with axis ratios of 1:0.97:0.73 (see also Statler *et al.* 2004) and surprisingly consists primarily of orbits that rotate around the short axis.




Figure 2. The cumulative mass per orbit-type as a function of radius. The modest fraction of stars on long axis tube orbits all have the same sense of rotation, and are responsible for the observed mean rotation.




Figure 3. Kinematics of the model up to a radius of $60''$. Only the part inside the rectangular box is constrained by the kinematic observations, the rest is extrapolated. The kinematics outside the box are thus not very reliable, but they do allow one to see the extended structure.

2. Orbit types

In our best-fit model the short axis tube orbits account for 75% of the mass inside $30''$ (Fig. 2) and hence dominate even outside the KDC. This seems in contradiction with the rotation seen around the major axis at large radius in the observations. However, the average rotation of the stars populating the short axis orbits is almost negligible in this region, causing the high dispersion measured along the major axis. To be able to visualise the structure of the model we show the extended kinematics (Fig. 3). The extended structure, including the sigma dip along the minor axis and the continuation of the unaligned zero velocity curve show more clearly that the galaxy is oblate/triaxial.

Acknowledgements

RvdB thanks the Leids Kerkhoven-Bosscha Fonds for travel support.

References

- Bacon R. *et al.* 2001, MNRAS, 326, 23
- Davies R. L., *et al.* 2001, ApJ, 548, L33
- Schwarzschild M. 1979, ApJ, 232, 2
- Surma P. & Bender R. 1995, A&A, 298, 405
- Statler T. S., Emsellem E., Peletier R. F. & Bacon R. 2004, MNRAS, 353, 1
- van den Bosch R. C. E., van de Ven G., Verolme E. K., Cappellari M. & de Zeeuw P. T. 2006, MNRAS, submitted
- van de Ven G., de Zeeuw P. T. & van den Bosch R. C. E. 2006, MNRAS, submitted