

time were more mentally stable and had no current somatic problems. Multidisciplinary teamwork is important for patients with deteriorating MMS scores.

Conclusion By illustrating the impact of somatic and psychiatric factors on dementia, the present study underlines the value of multidisciplinary professional care, the role of the family and the importance of long-stay wards.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.1703>

EV719

Depressive symptoms in older people in Greece and Cyprus

K. Argyropoulos^{1,*}, G. Panteli², G. Charalambous², A. Argyropoulou³, P. Gourzis⁴, E. Jelastopulu¹

¹ Medical School, University of Patras, Public Health, Patras, Greece

² Frederick University, Postgraduate Program Health Management, Nicosia, Cyprus

³ Health Center of Andravida, General Practice, Andravida, Greece

⁴ University Hospital of Patras, Psychiatry, Patras, Greece

* Corresponding author.

Introduction Depression is fast becoming a major public health problem with a very high prevalence rate in the 65 and over age group.

Objectives The aim of the present study was to estimate the prevalence of depression in Greeks and Cypriots older adults.

Methods A cross-sectional study was conducted among the 445 participants, 239 members of three day care centers for older people, in the municipality of Patras, West-Greece and 206 older adults (110 in the community, 65 in outpatient clinics, 31 in nursing homes) in Cyprus, aged >60 years. A questionnaire was administered including socio-demographic characteristics. Depression was assessed using the Greek version of Geriatric Depression Scale (GDS-15).

Results The overall prevalence of depression according to GDS-15 was 33% (28% moderate, 5% severe type). Depressive symptoms were more frequent in women (41,6% vs. 28,3%, $P < 0,001$), in not married (43,0% vs. 29,3%, $P < 0,001$), in elderly with chronic diseases (36,8% vs. 25,0%, $P = 0,007$), in older people dwellers of urban areas compared to rural (36,3% vs. 26,4%, $P = 0,028$) and in ages between 70 to 80 years old (38,7% vs. 31,6%, $P = 0,038$). Moreover, higher prevalence of depression was measured in Greeks compared to Cypriots (44,3% vs. 20,6%, $P < 0,001$). In a univariate analysis, the following variables were significantly associated with depression: female gender ($P < 0,001$), co-morbidity ($P = 0,004$), higher age group ($P = 0,018$), place of living ($P = 0,022$) and Greek nationality ($P < 0,001$).

Conclusions High prevalence and several risk factors are strongly associated with depression, whereas Greeks are in higher danger of developing depressive symptoms in late life, than Cypriots.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.1704>

EV720

Treatment of Charles Bonnet syndrome with continuous positive airway pressure in an older adult

M. Arts^{1,*}, P. Michielsen², S. Petrykiv³, L. de Jonge¹, R. Oude Voshaar¹

¹ UMCG, Old Age Psychiatry, Groningen, Netherlands

² GGZWNB, Psychiatry, Bergen op Zoom, Netherlands

³ GGZ Friesland, Emergency Psychiatry, Leeuwarden, Netherlands

* Corresponding author.

Introduction Charles Bonnet syndrome (CBS) is a disorder in older adults, and is characterized by a triad of recurrent vivid visual hallucinations, ocular pathology causing visual impairment due to lesions in central or peripheral visual pathways, and normal cognitive status. It is often misdiagnosed as a psychosis, early dementia or a drug related condition. Hypoxemia was anecdotally reported as a cause of CBS.

Objectives We present an older adult with CBS caused by severe obstructive sleep apnea syndrome.

Aims To report a case study, describing treatment of obstructive sleep apnea syndrome as a cause of CBS.

Methods A case study is presented and discussed.

Results An older male adult was admitted to hospital for persistent vivid visual hallucinations. There was no personal or family history of mental illness. Neurological examination was normal, except for visual impairment due to age related macular degeneration. The remainder of his physical examination was normal. Previous treatment with antipsychotics proved not to be effective. Severe hypoxemia (SaO₂ 79%) was diagnosed with overnight pulse oximetry and subsequent polysomnography revealed an obstructive sleep apnea syndrome. After three nights of nasal continuous positive airway pressure, the vivid hallucinations ceased.

Conclusion Physicians need to understand the underlying causes and mechanisms of CBS. One should be aware of the importance of a full clinical examination and sleep apnea research in elderly persons with visual impairment.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2016.01.1705>

EV721

Management of late-life insomnia

S. Bravo Herrero^{1,*}, C. Moreno Menguiano², R. Martín Aragón¹, M. Gutiérrez Rodríguez², J.F. Cruz Fourcade¹, N. Rodríguez Criado¹, P. Muñoz-Calero Franco¹, B. Sánchez Sánchez¹

¹ Hospital Universitario de Móstoles, UHB Psiquiatría, Móstoles, Spain

² Hospital Universitario de Móstoles, CSM Móstoles, Móstoles, Spain

* Corresponding author.

Introduction Insomnia is the most frequent sleep disorder in late life. Forty-two percent of elderly people in the United States often complain about difficulties to get or maintain sleep, or awakening too early. Insomnia is frequent in old people greatly due to frequency of concomitant medical illnesses and polypharmacy, rather than because of age.

Objectives The objective of our research was to revise the current state of knowledge about management of insomnia in people above 65 years of age.

Methodology For that, a bibliographical search through PubMed.gov has been made. From the obtained results, the 14 which best suited for our goals were selected, 10 of them dealing with people above 65 years and the rest with people above 75 or 80 years of age.

Results Based on the literature reviewed, the current options of management of late-life insomnia are based on behavioral or pharmacological therapy. The combination of behavioral therapies shows results and is currently considered as an option, especially given the possibility of medicine interaction and the secondary effects hypnotic and sedative medicines might produce. There is a paucity of long-term safety and efficacy data for the use of non-benzodiazepine sedative-hypnotics. There are no criteria for the use of antidepressant sedatives in elderly people without diagnosed depression, although they are still used in practice.

Conclusion Possibility of using behavioral therapy as first option. In case of polymedicated or multi-pathological patients, pay special attention when starting a pharmacological treatment, choose the most suitable one and supervise it closely.