

CORRIGENDA

Journal of Agricultural Science, Vol. 60 (1963)

Soil changes in ley-arable experiments

By A. J. LOW, F. J. PIPER AND P. ROBERTS

p. 231, Table 6 heading:

for blocks 1-5 *read* blocks 1, 2, 4 and 5

p. 233, column 2 fourth sentence in second paragraph to read as follows:

There was a good response to the nitrogen: the yield increased from 10.2 cwt. grain (no N fertilizer) to 25.7 cwt. with 1 ton of 'Nitro-Chalk' (15.5 % N)—about 13 % recovery of the nitrogen—the straw recovered another 5 %.

p. 233, Table 14 last line to read as follows:

After 8 years ley

CONTENTS

| | PAGE |
|---|------|
| <p> RIDLER, B., BROSTER, W. H. and FOOT, A. S. The growth rate of heifers in a dairy herd. (With Three Text-figures) </p> | 1 |
| <p> HOLME, D. W., COEY, W. E. and ROBINSON, K. L. The prediction of pig carcass composition from measurements of carcass density </p> | 9 |
| <p> CARROLL, M. A., PEARCE, J., BASSETT, J. M., GILLARD, P. B. and LAWRENCE, T. L. J. The effects of age of calves at castration on growth rate, carcass composition and feed efficiency </p> | 19 |
| <p> COWLISHAW, S. J. A study of the effects of two methods of grazing poultry on the persistence and palatability of seven herbage species. (With Two Text-figures) </p> | 27 |
| <p> ARNOLD, G. W. and DUDZINSKI, M. L. The use of faecal nitrogen as an index for estimating the consumption of herbage by grazing animals. (With Four Text-figures) </p> | 33 |
| <p> DONEY, J. M. The fleece of the Scottish Blackface sheep. III. The relative importance of the fleece components. (With Six Text-figures) </p> | 45 |
| <p> MARTIN, I. C. A. The influence of diluents and processing time after ejaculation on the revival of deep-frozen bull spermatozoa. (With Three Text-figures) </p> | 55 |
| <p> VAN EIJNATTEN, C. L. M. A study of the development of two varieties of maize at Ibadan, Nigeria. (With Two Text-figures) </p> | 65 |
| <p> NAGA, M. M. A. and EL-SHAZLY, K. The use of the <i>in vitro</i> fermentation technique to estimate the digestible energy content of some Egyptian forages. I. The <i>in vitro</i> digestion of cellulose as a criterion of energy content </p> | 73 |
| <p> HALE, R. W. and COEY, W. E. Genotype-environment interactions in a herd of bacon pigs </p> | 81 |
| <p> HERRIOTT, J. B. D. and WELLS, D. A. The grazing animal and sward productivity. (With One Text-figure) </p> | 89 |
| <p> BREDON, R. M., HARKER, K. W. and MARSHALL, B. The nutritive value of grasses grown in Uganda when fed to zebu cattle. I. The relation between the percentage of crude protein and other nutrients </p> | 101 |
| <p> BREDON, R. M., HARKER, K. W. and MARSHALL, B. The nutritive value of grasses grown in Uganda when fed to zebu cattle. II. The relation between crude fibre and nitrogen-free extract and other nutrients </p> | 105 |
| <p> EL-SHAZLY, K., ABOU AKKADA, A. R. and NAGA, M. M. A. The use of the <i>in vitro</i> fermentation technique to estimate the digestible energy content of some Egyptian forages. II. The <i>in vitro</i> production of total volatile fatty acids and organic acids as criteria of energy content. (With One Text-figure) </p> | 109 |
| <p> ENYI, B. A. C. The effect of time of irrigation on the growth and yield of transplanted rice plants (<i>Oryza sativa</i> L.) (With Three Text-figures) </p> | 115 |
| <p> BAYLEY, H. S. and LEWIS, D. The use of fats in pig rations </p> | 121 |
| <p> LUITINGH, H. C. The efficiency of beef production in terms of carcass-weight increase as influenced by the ration concentration and the age of steers. (With Three Text-figures) </p> | 127 |
| <p> MORIARTY, F. The decline of a pea root eelworm (<i>Heterodera göttingiana</i> Liebscher) population in the absence of host plants. (With Two Text-figures) </p> | 135 |
| <p> BREMNER, P. M., ECKERSALL, R. N. and SCOTT, R. K. The relative importance of embryo size and endosperm size in causing the effects associated with seed size in wheat. (With Four Text-figures) </p> | 139 |

The Journal of Agricultural Science is issued every other month in parts of about 136 pages with plates and figures, three parts constituting a volume. Two volumes yearly.

Quotations for available back volumes and parts may be obtained on application to the Cambridge University Press.

Papers for publication should be typewritten or written in a very legible hand, and may be sent to SIR JOHN HAMMOND, School of Agriculture, Cambridge, or to the associate Editors. Other communications should be addressed to the University Press, Cambridge.

If Plates in excess of one per 24 pages of text or coloured plates are submitted the author may be asked to bear the cost of such Plates.

The lettering on text figures should be clear of the illustration and indicating by blue pencilled lines the desired position.

Bibliography should be given under the heading of 'References' in the following form:

Surname of authors (in alphabetical order), initials, date of publication (in brackets), title of Journal (abbreviated according to the *World List of Scientific Periodicals*), volume and page of reference. In the text a reference should be quoted by the author's name and date (in brackets).

All names of Journals and books are set in italics; they should be indicated on the typescript by underlining.

Contributors will receive twenty-five copies of their papers free.

The subscription price, payable in advance, is 65s. net per volume (post free); single numbers 27s. 6d. net. Subscriptions may be sent to any bookseller, or to The Cambridge University Press, Bentley House, 200 Euston Road, London, N.W. 1.

Subscription price in the U.S.A. is \$12.50 per volume; single number \$4.75. Enquiries should be addressed to the Cambridge University Press, American Branch, 32 East 57th Street, New York 22, N.Y.

*Printed in Great Britain at the University Press, Cambridge
(Brooke Crutchley, University Printer)*