

equally accessible to all children in Maine. This may restrict their access to specific asthma therapies such as allergic desensitization and monoclonal antibodies.

248

HPV vaccine knowledge and beliefs among women in rural areas

Deborah Smith¹, Shreveport Cory Coehoorn¹, Shreveport Robbie A. Beyl², Yanna Xhari Alesafis de Koter², Emily Martinson Dantes¹ and Shreveport Jerry W. McLarty¹

¹Louisiana State University Health Science and ²Pennington Biomedical Research Center

OBJECTIVES/GOALS: To reduce cervical cancer in the USA, it is essential to identify the reasons underlying the low HPV vaccination rates. This study aims to identify knowledge and beliefs about HPV vaccination among women in rural areas. The knowledge gained from this study will directly lead to developing an education tool tailored specifically for women in rural areas. **METHODS/STUDY POPULATION:** We conducted a cross-sectional study from November 2022 to July 2023. We recruited women (n = 141) who visited a mobile health unit in rural North Louisiana. The inclusion criteria were women aged 25–64 years who spoke English, had not undergone a total hysterectomy, and had no history of cervical cancer. Data collected included sociodemographic characteristics, sexual history, awareness and knowledge of HPV infection, cervical cancer, genital warts, and HPV vaccination, the perceived risk of acquiring genital HPV infection and developing cervical cancer, and the willingness to receive an HPV vaccine. Descriptive statistics were used to evaluate participant responses. Written informed consent was obtained before completing the self-administered questionnaire. **RESULTS/ANTICIPATED RESULTS:** Our findings showed significant gaps in HPV vaccine knowledge and uptake. Approximately 40% of the participants were unaware of the HPV vaccine, 96.5% had never received it, and 91.4% had never been offered it. However, 42% indicated a willingness to consider vaccination if it were offered. Factors influencing their decision to vaccinate against HPV included a family history of cervical cancer (44.7%), having multiple sexual partners (48%), and engaging in unprotected sex (61.7%). Furthermore, there was significant uncertainty surrounding the vaccine, with 65.2% of participants unsure about its safety, 66% unaware of potential health risks, and 47.5% uncertain about its effectiveness in preventing HPV. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Our research emphasizes the need for tailored behavioral interventions to address knowledge gaps about the HPV vaccine. Low HPV vaccination rates in rural areas contribute to health disparities in cervical cancer. Implementing educational interventions in healthcare settings can enhance vaccination rates and mitigate the risk of cervical cancer.

250

Transplant center level variation in listing patients for liver transplant with initial inactive status

Sarah Huber¹, Jonathan Fridell¹, Rachel Patzer² and Katie Ross Driscoll¹

¹Indiana University School of Medicine, Dept of Surgery and

²Indiana University School of Medicine, Dept of Surgery, and Regenstrief Institute

OBJECTIVES/GOALS: Our objectives were to describe the characteristics of patients initially listed with inactive status on the liver

transplant waitlist to identify if disparities exist and compare the prevalence of initial inactive status listings across transplant centers. **METHODS/STUDY POPULATION:** This was a retrospective cohort study of candidates waitlisted for liver transplant between March 1, 2023 and February 12, 2023, utilizing the Scientific Registry of Transplant Recipients, a national database including all solid organ transplant candidates. 224,736 candidates were included in analysis, and covariates included race, ethnicity, sex, age, body mass index, primary payer, MELD at listing, and etiology of liver disease. **RESULTS/ANTICIPATED RESULTS:** Totally, 8,131 (3.62%) candidates were initially listed for liver transplant with inactive status. Although there were statistically significant differences between those listed initially with active status and those listed initially with inactive status in each covariate, these differences did not reach clinical significance. Of the 151 transplant centers, 128 listed any patients with an initial inactive status, with inactive status listings compromising 0–49.36% of total listings by transplant center. There is significant variation between listing centers in the practice of listing with initial inactive status by both liver disease etiology and across different eras of liver allocation policies. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Despite no significant clinical difference in the characteristics of patients listed with initial inactive status, there is significant variation across transplant centers of the prevalence of listing with initial inactive status. Subsequent investigations will focus on understanding these differences in listing practices between centers.

251

Ethical care systems for substance use disorder (SUD): Evaluating stigma and clinical decision-making

Jessica Olivares

The University of Texas Medical Branch at Galveston

OBJECTIVES/GOALS: This project (1) assesses provider behavior at the Galveston Secure Against the Fentanyl Epidemic (G-SAFE) Clinic in providing evidence-based care for underserved patients in our community and (2) implements evidence-based anti-stigma intervention in local health care. **METHODS/STUDY POPULATION:** The efficacy and impact of G-SAFE on healthcare utilization, health outcomes, and patient satisfaction will be determined. Evaluations assessing provider behavior will be made across at least 25 clinic patient visits, taking note of medical students, clinicians, and health providers' interactions with patients related to decision-making, care plan creation, and communication. Health provider discussion groups will provide essential data in understanding the successful management of SUDs, management of safety and risk, and how best to pilot and assess interventions that promote greater trust-building and a harm-reduction approach to clinical care for patient empowerment and positive outcomes. **RESULTS/ANTICIPATED RESULTS:** Observations in the clinic will be turned into interventions to improve diagnostics and therapeutics for SUD medical care and behavioral change. We will recruit 40 medical students (with/without exposure to UTMB's free clinics) as well as 20 clinicians in the G-SAFE Clinic, Emergency Medicine, and Primary Care. Baseline stigma will be measured with the Medical Condition Regard Scale and the Drug and Drug Problems Questionnaire alongside measures of care approaches. The data collected and the educational intervention for medical students and clinicians will make a positive impact on the lives of those affected by SUDs and serve as a model for future interventions on a larger scale. **DISCUSSION/SIGNIFICANCE OF IMPACT:** Health provider's