



COMPOSITIO MATHEMATICA

Odd zeta motive and linear forms in odd zeta values

Clément Dupont

Compositio Math. **154** (2018), 342–379.

[doi:10.1112/S0010437X17007588](https://doi.org/10.1112/S0010437X17007588)





Odd zeta motive and linear forms in odd zeta values

Clément Dupont

With a joint appendix with Don Zagier

ABSTRACT

We study a family of mixed Tate motives over \mathbb{Z} whose periods are linear forms in the zeta values $\zeta(n)$. They naturally include the Beukers–Rhin–Viola integrals for $\zeta(2)$ and the Ball–Rivoal linear forms in odd zeta values. We give a general integral formula for the coefficients of the linear forms and a geometric interpretation of the vanishing of the coefficients of a given parity. The main underlying result is a geometric construction of a minimal ind-object in the category of mixed Tate motives over \mathbb{Z} which contains all the non-trivial extensions between simple objects. In a joint appendix with Don Zagier, we prove the compatibility between the structure of the motives considered here and the representations of their periods as sums of series.

1. Introduction

1.1 Constructing linear forms in zeta values

The study of the values at integers $n \geq 2$ of the Riemann zeta function

$$\zeta(n) = \sum_{k \geq 1} \frac{1}{k^n}$$

goes back to Euler, who showed that the even zeta value $\zeta(2n)$ is a rational multiple of π^{2n} . Lindemann’s theorem thus implies that the even zeta values are transcendental numbers. It is conjectured that the odd zeta values $\zeta(3), \zeta(5), \zeta(7), \dots$ are algebraically independent over $\mathbb{Q}[\pi]$.

Many of the results in the direction of this conjecture use as a key ingredient certain families of period integrals which evaluate to linear combinations of 1 and zeta values:

$$\int_{\sigma} \omega = a_0 + a_2 \zeta(2) + \dots + a_n \zeta(n), \quad (1)$$

with $a_k \in \mathbb{Q}$ for every k . We can cite in particular the following results (see Fischler’s Bourbaki talk [Fis04] for a more complete survey).

- Apéry’s proof [Apé79] of the irrationality of $\zeta(2)$ and $\zeta(3)$ was simplified by Beukers [Beu79] by using a family of integrals evaluating to linear combinations $a_0 + a_2 \zeta(2)$ and $a_0 + a_3 \zeta(3)$.
- Ball and Rivoal’s proof [Riv00, BR01] that infinitely many odd zeta values are irrational relies on a family of integrals evaluating to linear combinations (1) for which all the even coefficients a_2, a_4, a_6, \dots vanish.

Received 21 November 2016, accepted in final form 28 July 2017, published online 30 October 2017.

2010 Mathematics Subject Classification 11M06, 14F42, 14G10 (primary).

Keywords: zeta values, mixed Tate motives.

This journal is © Foundation Compositio Mathematica 2017.

- Rhin and Viola’s irrationality measures [RV96, RV01] for $\zeta(2)$ and $\zeta(3)$ are built on generalizations of the Beukers integrals and precise estimates for the coefficients a_2 and a_3 .

In view of diophantine applications, it is crucial to have some control over the coefficients a_k appearing in linear combinations (1), in particular to be able to predict the vanishing of certain coefficients.

In the present article, we study the family of integrals

$$\int_{[0,1]^n} \omega \quad \text{with } \omega = \frac{P(x_1, \dots, x_n)}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n, \tag{2}$$

where $n \geq 1$ and $N \geq 0$ are integers and $P(x_1, \dots, x_n)$ is a polynomial with rational coefficients. This family contains the Beukers–Rhin–Viola integrals for $\zeta(2)$ and the Ball–Rivoal integrals. We say that an algebraic differential form ω as in (2) is *integrable* if the integral in (2) is absolutely convergent. Our first result is that such integrals evaluate to linear combinations of 1 and zeta values, with an integral formula for the coefficients.

THEOREM 1.1. *There exists a family $(\sigma_2, \dots, \sigma_n)$ of relative n -cycles with rational coefficients in $(\mathbb{C}^*)^n - \{x_1 \cdots x_n = 1\}$ such that for every integrable ω we have*

$$\int_{[0,1]^n} \omega = a_0(\omega) + a_2(\omega)\zeta(2) + \cdots + a_n(\omega)\zeta(n),$$

with $a_k(\omega)$ a rational number for every k , given for $k = 2, \dots, n$ by the formula

$$a_k(\omega) = (2\pi i)^{-k} \int_{\sigma_k} \omega. \tag{3}$$

The case $n = k = 2$ of this theorem is Rhin and Viola’s contour formula for $\zeta(2)$ [RV96, Lemma 2.6]. We note that in Theorem 1.1, the relative homology classes of the n -cycles σ_k are uniquely determined, see Theorem 4.9 for a precise statement. Furthermore, they are invariant, up to a sign, by the involution

$$\tau : (x_1, \dots, x_n) \mapsto (x_1^{-1}, \dots, x_n^{-1}), \tag{4}$$

which implies a general vanishing theorem for the coefficients $a_k(\omega)$, as follows.

THEOREM 1.2. *For $k = 2, \dots, n$ the relative cycle $\tau.\sigma_k$ is homologous to $(-1)^{k-1}\sigma_k$. Thus, for every integrable ω :*

- (i) *if $\tau.\omega = \omega$ then $a_k(\omega) = 0$ for $k \neq 0$ even;*
- (ii) *if $\tau.\omega = -\omega$ then $a_k(\omega) = 0$ for k odd.*

This allows us to construct families of integrals (2) which evaluate to linear combinations of 1 and odd zeta values, or 1 and even zeta values. This is the case for the integrals (see Corollary 5.6)

$$\int_{[0,1]^n} \frac{x_1^{u_1-1} \cdots x_n^{u_n-1} (1 - x_1)^{v_1-1} \cdots (1 - x_n)^{v_n-1}}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n,$$

where the integers $u_i, v_i \geq 1$ satisfy $2u_i + v_i = N + 1$ for every i . Depending on the parity of the product $(n + 1)(N + 1)$, the differential form is invariant or anti-invariant by τ and we get the vanishing of even or odd coefficients. This gives a geometric interpretation of the vanishing of the coefficients in the Ball–Rivoal integrals [Riv00, BR01], which correspond to special values of the parameters u_i, v_i .

The fact that the vanishing of certain coefficients in the Ball–Rivoal integrals could be explained by the existence of (anti-)invariant relative cycles was suggested to me by Rivoal during a visit at Institut Fourier, Grenoble, in October 2015. The special role played by the involution τ was first remarked by Deligne in a letter to Rivoal (17 February 2001, Princeton, NJ).

In an appendix written jointly with Don Zagier, we give an interpretation of the coefficients $a_k(\omega)$ appearing in Theorem 1.1 in elementary terms, that is in terms of the natural representations of the integrals in (2) as sums of series. This should be viewed as a geometric version of the dictionary between integrals and sums of series which is used in [Riv00, BR01]. It also gives an elementary proof of the vanishing properties of Theorem 1.2, which is essentially already present in the literature, see e.g. [Riv00, BR01], [Zud04, § 8] and [CFR08b, § 3.1].

The existence of the integral formulas (3) follows from the computation of certain motives, which are the central objects of the present article and that we now describe.

1.2 Constructing extensions in mixed Tate motives

Recall that the category $\text{MT}(\mathbb{Z})$ of mixed Tate motives over \mathbb{Z} is a (neutral) \mathbb{Q} -linear tannakian category defined in [DG05] and whose abstract structure is well understood. The only simple objects in $\text{MT}(\mathbb{Z})$ are the pure Tate objects $\mathbb{Q}(-k)$, for k an integer, and every object in $\text{MT}(\mathbb{Z})$ has a canonical weight filtration whose graded quotients are sums of pure Tate objects. The only non-zero extension groups between the pure Tate objects are given by

$$\text{Ext}_{\text{MT}(\mathbb{Z})}^1(\mathbb{Q}(-(2n + 1)), \mathbb{Q}(0)) \cong \mathbb{Q} \quad (n \geq 1). \tag{5}$$

Furthermore, a period matrix of the (essentially unique) non-trivial extension of $\mathbb{Q}(-(2n + 1))$ by $\mathbb{Q}(0)$ has the form

$$\begin{pmatrix} 1 & \zeta(2n + 1) \\ 0 & (2\pi i)^{2n+1} \end{pmatrix}.$$

The difficulty of constructing linear combinations (1) with many vanishing coefficients reflects the difficulty of constructing objects of $\text{MT}(\mathbb{Z})$ with many vanishing weight-graded quotients [Bro16, § 1.4]. In particular, the difficulty of constructing linear combinations involving only 1 and $\zeta(2n + 1)$ reflects the difficulty of giving a geometric construction of the extensions (5).

In this article, we construct a minimal ind-object \mathcal{Z}^{odd} in the category $\text{MT}(\mathbb{Z})$ which contains all the non-trivial extensions (5). The construction goes as follows. We first define, for every integer n , an object $\mathcal{Z}^{(n)} \in \text{MT}(\mathbb{Z})$ whose periods naturally include all the integrals (2). More precisely, any integrable form ω defines a class in the de Rham realization $\mathcal{Z}_{\text{dR}}^{(n)}$, and the unit n -cube $[0, 1]^n$ defines a class in the dual of the Betti realization $\mathcal{Z}_{\text{B}}^{(n),\vee}$, the pairing between these classes being the integral (2). The technical heart of this article is the computation of the full period matrix of $\mathcal{Z}^{(n)}$.

THEOREM 1.3. *We have a short exact sequence*

$$0 \rightarrow \mathbb{Q}(0) \rightarrow \mathcal{Z}^{(n)} \rightarrow \mathbb{Q}(-2) \oplus \cdots \oplus \mathbb{Q}(-n) \rightarrow 0$$

and $\mathcal{Z}^{(n)}$ has the following period matrix which is compatible with this short exact sequence:

$$\begin{pmatrix} 1 & \zeta(2) & \zeta(3) & \cdots & \cdots & \zeta(n-1) & \zeta(n) \\ & (2\pi i)^2 & & & & & \\ & & (2\pi i)^3 & & & 0 & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & 0 & & & & (2\pi i)^{n-1} & \\ & & & & & & (2\pi i)^n \end{pmatrix}.$$

Concretely, this theorem says that we can find a basis (v_0, v_2, \dots, v_n) of the de Rham realization $\mathcal{Z}_{\text{dR}}^{(n)}$ (which we will compute explicitly in terms of a special family of integrable forms) and a basis $(\varphi_0, \varphi_2, \dots, \varphi_n)$ of the dual of the Betti realization $\mathcal{Z}_{\text{B}}^{(n), \vee}$, such that the matrix of the integrals $\langle \varphi_i, v_j \rangle$ is the one given. The basis element φ_0 is the class of the unit n -cube $[0, 1]^n$. Expressing the class $[\omega] \in \mathcal{Z}_{\text{dR}}^{(n)}$ of an integrable form ω in the basis (v_0, v_2, \dots, v_n) as

$$[\omega] = a_0(\omega)v_0 + a_2(\omega)v_2 + \cdots + a_n(\omega)v_n$$

and pairing with the dual basis of the Betti realization gives the proof of Theorem 1.1, with the cycles $(\sigma_2, \dots, \sigma_n)$ chosen as representatives of the classes $(\varphi_2, \dots, \varphi_n)$.

The involution (4) plays an important role in the proof of Theorem 1.3. It induces a natural involution, still denoted by τ , on the quotient $\mathcal{Z}^{(n)}/\mathbb{Q}(0) \cong \mathbb{Q}(-2) \oplus \cdots \oplus \mathbb{Q}(-n)$.

THEOREM 1.4. *For $k = 2, \dots, n$, the involution τ acts on the direct summand $\mathbb{Q}(-k)$ of $\mathcal{Z}^{(n)}/\mathbb{Q}(0)$ by multiplication by $(-1)^{k-1}$.*

This readily implies Theorem 1.2. Now if we write

$$\mathcal{Z}^{(n)}/\mathbb{Q}(0) = (\mathcal{Z}^{(n)}/\mathbb{Q}(0))_+ \oplus (\mathcal{Z}^{(n)}/\mathbb{Q}(0))_-$$

for the decomposition into invariant and anti-invariants with respect to τ and write $p : \mathcal{Z}^{(n)} \rightarrow \mathcal{Z}^{(n)}/\mathbb{Q}(0)$ for the natural projection, we may set

$$\mathcal{Z}^{(n), \text{odd}} := p^{-1}((\mathcal{Z}^{(n)}/\mathbb{Q}(0))_+)$$

whose period matrix only contains odd zeta values in the first row. The objects $\mathcal{Z}^{(n), \text{odd}} \in \text{MT}(\mathbb{Z})$ form an inductive system, and the limit

$$\mathcal{Z}^{\text{odd}} := \varinjlim_n \mathcal{Z}^{(n), \text{odd}}$$

has an infinite period matrix

$$\begin{pmatrix} 1 & \zeta(3) & \zeta(5) & \zeta(7) & \cdots & \cdots & \cdots \\ & (2\pi i)^3 & & & & & \\ & & (2\pi i)^5 & & & 0 & \\ & & & (2\pi i)^7 & & & \\ & & & & \ddots & & \\ & 0 & & & & \ddots & \\ & & & & & & \ddots \end{pmatrix}. \tag{6}$$

We call \mathcal{Z}^{odd} the *odd zeta motive*.

1.3 Related work and open questions

This article follows the program initiated by Brown [Bro16], which aims at explaining and possibly producing irrationality proofs for zeta values by means of algebraic geometry. However, the motives that we are considering are different from the general motives considered by Brown, and in particular, easier to compute. It would be interesting to determine the precise relationship between our motives and those defined in [Bro16] in terms of the moduli spaces $\mathcal{M}_{0,n+3}$.

In another direction, an explicit description of the relative cycles defined in Theorem 1.1 could prove helpful in proving quantitative results on the irrationality measures of zeta values, in the spirit of [RV96, RV01].

It is also tempting to apply our methods to other families of integrals appearing in the literature, such as the Beukers integrals for $\zeta(3)$ and their generalizations. One should be able, for instance, to recover Rhin and Viola's contour integrals for $\zeta(3)$ [RV01, Theorem 3.1]. The symmetry properties studied by Cresson *et al.* [CFR08a] can probably be explained geometrically via finite group actions as in the present article. The *ad hoc* long exact sequences appearing here should be replaced by more systematic tools such as the Orlik–Solomon bi-complexes from [Dup17].

Finally, it should be possible to extend our results to a functional version of the periods (2), where one replaces $1 - x_1 \cdots x_n$ in the denominator by $1 - z x_1 \cdots x_n$, with z a complex parameter. Such functions have already been considered in [Riv00, BR01]. The relevant geometric objects are variations of mixed Hodge–Tate structures on $\mathbb{C} - \{0, 1\}$, or mixed Tate motives over $\mathbb{A}_{\mathbb{Q}}^1 - \{0, 1\}$.

1.4 Contents

In §2 we recall some general facts about the categories in which the objects that we will be considering live, and in particular the categories $\text{MT}(\mathbb{Z})$ and $\text{MT}(\mathbb{Q})$ of mixed Tate motives over \mathbb{Z} and \mathbb{Q} . In §3 we introduce the zeta motives and examine their Betti and de Rham realizations. In §4, which is more technical than the rest of the paper, we compute the full period matrix of the zeta motives, which allows us to define the odd zeta motives. In §5, we apply our results to proving Theorems 1.1 and 1.2 on the coefficients of linear forms in zeta values.

2. Mixed Tate motives and their period matrices

We recall the construction of the categories MHTS , $\text{MT}(\mathbb{Q})$ and $\text{MT}(\mathbb{Z})$, which sit as full subcategories of one another, as follows:

$$\text{MT}(\mathbb{Z}) \hookrightarrow \text{MT}(\mathbb{Q}) \hookrightarrow \text{MHTS}.$$

2.1 Mixed Hodge–Tate structures and their period matrices

DEFINITION 2.1. A *mixed Hodge–Tate structure* is a triple $H = (H_{\text{dR}}, H_{\text{B}}, \alpha)$ consisting of:

- a finite-dimensional \mathbb{Q} -vector space H_{B} , together with a finite increasing filtration indexed by even integers: $\cdots \subset W_{2(n-1)}H_{\text{B}} \subset W_{2n}H_{\text{B}} \subset \cdots \subset H_{\text{B}}$;
- a finite-dimensional \mathbb{Q} -vector space H_{dR} , together with a grading indexed by even integers: $H_{\text{dR}} = \bigoplus_n (H_{\text{dR}})_{2n}$;
- an isomorphism $\alpha : H_{\text{dR}} \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\simeq} H_{\text{B}} \otimes_{\mathbb{Q}} \mathbb{C}$;

which satisfy the following conditions.

- For every integer n , the isomorphism α sends $(H_{\text{dR}})_{2n} \otimes_{\mathbb{Q}} \mathbb{C}$ to $W_{2n}H_{\text{B}} \otimes_{\mathbb{Q}} \mathbb{C}$.
- For every integer n , it induces an isomorphism $\alpha_n : (H_{\text{dR}})_{2n} \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\cong} (W_{2n}H_{\text{B}}/W_{2(n-1)}H_{\text{B}}) \otimes_{\mathbb{Q}} \mathbb{C}$, which sends $(H_{\text{dR}})_{2n}$ to $(W_{2n}H_{\text{B}}/W_{2(n-1)}H_{\text{B}}) \otimes_{\mathbb{Q}} (2\pi i)^n \mathbb{Q}$.

We call H_{B} and H_{dR} the *Betti realization* and the *de Rham realization* of the mixed Hodge–Tate structure, and α the *comparison isomorphism*. The filtration W on H_{B} is called the *weight filtration*. The grading on H_{dR} is called the *weight grading*, and the corresponding filtration $W_{2n}H_{\text{dR}} := \bigoplus_{k \leq n} (H_{\text{dR}})_{2k}$ the *weight filtration*.

Remark 2.2. More classically, a mixed Hodge–Tate structure is defined to be a mixed Hodge structure [Del71, Del74] whose weight-graded quotients are of Tate type, i.e., of type (p, p) for some integer p . One passes from that classical definition to Definition 2.1 by setting $H_{\text{B}} := H$ and $H_{\text{dR}} := \bigoplus_n W_{2n}H/W_{2(n-1)}H$. The isomorphism α is induced by the inverses of the isomorphisms

$$(W_{2n}H/W_{2(n-1)}H) \otimes_{\mathbb{Q}} \mathbb{C} \xleftarrow{\cong} (W_{2n}H \otimes_{\mathbb{Q}} \mathbb{C}) \cap F^n(H \otimes_{\mathbb{Q}} \mathbb{C}) \tag{7}$$

(multiplied by $(2\pi i)^n$) which express the fact that the weight-graded quotients are of Tate type.

It is convenient to view the comparison isomorphism $\alpha : H_{\text{dR}} \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\cong} H_{\text{B}} \otimes_{\mathbb{Q}} \mathbb{C}$ as a pairing

$$H_{\text{B}}^{\vee} \otimes_{\mathbb{Q}} H_{\text{dR}} \longrightarrow \mathbb{C}, \quad \varphi \otimes v \mapsto \langle \varphi, v \rangle, \tag{8}$$

where $(\cdot)^{\vee}$ denotes the linear dual. The weight filtration on H_{B}^{\vee} is defined by

$$W_{-2n}H_{\text{B}}^{\vee} := (H_{\text{B}}/W_{2(n-1)}H_{\text{B}})^{\vee},$$

so that we have

$$W_{-2n}H_{\text{B}}^{\vee}/W_{-2(n+1)}H_{\text{B}}^{\vee} \cong (W_{2n}H_{\text{B}}/W_{2(n-1)}H_{\text{B}})^{\vee}.$$

The pairing (8) is compatible with the weight filtrations in that we have $\langle \varphi, v \rangle = 0$ for $\varphi \in W_{-2m}H_{\text{B}}^{\vee}$, $v \in W_{2n}H_{\text{dR}}$ and $m < n$.

If we choose bases for the \mathbb{Q} -vector spaces H_{dR} and H_{B} , then the matrix of α in these bases, or equivalently the matrix of the pairing (8), is called a *period matrix* of the mixed Hodge–Tate structure. We will always make the following assumptions on the choice of bases.

- The basis of H_{B} is compatible with the weight filtration.
- The basis of H_{dR} is compatible with the weight grading.
- For every n , the matrix of the comparison isomorphism α_n in the corresponding basis is $(2\pi i)^n$ times the identity.

This implies that any period matrix is block upper-triangular with successive blocks of $(2\pi i)^n \text{Id}$ on the diagonal. Conversely, any block upper-triangular matrix with successive blocks of $(2\pi i)^n \text{Id}$ on the diagonal is a period matrix of a mixed Hodge–Tate structure.

Example 2.3. Any matrix of the form

$$\begin{pmatrix} 1 & * & * & * & * \\ 0 & 2\pi i & 0 & * & * \\ 0 & 0 & 2\pi i & * & * \\ 0 & 0 & 0 & (2\pi i)^2 & 0 \\ 0 & 0 & 0 & 0 & (2\pi i)^2 \end{pmatrix}$$

defines a mixed Hodge–Tate structure H such that $H_{\text{dR}} = (H_{\text{dR}})_0 \oplus (H_{\text{dR}})_2 \oplus (H_{\text{dR}})_4$ has graded dimension $(1, 2, 2)$.

2.2 The category of mixed Hodge–Tate structures

We denote by MHTS the category of mixed Hodge–Tate structures. It is a neutral tannakian category over \mathbb{Q} , which means in particular that it is an abelian \mathbb{Q} -linear category equipped with a \mathbb{Q} -linear tensor product \otimes . We note that an object $H \in \text{MHTS}$ is endowed with a canonical weight filtration W by subobjects such that the morphisms in MHTS are strictly compatible with W . We have two natural fiber functors

$$\omega_{\text{B}} : \text{MHTS} \rightarrow \text{Vect}_{\mathbb{Q}} \quad \text{and} \quad \omega_{\text{dR}} : \text{MHTS} \rightarrow \text{Vect}_{\mathbb{Q}} \tag{9}$$

from MHTS to the category of finite-dimensional vector spaces over \mathbb{Q} , which only remember the Betti realization H_{B} and the de Rham realization H_{dR} respectively. We note that the de Rham realization functor ω_{dR} factors through the category of finite-dimensional graded vector spaces. The comparison isomorphisms α gives an isomorphism between the complexifications of the two fiber functors:

$$\omega_{\text{dR}} \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\cong} \omega_{\text{B}} \otimes_{\mathbb{Q}} \mathbb{C}. \tag{10}$$

For an integer n , we denote by $\mathbb{Q}(-n)$ the mixed Hodge–Tate structure whose period matrix is the 1×1 matrix $((2\pi i)^n)$. Its weight grading and filtration are concentrated in weight $2n$, hence we call it the *pure Tate structure* of weight $2n$. For H a mixed Hodge–Tate structure, the tensor product $H \otimes \mathbb{Q}(-n)$ is simply denoted by $H(-n)$ and called the *n th Tate twist* of H . A period matrix of $H(-n)$ is obtained by multiplying a period matrix of H by $(2\pi i)^n$. The weight grading and filtration of $H(-n)$ are those of H , shifted by $2n$.

2.3 Extensions between pure Tate structures

The pure Tate structures $\mathbb{Q}(-n)$ are the only simple objects of the category MHTS . The extensions between them are easily described. Up to a Tate twist, it is enough to describe the extensions of $\mathbb{Q}(-n)$ by $\mathbb{Q}(0)$ for some integer n . The corresponding extension group is given by

$$\text{Ext}_{\text{MHTS}}^1(\mathbb{Q}(-n), \mathbb{Q}(0)) = \begin{cases} \mathbb{C}/(2\pi i)^n \mathbb{Q} & \text{if } n > 0, \\ 0 & \text{otherwise.} \end{cases}$$

More concretely, the extension corresponding to a number $z \in \mathbb{C}/(2\pi i)^n \mathbb{Q}$ has a period matrix

$$\begin{pmatrix} 1 & z \\ 0 & (2\pi i)^n \end{pmatrix}.$$

We note that the higher extension groups vanish: $\text{Ext}_{\text{MHTS}}^r(H, H') = 0$ for $r \geq 2$ and H, H' two mixed Hodge–Tate structures.

Example 2.4. For a complex number $a \in \mathbb{C} - \{0, 1\}$, the cohomology group $H^1(\mathbb{C}^*, \{1, a\})$ is an extension of $\mathbb{Q}(-1)$ by $\mathbb{Q}(0)$ corresponding to $z = \log(a) \in \mathbb{C}/(2\pi i)\mathbb{Q}$. It is called the *Kummer extension* of parameter a .

2.4 Mixed Tate motives over \mathbb{Q}

Let $DM(\mathbb{Q})$ denote Voevodsky’s triangulated category of motives over \mathbb{Q} [Voe00]. It is a \mathbb{Q} -linear triangulated tensor category whose objects can be described in terms of complexes of varieties and whose morphisms can be described in terms of algebraic cycles (in particular, in terms of Bloch’s higher Chow groups). There are invertible objects $\mathbb{Q}(-n) \in DM(\mathbb{Q})$, where $\mathbb{Q}(-1)$ is the reduced motive of the multiplicative group \mathbb{G}_m , shifted by -1 (we work with cohomological conventions). The triangulated subcategory of $DM(\mathbb{Q})$ generated by these objects is denoted by $DMT(\mathbb{Q})$. By using the relation between higher Chow groups and rational K -theory [Blo86] and Borel’s computation of the rational K -theory of number fields [Bor77], Levine defined a natural t -structure on $DMT(\mathbb{Q})$ [Lev93]. The heart of this t -structure is denoted by $MT(\mathbb{Q})$ and called the category of *mixed Tate motives* over \mathbb{Q} . It is a (neutral) tannakian \mathbb{Q} -linear category which contains the objects $\mathbb{Q}(-n)$.

There is a faithful and exact functor

$$MT(\mathbb{Q}) \rightarrow MHTS \tag{11}$$

from $MT(\mathbb{Q})$ to the category $MHTS$ of mixed Hodge–Tate structures, which is called the *Hodge realization functor* ([DG05, § 2.13], see also [Hub00, Hub04]). It sends the object $\mathbb{Q}(-n) \in MT(\mathbb{Q})$ to the object $\mathbb{Q}(-n) \in MHTS$. Composing (11) with the fiber functors (9) gives the Betti and de Rham realization functors, still denoted by

$$\omega_B : MT(\mathbb{Q}) \rightarrow \text{Vect}_{\mathbb{Q}} \quad \text{and} \quad \omega_{dR} : MT(\mathbb{Q}) \rightarrow \text{Vect}_{\mathbb{Q}}, \tag{12}$$

and we still have a comparison isomorphism (10). We note that any object in $MT(\mathbb{Q})$ is endowed with a canonical weight filtration W by subobjects such that the morphisms in $MT(\mathbb{Q})$ are strictly compatible with W . The realization morphisms are compatible with the weight filtrations.

Remark 2.5. The functors (12) are fiber functors for the tannakian category $MT(\mathbb{Q})$. In particular, they are conservative.

The extension groups between the objects $\mathbb{Q}(-n)$ are computed by the rational K -theory of \mathbb{Q} [Lev93, § 4] and hence given, after Borel [Bor77], by

$$\text{Ext}_{MT(\mathbb{Q})}^1(\mathbb{Q}(-n), \mathbb{Q}(0)) = \begin{cases} \bigoplus_{p \text{ prime}} \mathbb{Q} & \text{if } n = 1, \\ \mathbb{Q} & \text{if } n \text{ is odd } \geq 3, \\ 0 & \text{otherwise.} \end{cases} \tag{13}$$

As in the category $MHTS$, the higher extension groups vanish in $MT(\mathbb{Q})$. The morphisms

$$\text{Ext}_{MT(\mathbb{Q})}^1(\mathbb{Q}(-n), \mathbb{Q}(0)) \longrightarrow \text{Ext}_{MHTS}^1(\mathbb{Q}(-n), \mathbb{Q}(0)) \cong \mathbb{C}/(2\pi i)^n \mathbb{Q} \tag{14}$$

induced by (11) are easy to describe. For $n = 1$, the image of the direct summand indexed by a prime p is the line spanned by the class of the Kummer extension of parameter p , i.e., by $\log(p) \in \mathbb{C}/(2\pi i)\mathbb{Q}$. For $n \geq 3$ odd, the image is the line spanned by $\zeta(n) \in \mathbb{C}/(2\pi i)^n \mathbb{Q}$. Thus, the morphism (14) is injective for every n . This implies the following theorem [DG05, Proposition 2.14].

THEOREM 2.6. *The realization functor (11) is fully faithful.*

This theorem is very helpful, since it allows one to compute in the category $MT(\mathbb{Q})$ with period matrices; in other words, a mixed Tate motive over \mathbb{Q} is uniquely determined by its period matrix.

2.5 Mixed Tate motives over \mathbb{Z}

Let $\text{MT}(\mathbb{Z})$ denote the category of mixed Tate motives over \mathbb{Z} , as defined in [DG05]. By definition, it is a full tannakian subcategory

$$\text{MT}(\mathbb{Z}) \hookrightarrow \text{MT}(\mathbb{Q})$$

of the category of mixed Tate motives over \mathbb{Q} , which contains the pure Tate motives $\mathbb{Q}(-n)$ for every integer n . An object of $\text{MT}(\mathbb{Q})$ is in $\text{MT}(\mathbb{Z})$ if and only if it has no subquotient isomorphic to a non-split extension of $\mathbb{Q}(-n)$ by $\mathbb{Q}(-n + 1)$.

The extension groups in the category $\text{MT}(\mathbb{Z})$ satisfy the following properties:

- (1) $\text{Ext}_{\text{MT}(\mathbb{Z})}^1(\mathbb{Q}(-1), \mathbb{Q}(0)) = 0$;
- (2) the natural morphism $\text{Ext}_{\text{MT}(\mathbb{Z})}^1(\mathbb{Q}(-n), \mathbb{Q}(0)) \rightarrow \text{Ext}_{\text{MT}(\mathbb{Q})}^1(\mathbb{Q}(-n), \mathbb{Q}(0))$ is an isomorphism for $n \neq 1$.

As in the categories MHTS and $\text{MT}(\mathbb{Q})$, the higher extension groups vanish in $\text{MT}(\mathbb{Z})$.

For $n \geq 3$ odd, there is an essentially unique non-trivial extension of $\mathbb{Q}(-n)$ by $\mathbb{Q}(0)$ in the category $\text{MT}(\mathbb{Q})$, which actually lives in $\text{MT}(\mathbb{Z})$. A period matrix for such an extension is

$$\begin{pmatrix} 1 & \zeta(n) \\ 0 & (2\pi i)^n \end{pmatrix}.$$

Apart from the case $n = 3$ (see [Bro16, Corollary 11.3] or Proposition 4.11 below), we do not know of any *geometric* construction of these extensions.

3. Definition of the zeta motives $\mathcal{Z}^{(n)}$

We define the zeta motives $\mathcal{Z}^{(n)}$ and explain how to define elements of their Betti and de Rham realizations. In particular, we define the classes of the Eulerian differential forms, which are elements of the de Rham realization $\mathcal{Z}_{\text{dR}}^{(n)}$ constructed out of the family of Eulerian polynomials. We also note that the zeta motives fit into an inductive system $\dots \rightarrow \mathcal{Z}^{(n-1)} \rightarrow \mathcal{Z}^{(n)} \rightarrow \dots$ which is compatible with the Eulerian differential forms.

3.1 The definition

Let $n \geq 1$ be an integer. In the affine n -space $X_n = \mathbb{A}_{\mathbb{Q}}^n$ we consider the hypersurfaces

$$A_n = \{x_1 \cdots x_n = 1\} \quad \text{and} \\ B_n = \bigcup_{1 \leq i \leq n} \{x_i = 0\} \cup \bigcup_{1 \leq i \leq n} \{x_i = 1\}.$$

The union $A_n \cup B_n$ is almost a normal crossing divisor inside X_n : around the point $P_n = (1, \dots, 1)$, it looks like $z_1 \cdots z_n(z_1 + \cdots + z_n) = 0$ (set $x_i = \exp(z_i)$). Let

$$\pi_n : \tilde{X}_n \rightarrow X_n$$

be the blow-up at P_n , and $E_n = \pi_n^{-1}(P_n)$ be the exceptional divisor. We denote respectively by \tilde{A}_n and \tilde{B}_n the strict transforms of A_n and B_n along π_n . The union $\tilde{A}_n \cup \tilde{B}_n \cup E_n$ is a simple normal crossing divisor inside \tilde{X}_n .

There is an object $\mathcal{Z}^{(n)} \in \text{MT}(\mathbb{Q})$, which we may abusively denote by

$$\mathcal{Z}^{(n)} = H^n(\tilde{X}_n - \tilde{A}_n, (\tilde{B}_n \cup E_n) - (\tilde{B}_n \cup E_n) \cap \tilde{A}_n),$$

such that its Betti and de Rham realizations (12) are ($? \in \{B, dR\}$)

$$\mathcal{Z}_?^{(n)} = H_?^n(\tilde{X}_n - \tilde{A}_n, (\tilde{B}_n \cup E_n) - (\tilde{B}_n \cup E_n) \cap \tilde{A}_n).$$

We now give the precise definition of $\mathcal{Z}^{(n)}$, along the lines of [Gon02, Proposition 3.6]. Let us write $Y = \tilde{X}_n - \tilde{A}_n$ and $\partial Y = (\tilde{B}_n \cup E_n) - (\tilde{B}_n \cup E_n) \cap \tilde{A}_n$, viewed as schemes defined over \mathbb{Q} . We have a decomposition into smooth irreducible components $\partial Y = \bigcup_i \partial_i Y$, where i runs in a set of cardinality $2n + 1$. For a set $I = \{i_1, \dots, i_r\}$ of indices, we denote by $\partial_I Y = \partial_{i_1} Y \cap \dots \cap \partial_{i_r} Y$ the corresponding intersection; it is either empty or a smooth subvariety of X of codimension r .

We thus get an object

$$\dots \rightarrow \bigsqcup_{|I|=3} \partial_I Y \rightarrow \bigsqcup_{|I|=2} \partial_I Y \rightarrow \bigsqcup_{|I|=1} \partial_I Y \rightarrow Y \rightarrow 0 \tag{15}$$

in Voevodsky’s triangulated category $DM(\mathbb{Q})$, see §2.4. The differentials are the alternating sums of the natural closed immersions. One readily checks that the complex (15) lives in the triangulated subcategory $DMT(\mathbb{Q})$. By definition, the object $\mathcal{Z}^{(n)}$ in $MT(\mathbb{Q})$ is the n th cohomology group of the complex (15) with respect to the t -structure.

DEFINITION 3.1. For $n \geq 1$, we call $\mathcal{Z}^{(n)} \in MT(\mathbb{Q})$ the n th zeta motive.

Remark 3.2. For $n = 1$, the blow-up map $\pi_1 : \tilde{X}_1 \rightarrow X_1$ is an isomorphism and $\tilde{A}_1 = \emptyset$, so that we get $\mathcal{Z}^{(1)} = H^1(\mathbb{A}_{\mathbb{Q}}^1, \{0, 1\})$. We have a long exact sequence in relative cohomology

$$0 \rightarrow H^0(\mathbb{A}_{\mathbb{Q}}^1, \{0, 1\}) \rightarrow H^0(\mathbb{A}_{\mathbb{Q}}^1) \rightarrow H^0(\{0\}) \oplus H^0(\{1\}) \rightarrow \mathcal{Z}^{(1)} \rightarrow 0,$$

which shows that $H^0(\mathbb{A}_{\mathbb{Q}}^1, \{0, 1\}) = 0$ and that we have an isomorphism $\mathcal{Z}^{(1)} \simeq \mathbb{Q}(0)$.

Remark 3.3. We will prove in Proposition 4.12 that $\mathcal{Z}^{(n)}$ is actually an object of the full subcategory $MT(\mathbb{Z}) \hookrightarrow MT(\mathbb{Q})$. It would be possible, but a little technical, to prove it directly from the definition by using the criterion [GM04, Proposition 4.3] on some compactification of $\tilde{X}_n - \tilde{A}_n$.

3.2 Betti and de Rham realizations, 1

We now give a first description of the Betti and de Rham realizations of the zeta motive $\mathcal{Z}^{(n)}$.

We let C_{\bullet} denote the functor which assigns to a topological space the complex of singular chains with rational coefficients. By definition, the dual of the Betti realization $\mathcal{Z}_B^{(n), \vee}$ is the n th homology group of the total complex of the double complex

$$\begin{array}{ccccc}
 \cdots \rightarrow & \bigoplus_{|I|=2} C_0(\partial_I Y(\mathbb{C})) & \longrightarrow & \bigoplus_{|I|=1} C_0(\partial_I Y(\mathbb{C})) & \longrightarrow & C_0(Y(\mathbb{C})) \\
 & \uparrow & & \uparrow & & \uparrow \\
 & \cdots \rightarrow & \bigoplus_{|I|=1} C_1(\partial_I Y(\mathbb{C})) & \longrightarrow & C_1(Y(\mathbb{C})) & \\
 & & \uparrow & & \uparrow & \\
 & & \cdots \rightarrow & C_2(Y(\mathbb{C})) & & \\
 & & & \uparrow & & \\
 & & & \cdots & &
 \end{array} \tag{16}$$

obtained by applying the functor C_\bullet to the complex (15). One readily verifies that this complex is quasi-isomorphic to the quotient complex $C_\bullet(Y(\mathbb{C}))/C_\bullet(\partial Y(\mathbb{C}))$, classically used to define the relative homology groups $H_\bullet^B(Y, \partial Y) = H_\bullet^{\text{sing}}(Y(\mathbb{C}), \partial Y(\mathbb{C}))$.

We let $\Omega_{\partial_I Y}^\bullet$ denote the complex of sheaves of algebraic differential forms on the smooth variety $\partial_I Y$, extended by zero to Y . By definition, the de Rham realization $\mathcal{Z}_{\text{dR}}^{(n)}$ is the hypercohomology of the total complex of the double complex of sheaves

$$\begin{array}{ccccc}
 \cdots & \bigoplus_{|I|=2} \Omega_{\partial_I Y}^0 & \longleftarrow & \bigoplus_{|I|=1} \Omega_{\partial_I Y}^0 & \longleftarrow & \Omega_Y^0 \\
 & \downarrow & & \downarrow & & \downarrow \\
 & \cdots & & \bigoplus_{|I|=1} \Omega_{\partial_I Y}^1 & \longleftarrow & \Omega_Y^1 \\
 & & & \downarrow & & \downarrow \\
 & & & \cdots & & \Omega_Y^2 \\
 & & & & & \downarrow \\
 & & & & & \cdots
 \end{array} \tag{17}$$

where the vertical arrows are the exterior derivatives and the horizontal arrows are the alternating sums of the natural restriction maps as in the complex (15).

The comparison morphism between the Betti and de Rham realizations of $\mathcal{Z}^{(n)}$ is induced, after complexification, by the morphism from the double complex (17) to the double complex (16) given by integration. Note that one first has to replace (16) by the double complex of sheaves of singular cochains.

3.3 Betti and de Rham realizations, 2

We now give descriptions of the Betti and de Rham realizations of $\mathcal{Z}^{(n)}$ that allow one to work directly in the affine space X_n and do not require to work in the blow-up \tilde{X}_n . The justification of the blow-up process goes as follows. Suppose that one wants to find a motive whose periods include all absolutely convergent integrals of the form

$$\int_{[0,1]^n} \frac{P(x_1, \dots, x_n)}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n, \tag{18}$$

where $P(x_1, \dots, x_n)$ is a polynomial with rational coefficients, and $N \geq 0$ is an integer. On the Betti side, we note that the boundary of $[0, 1]^n$ intersects the divisor $A_n(\mathbb{C})$ of poles of the differential forms at the point $P_n(\mathbb{C})$. The blow-up process is thus required in order to have a class that represents the integration domain. On the de Rham side, the blow-up process is required in order to only consider *absolutely convergent* integrals of the form (18). This is made precise by Propositions 3.4 and 3.6 below.

We start with the Betti realization. Let us write $\mathring{A}_n = A_n - P_n$ and note that this is not a closed subset, but only a locally closed subset, of X_n .

PROPOSITION 3.4. *The blow-up morphism $\pi_n : \tilde{X}_n \rightarrow X_n$ induces an isomorphism*

$$\mathcal{Z}_B^{(n), \vee} \xrightarrow{\cong} H_n^{\text{sing}}(X_n(\mathbb{C}) - \mathring{A}_n(\mathbb{C}), B_n(\mathbb{C}) - B_n(\mathbb{C}) \cap \mathring{A}_n(\mathbb{C})).$$

Proof. The blow-up morphism π_n is the contraction of the exceptional divisor E_n onto the point P_n . Thus, this is a consequence of the classical excision theorem in singular homology, see for instance [Hat02, Proposition 2.22]. \square

As a consequence of Proposition 3.4, we see that the unit n -square $\square^n = [0, 1]^n \subset X_n(\mathbb{C}) - \mathring{A}_n(\mathbb{C})$ defines a class

$$[\square^n] \in \mathcal{Z}_{\mathbb{B}}^{(n), \vee}.$$

When viewed in $\tilde{X}_n(\mathbb{C}) - \tilde{A}_n(\mathbb{C})$, it is the class of the strict transform $\tilde{\square}^n$, which has the combinatorial structure of an n -cube truncated at one of its vertices.

We now turn to a description of the de Rham realization of $\mathcal{Z}^{(n)}$. Instead of giving a general description in terms of algebraic differential forms on $X_n - A_n$, we will only give a way of defining many classes in $\mathcal{Z}_{\text{dR}}^{(n)}$, which will turn out to be enough for our purposes.

DEFINITION 3.5. An algebraic differential n -form on $X_n - A_n$ is said to be *integrable* if it can be written as a linear combination of forms of the type

$$\omega = \frac{(1 - x_1)^{v_1 - 1} \cdots (1 - x_n)^{v_n - 1} f(x_1, \dots, x_n)}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n \tag{19}$$

with $v_1, \dots, v_n \geq 1$ and $N \geq 0$ integers such that $v_1 + \cdots + v_n \geq N + 1$, and $f(x_1, \dots, x_n)$ a polynomial with rational coefficients.

The terminology is justified by the following proposition.

PROPOSITION 3.6. *Let ω be an algebraic differential n -form on $X_n - A_n$. If ω is integrable, then $\pi_n^*(\omega)$ does not have a pole along E_n , and thus defines a class in $\mathcal{Z}_{\text{dR}}^{(n)}$. In particular, the integral*

$$\int_{\tilde{\square}^n} \pi_n^*(\omega) = \int_{\square^n} \omega$$

is absolutely convergent and is a period of $\mathcal{Z}^{(n)}$.

Proof. We write ω as in (19). We note that the only problem for absolute convergence is around the point $(1, \dots, 1)$. Let us thus make the change of variables $y_i = 1 - x_i$ for $i = 1, \dots, n$, and $g(y_1, \dots, y_n) = (-1)^n f(x_1, \dots, x_n)$. We write $h(y_1, \dots, y_n) = 1 - (1 - y_1) \cdots (1 - y_n)$ so that we have

$$\omega = \frac{y_1^{v_1 - 1} \cdots y_n^{v_n - 1} g(y_1, \dots, y_n)}{h(y_1, \dots, y_n)^N} dy_1 \cdots dy_n.$$

There are n natural affine charts for the blow-up $\pi_n : \tilde{X}_n \rightarrow X_n$ of the point $(0, \dots, 0)$, and by symmetry it is enough to work in the first one. We then have local coordinates (z_1, \dots, z_n) on \tilde{X}_n , which are linked to the coordinates $(y_1, \dots, y_n) = \pi_n(z_1, \dots, z_n)$ by the formula

$$(y_1, \dots, y_n) = (z_1, z_1 z_2, \dots, z_1 z_n).$$

The problem of convergence occurs in the neighborhood of the exceptional divisor E_n , which is defined by the equation $z_1 = 0$. Since $h(0, \dots, 0) = 0$, we may write

$$h(z_1, z_1 z_2, \dots, z_1 z_n) = z_1 \tilde{h}(z_1, \dots, z_n)$$

with $\tilde{h}(z_1, \dots, z_n)$ a polynomial such that $\tilde{h}(0, \dots, 0) = 1$. The strict transform \tilde{A}_n of A_n is thus defined by the equation $\tilde{h}(z_1, \dots, z_n) = 0$. We note that we have $dy_1 \cdots dy_n = z_1^{n-1} dz_1 \cdots dz_n$, so that we can write

$$\pi_n^*(\omega) = \frac{z_1^{v_1-1} (z_1 z_2)^{v_2-1} \cdots (z_1 z_n)^{v_n-1} g(z_1, z_1 z_2, \dots, z_1 z_n)}{z_1^N \tilde{h}(z_1, \dots, z_n)^N} z_1^{n-1} dz_1 \cdots dz_n = z_1^{v_1+\cdots+v_n-N-1} \Omega,$$

where Ω has a pole along \tilde{A}_n but not along E_n . The claim follows. □

We make an abuse of notation and denote by

$$[\omega] \in \mathcal{Z}_{\text{dR}}^{(n)}$$

the class of the pullback $\pi_n^*(\omega)$ for ω integrable, so that the comparison isomorphism reads

$$\langle [\square^n], [\omega] \rangle = \int_{\square^n} \omega.$$

We note the converse of Proposition 3.6, which we will not use.

PROPOSITION 3.7. *Let ω be an algebraic differential n -form on $X_n - A_n$. If the integral $\int_{\square^n} \omega$ is absolutely convergent, then ω is integrable.*

Proof. In the coordinates $y_i = 1 - x_i$, we write

$$\omega = \frac{P(y_1, \dots, y_n)}{h(y_1, \dots, y_n)^N} dy_1 \cdots dy_n$$

with $P(y_1, \dots, y_n)$ a polynomial with rational coefficients. If the integral $\int_{\square^n} \omega$ is absolutely convergent in the neighborhood of the point $(0, \dots, 0)$, then after the change of variables

$$\phi(z_1, \dots, z_n) = (z_1, z_1 z_2, \dots, z_1 z_n)$$

we get an absolutely convergent integral in the neighborhood of $z_1 = 0$. We write, as in the proof of Proposition 3.6,

$$\phi^*(\omega) = \frac{P(z_1, z_1 z_2, \dots, z_1 z_n)}{z_1^{N-n+1} \tilde{h}(z_1, \dots, z_n)^N} dz_1 \cdots dz_n.$$

Let us write

$$P(y_1, \dots, y_n) = \sum_{\underline{a}} \lambda_{\underline{a}} y_1^{a_1-1} \cdots y_n^{a_n-1}$$

with $\lambda_{\underline{a}} \in \mathbb{Q}$ for every multi-index $\underline{a} = (a_1, \dots, a_n)$. We then have

$$P(z_1, z_1 z_2, \dots, z_1 z_n) = \sum_{\underline{a}} \lambda_{\underline{a}} z_1^{a_1+\cdots+a_n-n} z_2^{a_2-1} \cdots z_n^{a_n-1}.$$

Let v denote the smallest integer such that there exists a multi-index \underline{a} with $|\underline{a}| := a_1 + \cdots + a_n = v$. We then have an equivalence

$$P(z_1, z_1 z_2, \dots, z_1 z_n) \sim_{z_1 \rightarrow 0} z_1^{v-n} Q(z_2, \dots, z_n),$$

where $Q(z_2, \dots, z_n) = \sum_{|\underline{a}|=v} \lambda_{\underline{a}} z_2^{a_2-1} \cdots z_n^{a_n-1}$. We also have the equivalence

$$\tilde{h}(z_1, \dots, z_n) \sim_{z_1 \rightarrow 0} 1 + z_2 + \cdots + z_n,$$

from which we deduce

$$\phi^*(\omega) \sim_{z_1 \rightarrow 0} z_1^{v-N-1} dz_1 \frac{Q(z_2, \dots, z_n)}{(1 + z_2 + \cdots + z_n)^N} dz_2 \cdots dz_n.$$

This gives an absolutely convergent integral in the neighborhood of $z_1 = 0$ if and only if $v \geq N+1$, which is exactly the integrability condition. □

3.4 The Eulerian differential forms

Recall that the family of *Eulerian polynomials* $E_r(x)$, $r \geq 0$, is defined by the equation

$$\frac{E_r(x)}{(1-x)^{r+1}} = \sum_{j \geq 0} (j+1)^r x^j. \tag{20}$$

We refer to [Foa10] for a survey on Eulerian polynomials. If $r \geq 1$, then (20) is equivalent to

$$\frac{E_r(x)}{(1-x)^{r+1}} = \frac{1}{x} \left(x \frac{d}{dx} \right)^r \frac{1}{1-x}.$$

For instance, we have $E_0(x) = E_1(x) = 1$, $E_2(x) = 1 + x$, $E_3(x) = 1 + 4x + x^2$. The Eulerian polynomials satisfy the recurrence relation

$$E_{r+1}(x) = x(1-x)E'_r(x) + (1+rx)E_r(x). \tag{21}$$

For integers $n \geq 2$ and $k = 2, \dots, n$, we define a differential form

$$\omega_k^{(n)} = \frac{E_{n-k}(x_1 \cdots x_n)}{(1-x_1 \cdots x_n)^{n-k+1}} dx_1 \cdots dx_n.$$

Note that we have $\omega_n^{(n)} = dx_1 \cdots dx_n / (1-x_1 \cdots x_n)$.

LEMMA 3.8. For $k = 2, \dots, n$, the form $\omega_k^{(n)}$ defines a class $[\omega_k^{(n)}] \in \mathcal{Z}_{\text{dR}}^{(n)}$ and we have

$$\langle [\square^n], [\omega_k^{(n)}] \rangle = \int_{\square^n} \omega_k^{(n)} = \zeta(k). \tag{22}$$

Proof. The first statement follows from Proposition 3.6. The computation of the period is then straightforward using the definition (20) of the Eulerian polynomials:

$$\int_{\square^n} \omega_k^{(n)} = \sum_{j \geq 0} (j+1)^{n-k} \int_{[0,1]^n} (x_1 \cdots x_n)^j dx_1 \cdots dx_n = \sum_{j \geq 0} (j+1)^{-k} = \zeta(k). \quad \square$$

For every $n \geq 0$, we define $\omega_0^{(n)} = dx_1 \cdots dx_n$; we also have the class $[\omega_0^{(n)}] \in \mathcal{Z}_{n,\text{dR}}$, whose pairing with the class $[\square^n]$ is

$$\langle [\square^n], [\omega_0^{(n)}] \rangle = \int_{\square^n} \omega_0^{(n)} = 1.$$

We call the differential forms $\omega_k^{(n)}$, for $k = 0, 2, \dots, n$, the *Eulerian differential forms*.

3.5 An inductive system

For $n \geq 2$ there are natural morphisms

$$i^{(n)} : \mathcal{Z}^{(n-1)} \rightarrow \mathcal{Z}^{(n)} \tag{23}$$

in the category $\text{MT}(\mathbb{Q})$, that we now define. We fix the identification $X_{n-1} = \{x_n = 1\} \subset X_n$, which implies the equality $A_{n-1} = A_n \cap X_{n-1}$. Let us set

$$B'_n = \bigcup_{1 \leq i \leq n} \{x_i = 0\} \cup \bigcup_{1 \leq i \leq n-1} \{x_i = 1\},$$

so that we have $B_n = B'_n \cup X_{n-1}$, and $B_{n-1} = B'_n \cap X_{n-1}$.

In the blow-up \tilde{X}_n , we thus get an embedding $\tilde{X}_{n-1} \subset \tilde{X}_n$ and identifications $\tilde{A}_{n-1} = \tilde{A}_n \cap \tilde{X}_{n-1}$, $\tilde{B}_{n-1} = \tilde{B}'_n \cap \tilde{X}_{n-1}$ and $E_{n-1} = E_n \cap \tilde{X}_{n-1}$. Thus, the complex in $\text{DM}(\mathbb{Q})$ that we have used to define $\mathcal{Z}^{(n-1)}$ is the subcomplex

$$\cdots \rightarrow \bigsqcup_{\substack{|I|=3 \\ \partial_I Y \subset \tilde{X}_{n-1}}} \partial_I Y \rightarrow \bigsqcup_{\substack{|I|=2 \\ \partial_I Y \subset \tilde{X}_{n-1}}} \partial_I Y \rightarrow \tilde{X}_{n-1} \rightarrow 0 \rightarrow 0 \tag{24}$$

of the complex (15) that we have used to define $\mathcal{Z}^{(n)}$, shifted by 1. Taking the n th cohomology groups with respect to the t -structure gives the morphism (23).

In the Betti and the de Rham realizations, the morphism (23) is also induced by the inclusion of double subcomplexes of (16) and (17).

We define the ind-motive

$$\mathcal{Z} = \varinjlim_n \mathcal{Z}^{(n)},$$

viewed as an ind-object in the category $\text{MT}(\mathbb{Q})$, and simply call it the *zeta motive*.

The map $i_{\mathbb{B}}^{(n),\vee} : \mathcal{Z}_{\mathbb{B}}^{(n),\vee} \rightarrow \mathcal{Z}_{\mathbb{B}}^{(n-1),\vee}$ given by the transpose of the Betti realization of $i^{(n)}$ satisfies

$$i_{\mathbb{B}}^{(n),\vee}([\square^{n-1}]) = [\square^n]. \tag{25}$$

More generally and loosely speaking, if σ is a chain on $\tilde{X}_n(\mathbb{C}) - \tilde{A}_n(\mathbb{C})$ whose boundary is on $\tilde{B}_n(\mathbb{C}) \cup E_n(\mathbb{C})$, then $i_{\mathbb{B}}^{(n),\vee}([\sigma])$ is the class of ‘the component of the boundary of σ that lives on $\tilde{X}_{n-1}(\mathbb{C})$ ’. According to Proposition 3.4, one can also work with chains on $X_n(\mathbb{C}) - \mathring{A}_n(\mathbb{C})$. We note that (25) allows us to define a class

$$[\square] \in \mathcal{Z}_{\mathbb{B}}^{\vee} := \varinjlim_n \mathcal{Z}_{\mathbb{B}}^{(n),\vee}.$$

Remark 3.9. There are (alternating) signs in the differentials of the complexes (15)–(17), that we leave to the reader. This also induces signs on the different components of the inclusions of subcomplexes such as (24); these signs are fixed once and for all by (25).

The next proposition shows that the Eulerian differential forms $\omega_k^{(n)}$ are compatible with the inductive structure on the zeta motives.

PROPOSITION 3.10. *For integers $n \geq 2$ and $k = 0, 2, \dots, n - 1$, the map $i_{\text{dR}}^{(n)} : \mathcal{Z}_{\text{dR}}^{(n-1)} \rightarrow \mathcal{Z}_{\text{dR}}^{(n)}$ sends the class $[\omega_k^{(n-1)}]$ to the class $[\omega_k^{(n)}]$.*

Proof. Since all the differential forms that we are manipulating have no poles along the exceptional divisors E_{n-1} and E_n , it is safe to do the computations in the affine spaces X_{n-1} and X_n ; we leave it to the reader to turn them into computations in \tilde{X}_{n-1} and \tilde{X}_n by working in local charts as in the proof of Proposition 3.6. Let us first assume that $k \in \{2, \dots, n - 1\}$. We put

$$\eta_k^{(n-1)} = \frac{x_n E_{n-1-k}(x_1 \cdots x_n)}{(1 - x_1 \cdots x_n)^{n-k}} dx_1 \cdots dx_{n-1},$$

viewed as a form on X_n . Then we have $(\eta_k^{(n-1)})|_{X_{n-1}} = \omega_k^{(n-1)}$ and $(\eta_k^{(n-1)})|_{B'_{n-1}} = 0$. A diagram chase in the double complex (17) shows that $i_{\text{dR}}^{(n)}([\omega_k^{(n-1)}])$ is the class of

$$(-1)^{n-1} (d(\eta_k^{(n-1)}))$$

(the sign is here to be consistent with the Betti version, see Remark 3.9). We have

$$(-1)^{n-1} d(\eta_k^{(n-1)}) = \frac{\partial}{\partial x_n} \left(\frac{x_n E_{n-1-k}(x_1 \cdots x_n)}{(1 - x_1 \cdots x_n)^{n-k}} \right) dx_1 \cdots dx_n$$

and one easily sees that setting $x = x_1 \cdots x_n$ we have

$$\frac{\partial}{\partial x_n} \left(\frac{x_n E_{n-1-k}(x_1 \cdots x_n)}{(1 - x_1 \cdots x_n)^{n-k}} \right) = \frac{x(1-x)E'_{n-1-k}(x) + (1+(n-1-k)x)E_{n-1-k}(x)}{(1-x)^{n-k+1}}.$$

Using the recurrence relation (21), one then concludes that

$$(-1)^{n-1} d(\eta_k^{(n-1)}) = \frac{E_{n-k}(x_1 \cdots x_n)}{(1 - x_1 \cdots x_n)^{n-k+1}} dx_1 \cdots dx_n = \omega_k^{(n)}.$$

For $k = 0$, this is the same computation with $\eta_0^{(n)} = x_n dx_1 \cdots dx_{n-1}$ and

$$(-1)^{n-1} d(\eta_0^{(n-1)}) = dx_1 \cdots dx_n = \omega_0^{(n)}. \quad \square$$

Proposition 3.10 allows us to unambiguously define classes

$$[\omega_k] \in \mathcal{Z}_{\text{dR}}$$

for $k = 0, 2, 3, \dots$, whose pairing with the class $[\square] \in \mathcal{Z}_{\text{B}}^\vee$ is

$$\langle [\square], [\omega_0] \rangle = 1 \quad \text{and} \quad \langle [\square], [\omega_k] \rangle = \zeta(k) \quad (k \geq 2).$$

Remark 3.11. The proof of Proposition 3.10 can be thought of as a cohomological version of the relation

$$\int_{\square^n} \omega_k^{(n)} = \int_{\square^{n-1}} \omega_k^{(n-1)},$$

which may be proved using Stokes’s theorem and the recurrence relation (21).

PROPOSITION 3.12. *For integers $n \geq 1$ and $k = 0, 2, \dots, n$, the class $[\omega_k^{(n)}]$ lives in the pure weight $2k$ component of $\mathcal{Z}_{\text{dR}}^{(n)}$.*

Proof. For $k = 0$, Proposition 3.10 and the fact that the maps $i_{\text{dR}}^{(n)}$ are compatible with the weight gradings implies that it is enough to do the proof for $n = 1$; this case is easy since $\mathcal{Z}^{(1)} \cong \mathbb{Q}(0)$ only has weight 0. We now turn to the case $k = 2, \dots, n$. Thanks to Proposition 3.10 and the fact that the maps $i_{\text{dR}}^{(n)}$ are compatible with the weight gradings, it is enough to check it for $k = n$.

By (7), we need to prove that the class of $\omega_n^{(n)}$ is in $F^n \mathcal{Z}_{\text{dR}}^{(n)}$. Let Y be a smooth projective variety of dimension n , D be a normal crossing divisor inside Y , and Z be a closed subvariety of Y of dimension $\leq n - 1$. Then we have

$$F^n H_{\text{dR}}^n(Y - D, Z - Z \cap D) = \text{Im}(H^0(\Omega_Y^n(\log D)) \longrightarrow H_{\text{dR}}^n(Y - D, Z - Z \cap D)).$$

Thus, it is enough to prove that there is a compactification Y of $\tilde{X}_n - \tilde{A}_n$ such that $Y - (\tilde{X}_n - \tilde{A}_n)$ is a normal crossing divisor D , and such that $\omega_n^{(n)}$ has logarithmic singularities along D . Since $\omega_n^{(n)}$ does not have poles along E_n , we can work on $X_n - A_n$ instead. Let us start with $Y_1 = (\mathbb{P}^1)^n$ with coordinates $((x_1 : y_1), \dots, (x_n : y_n))$, and D_1 the divisor given by the union of the subvarieties $\{y_i = 0\}$ for $i = 1, \dots, n$, and the subvariety $\{x_1 \cdots x_n = y_1 \cdots y_n\}$ (this is the closure of A_n). This is not enough since D_1 is not a normal crossing divisor. We then finish thanks to the following lemma. □

LEMMA 3.13. Let $\phi : Y \rightarrow Y_1$ be the iterated blow-up of all the codimension-2 subvarieties $Z_{i,j} = \{y_i = x_j = 0\}$, $i \neq j$, in any order. Then $D = \phi^{-1}(D_1)$ is a normal crossing divisor inside Y and $\phi^*(\omega_n^{(n)})$ has logarithmic singularities along D .

Proof. This is checked locally on the standard affine cover of Y , consisting of 2^n affine spaces. By symmetry of the variables, it is enough to look at the charts U_r with affine coordinates

$$(y_1, \dots, y_r, x_{r+1}, \dots, x_n),$$

for $r = 0, \dots, n$. We note that in the chart U_r the divisor D_1 is the union of the subvarieties $\{y_i = 0\}$ for $i = 1, \dots, r$, and the subvariety $\{y_1 \cdots y_r = x_{r+1} \cdots x_n\}$. In that chart the differential form that we are looking at is (up to a sign)

$$\omega = \frac{dy_1 \cdots dy_r dx_{r+1} \cdots dx_n}{y_1 \cdots y_r (y_1 \cdots y_r - x_{r+1} \cdots x_n)}.$$

We proceed by induction on r . For $r = 0$, D_1 only consists of $\{x_1 \cdots x_n = 1\}$, which is a normal crossing divisor, and ω has logarithmic singularities along D_1 . The subvarieties $Z_{i,j}$ do not intersect U_0 , so the blow-ups do not change anything. For a given $r = 1, \dots, n$, let us look at the blow-up of $Z_{1,n} = \{y_1 = x_n = 0\}$ (this is enough for reasons of symmetry) in the chart U_r . There are two natural affine charts $\mathbb{A}^n \rightarrow U_r$ for the blow-up.

(i) On the first chart, the blow-up map is given by

$$(y_1, \dots, y_r, x_{r+1}, \dots, x_n) = (v_1, \dots, v_r, u_{r+1}, \dots, u_{n-1}, v_1 u_n).$$

The preimage of D_1 consists of the subvarieties $\{v_i = 0\}$ for $i = 1, \dots, r$, and the subvariety $\{v_2 \cdots v_r = u_{r+1} \cdots u_n\}$. The pullback of ω is

$$\tilde{\omega} = \frac{dv_1 \cdots dv_r du_{r+1} \cdots du_n}{v_1 \cdots v_r (v_2 \cdots v_r - u_{r+1} \cdots u_n)} = \frac{dv_1}{v_1} \wedge \omega' \quad \text{with } \omega' = \frac{dv_2 \cdots dv_r du_{r+1} \cdots du_n}{v_2 \cdots v_r (v_2 \cdots v_r - u_{r+1} \cdots u_n)}.$$

We note that $\{v_1 = 0\}$ is the exceptional divisor and that the total transforms of the subvarieties $Z_{1,j}$ are empty in this chart. By the induction hypothesis (with n replaced by $n - 1$), the pullback of ω' by the successive blow-up of the subvarieties $Z_{i,j}$ with $i \neq 1$ has logarithmic singularities. Since dv_1/v_1 has logarithmic singularities along $\{v_1 = 0\}$, we are done.

(ii) On the second chart, the blow-up map is given by

$$(y_1, \dots, y_r, x_{r+1}, \dots, x_n) = (v_1 u_n, v_2, \dots, v_r, u_{r+1}, \dots, u_n).$$

The same argument as in the first chart applies. □

3.6 A long exact sequence

We now show that the morphism $i^{(n)} : \mathcal{Z}^{(n-1)} \rightarrow \mathcal{Z}^{(n)}$ fits into a long exact sequence. We first define objects of $\text{MT}(\mathbb{Q})$:

$$\mathcal{Z}^{(n),r} = H^r(\tilde{X}_n - \tilde{A}_n, (\tilde{B}_n \cup E_n) - (\tilde{B}_n \cup E_n) \cap \tilde{A}_n)$$

and

$$'\mathcal{Z}^{(n),r} = H^r(\tilde{X}_n - \tilde{A}_n, (\tilde{B}'_n \cup E_n) - (\tilde{B}'_n \cup E_n) \cap \tilde{A}_n),$$

so that $\mathcal{Z}^{(n)} = \mathcal{Z}^{(n),n}$. We leave it to the reader to fill in the technical definitions of these objects by mimicking that of $\mathcal{Z}^{(n)}$ from §3.1.

PROPOSITION 3.14. For $n \geq 2$, we have a long exact sequence in $\text{MT}(\mathbb{Q})$:

$$\dots \rightarrow \mathcal{Z}^{(n-1),r-1} \rightarrow \mathcal{Z}^{(n),r} \rightarrow {}'\mathcal{Z}^{(n),r} \rightarrow \mathcal{Z}^{(n-1),r} \rightarrow \mathcal{Z}^{(n),r+1} \rightarrow \dots \tag{26}$$

Proof. The objects $\mathcal{Z}^{(n-1),\bullet}$, $\mathcal{Z}^{(n),\bullet}$ and ${}'\mathcal{Z}^{(n),\bullet}$ are defined via objects in $\text{DMT}(\mathbb{Q})$ that we denote by $C^{(n-1)}$, $C^{(n)}$ and ${}'C^{(n)}$ respectively, $C^{(n)}$ being the complex (15) and $C^{(n-1)}$ the subcomplex (24). Now there is an obvious exact triangle

$$C^{(n-1)}[-1] \longrightarrow C^{(n)} \longrightarrow {}'C^{(n)} \xrightarrow{+1},$$

in $\text{DMT}(\mathbb{Q})$, which gives the desired long exact sequence after taking the cohomology with respect to the t -structure. \square

We note that the map $\mathcal{Z}^{(n-1),n-1} \rightarrow \mathcal{Z}^{(n),n}$ in the long exact sequence (26) is exactly $i^{(n)}$.

4. Computation of the zeta motives $\mathcal{Z}^{(n)}$

This section is the technical heart of this article, where we compute (Theorem 4.9) the full period matrix of the zeta motives $\mathcal{Z}^{(n)}$. The main difficulty is showing that the motives $\mathcal{T}^{(n)}$, introduced below, are semi-simple. For that we use the involution τ defined in the introduction and the computation of the extension groups in the category $\text{MT}(\mathbb{Q})$. We then define the odd zeta motive and compute its period matrix. We finish with an elementary (Hodge-theoretic) proof that the motives $\mathcal{T}^{(n)}$ are semi-simple.

4.1 The Gysin long exact sequence

Since the divisor A_n is smooth, it is natural to decompose the motives $\mathcal{Z}^{(n),r}$ thanks to a Gysin long exact sequence. In the next proposition, the definition of the objects $H^\bullet(X_n, B_n)$ and $H^\bullet(A_n, B_n \cap A_n)$ of $\text{MT}(\mathbb{Q})$ is similar to that of $\mathcal{Z}^{(n)}$ from § 3.1.

PROPOSITION 4.1. For $n \geq 1$, we have a long exact sequence in $\text{MT}(\mathbb{Q})$:

$$\dots \rightarrow H^r(X_n, B_n) \rightarrow \mathcal{Z}^{(n),r} \rightarrow H^{r-1}(A_n, B_n \cap A_n)(-1) \rightarrow H^{r+1}(X_n, B_n) \rightarrow \mathcal{Z}^{(n),r+1} \rightarrow \dots \tag{27}$$

Proof. Recall from [Voe00, (3.5.4)] the existence of a Gysin exact triangle in the category $\text{DM}(\mathbb{Q})$. For the pair $(\tilde{X}_n, \tilde{A}_n)$, it reads (with cohomological conventions)

$$\tilde{X}_n \longrightarrow \tilde{X}_n - \tilde{A}_n \longrightarrow \tilde{A}_n(-1)[-1] \xrightarrow{+1}$$

and is an exact triangle in the category $\text{DMT}(\mathbb{Q})$. Applying this triangle to every pair $(\partial_I Y, \partial_I Y \cap \tilde{A}_n)$ in the complex (15) and taking the cohomology with respect to the t -structure leads to a long exact sequence

$$\dots \rightarrow H^r(\tilde{X}_n, \tilde{B}_n \cup E_n) \rightarrow H^r(\tilde{X}_n - \tilde{A}_n, (\tilde{B}_n \cup E_n) - \tilde{A}_n) \rightarrow H^{r-1}(\tilde{A}_n, (\tilde{B}_n \cup E_n) \cap \tilde{A}_n)(-1) \rightarrow \dots$$

in $\text{MT}(\mathbb{Q})$. One finishes with the fact that the natural morphisms

$$H^r(\tilde{X}_n, \tilde{B}_n \cup E_n) \rightarrow H^r(X_n, B_n) \quad \text{and} \quad H^{r-1}(\tilde{A}_n, (\tilde{B}_n \cup E_n) \cap \tilde{A}_n) \rightarrow H^{r-1}(A_n, B_n \cap A_n)$$

are isomorphisms. This can be checked in the Betti realization (see Remark 2.5), where it is a consequence of the excision theorem as in the proof of Proposition 3.4. \square

4.2 The motives $H^\bullet(X_n, B_n)$

The computation of the motives $H^\bullet(X_n, B_n)$ appearing in the long exact sequence (27) is relatively easy.

PROPOSITION 4.2. (i) We have $H^r(X_n, B_n) = 0$ for $r \neq n$, and an isomorphism $H^n(X_n, B_n) \cong \mathbb{Q}(0)$.

- (ii) A basis for the de Rham realization $H_{\text{dR}}^n(X_n, B_n)$ is the class of the form $dx_1 \cdots dx_n$.
- (iii) A basis for the Betti realization $H_n^{\text{B}}(X_n, B_n)$ is the class of the unit n -cube $\square^n = [0, 1]^n$.

Proof. By the relative Künneth formula we have $H^\bullet(X_n, B_n) \cong H^\bullet(X_1, B_1)^{\otimes n}$ so that it is enough to prove the proposition for $n = 1$. This has already been done in Remark 3.2. □

4.3 The motives $H^\bullet(A_n, B_n \cap A_n)$

For $n \geq 1$, we realize the n -torus as $T^n = \{x_1 \cdots x_{n+1} = 1\}$, and we have subtori $T_i^{n-1} = \{x_i = 1\} \subset T^n$ for $i = 1, \dots, n + 1$. We define

$$\mathcal{T}^{(n),r} = H^r\left(T^n, \bigcup_{1 \leq i \leq n+1} T_i^{n-1}\right) \quad \text{and} \quad {}'\mathcal{T}^{(n),r} = H^r\left(T^n, \bigcup_{1 \leq i \leq n} T_i^{n-1}\right),$$

which are objects in $\text{MT}(\mathbb{Q})$ (whose definition is similar to that of $\mathcal{Z}^{(n)}$ from §3.1) and write

$$\mathcal{T}^{(n)} = \mathcal{T}^{(n),n} \quad \text{and} \quad {}'\mathcal{T}^{(n)} = {}'\mathcal{T}^{(n),n}.$$

We then have

$$H^{r-1}(A_n, B_n \cap A_n) \cong \mathcal{T}^{(n-1),r-1}.$$

By mimicking the proof of Proposition 3.14, one produces a long exact sequence in $\text{MT}(\mathbb{Q})$:

$$\dots \rightarrow \mathcal{T}^{(n-1),r-1} \rightarrow \mathcal{T}^{(n),r} \rightarrow {}'\mathcal{T}^{(n),r} \rightarrow \mathcal{T}^{(n-1),r} \rightarrow \mathcal{T}^{(n),r+1} \rightarrow \dots \tag{28}$$

PROPOSITION 4.3. (i) We have ${}'\mathcal{T}^{(n),r} = 0$ for $r \neq n$, and an isomorphism ${}'\mathcal{T}^{(n)} \cong H^n(T^n) \cong \mathbb{Q}(-n)$.

- (ii) We have $\mathcal{T}^{(n),r} = 0$ for $r \neq n$, and short exact sequences in $\text{MT}(\mathbb{Q})$:

$$0 \rightarrow \mathcal{T}^{(n-1)} \xrightarrow{j^{(n)}} \mathcal{T}^{(n)} \rightarrow H^n(T^n) \rightarrow 0. \tag{29}$$

Proof. If (i) is proved, then (ii) follows from the long exact sequence (28). By choosing coordinates (x_1, \dots, x_n) on T^n we see that we have

$${}'\mathcal{T}^{(n),\bullet} = H^\bullet\left(\left(\mathbb{A}_{\mathbb{Q}}^1 - \{0\}\right)^n, \bigcup_{1 \leq i \leq n} \{x_i = 1\}\right) \cong H^\bullet(\mathbb{A}_{\mathbb{Q}}^1 - \{0\}, \{1\})^{\otimes n} = ({}'\mathcal{T}^{(1),\bullet})^{\otimes n},$$

where we have used the relative Künneth formula. Thus, it is enough to prove (i) for $n = 1$, which is easy since ${}'\mathcal{T}^{(1),\bullet}$ is nothing but the reduced cohomology of $\mathbb{A}_{\mathbb{Q}}^1 - \{0\}$. □

Remark 4.4. We note that the morphism $j^{(n)} : \mathcal{T}^{(n-1)} \rightarrow \mathcal{T}^{(n)}$ in (29) is defined in the same way as the morphism $i^{(n)} : \mathcal{Z}^{(n-1)} \rightarrow \mathcal{Z}^{(n)}$ from §3.5.

We note that we have $\mathcal{T}^{(0)} = H^0(\text{pt}, \text{pt}) = 0$, so that Proposition 4.3 implies that we have

$$\text{gr}_{2k}^W \mathcal{T}^{(n)} = \begin{cases} \mathbb{Q}(-k) & \text{if } k \in \{1, \dots, n\}, \\ 0 & \text{otherwise.} \end{cases}$$

In the next proposition, we will prove that the weight filtration of $\mathcal{T}^{(n)}$ actually splits in $\text{MT}(\mathbb{Q})$. For that we introduce the involution τ which acts on the tori T^n by

$$\tau : (x_1, \dots, x_{n+1}) \mapsto (x_1^{-1}, \dots, x_{n+1}^{-1}).$$

This induces an involution, still denoted by τ , on the objects $\mathcal{T}^{(n),r}$ and ${}'\mathcal{T}^{(n),r}$ of $\text{MT}(\mathbb{Q})$, such that all the maps in the long exact sequence (28) commute with τ .

PROPOSITION 4.5. (i) *The short exact sequences (29) split in $\text{MT}(\mathbb{Q})$, hence we have isomorphisms:*

$$\mathcal{T}^{(n)} \cong \mathbb{Q}(-1) \oplus \mathbb{Q}(-2) \oplus \dots \oplus \mathbb{Q}(-n).$$

Thus, a period matrix for $\mathcal{T}^{(n)}$ is the diagonal matrix $\text{Diag}(2\pi i, (2\pi i)^2, \dots, (2\pi i)^n)$.

(ii) *The involution τ acts on the direct summand $\mathbb{Q}(-k)$ of $\mathcal{T}^{(n)}$ by multiplication by $(-1)^k$.*

Proof. We first note that τ acts on $H^1(T^1)$ by multiplication by -1 . It is enough to prove it in the de Rham realization, where it follows from $\tau.\text{dlog}(x_1) = -\text{dlog}(x_1)$. Thus, τ acts on $\text{gr}_{2n}^W \mathcal{T}^{(n)} \cong H^n(T^n) \cong H^1(T^1)^{\otimes n}$ by multiplication by $(-1)^n$, and we are left with proving (i). We denote by $\mathcal{T}^{(n)} = \mathcal{T}_+^{(n)} \oplus \mathcal{T}_-^{(n)}$ the direct sum decomposition of $\mathcal{T}^{(n)}$ into its invariant and anti-invariant parts with respect to τ . We have to prove that we have isomorphisms

$$\mathcal{T}_+^{(2n)} \cong \mathcal{T}_+^{(2n+1)} \cong \mathbb{Q}(-2) \oplus \mathbb{Q}(-4) \oplus \dots \oplus \mathbb{Q}(-2n)$$

and

$$\mathcal{T}_-^{(2n+1)} \cong \mathcal{T}_-^{(2n+2)} \cong \mathbb{Q}(-1) \oplus \mathbb{Q}(-3) \oplus \dots \oplus \mathbb{Q}(-(2n+1)).$$

We only prove the statements corresponding to the invariant parts, the statements corresponding to the anti-invariant parts being proved similarly. We use induction on n , the case $n = 0$ being trivial: $\mathcal{T}_+^{(0)} = \mathcal{T}_+^{(1)} = 0$. The short exact sequences (29) imply that we have short exact sequences

$$0 \rightarrow \mathcal{T}_+^{(2n+1)} \rightarrow \mathcal{T}_+^{(2n+2)} \rightarrow \mathbb{Q}(-(2n+2)) \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathcal{T}_+^{(2n+2)} \rightarrow \mathcal{T}_+^{(2n+3)} \rightarrow 0 \rightarrow 0.$$

Using the induction hypothesis we see that we have

$$\begin{aligned} \text{Ext}_{\text{MT}(\mathbb{Q})}^1(\mathbb{Q}(-(2n+2)), \mathcal{T}_+^{(2n+1)}) &\cong \text{Ext}_{\text{MT}(\mathbb{Q})}^1(\mathbb{Q}(-(2n+2)), \mathbb{Q}(-2) \oplus \mathbb{Q}(-4) \oplus \dots \oplus \mathbb{Q}(-2n)) \\ &\cong \bigoplus_{1 \leq k \leq n} \text{Ext}_{\text{MT}(\mathbb{Q})}^1(\mathbb{Q}(-2k), \mathbb{Q}(0)) \\ &= 0, \end{aligned}$$

where we have used (13). Thus, the first short exact sequence splits. The second short exact sequence then completes the induction. □

Remark 4.6. From the short exact sequences (29) it is clear that, for every n , $\mathcal{T}_{\text{dR}}^{(n)}$ has a basis $(w_1^{(n)}, \dots, w_n^{(n)})$ which is compatible with the weight grading, such that $w_n^{(n)}$ is the class of the form $\text{dlog}(x_1) \wedge \dots \wedge \text{dlog}(x_n)$, and such that these bases are compatible with the short exact sequences (29).

4.4 The structure of the zeta motives

We can now determine the structure of the zeta motives $\mathcal{Z}^{(n)}$, for $n \geq 1$.

THEOREM 4.7. (i) We have a short exact sequence in $\text{MT}(\mathbb{Q})$:

$$0 \rightarrow \mathbb{Q}(0) \rightarrow \mathcal{Z}^{(n)} \xrightarrow{p^{(n)}} \mathcal{T}^{(n-1)}(-1) \rightarrow 0, \tag{30}$$

with $\mathcal{T}^{(n-1)}(-1) \cong \mathbb{Q}(-2) \oplus \dots \oplus \mathbb{Q}(-n)$.

(ii) We have a short exact sequence in $\text{MT}(\mathbb{Q})$:

$$0 \rightarrow \mathcal{Z}^{(n-1)} \xrightarrow{i^{(n)}} \mathcal{Z}^{(n)} \rightarrow \mathbb{Q}(-n) \rightarrow 0. \tag{31}$$

(iii) These short exact sequences fit into a commutative diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{Q}(0) & \longrightarrow & \mathcal{Z}^{(n-1)} & \xrightarrow{p^{(n-1)}} & \mathcal{T}^{(n-2)}(-1) \longrightarrow 0 \\
 & & \downarrow = & & \downarrow i^{(n)} & & \downarrow j^{(n-1)} \\
 0 & \longrightarrow & \mathbb{Q}(0) & \longrightarrow & \mathcal{Z}^{(n)} & \xrightarrow{p^{(n)}} & \mathcal{T}^{(n-1)}(-1) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & 0 & \longrightarrow & \mathbb{Q}(-n) & \xrightarrow{=} & \mathbb{Q}(-n) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array} \tag{32}$$

where all rows and columns are exact.

Proof. Assertion (1) follows from Propositions 4.1, 4.2 and 4.5. The commutativity of (32) follows from the compatibility of the long exact sequences (26) and (28). A diagram chase implies that (31) is exact. \square

Remark 4.8. The difference between the sign $(-1)^k$ in Proposition 4.5(2) and the sign $(-1)^{k-1}$ in Theorem 1.4 comes from the Tate twist (-1) in the short exact sequence (30).

THEOREM 4.9. (i) The classes

$$v_k^{(n)} := [\omega_k^{(n)}] \quad (k = 0, 2, \dots, n)$$

of the Eulerian differential forms provide a basis $(v_0^{(n)}, v_2^{(n)}, \dots, v_n^{(n)})$ of the de Rham realization $\mathcal{Z}_{\text{dR}}^{(n)}$ which is compatible with the weight grading.

(ii) There exists a unique basis $(\varphi_0^{(n)}, \varphi_2^{(n)}, \dots, \varphi_n^{(n)})$ for the dual of the Betti realization $\mathcal{Z}_B^{(n), \vee}$ which is compatible with the weight filtration and such that the period matrix for $\mathcal{Z}^{(n)}$ in the v -basis and the φ -basis is

$$\begin{pmatrix} 1 & \zeta(2) & \zeta(3) & \cdots & \cdots & \zeta(n-1) & \zeta(n) \\ & (2\pi i)^2 & & & & & \\ & & (2\pi i)^3 & & & 0 & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & 0 & & & & (2\pi i)^{n-1} & \\ & & & & & & (2\pi i)^n \end{pmatrix}. \tag{33}$$

Proof. (i) Proposition 3.12 says that $v_k^{(n)}$ is in the pure weight $2k$ component of $\mathcal{Z}_{\text{dR}}^{(n)}$. Thus, it is enough to show that it is non-zero, which is a consequence of the equalities $\langle [\square^n], v_0^{(n)} \rangle = 1 \neq 0$ and $\langle [\square^n], v_k^{(n)} \rangle = \zeta(k) \neq 0$ for $k = 2, \dots, n$.

(ii) We put $\varphi_0^{(n)} = [\square^n]$. Let $(\psi_1^{(n-1)}, \dots, \psi_{n-1}^{(n-1)})$ be a basis of $\mathcal{T}_B^{(n-1), \vee}$ for which the period matrix is diagonal, as in Proposition 4.5. Let $p^{(n)}$ denote the morphism $\mathcal{Z}^{(n)} \rightarrow \mathcal{T}^{(n-1)}(-1)$, and let us consider the transpose of its Betti realization $p_B^{(n), \vee} : \mathcal{T}_B^{(n-1), \vee} \rightarrow \mathcal{Z}_B^{(n), \vee}$. Then we can put $\varphi_k^{(n)} = p_B^{(n), \vee}(\psi_{k-1}^{(n-1)})$ for $k = 2, \dots, n$. The fact that this gives a basis of $\mathcal{Z}_B^{(n), \vee}$ is a consequence of the short exact sequence (30). The fact that the period matrix is as required follows from Lemma 3.8 and Proposition 4.5. The uniqueness statement is obvious. \square

We have already noted that the classes $v_k^{(n)}$ are compatible with the inductive system of the zeta motives. By the uniqueness statement in Theorem 4.9, this is also the case for the classes $\varphi_k^{(n)}$, and the zeta motive \mathcal{Z} has an infinite period matrix

$$\begin{pmatrix} 1 & \zeta(2) & \zeta(3) & \zeta(4) & \cdots & \cdots & \cdots \\ & (2\pi i)^2 & & & & & \\ & & (2\pi i)^3 & & & 0 & \\ & & & (2\pi i)^4 & & & \\ & & & & \ddots & & \\ & 0 & & & & \ddots & \\ & & & & & & \ddots \end{pmatrix}.$$

4.5 The odd zeta motive

Let us write $\mathcal{T}^{(n-1)} = \mathcal{T}_+^{(n-1)} \oplus \mathcal{T}_-^{(n-1)}$ for the direct sum decomposition into its invariant and anti-invariant parts with respect to τ , and let us write $p^{(n)} : \mathcal{Z}^{(n)} \rightarrow \mathcal{T}^{(n-1)}(-1)$ for the surjection appearing in the short exact sequence (30).

DEFINITION 4.10. The n th odd zeta motive $\mathcal{Z}^{(n),\text{odd}}$ is the object of $\text{MT}(\mathbb{Q})$ defined by

$$\mathcal{Z}^{(n),\text{odd}} := (p^{(n)})^{-1}(\mathcal{T}_+^{(n-1)}(-1)).$$

We obviously have a short exact sequence

$$0 \rightarrow \mathbb{Q}(0) \rightarrow \mathcal{Z}^{(n),\text{odd}} \rightarrow \mathcal{T}_+^{(n-1)}(-1) \rightarrow 0 \tag{34}$$

with

$$\mathcal{T}_+^{(n-1)}(-1) \cong \bigoplus_{3 \leq 2k+1 \leq n} \mathbb{Q}(-(2k+1)).$$

We note that there are morphisms

$$i^{(n),\text{odd}} : \mathcal{Z}^{(n-1),\text{odd}} \rightarrow \mathcal{Z}^{(n),\text{odd}}$$

such that $i^{(2n),\text{odd}}$ is an isomorphism for every integer n . The limit

$$\mathcal{Z}^{\text{odd}} := \varinjlim_n \mathcal{Z}^{(n),\text{odd}}$$

is an ind-object in $\text{MT}(\mathbb{Q})$ that we simply call the *odd zeta motive*.

PROPOSITION 4.11. (i) We have a direct sum decomposition

$$\mathcal{Z}^{(n)} \cong \mathcal{Z}^{(n),\text{odd}} \oplus \bigoplus_{2 \leq 2k \leq n} \mathbb{Q}(-2k). \tag{35}$$

(ii) A period matrix for $\mathcal{Z}^{(2n+1),\text{odd}} \cong \mathcal{Z}^{(2n+2),\text{odd}}$ is

$$\begin{pmatrix} 1 & \zeta(3) & \zeta(5) & \cdots & \cdots & \zeta(2n-1) & \zeta(2n+1) \\ & (2\pi i)^3 & & & & & \\ & & (2\pi i)^5 & & & 0 & \\ & & & \ddots & & & \\ & & & & \ddots & & \\ & 0 & & & & (2\pi i)^{2n-1} & \\ & & & & & & (2\pi i)^{2n+1} \end{pmatrix}. \tag{36}$$

Proposition 4.11 implies that the odd zeta motive \mathcal{Z}^{odd} has an infinite period matrix (6). In particular, $\mathcal{Z}^{(3),\text{odd}}$ is the essentially unique non-trivial extension of $\mathbb{Q}(-3)$ by $\mathbb{Q}(0)$ in $\text{MT}(\mathbb{Q})$.

Proof. This is a consequence of the short exact sequence (30) and the vanishing of the extension groups $\text{Ext}_{\text{MT}(\mathbb{Q})}^1(\mathbb{Q}(-2k), \mathbb{Q}(0))$, see (13). An alternative proof which does not use the computation of extension groups goes as follows. A basis for $\mathcal{Z}_{\text{dR}}^{(n),\text{odd}}$ is given by $v_0^{(n)}$ and the $v_{2k+1}^{(n)}$, for $3 \leq 2k+1 \leq n$, and a basis for $\mathcal{Z}_{\text{B}}^{(n),\text{odd},\vee}$ is given by $\varphi_0^{(n)}$ and the $\varphi_{2k+1}^{(n)}$, for $3 \leq 2k+1 \leq n$. This gives the desired shape for the period matrix (36). Now, Euler’s solution to the Basel problem implies that we have $\zeta(2k) = \lambda_{2k}(2\pi i)^{2k}$ for every integer $k \geq 1$,

with $\lambda_{2k} = -B_{2k}/2(2k)! \in \mathbb{Q}$. Thus, we may replace the basis $(\varphi_0^{(n)}, \varphi_2^{(n)}, \dots, \varphi_n^{(n)})$ of Theorem 4.9 by the basis $(\prime\varphi_0^{(n)}, \varphi_2^{(n)}, \dots, \varphi_n^{(n)})$ with

$$\prime\varphi_0^{(n)} = \varphi_0^{(n)} - \sum_{2 \leq 2k \leq n} \lambda_{2k} \varphi_{2k}^{(n)}$$

to get a period matrix similar to (33) where the even zeta values $\zeta(2k)$ in the first row are replaced by 0. This implies the direct sum decomposition (35). □

We finish by proving that all the objects in $\text{MT}(\mathbb{Q})$ considered earlier actually live in the full subcategory $\text{MT}(\mathbb{Z})$.

PROPOSITION 4.12. *The zeta motives $\mathcal{Z}^{(n)}$ and the odd zeta motives $\mathcal{Z}^{(n),\text{odd}}$ are objects of the category $\text{MT}(\mathbb{Z})$.*

Proof. Thanks to the direct sum decomposition (35), it is enough to prove it for the odd zeta motives. Let us recall the definition [DG05, Définition 1.4] of the category $\text{MT}(\mathbb{Z})$. According to the tannakian formalism, the de Rham realization functor $\text{MT}(\mathbb{Q}) \rightarrow \text{grVect}_{\mathbb{Q}}$ induces an equivalence of categories

$$\text{MT}(\mathbb{Q}) \cong \text{grRep}(\mathfrak{g}_{\text{dR}}^{\mathbb{Q}})$$

between $\text{MT}(\mathbb{Q})$ and the category of graded finite-dimensional representations of a graded Lie algebra $\mathfrak{g}_{\text{dR}}^{\mathbb{Q}}$. The degree in $\mathfrak{g}_{\text{dR}}^{\mathbb{Q}}$ is half the weight. This Lie algebra is non-positively graded. The category $\text{MT}(\mathbb{Z})$ is defined as the full subcategory of $\text{MT}(\mathbb{Q})$ consisting of objects H such that the degree -1 component of $\mathfrak{g}_{\text{dR}}^{\mathbb{Q}}$ acts trivially on H_{dR} . This is obviously the case for $\mathcal{Z}^{(n),\text{odd}}$, which is concentrated in weights 0 and $2(2k + 1)$ with $2k + 1 \geq 3$ by the short exact sequence (34). □

Remark 4.13. A tannakian interpretation of the odd zeta motive goes as follows. Let $\mathfrak{g}^{\mathbb{Z},\vee}$ be the graded dual of the fundamental Lie algebra $\mathfrak{g}^{\mathbb{Z}}$ of the tannakian category $\text{MT}(\mathbb{Z})$. It is an ind-object in $\text{MT}(\mathbb{Z})$, independent of the choice of a fiber functor [Del89, Définition 6.1]. Then one has a short exact sequence

$$0 \rightarrow \mathbb{Q}(0) \rightarrow \mathfrak{g}^{\mathbb{Z},\vee} \rightarrow \mathfrak{u}^{\mathbb{Z},\vee} \rightarrow 0,$$

where $\mathfrak{u}^{\mathbb{Z}}$ is the pro-unipotent radical of $\mathfrak{g}^{\mathbb{Z}}$. One views \mathcal{Z}^{odd} inside the exact subsequence

$$0 \rightarrow \mathbb{Q}(0) \rightarrow \mathcal{Z}^{\text{odd}} \rightarrow \mathfrak{u}^{\mathbb{Z},\text{ab},\vee} \rightarrow 0,$$

where $\mathfrak{u}^{\mathbb{Z},\text{ab},\vee} \cong \bigoplus_{k \geq 1} \mathbb{Q}(-(2k + 1))$ is the graded dual of the abelianization of $\mathfrak{u}^{\mathbb{Z}}$.

4.6 An elementary computation of the motives $\mathcal{T}^{(n)}$

We give an elementary proof of Proposition 4.5, which only uses basic algebraic topology. The proof is Hodge-theoretic, and the only drawback is that we have to use the full faithfulness of the Hodge realization (Theorem 2.6). Let us consider the relative homology group

$$\mathcal{T}_{\mathbb{B}}^{(n),\vee} = H_n^{\text{sing}} \left((\mathbb{C}^*)^n, \bigcup_{1 \leq i \leq n} \{x_i = 1\} \cup \{x_1 \cdots x_n = 1\} \right).$$

By homotopy invariance, one may replace every \mathbb{C}^* by the unit circle $S^1 = \{|x| = 1\} \hookrightarrow \mathbb{C}^*$ and the divisor $\{x_1 \cdots x_n = 1\}$ by its intersection with $(S^1)^n$, and we get

$$\mathcal{T}_B^{(n),\vee} \cong H_n^{\text{sing}}\left((S^1)^n, \bigcup_{1 \leq i \leq n} \{x_i = 1\} \cup \{x_1 \cdots x_n = 1\}\right).$$

Let us look at the projection $[0, 1]^n \rightarrow (S^1)^n, (t_1, \dots, t_n) \mapsto (e^{2\pi i t_1}, \dots, e^{2\pi i t_n})$. Then by excision we can write

$$\mathcal{T}_B^{(n),\vee} \cong H_n^{\text{sing}}\left([0, 1]^n, \bigcup_{1 \leq i \leq n} \{t_i \in \mathbb{Z}\} \cup \{t_1 + \cdots + t_n \in \mathbb{Z}\}\right).$$

This is simply the singular homology of the unit hypercube $[0, 1]^n$ relative to the union of its faces $\{t_i = 0\}$ and $\{t_i = 1\}$, for $1 \leq i \leq n$, and the hyperplanes $\{t_1 + \cdots + t_n = k\}$ for $k = 0, 1, \dots, n$. We note that these hyperplanes cut the unit hypercube into polytopes

$$\Delta(n, k) = \{(t_1, \dots, t_n) \in [0, 1]^n \mid k \leq t_1 + \cdots + t_n \leq k + 1\},$$

for $k = 0, \dots, n - 1$. We note that $\Delta(n, 0)$ is the usual n -simplex; the polytopes $\Delta(n, k)$ are usually called *hypersimplices*.

LEMMA 4.14. (i) *The classes $[\Delta(n, k)]$, for $k = 0, \dots, n - 1$, form a basis of $\mathcal{T}_B^{(n),\vee}$.*

(ii) *The morphism $j_B^{(n),\vee} : \mathcal{T}_B^{(n),\vee} \rightarrow \mathcal{T}_B^{(n-1),\vee}$ sends*

- (a) $[\Delta(n, 0)]$ to $[\Delta(n - 1, 0)]$;
- (b) $[\Delta(n, k)]$ to $[\Delta(n - 1, k)] - [\Delta(n - 1, k - 1)]$ for $k = 1, \dots, n - 2$;
- (c) $[\Delta(n, n - 1)]$ to $-[\Delta(n - 1, n - 2)]$.

Proof. (i) This is clear by excision, since collapsing the boundary of $[0, 1]^n$ and the hyperplanes $\{t_1 + \cdots + t_n = k\}$ onto a point creates a wedge sum of n spheres of dimension n , one for each hypersimplex.

(ii) Recall (see Remark 4.4 and § 3.5) that $j_B^{(n),\vee}$ computes ‘the component of the boundary that lives on $\{x_n = 1\}$ ’. In the t -coordinates, $\{x_n = 1\}$ corresponds to $\{t_n = 0\}$ (counted positively) and $\{t_n = 1\}$ (counted negatively). In case (b), the intersection of $\Delta(n, k)$ with $\{t_n = 0\}$ is $\Delta(n - 1, k)$ and its intersection with $\{t_n = 1\}$ is $\Delta(n - 1, k - 1)$, which proves the claim. Cases (a) and (c) are similar. \square

Remark 4.15. One may check that the sum of the classes $[\Delta(n, k)]$, for $k = 0, \dots, n - 1$, is sent to 0 by the morphism $j_B^{(n),\vee}$. This is because this sum is represented by the unit square $[0, 1]^n$ in the t -coordinates, or by the compact n -torus $(S^1)^n \subset (\mathbb{C}^*)^n$ in the x -coordinates, which has empty boundary.

The *Eulerian numbers* are the coefficients of the Eulerian polynomials and are denoted by symbols $\langle n \rangle_k$:

$$E_n(x) = \sum_{k=0}^{n-1} \langle n \rangle_k x^k.$$

They satisfy many beautiful identities, including the following recursion, which can be deduced from (21):

$$\langle n \rangle_k = (n - k) \langle n - 1 \rangle_{k-1} + (k + 1) \langle n - 1 \rangle_k.$$

The following lemma is a classical result due to Laplace [Foa77, § 2].

LEMMA 4.16. For $k = 0, \dots, n - 1$, the volume of the hypersimplex $\Delta(n, k)$ is the ratio $\langle n \rangle_k / n!$.

Recall from Remark 4.6 that, for every integer $n \geq 1$, $\mathcal{T}_{\text{dR}}^{(n)}$ has a basis $(w_1^{(n)}, \dots, w_n^{(n)})$ which is compatible with the weight grading and with the morphisms $j_{\text{dR}}^{(n)} : \mathcal{T}_{\text{dR}}^{(n-1)} \rightarrow \mathcal{T}_{\text{dR}}^{(n)}$. We let P_n be the period matrix of $\mathcal{T}^{(n)}$ with respect to the w -basis and the Δ -basis from Lemma 4.14. The first period matrix P_1 is simply the 1×1 matrix $(2\pi i)$. Let us introduce the following $n \times n$ integer matrix encoding the family of Eulerian numbers:

$$A_n = \begin{pmatrix} 1 & & & & & \langle n \rangle_0 \\ -1 & 1 & & & & \langle n \rangle_1 \\ & -1 & 1 & & & \langle n \rangle_2 \\ & & \ddots & \ddots & & \\ & & & \ddots & \ddots & \\ 0 & & & -1 & 1 & \langle n \rangle_{n-2} \\ & & & & -1 & \langle n \rangle_{n-1} \end{pmatrix}.$$

PROPOSITION 4.17. The period matrices P_n satisfy the recurrence relation

$$P_n = A_n \left(\begin{array}{cccc|c} & & & & 0 \\ & & & & 0 \\ & & & & \vdots \\ & & & & 0 \\ \hline 0 & 0 & \dots & 0 & \frac{(2\pi i)^n}{n!} \end{array} \right).$$

Proof. Recall the short exact sequence (29)

$$0 \rightarrow \mathcal{T}^{(n-1)} \xrightarrow{j^{(n)}} \mathcal{T}^{(n)} \rightarrow H^n(T^n) \rightarrow 0$$

and the fact (see Remark 4.6) that the morphism $j^{(n)}$ is compatible with the w -bases. Then Lemma 4.14 shows that the first $(n - 1)$ columns of P_n are as stated. It only remains to compute the entries in the last column, i.e., compute the integral of the n -form $dx_1/x_1 \wedge \dots \wedge dx_n/x_n$ on a hypersimplex $\Delta(n, k)$. After the change of variables $(x_1, \dots, x_n) = (e^{2\pi i t_1}, \dots, e^{2\pi i t_n})$, one sees that this integral is simply $(2\pi i)^n$ times the volume of $\Delta(n, k)$, and completes the proof thanks to Lemma 4.16. \square

We note that the period matrices P_n are not block upper-triangular. This is because the Δ -basis is not compatible with the weight filtration. We thus have to introduce a change of basis.

Let $(Q_n)_{n \geq 1}$ be the family of matrices (with rational entries) defined by $Q_1 = (1)$ and the recurrence relation

$$Q_n = \left(\begin{array}{cccc|c} & & & & 0 \\ & & & & 0 \\ & & & & \vdots \\ & & & & 0 \\ \hline 0 & 0 & \dots & 0 & n! \end{array} \right) A_n^{-1}.$$

The first terms are

$$Q_1 = (1), \quad Q_2 = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ 1 & 1 \end{pmatrix}, \quad Q_3 = \begin{pmatrix} \frac{1}{3} & -\frac{1}{6} & \frac{1}{3} \\ 1 & 0 & -1 \\ 1 & 1 & 1 \end{pmatrix}, \quad Q_4 = \begin{pmatrix} \frac{1}{4} & -\frac{1}{12} & \frac{1}{12} & -\frac{1}{4} \\ \frac{11}{12} & -\frac{1}{12} & -\frac{1}{12} & \frac{11}{12} \\ \frac{3}{2} & \frac{1}{2} & -\frac{1}{2} & -\frac{3}{2} \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

Let us put

$$\begin{pmatrix} \Sigma_1^{(n)} \\ \Sigma_2^{(n)} \\ \vdots \\ \Sigma_n^{(n)} \end{pmatrix} = Q_n \begin{pmatrix} \Delta(n, 0) \\ \Delta(n, 1) \\ \vdots \\ \Delta(n, n-1) \end{pmatrix}.$$

We view $\Sigma_k^{(n)}$ as a relative cycle with rational coefficients. The change of indexing is here to remind the reader that $\Sigma_k^{(n)}$ lives in weight $\leq 2k$. We have thus proved the following result.

PROPOSITION 4.18. *The classes $[\Sigma_k^{(n)}]$, for $k = 1, \dots, n$, form a basis of $\mathcal{T}_B^{(n), \vee}$ and the period matrix of $\mathcal{T}^{(n)}$ in the w -basis and the Σ -basis is the diagonal matrix $\text{Diag}(2\pi i, \dots, (2\pi i)^n)$.*

Proof. This amounts to saying that the product $Q_n P_n$ is the matrix $\text{Diag}(2\pi i, \dots, (2\pi i)^n)$, which is easily proved by induction on n using Proposition 4.17. □

By using Theorem 2.6, we thus get an alternate (Hodge-theoretic) proof of Proposition 4.5.

Remark 4.19. Proposition 4.18 implies that we can choose $(\Sigma_1^{(n-1)}, \dots, \Sigma_{n-1}^{(n-1)})$ as representatives for the classes $(\psi_1^{(n-1)}, \dots, \psi_{n-1}^{(n-1)})$ from the proof of Theorem 4.9.

Remark 4.20. One can easily prove that the last row of the matrix Q_n is filled with 1s, which means that $\Sigma_n^{(n)}$ is homologous to the unit hypercube $[0, 1]^n$. In the x -coordinates, it is homologous to the compact n -torus $(S^1)^n \subset (\mathbb{C}^*)^n$.

5. Linear forms in zeta values

We apply our results from the previous section to prove Theorems 1.1 and 1.2 from the Introduction.

5.1 Integral formulas for the coefficients

THEOREM 5.1. *For ω an integrable algebraic differential form on $X_n - A_n$, we have*

$$\int_{[0,1]^n} \omega = a_0(\omega) + a_2(\omega)\zeta(2) + \dots + a_n(\omega)\zeta(n) \tag{37}$$

with $a_k(\omega)$ a rational number for every k , given for $k = 2, \dots, n$ by the formula

$$a_k(\omega) = (2\pi i)^{-k} \langle \varphi_k^{(n)}, [\omega] \rangle. \tag{38}$$

Proof. According to Proposition 3.6, the class $[\omega]$ defines an element in $\mathcal{Z}_{n,dR}$, hence we may write

$$[\omega] = a_0(\omega)v_0 + a_2(\omega)v_2 + \dots + a_n(\omega)v_n$$

with $a_k(\omega) \in \mathbb{Q}$ for every k . Pairing with the class $\varphi_0^{(n)} = [\square^n]$ gives the equality (37), and pairing with the class $\varphi_k^{(n)}$, $k = 2, \dots, n$, gives the equality (38). \square

Remark 5.2. If we represent the class $\varphi_k^{(n)}$ by a relative cycle $\sigma_k^{(n)}$, then (38) becomes

$$a_k(\omega) = (2\pi i)^{-k} \int_{\sigma_k^{(n)}} \omega.$$

Here we will not give explicit representatives for the classes $\varphi_k^{(n)}$. Recall from the proof of Theorem 4.9 that the class $\varphi_k^{(n)}$ is the image by the map $p_B^{(n),\vee} : \mathcal{T}_B^{(n-1),\vee} \rightarrow \mathcal{Z}_B^{(n),\vee}$ of an element $\psi_{k-1}^{(n-1)}$, which by Remark 4.19 can be represented by the cycle $\Sigma_{k-1}^{(n-1)}$. The question is then how to compute the map $p_B^{(n),\vee}$ at the level of cycles. Such a task would involve the following ingredients. Let $T \subset \mathbb{C}^n$ be a tubular neighborhood of $A_n(\mathbb{C})$ in \mathbb{C}^n . Let us denote by $\rho : T \rightarrow A_n(\mathbb{C})$ the corresponding projection, and by $\partial\rho : \partial T \rightarrow A_n(\mathbb{C})$ the projection corresponding to the boundary of the tubular neighborhood; it is an S^1 -bundle. The natural map $H_r^{\text{sing}}(A_n(\mathbb{C})) \rightarrow H_{r+1}^{\text{sing}}(\mathbb{C}^n - A_n(\mathbb{C}))$ can be computed at the level of singular chains by mapping an r -cycle σ to the $(r+1)$ -cycle $(\partial\rho)^{-1}(\sigma)$. We note that since $A_n(\mathbb{C})$ does not intersect the hyperplanes $\{x_i = 0\}$, we can do the computation with a tubular neighborhood inside $(\mathbb{C}^*)^n$ and get representatives in $(\mathbb{C}^*)^n$. Now if we want to play this game for the relative homology groups $\mathcal{Z}_B^{(n),\vee}$, we need the tubular neighborhood to be ‘compatible’ with the subvariety $B_n(\mathbb{C})$, in the sense that ρ should pull back $A_n(\mathbb{C}) \cap B_n(\mathbb{C})$ to $B_n(\mathbb{C})$. At this point, it is probably easier to ask for something weaker than a tubular neighborhood, i.e., something that is a tubular neighborhood on a dense open subset of $A_n(\mathbb{C})$ (this does not change anything for the integral formulas). We will not try to give formulas here and postpone this discussion to a future article. Nevertheless, we can give more explicit formulas than (38) in two situations.

5.1.1 The highest weight coefficient. Let us fix real numbers $\rho_1, \dots, \rho_{n-1}, \rho_n > 0$ and let us introduce the cycle $S^{(n)} \subset \mathbb{C}^n - A_n(\mathbb{C})$ defined by the conditions

$$|x_1| = \rho_1, \dots, |x_{n-1}| = \rho_{n-1}, \quad \left| x_n - \frac{1}{x_1 \dots x_{n-1}} \right| = \rho_n.$$

PROPOSITION 5.3. *Let ω be an integrable differential form on $X_n - A_n$. Then the highest weight coefficient $a_n(\omega)$ from Theorem 5.1 is given by the integral formula*

$$a_n(\omega) = (2\pi i)^{-n} \int_{S^{(n)}} \omega.$$

Proof. The integral formula is obviously independent of the choice of $\rho_1, \dots, \rho_{n-1}, \rho_n$ and we can assume that we have $\rho_1 = \dots = \rho_{n-1} = \rho_n = 1$. We have noted in Remark 4.20 that the highest weight basis vector $\psi_{n-1}^{(n-1)}$ of $\mathcal{T}_B^{(n-1), \vee}$ can be represented by the $(n-1)$ -torus $\{|x_1| = \dots = |x_{n-1}| = 1\}$. Since this has an empty boundary we can make the computation explained in Remark 5.2 with the choice of any tubular neighborhood of $A_n(\mathbb{C})$ in \mathbb{C}^n , for instance the one defined by $|x_n - 1/(x_1 \cdots x_{n-1})| \leq 1$, with projection map $\rho(x_1, \dots, x_n) = (x_1, \dots, x_{n-1}, 1/(x_1 \cdots x_{n-1}))$. The pullback of the $(n-1)$ -torus by the projection $\partial\rho$ is exactly $S^{(n)}$. \square

The case $n = 2$ is Rhin and Viola’s contour integral for $\zeta(2)$ [RV96, Lemma 2.6].

5.1.2 *The case of forms with simple poles.* We say that a differential form on $X_n - A_n$ has a simple pole along A_n if it can be written as

$$\omega = \alpha + \text{dlog}(1 - x_1 \cdots x_n) \wedge \beta,$$

where α and β do not have poles along A_n . The residue of such a form along A_n is the restriction

$$\text{Res}(\omega) = \beta|_{A_n}.$$

Recall that the relative cycles $\Sigma_{k-1}^{(n-1)}$ were defined in § 4.6.

PROPOSITION 5.4. *Let ω be an integrable differential form on $X_n - A_n$ which has a simple pole along A_n . Then the coefficients $a_k(\omega)$, $k = 2, \dots, n$, from Theorem 5.1 are given by the integral formulas*

$$a_k(\omega) = (2\pi i)^{-k+1} \int_{\Sigma_{k-1}^{(n-1)}} \text{Res}(\omega).$$

Proof. Recall from the proof of Theorem 4.9 that we have defined

$$\varphi_k^{(n)} = p_B^{(n), \vee}(\psi_{k-1}^{(n-1)}),$$

where $(\psi_1^{(n-1)}, \dots, \psi_{n-1}^{(n-1)})$ is a basis of $\mathcal{T}_B^{(n-1), \vee}$ for which the period matrix is diagonal. In the light of Remark 4.19 we see that $\psi_{k-1}^{(n-1)}$ is the class of the cycle $\Sigma_{k-1}^{(n-1)}$, hence we get

$$a_k(\omega) = (2\pi i)^{-k} \langle p_B^{(n), \vee}([\Sigma_{k-1}^{(n-1)}]), [\omega] \rangle = (2\pi i)^{-k+1} \langle [\Sigma_{k-1}^{(n-1)}], p_{\text{dR}}^{(n)}([\omega]) \rangle,$$

where the extra $2\pi i$ comes from the Tate twist at the target of $p^{(n)}$. Since ω has a simple pole, $p_{\text{dR}}^{(n)}([\omega])$ is simply the class of $\text{Res}(\omega)$, hence the result. \square

5.1.3 *Vanishing of coefficients.*

THEOREM 5.5. *For ω an integrable algebraic differential form on $X_n - A_n$, we have:*

- (i) if $\tau.\omega = \omega$ then $a_k(\omega) = 0$ for $k \neq 0$ even;
- (ii) if $\tau.\omega = -\omega$ then $a_k(\omega) = 0$ for k odd.

Proof. Let us assume that we have $\tau.\omega = \omega$, and let us write x for the image of $[\omega]$ in $\mathcal{T}_{\text{dR}}^{(n-1)}$. Then we have $\tau.x = x$; according to Proposition 4.5, this implies that x only has components of weights $2k$ with k even. Thus, $[\omega] \in \mathcal{Z}_{\text{dR}}^{(n)}$ only has components in weight 0 and $2k$ with k odd, which implies that we have $a_k(\omega) = 0$ for $k \neq 0$ even. The second case is similar. \square

Let us write an integrable form as

$$\omega = \frac{P(x_1, \dots, x_n)}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n \tag{39}$$

with $P(x_1, \dots, x_n)$ a polynomial with rational coefficients and $N \geq 0$ an integer. Then we have

$$\tau.\omega = \pm\omega \iff P(x_1, \dots, x_n) = \pm(-1)^{N+n}(x_1 \cdots x_n)^{N-2}P(x_1^{-1}, \dots, x_n^{-1}). \tag{40}$$

5.2 The Ball–Rivoal integrals

We apply Theorems 5.1 and 5.5 to a special family of integrals.

COROLLARY 5.6. *Let $u_1, \dots, u_n, v_1, \dots, v_n \geq 1$ and $N \geq 0$ be integers such that $v_1 + \dots + v_n \geq N + 1$. Then the integral*

$$\int_{[0,1]^n} \frac{x_1^{u_1-1} \cdots x_n^{u_n-1} (1 - x_1)^{v_1-1} \cdots (1 - x_n)^{v_n-1}}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_n \tag{41}$$

is absolutely convergent and evaluates to a linear combination

$$a_0 + a_2\zeta(2) + a_3\zeta(3) + \cdots + a_n\zeta(n)$$

with a_k a rational number for every k . If furthermore we have $2u_i + v_i = N + 1$ for every i , then we get:

- (i) if $(n + 1)(N + 1)$ is odd then $a_k = 0$ for $k \neq 0$ even;
- (ii) if $(n + 1)(N + 1)$ is even then $a_k = 0$ for k odd.

Proof. This is a direct application of Theorem 5.5. The polynomial

$$P(x_1, \dots, x_n) = x_1^{u_1-1} \cdots x_n^{u_n-1} (1 - x_1)^{v_1-1} \cdots (1 - x_n)^{v_n-1}$$

satisfies

$$P(x_1, \dots, x_n) = (-1)^{n+v_1+\dots+v_n} x_1^{2u_1+v_1-3} \cdots x_n^{2u_n+v_n-3} P(x_1^{-1}, \dots, x_n^{-1}).$$

Let us assume that we have $2u_i + v_i = N + 1$ for every i , then $v_1 + \dots + v_n \equiv n(N + 1) \pmod{2}$ and we get

$$P(x_1, \dots, x_n) = -(-1)^{(n+1)(N+1)} (-1)^{N+n} (x_1 \cdots x_n)^{N-2} P(x_1^{-1}, \dots, x_n^{-1}),$$

hence the result, in view of (40). □

Corollary 5.6 applies in particular to the special case

$$N = (2r + 1)m + 2, \quad u_i = rm + 1, \quad v_i = m + 1$$

for some integer parameters $r, m \geq 0$ satisfying $n(m + 1) \geq (2r + 1)m + 3$. We then recover the integrals considered by Ball and Rivoal [BR01, Lemme 2]. The vanishing of the coefficients is [BR01, Lemme 1]. The notations (a, n, r) in [BR01] correspond to our notations $(n - 1, m, r)$.

The integrals (41) can be expressed as generalized hypergeometric series

$$\begin{aligned} & \left(\prod_{i=1}^n \frac{(u_i - 1)!(v_i - 1)!}{(u_i + v_i - 1)!} \right) {}_{n+1}F_n \left(\begin{matrix} u_1, \dots, u_n, N \\ u_1 + v_1, \dots, u_n + v_n \end{matrix}; 1 \right) \\ &= \frac{\prod_{i=1}^n (v_i - 1)!}{(N - 1)!} \sum_{k \geq 0} \frac{(k)_{u_1} \cdots (k)_{u_n} (k + 1)_{N-1}}{(k)_{u_1+v_1} \cdots (k)_{u_n+v_n}}. \end{aligned} \tag{42}$$

If $2u_i + v_i = N + 1$, then the corresponding generalized hypergeometric series is said to be *well-poised*.

5.3 Weight drop

In the context of Theorem 5.1, we say that the integral $\int_{[0,1]^n} \omega$ has *weight drop* if the highest weight coefficient $a_n(\omega)$ vanishes. This amounts to saying that the class $[\omega]$ actually lives in the step $W_{2(n-1)}\mathcal{Z}_{n,\text{dR}}$ of the weight filtration, hence the terminology. We give a sufficient condition for this phenomenon to happen.

LEMMA 5.7. *Let $u, v \geq 1$ and $N \geq 0$ be integers such that $u + v \leq N$. Then there exists a polynomial $P(t)$ with rational coefficients such that*

$$\int_0^1 \frac{x^{u-1}(1-x)^{v-1}}{(1-tx)^N} dx = \frac{P(t)}{(1-t)^{N-v}}$$

for every $0 \leq t < 1$.

Proof. We can write

$$x^{u-1}(1-x)^{v-1} = \sum_{k=0}^{u+v-2} a_k(t)(1-tx)^k$$

with $a_k(t)$ a Laurent polynomial with rational coefficients for every k . We then have

$$\frac{x^{u-1}(1-x)^{v-1}}{(1-tx)^N} = \sum_{k=0}^{u+v-2} \frac{a_k(t)}{(1-tx)^{N-k}}$$

and all the powers of $(1-tx)$ appearing in the denominators are $\geq N-(u+v-2) \geq N-u-v+2 \geq 2$. Thus, we may integrate and get

$$\int_0^1 \frac{x^{u-1}(1-x)^{v-1}}{(1-tx)^N} dx = \frac{Q(t)}{(1-t)^{N-1}}$$

with $Q(t)$ a Laurent polynomial with rational coefficients. The left-hand side has a limit when t tends to 0, so $Q(t)$ has to be a polynomial. To finish, it is enough to show that

$$(1-t)^{N-v} \int_0^1 \frac{x^{u-1}(1-x)^{v-1}}{(1-tx)^N} dx$$

is bounded when t approaches 1. We make the change of variables $s = 1 - t$, $y = 1 - x$, and consider integrals

$$s^{N-v} \int_0^1 \frac{(1-y)^{u-1}y^{v-1}}{(y+s-ys)^N} dy$$

with s approaching 0. Since $(1-y)^{u-1} \leq 1$ and $y+s-ys \geq \frac{1}{2}(y+s)$, it is enough to prove that the quantities

$$s^{N-v} \int_0^1 \frac{y^{v-1}}{(y+s)^N} dy$$

are bounded when s approaches 0. This equals

$$s^{N-v} \int_0^1 \left(\frac{y}{y+s}\right)^{v-1} \frac{dy}{(y+s)^{N-v+1}} \leq s^{N-v} \int_0^1 \frac{dy}{(y+s)^{N-v+1}} = \frac{1}{N-v} \left(1 - \left(\frac{s}{1+s}\right)^{N-v}\right)$$

and we are done. □

PROPOSITION 5.8. Let $u_1, \dots, u_n, v_1, \dots, v_n \geq 1$ and $N \geq 0$ be integers such that $v_1 + \dots + v_n \geq N + 1$. Let us assume that there exists an index $i \in \{1, \dots, n\}$ such that

$$u_i + v_i \leq N.$$

Then the integral

$$\int_{[0,1]^n} \frac{x_1^{u_1-1} \dots x_n^{u_n-1} (1-x_1)^{v_1-1} \dots (1-x_n)^{v_n-1}}{(1-x_1 \dots x_n)^N} dx_1 \dots dx_n$$

is absolutely convergent and evaluates to a linear combination

$$a_0 + a_2\zeta(2) + a_3\zeta(3) + \dots + a_{n-1}\zeta(n-1)$$

with $a_i \in \mathbb{Q}$ for every i .

Proof. By symmetry, we can assume that $u_n + v_n \leq N$. Therefore, applying Lemma 5.7 to the variables $x = x_n$ and $t = x_1 \dots x_{n-1}$ in the integral leads to the $(n-1)$ -dimensional integral

$$\int_{[0,1]^{n-1}} \frac{x_1^{u_1-1} \dots x_{n-1}^{u_{n-1}-1} (1-x_1)^{v_1-1} \dots (1-x_{n-1})^{v_{n-1}-1} P(x_1 \dots x_{n-1})}{(1-x_1 \dots x_{n-1})^{N-v_n}} dx_1 \dots dx_{n-1}.$$

Since $v_1 + \dots + v_{n-1} \geq N - v_n + 1$, one can then finish thanks to Theorem 5.1. □

Note that Proposition 5.8 applies in particular if for every i , $2u_i + v_i = N + 1$. This gives in particular a geometric interpretation of the weight drop in the Ball–Rivoal integrals [Riv00, BR01], which comes from the representations as hypergeometric series (42). Note that a careful analysis of the degree of the polynomial $P(t)$ in Lemma 5.7 can lead to sufficient conditions for the vanishing of the subleading coefficients.

ACKNOWLEDGEMENTS

Many thanks to Francis Brown, Pierre Cartier, Tanguy Rivoal and Don Zagier for fruitful discussions as well as comments and corrections on a preliminary version.

Appendix. An approach via series

Written jointly with Don Zagier

The aim of this appendix is to give an elementary construction of the coefficients $a_k(\omega)$ from Theorem 5.1. The dictionary between integrals and sums of series leads to an interpretation of the (de Rham realization of the) zeta motive \mathcal{Z} , modulo weight 0, in terms of rational functions in one variable.

A.1 Series, integrals, and zeta values

A.1.1 *Series of rational functions and zeta values.* We denote by $\mathbb{Q}(k)$ the field of rational functions in the variable k with rational coefficients. Let V denote the subspace of $\mathbb{Q}(k)$ consisting of rational functions with poles in $\{-1, -2, -3, \dots\}$ and V_0 be the subspace of functions vanishing at ∞ . Then $V = V_0 \oplus \mathbb{Q}[k]$ and the set of functions $(k+j)^{-r}$, with $j, r \geq 1$ integers, is a basis of V_0 . The forward difference operator $\Delta : \mathbb{Q}(k) \rightarrow \mathbb{Q}(k)$ defined by $\Delta R(k) = R(k+1) - R(k)$ preserves

the spaces V and V_0 and one has direct sum decompositions $V = \Delta(V) \oplus B$ and $V_0 = \Delta(V_0) \oplus B$, where B is the space spanned by the functions $(k + 1)^{-r}$, for $r \geq 1$ integers. We thus have an identification $V_0/\Delta(V_0) \cong V/\Delta(V)$ and an isomorphism

$$\beta : V/\Delta(V) \xrightarrow{\cong} \bigoplus_{r \geq 1} \mathbb{Q}, \quad R \mapsto (\beta_1(R), \beta_2(R), \dots), \tag{A.1}$$

where the numbers $\beta_r(R) \in \mathbb{Q}$, for $R \in V$, are defined by

$$R(k) \equiv \sum_{r \geq 1} \frac{\beta_r(R)}{(k + 1)^r} \pmod{\Delta(V)}.$$

For $R \in V_0$ we can write

$$R(k) = \sum_{r \geq 1} \frac{\beta_r(R)}{(k + 1)^r} - \Delta R_0(k),$$

for some $R_0 \in V_0$, which is unique because $\Delta : V_0 \rightarrow V_0$ is injective. Thus, the sum $\sum_{k=0}^{\infty} R(k)$ is absolutely convergent if and only if $R \in V_0$ and $\beta_1(R) = 0$, and in this case we have

$$\sum_{k=0}^{\infty} R(k) = R_0(0) + \sum_{r \geq 2} \beta_r(R) \zeta(r) \in \mathbb{Q} + \sum_{r \geq 2} \mathbb{Q} \zeta(r). \tag{A.2}$$

A.1.2 From differential forms to rational functions. For $n \geq 1$ an integer, we define

$$\Omega_n = \mathbb{Q}[x_1, \dots, x_n, (1 - x_1 \cdots x_n)^{-1}]$$

and we interpret an element $F \in \Omega_n$ as the algebraic differential n -form $\omega = F dx_1 \cdots dx_n$.

LEMMA A.1. *The formula*

$$\Phi_n \left(\frac{x_1^{a_1-1} \cdots x_n^{a_n-1}}{(1 - x_1 \cdots x_n)^N} \right) = \begin{cases} 0 & \text{if } N = 0, \\ \binom{k + N - 1}{N - 1} \frac{1}{(k + a_1) \cdots (k + a_n)} & \text{if } N \geq 1, \end{cases} \tag{A.3}$$

for $a_1, \dots, a_n \geq 1$ and $N \geq 0$ integers, defines a linear map $\Phi_n : \Omega_n \rightarrow V/\Delta(V)$.

Proof. If we rewrite

$$x_1^{a_1-1} \cdots x_n^{a_n-1} \quad \text{as} \quad \frac{x_1^{a_1-1} \cdots x_n^{a_n-1} - x_1^{a_1} \cdots x_n^{a_n}}{1 - x_1 \cdots x_n},$$

then its image by Φ_n is

$$\begin{aligned} \frac{1}{(k + a_1) \cdots (k + a_n)} - \frac{1}{(k + a_1 + 1) \cdots (k + a_n + 1)} &= \Delta \left(-\frac{1}{(k + a_1) \cdots (k + a_n)} \right) \\ &\equiv 0 \pmod{\Delta(V)}. \end{aligned}$$

For $N \geq 1$, if we rewrite

$$\frac{x_1^{a_1-1} \cdots x_n^{a_n-1}}{(1 - x_1 \cdots x_n)^N} \quad \text{as} \quad \frac{x_1^{a_1-1} \cdots x_n^{a_n-1} - x_1^{a_1} \cdots x_n^{a_n}}{(1 - x_1 \cdots x_n)^{N+1}},$$

then we replace the function

$$R(k) = \binom{k + N - 1}{N - 1} \frac{1}{(k + a_1) \cdots (k + a_n)}$$

by the function

$$\begin{aligned} R^*(k) &= \binom{k + N}{N} \left(\frac{1}{(k + a_1) \cdots (k + a_n)} - \frac{1}{(k + a_1 + 1) \cdots (k + a_n + 1)} \right) \\ &= R(k) + \frac{kR(k) - (k + 1)R(k + 1)}{N} \\ &\equiv R(k) \pmod{\Delta(V)}. \end{aligned}$$

This shows that the definition of $\Phi_n(F)$ for $F \in (1 - x_1 \cdots x_n)^{-N} \mathbb{Q}[x_1, \dots, x_n]$ is independent of the choice of N . □

Combining with (A.1), we get well-defined maps

$$b_r : \Omega_n \rightarrow \mathbb{Q}, \quad \omega \mapsto \beta_r(\Phi_n(\omega)).$$

Note that this is zero for $r > n$ for degree reasons. We denote by $\Omega_n^{\text{int}} \subset \Omega_n$ the subspace of integrable differential forms, which are the forms ω such that the integral $\int_{[0,1]^n} \omega$ is absolutely convergent (see Definition 3.5 and Propositions 3.6 and 3.7).

PROPOSITION A.2. *For every $\omega \in \Omega_n^{\text{int}}$ we have $b_1(\omega) = 0$ and*

$$\int_{[0,1]^n} \omega = b_2(\omega)\zeta(2) + \cdots + b_n(\omega)\zeta(n) \pmod{\mathbb{Q}}.$$

Proof. Let us write $\omega = P(x_1, \dots, x_n)/(1 - x_1 \cdots x_n)^N$ with $P(x_1, \dots, x_n)$ a polynomial with rational coefficients and $N \geq 1$ an integer. Let $R \in V$ be the representative of $\Phi_n(\omega)$ obtained by applying (A.3) to every monomial in $P(x_1, \dots, x_n)$ and using linearity. Then the formula

$$\frac{1}{(1 - x)^N} = \sum_{k=0}^{\infty} \binom{k + N - 1}{N - 1} x^k$$

implies that we have

$$\int_{[0,1]^n} \omega = \sum_{k=0}^{\infty} R(k).$$

Thus, the sum $\sum_{k=0}^{\infty} R(k)$ is convergent, which implies that we have $R \in V_0$ and $\beta_1(R) = 0$. The claim then follows from (A.2). □

Proposition A.2 implies that there is a well-defined map $b_0 : \Omega_n^{\text{int}} \rightarrow \mathbb{Q}$ such that for every $\omega \in \Omega_n^{\text{int}}$ we have

$$\int_{[0,1]^n} \omega = b_0(\omega) + b_2(\omega)\zeta(2) + \cdots + b_n(\omega)\zeta(n). \tag{A.4}$$

We note that applying Φ_n to the integrals (41) leads to the hypergeometric series representations (42).

A.1.3 *Parity.* Let us recall that τ denotes the involution $(x_1, \dots, x_n) \mapsto (x_1^{-1}, \dots, x_n^{-1})$. The following proposition is nothing but a generalization of the classical well-posed symmetry of the hypergeometric series (42), and is similar to the parity considerations in [Zud04, § 8] and [CFR08b, § 3.1].

PROPOSITION A.3. *Let $\omega \in \Omega_n$ be a differential form such that $\tau.\omega$ belongs to Ω_n . We have, for every integer $r \geq 1$,*

$$b_r(\tau.\omega) = (-1)^{r-1}b_r(\omega).$$

In particular, we have:

- (i) if $\tau.\omega = \omega$ then $b_r(\omega) = 0$ for $r \neq 0$ even;
- (ii) if $\tau.\omega = -\omega$ then $b_r(\omega) = 0$ for r odd.

Proof. Let R and S be representatives of $\Phi_n(\omega)$ and $\Phi_n(\tau.\omega)$ respectively, constructed as in the proof of Proposition A.2. The involution τ acts on differential forms by the formula

$$\frac{x_1^{a_1-1} \dots x_n^{a_n-1}}{(1-x_1 \dots x_n)^N} dx_1 \dots dx_n \mapsto (-1)^{N+n} \frac{x_1^{N-a_1-1} \dots x_n^{N-a_n-1}}{(1-x_1 \dots x_n)^N} dx_1 \dots dx_n.$$

Thus, by looking at the formula for Φ_n , we see that we have $S(k) = -R(-N-k)$. This implies, for every integer $r \geq 1$, the equality,

$$\beta_r(S) = (-1)^{r-1}\beta_r(R),$$

and the claim follows. □

A.2 Comparison of the coefficients

The aim of this section is to prove the following theorem.

THEOREM A.4. *For every $\omega \in \Omega_n^{\text{int}}$ and every integer $r = 0, 2, \dots, n$ we have $a_r(\omega) = b_r(\omega)$.*

Note that this theorem would follow from the conjecture that 1 and the zeta values $\zeta(n)$, $n \geq 2$, are linearly independent over \mathbb{Q} , by looking at (37) and (A.4).

A.2.1 *Inductive structure on the motives $\mathcal{Z}^{(n)}$.* Let us recall from § 3.5 the morphisms $i_{\text{dR}}^{(n)} : \mathcal{Z}_{\text{dR}}^{(n-1)} \rightarrow \mathcal{Z}_{\text{dR}}^{(n)}$, which come from the identification $X_{n-1} = \{x_n = 1\} \subset X_n$. Let us consider an $(n-1)$ -form of the type

$$\eta = \frac{P(x_1, \dots, x_n)}{(1-x_1 \dots x_n)^N} dx_1 \dots dx_{n-1},$$

with $P(x_1, \dots, x_n)$ a polynomial with rational coefficients and $N \geq 0$ an integer. We say that such a form is integrable if the pullback $\pi_n^*(\eta)$ does not have a pole along the exceptional divisor E_n (π_n and E_n are introduced in § 3.1). This can be characterized in the same way as in Propositions 3.6 and 3.7, but we will not need such a characterization. If η is integrable, then its derivative $d\eta$ is integrable in the sense of Definition 3.5, and the restriction $\eta|_{x_n=1}$, viewed as a form on X_{n-1} , is also integrable. We then have classes $[d\eta] \in \mathcal{Z}_{\text{dR}}^{(n)}$ and $[\eta|_{x_n=1}] \in \mathcal{Z}_{\text{dR}}^{(n-1)}$. They are related by the formula

$$i_{n,\text{dR}}([\eta|_{x_n=1}]) \equiv (-1)^{n-1}[d\eta] \pmod{W_0\mathcal{Z}_{\text{dR}}^{(n)}},$$

which is proved as in the proof of Proposition 3.10, by noticing that $\eta|_{x_n=0}$ is a polynomial, hence has weight zero. This formula is the de Rham-theoretic incarnation of Stokes’s formula

$$(-1)^{n-1} \int_{[0,1]^n} d\eta = \left(\int_{[0,1]^{n-1}} \eta|_{x_n=1} - \int_{[0,1]^{n-1}} \eta|_{x_n=0} \right) \equiv \int_{[0,1]^{n-1}} \eta|_{x_n=1} \pmod{\mathbb{Q}}.$$

If we now choose to make the identification $X_{n-1} = \{x_j = 1\} \subset X_n$, for some index $j = 1, \dots, n$, then we get a morphism $i_{\text{dR}}^{(n),j} : \mathcal{Z}_{\text{dR}}^{(n-1)} \rightarrow \mathcal{Z}_{\text{dR}}^{(n)}$, such that $i_{\text{dR}}^{(n)} = i_{\text{dR}}^{(n),n}$. They satisfy the equation

$$i_{\text{dR}}^{(n),j}([\eta|_{x_j=1}]) \equiv (-1)^{j-1}[d\eta] \pmod{W_0\mathcal{Z}_{\text{dR}}^{(n)}}, \tag{A.5}$$

for η an integrable $(n - 1)$ -form of the type

$$\frac{P(x_1, \dots, x_n)}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots \widehat{dx_j} \cdots dx_n. \tag{A.6}$$

One easily notes that the morphism $i_{\text{dR}}^{(n),j}$ does not depend on the index j , for instance by proving that Proposition 3.10 is valid for any choice of j : for every $d = 0, 2, \dots, n - 1$, the map $i_{\text{dR}}^{(n),j}$ sends the class $[\omega_d^{(n-1)}]$ to the class $[\omega_d^{(n)}]$. We nevertheless keep the notation $i_{\text{dR}}^{(n),j}$ since these morphisms have different geometric interpretations.

A.2.2 Compatibility of Φ_n with the induction. The crucial point is that the morphisms Φ_n are compatible with the inductive structure (A.5) on the motives $\mathcal{Z}_{\text{dR}}^{(n)}$, in the sense of the following lemma.

LEMMA A.5. For every $j = 1, \dots, n$ and every differential $(n - 1)$ -form η of type (A.6) we have

$$\Phi_n(d\eta) \equiv (-1)^{j-1}\Phi_{n-1}(\eta|_{x_j=1}) \pmod{\Delta(V)}.$$

Proof. We do the case $j = n$, the general case being similar. It is enough to do the proof for a monomial

$$\eta = \frac{x_1^{a_1-1} \cdots x_{n-1}^{a_{n-1}-1} x_n^{a_n}}{(1 - x_1 \cdots x_n)^N} dx_1 \cdots dx_{n-1},$$

with $a_1, \dots, a_{n-1} \geq 1, a_n \geq 0$ and $N \geq 1$. We have

$$(-1)^{n-1}d\eta = \left(a_n \frac{x_1^{a_1-1} \cdots x_n^{a_n-1}}{(1 - x_1 \cdots x_n)^N} + N \frac{x_1^{a_1} \cdots x_n^{a_n}}{(1 - x_1 \cdots x_n)^{N+1}} \right) dx_1 \cdots dx_n,$$

and thus $(-1)^{n-1}\Phi_n(d\eta)$ equals

$$a_n \binom{k + N - 1}{N - 1} \frac{1}{(k + a_1) \cdots (k + a_n)} + N \binom{k + N}{N} \frac{1}{(k + a_1 + 1) \cdots (k + a_n + 1)}.$$

By writing $a_n/(k + a_n) = 1 - k/(k + a_n)$ and $N \binom{k+N}{N} = (k + 1) \binom{k+N}{N-1}$, we get

$$(-1)^{n-1}\Phi_n(d\eta) \equiv \binom{k + N - 1}{N - 1} \frac{1}{(k + a_1) \cdots (k + a_{n-1})} = \Phi_{n-1}(\eta|_{x_n=1}) \pmod{\Delta(V)}. \quad \square$$

A.2.3 *Proof of Theorem A.4.* We prove Theorem A.4 by induction on n . The case $n = 1$ is trivial since in this case we have $a_0(\omega) = b_0(\omega) = \int_0^1 \omega$. Let us then assume that $n \geq 2$ and that the theorem is proved for $n - 1$. Recall the notation

$$\omega_n^{(n)} = \frac{dx_1 \cdots dx_n}{1 - x_1 \cdots x_n}$$

for the representative of the highest weight basis element in $\mathcal{Z}_{n,\text{dR}}$; it satisfies $\Phi_n(\omega_n^{(n)}) = (k + 1)^{-n}$. The short exact sequence (31) implies that, for every $\omega \in \Omega_n^{\text{int}}$, we may write

$$\omega = a_n(\omega)\omega_n^{(n)} + \sum_{j=1}^n d\eta_j$$

with η_j an integrable $(n - 1)$ -form of type (A.6), for every $j = 1, \dots, n$. The short exact sequence (31) actually implies that in addition we can assume that the classes of $d\eta_1, \dots, d\eta_{n-1}$ are zero, but we will not need it here. By using (A.5) we may write

$$[\omega] = a_n(\omega)[\omega_n^{(n)}] + \sum_{j=1}^n (-1)^{j-1} i_{\text{dR}}^{(n),j}([\eta_j]_{|x_j=1}) \pmod{W_0 \mathcal{Z}_{\text{dR}}^{(n)}}.$$

Now Lemma A.5 implies the formula

$$\Phi_n(\omega) \equiv \frac{a_n(\omega)}{(k + 1)^n} + \sum_{j=1}^n (-1)^{j-1} \Phi_{n-1}([\eta_j]_{|x_j=1}) \pmod{\Delta(V)}.$$

By using the induction hypothesis on the forms $(\eta_j)_{|x_j=1}$ and the fact that the morphisms $i_{\text{dR}}^{(n),j}$ are compatible with the bases, this implies that we have

$$\Phi_n(\omega) \equiv \sum_{r=2}^n \frac{a_r(\omega)}{(k + 1)^r} \pmod{\Delta(V)},$$

which completes the proof.

We note that a restatement of Theorem A.4 is that the morphisms Φ_n induce an isomorphism of graded vector spaces

$$\Phi : \mathcal{Z}_{\text{dR}}/W_0 \mathcal{Z}_{\text{dR}} \xrightarrow{\cong} (V/\Delta(V))_{\geq 2},$$

where $(V/\Delta(V))_{\geq 2}$ is the subspace of $V/\Delta(V)$ characterized by the condition $\beta_1 = 0$ and is graded by the morphisms $\beta_n, n \geq 2$.

REFERENCES

Apé79 R. Apéry, *Irrationalité de $\zeta(2)$ et $\zeta(3)$* , Astérisque **61** (1979), 11–13.
 BR01 K. Ball and T. Rivoal, *Irrationalité d’une infinité de valeurs de la fonction zêta aux entiers impairs*, Invent. Math. **146** (2001), 193–207.
 Beu79 F. Beukers, *A note on the irrationality of $\zeta(2)$ and $\zeta(3)$* , Bull. Lond. Math. Soc. **11** (1979), 268–272.
 Blo86 S. Bloch, *Algebraic cycles and higher K-theory*, Adv. Math. **61** (1986), 267–304.
 Bor77 A. Borel, *Cohomologie de SL_n et valeurs de fonctions zeta aux points entiers*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) **4** (1977), 613–636.

- Bro16 F. Brown, *Irrationality proofs for zeta values, moduli spaces and dinner parties*, Mosc. J. Comb. Number Theory **6** (2016), 102–165.
- CFR08a J. Cresson, S. Fischler and T. Rivoal, *Phénomènes de symétrie dans des formes linéaires en polyzêtas*, J. Reine Angew. Math. **617** (2008), 109–151.
- CFR08b J. Cresson, S. Fischler and T. Rivoal, *Séries hypergéométriques multiples et polyzêtas*, Bull. Soc. Math. France **136** (2008), 97–145.
- Del71 P. Deligne, *Théorie de Hodge II*, Publ. Math. Inst. Hautes Études Sci. **40** (1971), 5–57.
- Del74 P. Deligne, *Théorie de Hodge III*, Publ. Math. Inst. Hautes Études Sci. **44** (1974), 5–77.
- Del89 P. Deligne, *Le groupe fondamental de la droite projective moins trois points*, in *Galois groups over \mathbf{Q}* , Berkeley, CA, 1987, Mathematical Sciences Research Institute Publications, vol. 16 (Springer, New York, NY, 1989), 79–297.
- DG05 P. Deligne and A. B. Goncharov, *Groupes fondamentaux motiviques de Tate mixte*, Ann. Sci. Éc. Norm. Supér. (4) **38** (2005), 1–56.
- Dup17 C. Dupont, *Relative cohomology of bi-arrangements*, Trans. Amer. Math. Soc. **369** (2017), 8105–8160.
- Fis04 S. Fischler, *Irrationalité de valeurs de zêta (d’après Apéry, Rivoal, . . .)*, Astérisque **294** (2004), vii, 27–62.
- Foa77 D. Foata, *Distributions eulériennes et mahoniennes sur le groupe des permutations*, in *Higher combinatorics: Proc. NATO Advanced Study Institute, Berlin, 1976*, NATO Advanced Study Institutes Series, Series C, vol. 31 (Reidel, Dordrecht, 1977), 27–49.
- Foa10 D. Foata, *Eulerian polynomials: from Euler’s time to the present*, in *The legacy of Alladi Ramakrishnan in the mathematical sciences* (Springer, New York, 2010), 253–273.
- Gon02 A. B. Goncharov, *Periods and mixed motives*, Preprint (2002), [arXiv:math.AG/0202154](https://arxiv.org/abs/math/0202154).
- GM04 A. B. Goncharov and Yu. I. Manin, *Multiple ζ -motives and moduli spaces $\overline{\mathcal{M}}_{0,n}$* , Compos. Math. **140** (2004), 1–14.
- Hat02 A. Hatcher, *Algebraic topology* (Cambridge University Press, Cambridge, 2002).
- Hub00 A. Huber, *Realization of Voevodsky’s motives*, J. Algebraic Geom. **9** (2000), 755–799.
- Hub04 A. Huber, *Corrigendum to: “Realization of Voevodsky’s motives” [J. Algebraic Geom.] 9 (2000) 755–799; mr1775312*, J. Algebraic Geom. **13** (2004), 195–207.
- Lev93 M. Levine, *Tate motives and the vanishing conjectures for algebraic K-theory*, in *Algebraic K-theory and algebraic topology Lake Louise, AB, 1991*, NATO Advanced Study Institutes Series, Series C, vol. 407 (Kluwer Academic, Dordrecht, 1993), 167–188.
- Riv00 T. Rivoal, *La fonction zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs*, C. R. Acad. Sci. Paris Sér. I Math. **331** (2000), 267–270.
- RV96 G. Rhin and C. Viola, *On a permutation group related to $\zeta(2)$* , Acta Arith. **77** (1996), 23–56.
- RV01 G. Rhin and C. Viola, *The group structure for $\zeta(3)$* , Acta Arith. **97** (2001), 269–293.
- Voe00 V. Voevodsky, *Triangulated categories of motives over a field*, in *Cycles, transfers, and motivic homology theories*, Annals of Mathematics Studies, vol. 143 (Princeton University Press, Princeton, NJ, 2000), 188–238.
- Zud04 W. Zudilin, *Arithmetic of linear forms involving odd zeta values*, J. Théor. Nombres Bordeaux **16** (2004), 251–291.

Clément Dupont clement.dupont@umontpellier.fr
 Institut Montpellierain Alexander Grothendieck, CNRS, Univ. Montpellier, France

Don Zagier dbz@mpim-bonn.mpg.de
 Max-Planck-Institut für Mathematik, Vivatsgasse, 7, 53111 Bonn, Germany