

The Erdős–Rado Arrow for Singular Cardinals

Saharon Shelah

Abstract. We prove in ZFC that if $\text{cf}(\lambda) > \aleph_0$ and $2^{\text{cf}(\lambda)} < \lambda$, then $\lambda \rightarrow (\lambda, \omega + 1)^2$.

1 Introduction

For every finite cardinal κ , the Erdős–Dushnik–Miller theorem, [1, Theorem 11.1], states that $\kappa \rightarrow (\kappa, \omega)^2$. Erdős, Hajnal, Maté, and Rado proved that $\kappa \rightarrow (\kappa, \omega + 1)$ for every regular uncountable κ , (see [1, Theorem 11.3]). For singular cardinals, κ , they were only able to obtain the weaker result in [1, Theorem 11.1] that $\kappa \rightarrow (\kappa, \omega)^2$. It is not hard to see that if $\text{cf}(\kappa) = \omega$, then $\kappa \not\rightarrow (\kappa, \omega + 1)^2$. If $\text{cf}(\kappa) > \omega$ and κ is a strong limit cardinal, then it follows from the General Canonization Lemma, [1, Lemma 28.1], that $\kappa \rightarrow (\kappa, \omega + 1)^2$. Question 11.4 of [1] is whether this holds without the assumption that κ is a strong limit cardinal, *e.g.*, whether, in ZFC,

$$\aleph_{\omega_1} \rightarrow (\aleph_{\omega_1}, \omega + 1)^2.$$

In [5] it was proved that $\lambda \rightarrow (\lambda, \omega + 1)^2$ if $2^{\text{cf}(\lambda)} < \lambda$ and there is a nice filter on κ (see [3, Ch.V]; it follows from suitable failures of SCH). Also proved there are consistency results when $2^{\text{cf}(\lambda)} > \lambda$.

Here, continuing [5] but not relying on it, we eliminate the extra assumption, *i.e.*, we prove the following (in ZFC).

Theorem 1.1 *If $\aleph_0 < \kappa = \text{cf}(\lambda)$ and $2^\kappa < \lambda$ then $\lambda \rightarrow (\lambda, \omega + 1)^2$.*

Before starting the proof, let us recall the well-known definition.

Definition 1.2 Let D be an \aleph_1 -complete filter on Y , $f \in {}^Y\text{Ord}$, and $\alpha \in \text{Ord} \cup \{\infty\}$.

We define $\text{rk}_D(f) = \alpha$ by induction on α (it is well known that $\text{rk}_D(f) < \infty$): $\text{rk}_D(f) = \alpha$ if and only if $\beta < \alpha \Rightarrow \text{rk}_D(f) \neq \beta$ and for every $g \in {}^Y\text{Ord}$ satisfying $g <_D f$, there is $\beta < \alpha$ such that $\text{rk}_D(g) = \beta$.

Notice that we will use normal filters on $\kappa = \text{cf}(\kappa) > \aleph_0$, so the demand for \aleph_1 -completeness in the definition is satisfied.

Recall also the following definition.

Received by the editors May 15, 2006; revised February 20, 2007.

Research supported by the United States-Israel Binational Science Foundation. Publication 881

AMS subject classification: 03E20.

Keywords: set theory, partition calculus.

©Canadian Mathematical Society 2009.

Definition 1.3 Assume Y, D, f are as in Definition 1.2.

$$J[f, D] = \{Z \subseteq Y : Y \setminus Z \in D \text{ or } \text{rk}_{D+Z}(f) > \text{rk}_D(f)\}$$

Lastly, we quote the next claim (Definition 1.3 and Claim 1.4 are from [2], and explicitly [4, 5.8(2), 5.9]).

Claim 1.4 Assume $\kappa > \aleph_0$ is regular, and D is a κ -complete (resp. normal) filter on Y .

Then for any $f \in {}^Y \text{Ord}$, $J[f, D]$ is a κ -complete (resp. normal) ideal on Y disjoint to D .

2 The Proof

In this section we prove Theorem 1.1, which, for convenience, we now restate.

Theorem 2.1 If $\aleph_0 < \kappa = \text{cf}(\lambda)$, $2^\kappa < \lambda$ then $\lambda \rightarrow (\lambda, \omega + 1)^2$.

Proof

Stage A. Given that $\aleph_0 < \kappa = \text{cf}(\lambda) < \lambda$, $2^\kappa < \lambda$, we will show that $\lambda \rightarrow (\lambda, \omega + 1)^2$. So, towards a contradiction, suppose that

(i) $c: [\lambda]^2 \rightarrow \{\text{red}, \text{green}\}$ but has no red set of cardinality λ and no green set of order type $\omega + 1$.

Choose $\bar{\lambda}$ such that:

(ii) $\bar{\lambda} = \langle \lambda_i : i < \kappa \rangle$ is increasing and continuous with limit λ , and for $i = 0$ or i a successor ordinal, λ_i is a successor cardinal. We also let $\Delta_0 = \lambda_0$ and for $i < \kappa$, $\Delta_{1+i} = [\lambda_i, \lambda_{i+1})$. For $\alpha < \lambda$ we will let $\mathbf{i}(\alpha)$ be the unique $i < \kappa$ such that $\alpha \in \Delta_i$.

We can clearly assume, in addition, that

(iii) $\lambda_0 > 2^\kappa$, for $i < \kappa$, $\lambda_{i+1} \geq \lambda_i^{++}$, and each Δ_i is homogeneously red for c .

The last is justified by the Erdős–Hajnal–Maté–Rado theorem for λ_{i+1} , i.e., as $\lambda_{i+1} \rightarrow (\lambda_{i+1}, \omega + 1)^2$ because λ_{i+1} is regular.

Stage B. For $0 < i < \kappa$, we define Seq_i to be

$$\{\langle \alpha_0, \dots, \alpha_{n-1} \rangle : \mathbf{i}(\alpha_0) < \dots < \mathbf{i}(\alpha_{n-1}) < i\}.$$

For $\zeta \in \Delta_i$ and $\langle \alpha_0, \dots, \alpha_{n-1} \rangle = \bar{\alpha} \in \text{Seq}_i$, we say $\bar{\alpha} \in \mathcal{T}^\zeta$ if and only if $\{\alpha_0, \dots, \alpha_{n-1}, \zeta\}$ is homogeneously green for c . Note that an infinite \triangleleft -increasing branch in \mathcal{T}^ζ violates the non-existence of a green set of order type $\omega + 1$, so,

(iv) \mathcal{T}^ζ is well-founded, that is we cannot find $\eta_0 \triangleleft \eta_1 \triangleleft \dots \triangleleft \eta_n \triangleleft \dots$.

Therefore the following definition of a rank function, rk^ζ , on Seq_i can be carried out. If $\eta \in \text{Seq}_i \setminus \mathcal{T}^\zeta$ then $\text{rk}^\zeta(\eta) = -1$. We define $\text{rk}^\zeta: \text{Seq}_i \rightarrow \text{Ord} \cup \{-1\}$ by induction on the ordinal ξ as follows. We have $\text{rk}^\zeta(\bar{\alpha}) = \xi$ if and only if for all $\epsilon < \xi$, $\text{rk}^\zeta(\bar{\alpha})$ was not defined as ϵ but there is a β such that $\text{rk}^\zeta(\bar{\alpha} \widehat{\langle \beta \rangle}) \geq \epsilon$. Of course,

if ξ is a successor ordinal, it is enough to check for $\epsilon = \xi - 1$, and for a limit ordinal, δ , if for all $\xi < \delta$, $\text{rk}^\zeta(\bar{\alpha}) \geq \xi$, then $\text{rk}^\zeta(\bar{\alpha}) \geq \delta$. In fact, it is clear that the range of rk^ζ is a proper initial segment of μ_i^+ , where $\mu_i := \text{card}(\bigcup \{\Delta_\epsilon : \epsilon < i\})$, and so, in particular, the range of rk^ζ has cardinality at most λ_i . Note that $\lambda_{i+1} \geq \lambda_i^{++} > \mu_i^+$.

Now we can choose B_i , an end-segment of Δ_i such that for all $\bar{\alpha} \in \text{Seq}_i$ and all $-1 \leq \gamma < \mu_i^+$, if there is $\zeta \in B_i$ such that $\text{rk}^\zeta(\bar{\alpha}) = \gamma$, then there are λ_{i+1} such ζ . Recall that Δ_i and therefore also B_i are of order type λ_{i+1} , which is a successor cardinal $> \mu_i^+ > |\text{Seq}_i|$, hence such B_i exists. Everything is now in place for the main definition.

Stage C. $(\bar{\alpha}, Z, D, f) \in K$ if and only if

- (a) D is a normal filter on κ ,
- (b) $f: \kappa \rightarrow \text{Ord}$,
- (c) $Z \in D$
- (d) for some $0 < i < \kappa$ we have $\bar{\alpha} \in \text{Seq}_i$, Z is disjoint to $i + 1$ and for every $j \in Z$ (hence $j > i$) there is $\zeta \in B_j$ such that $\text{rk}^\zeta(\bar{\alpha}) = f(j)$ (so, in particular, $\bar{\alpha} \in \mathcal{T}^\zeta$).

Stage D. Note that $K \neq \emptyset$, since if we choose $\zeta_j \in B_j$, for $j < \kappa$, take $Z = \kappa \setminus \{0\}$, $\bar{\alpha} =$ the empty sequence, choose D to be any normal filter on κ and define f by $f(j) = \text{rk}^{\zeta_j}(\bar{\alpha})$, then $(\bar{\alpha}, Z, D, f) \in K$.

Now clearly by Definition 1.2, among the quadruples $(\bar{\alpha}, Z, D, f) \in K$, there is one with $\text{rk}_D(f)$ minimal. So, fix one such quadruple, and denote it by $(\bar{\alpha}^*, Z^*, D^*, f^*)$. Let D_1^* be the filter on κ dual to $J[f^*, D^*]$; so by Claim 1.4 it is a normal filter on κ extending D^* .

For $j \in Z^*$, set $C_j = \{\zeta \in B_j : \text{rk}^\zeta(\bar{\alpha}^*) = f^*(j)\}$. Thus by the choice of B_j we know that $\text{card}(C_j) = \lambda_{j+1}$, and for every $\zeta \in C_j$ the set $(\text{Rang}(\bar{\alpha}^*) \cup \{\zeta\})$ is homogeneously green under the colouring c . Now suppose $j \in Z^*$. For every $\Upsilon \in Z^* \setminus (j + 1)$ and $\zeta \in C_j$, let $C_\Upsilon^+(\zeta) = \{\xi \in C_\Upsilon : c(\{\zeta, \xi\}) = \text{green}\}$. Also, let $Z^+(\zeta) = \{\Upsilon \in Z^* \setminus (j + 1) : \text{card}(C_\Upsilon^+(\zeta)) = \lambda_{\Upsilon+1}\}$.

Stage E. For $j \in Z^*$ and $\zeta \in C_j$, let $Y(\zeta) = Z^* \setminus Z^+(\zeta)$. Since $\lambda_0 > 2^\kappa$ and $\lambda_{j+1} > \lambda_0$ is regular, for each $j \in Z^*$ there are $Y = Y_j \subseteq \kappa$ and $C'_j \subseteq C_j$ with $\text{card}(C'_j) = \lambda_{j+1}$ such that $\zeta \in C'_j \Rightarrow Y(\zeta) = Y_j$.

Let $\hat{Z} = \{j \in Z^* : Y_j \in D_1^*\}$. Now the proof splits into two cases.

Case 1. $\hat{Z} \neq \emptyset \pmod{D_1^*}$

Define $Y^* = \{j \in \hat{Z} : \text{for every } i \in \hat{Z} \cap j, \text{ we have } j \in Y_i\}$. Notice that Y^* is the intersection of \hat{Z} with the diagonal intersection of κ sets from D_1^* (since $i \in \hat{Z} \Rightarrow Y_i \in D_1^*$), hence (by the normality of D_1^*) $Y^* \neq \emptyset \pmod{D_1^*}$. But then, as we will see soon, by shrinking the C'_j for $j \in Y^*$, we can get a homogeneous red set of cardinality λ , which is contrary to the assumption toward contradiction.

We define \hat{C}_j for $j \in Y^*$ by induction on j such that \hat{C}_j is a subset of C'_j of cardinality λ_{j+1} . Now, for $j \in Y^*$, let \hat{C}_j be the set of $\xi \in C'_j$ such that for every $i \in Y^* \cap j$ and every $\zeta \in \hat{C}_i$ we have $\xi \notin C_\zeta^+(\zeta)$. So, in fact, \hat{C}_j has cardinality λ_{j+1} , as it is the result of removing $< \lambda_{j+1}$ elements from C'_j where $|C'_j| = \lambda_{j+1}$ by its choice. Indeed, the number of such pairs (i, ζ) is $\leq \lambda_j$ and for $i \in Y^* \cap j$ and $\zeta \in \hat{C}_i$

- (a) $j \in Y_i$ [by the definition of Y^* as $j \in Y^*$].

- (b) $\zeta \in C'_i$ [as $\zeta \in \hat{C}_i$ and $\hat{C}_i \subseteq C'_i$ by the induction hypothesis].
- (c) $Y(\zeta) = Y_i$ [as by (b) we have $\zeta \in C'_i$ and the choice of C'_i].
- (d) $j \in Y(\zeta)$ [by (a)+(c)].
- (e) $j \notin Z^+(\zeta)$ [by (d) and the choice of $Y(\zeta)$ as $Z^* \setminus Z^+(\zeta)$].
- (f) $C_j^+(\zeta)$ has cardinality $< \lambda_{j+1}$ [by (e) and the choice of $Z^+(\zeta)$, as $j \in \hat{Z} \subseteq Z^*$].

So \hat{C}_j is a well defined subset of C'_j of cardinality λ_{j+1} for every $j \in Y^*$. But then, clearly the union of the \hat{C}_j for $j \in Y^*$, call it \hat{C} , satisfies the following.

- (a) it has cardinality λ [as $j \in Y^* \Rightarrow |\hat{C}_j| = \lambda_{j+1}$ and $\sup(Y^*) = \kappa$ as $Y^* \neq \emptyset \text{ mod } D_1^*$],
- (b) $c \upharpoonright [\hat{C}_j]^2$ is constantly red [as we are assuming (iii)],
- (c) if $i < j$ are from Y^* and $\zeta \in \hat{C}_i, \xi \in \hat{C}_j$ then $c\{\zeta, \xi\} = \text{red}$ [as $\xi \notin C_j^+(\zeta)$].

So \hat{C} has cardinality λ and is homogeneously red. This concludes the proof of Case 1.

Case 2. $\hat{Z} = \emptyset \text{ mod } D_1^*$.

In this case there are $i \in Z^*, \beta \in C_i$ such that $Z^+(\beta) \neq \emptyset \text{ mod } D_1^*$
 [Because $Z^* \in D^* \subseteq D_1^*$ and $\hat{Z} = \emptyset \text{ mod } D_1^*$, hence $Z^* \setminus \hat{Z} \neq \emptyset$. Choose $i \in Z^* \setminus \hat{Z}$. By the definition of $\hat{Z}, Y_i \notin D_1^*$. So, if $\beta \in C'_i$ then $Y(\beta) = Y_i \notin D_1^*$ and choose $\beta \in C'_i$, so $Y(\beta) \notin D_1^*$ hence by the definition of $Y(\beta)$ we have $Z^* \setminus Z^+(\beta) = Y(\beta) \notin D_1^*$. Since $Z^* \in D_1^*$, we conclude that $Z^+(\beta) \neq \emptyset \text{ mod } D_1^*$].

Let $\bar{\alpha}' = \bar{\alpha}^* \frown \langle \beta \rangle, Z' = Z^+(\beta), D' = D^* + Z'$. It is a normal filter Claim 1.4, and by the previous sentence which makes sure that $Z' \neq \emptyset$, as $D^* \subseteq D_1^*$. Lastly we define $f' \in {}^\kappa \text{Ord}$ by

- (a) if $j \in Z'$ then $f'(j) = \text{Min}\{\text{rk}^\gamma(\bar{\alpha}') : \gamma \in C_j^+(\beta) \subseteq B_j\}$,
- (b) otherwise $f'(j) = 0$.

Clearly

- (a) $(\bar{\alpha}', Z', D', f') \in K$, and
- (b) $f' <_{D'} f^*$

[Because, as $Z' \in D'$ and if $j \in Z'$ then for some $\gamma \in C_j^+(\beta)$ we have $f'(j) = \text{rk}^\gamma(\bar{\alpha}') = \text{rk}^\gamma(\bar{\alpha}^* \frown \langle \beta \rangle)$ which by the definition of rk^γ is $< \text{rk}^\gamma(\bar{\alpha}^*) = f^*(j)$, recalling (d) from Stage C.]

hence

- (c) $\text{rk}_{D'}(f') < \text{rk}_{D'}(f^*)$
 [By Definition 1.2].

But $\text{rk}_{D'}(f^*) = \text{rk}_{D^*}(f^*)$ as $Z' = Z^+(\beta) \neq \emptyset \text{ mod } D_1^*$ by the definition of D_1^* as extending the filter dual to $J[f^*, D^*]$, see Definition 1.3. Hence $\text{rk}_{D'}(f') < \text{rk}_{D^*}(f^*)$, so we get a contradiction to the choice of $(\bar{\alpha}^*, Z^*, D^*, f^*)$.

Clearly at least one of the two cases holds, so we are done. ■

References

- [1] P. Erdős, A. Hajnal, A. Maté, and R. Rado, *Combinatorial set theory: partition relations for cardinals*. Studies in Logic and the Foundations of Mathematics 106, North Holland Publishing Co, Amsterdam, 1984.
- [2] S. Shelah, *A note on cardinal exponentiation*. J. Symbolic Logic 45(1980), no. 1, 56–66.
- [3] ———, *Cardinal Arithmetic*. Oxford Logic Guides 29, Oxford University Press, New York, 1994.

- [4] ———, *Applications of PCF theory*. J. Symbolic Logic **65**(2000), no. 4, 1624–1674.
- [5] S. Shelah and L. Stanley, *Filters, Cohen sets and consistent extensions of the Erdős–Dushnik–Miller theorem*. J. Symbolic Logic **65**(2000), no. 1, 259–271.

Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel, and Department of Mathematics, Rutgers University, New Brunswick, NJ 08854, USA
e-mail: shelah@math.huji.ac.il