

L^2 -BOUNDEDNESS OF THE CAUCHY TRANSFORM ON SMOOTH NON-LIPSCHITZ CURVES

HYEONBAE KANG* AND JIN KEUN SEO*

1. Introduction and statements of results

Let Γ be a curve defined by $y = A(x)$ in \mathbf{R}^2 . The Cauchy transform \mathcal{C}_A on the curve Γ is a singular integral operator defined by the singular integral kernel

$$(1.1) \quad K(x, y) = \frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))}.$$

If A is a Lipschitz function, i.e., $\|A'\|_\infty < \infty$, then \mathcal{C}_A makes a very significant example of non-convolution type singular integral operators. The problem of L^2 -boundedness of the Cauchy transform was raised and solved when $\|A'\|_\infty$ is small by A. P. Calderón in relation to the Dirichlet problem on Lipschitz domains [Cal1, Cal2]. Since then, it has been a central problem in the theory of singular integral operators and several significant techniques has been developed to deal with this problem. Among them are the $T(1)$ -Theorem of David and Journé, the technique of Coifman, McIntosh, and Meyer, and the technique of Coifman, Jones, and Semmes [D.J, C.M.M, C.J.S]. We refer to [Chi, Mur] for a history of development in the last decades on the theory of the Cauchy transform.

If $\|A'\|_\infty = \infty$, then the Cauchy kernel $K(x, y)$ given in (1.1) is not a standard kernel. An integral kernel on the line is called a standard kernel if it satisfies $|K(x, y)| \leq C|x - y|^{-1}$ and $|\nabla_{x,y} K(x, y)| \leq C|x - y|^{-2}$. If $\|A'\|_\infty = \infty$, then the Cauchy kernel does not satisfy both estimates. So, the theory of the singular integral operators may not be applied directly. Nevertheless, the question of L^2 -boundedness of \mathcal{C}_A is still an interesting one. In this paper, we deal with L^2 -boundedness of \mathcal{C}_A when A is smooth and $\|A'\|_\infty = \infty$.

Received June 22, 1992.

*Both authors are partially supported by GARC-KOSEF and NON DIRECTED RESEARCH FUND, Korea Research Foundation, 1992.

We first find two examples of curves on which the Cauchy transforms are not L^2 -bounded. Those are curves defined by $A'(x) = x \sin x$ and $A(x) = \exp(x^2)$. In the first example, A' has too many zeros while the derivative of the second A grows too fast relatively to A . The fact that both $\log |x \sin x|$ and $\log \exp x^2$ are *not* BMO functions is relevant. We then consider the case when A is a polynomial. If A is a polynomial, then A' has only finitely many zeros and $|A'(x)/A(x)|$ behaves like $|1/x|$ as $x \rightarrow \infty$. In fact, if A is a polynomial, then $\log |A(x)|$ is a BMO function. In this paper, we prove the following theorem.

MAIN THEOREM. *Let $A(x)$ be a polynomial of the form*

$$(1.2) \quad \begin{cases} A(x) \text{ is any polynomial if } d \text{ is an odd integer,} \\ A(x) = \sum_{i=1}^n a_i x^{2i} \text{ if } d = 2n \text{ is an even integer.} \end{cases}$$

Then, the Cauchy transform \mathcal{C}_A is bounded on L^p for any $1 < p < \infty$.

Among the polynomial which are not covered in (1.2) is $A(x) = x^4 - x^3$. This polynomial does not satisfy the estimate $|A(x) - A(y)| \approx |x - y| |x + y| (|x|^2 + |y|^2)$ when $|x| + |y|$ is large which is a crucial estimate for our proofs. However, polynomials in (1.2) include a significantly large class of polynomials.

In order to explain ideas of proofs in this paper, let us consider an example. If $A(x) = x^2$, the kernel given in (1.1) does not satisfy the standard estimates. But, the kernel can be decomposed as

$$(1.3) \quad K(x, y) = \frac{1 + 2iy}{(x - y) + i(x^2 - y^2)} = \frac{1}{x - y} + \frac{-i}{1 + (x + y)}.$$

The first kernel of the right hand side is the Hilbert kernel while the second one is a kernel of Poisson type. So, if $A(x) = x^2$, then \mathcal{C}_A is bounded on L^2 . It turns out this decomposition can be performed for general kernels by using a proper cut-off function. Then, each one of the decomposed kernels is a standard kernel and we can apply the $T(1)$ -Theorem to it.

This paper is organized as follows; In section 2, we give a sufficient condition for a function to belong to the BMO. In section 3, we collect some estimates on polynomials which will be used in later sections. In section 4, we decompose the kernel $K(x, y)$ into two standard kernels and show that both of them satisfy all the conditions of $T(1)$ -Theorem. In section 5, we show that if $A'(x) = x \sin x$ or $A(x) = \exp(x^2)$, then \mathcal{C}_A is *not* bounded on L^2 .

Throughout this paper constants C may differ in each occurrence and $A \approx B$

means that there are positive constants c and C such that $c \leq A/B \leq C$.

2. Preliminary lemma on BMO

Showing that a function is in BMO is a fairly hard task. One of the reasons is that being a BMO function is not just a size condition. For example, even if $|f| \in \text{BMO}$, f may not be a BMO function. It can also be shown easily that even if $0 \leq f \leq g$ and $g \in \text{BMO}$, f may not be a BMO function. In particular, that $f(x) = O(\log|x|)$ as $x \rightarrow \infty$ does not imply $f \in \text{BMO}$. In this section we obtain a sufficient condition for a function to belong to BMO which will be used repeatedly in section 4. We show that if $f'(x) = O(|x|^{-1})$ as $x \rightarrow \infty$, then $f \in \text{BMO}$.

LEMMA 2.1. *Suppose that there exists a positive number m such that f is bounded on $[-m, m]$ and f is continuously differentiable if $|x| \geq m$. If $|f'(x)| = O(|x|^{-1})$ as $x \rightarrow \infty$, then $f \in \text{BMO}$.*

Proof. By the assumption, there are large constants L and C such that $|f'(x)| \leq C|x|^{-1}$ if $|x| > L$. We write

$$f = f\chi_{(-\infty, -L)} + f\chi_{[-L, L]} + f\chi_{(L, \infty)} = f_1 + f_2 + f_3.$$

It is enough to show that $f_3 \in \text{BMO}$ since f_2 is bounded and that $f_1 \in \text{BMO}$ can be proved in the same way. For notational simplicity, we put $g = f_3$. We need to show that if $0 < a < b$, then

$$\frac{1}{b-a} \int_a^b |g(x) - g(b)| dx \leq C.$$

We may assume $L < a < b$. If $b \leq 2a$, then

$$\frac{1}{b-a} \int_a^b |g(x) - g(b)| dx \leq C \int_a^b |g'(x)| dx \leq C.$$

Suppose that $b \geq 2a$. Choose an integer N so that $2^{-N}b \leq a \leq 2^{-N+1}b$. Then,

$$\frac{1}{b-a} \int_a^b |g(x) - g(b)| dx \leq \frac{1}{b-a} \sum_{j=1}^N \int_{2^{-j}b}^{2^{-j+1}b} |g(x) - g(b)| dx.$$

And we have, for each j ,

$$\int_{2^{-j}b}^{2^{-j+1}b} |g(x) - g(b)| dx$$

$$\begin{aligned}
&\leq C \int_{2^{-j}b}^{2^{-j+1}b} |g(x) - g(2^{-j}b)| dx + 2^{-j}b |g(b) - g(2^{-j}b)| \\
&\leq C 2^{-j}b \int_{2^{-j}b}^{2^{-j+1}b} |g'(x)| dx + 2^{-j}b \sum_{j=1}^j |g(2^{-i+1}b) - g(2^{-j}b)| \\
&\leq C 2^{-j}b(\log 2 + j).
\end{aligned}$$

Therefore we obtain

$$\begin{aligned}
\frac{1}{b-a} \int_a^b |g(x) - g(b)| dx &\leq C \frac{1}{b-a} \sum_{j=1}^N 2^{-j} b(\log 2 + j) \\
&\leq C \frac{b}{b-a} \leq C.
\end{aligned}$$

This completes the proof.

3. Estimates on polynomials

In this section we collect estimates on polynomials which will be used in later sections. Let $A(x)$ be d -th degree polynomial of the form:

$$(3.1) \quad \begin{cases} A(x) \text{ is any polynomial if } d \text{ is an odd integer} \\ A(x) = \sum_{i=1}^n a_i x^{2i} \text{ if } d = 2n \text{ is an even integer.} \end{cases}$$

For these polynomials we have the following elementary but significant estimates.

LEMMA 3.1. *Let $A(x)$ be a polynomial of degree d as in (3.1). Then,*

(1) *If d is odd, then*

$$|A(x) - A(y)| \approx |x - y| (|x|^{d-1} + |y|^{d-1}).$$

Moreover, there exists a positive number M such that

(2) *If $|x| \geq M$, then $|A(x)| \approx |x|^d$, $|A'(x)| \approx |x|^{d-1}$, and $|A''(x)| \approx |x|^{d-2}$.*

(3) *If d is even and if either $|x| \geq M$ or $|y| \geq M$, then*

$$|A(x) - A(y)| \approx |x - y| |x + y| (|x|^{d-2} + |y|^{d-2}).$$

(4) *If $|x| < M$ and $|y| > 2M$, then*

$$|A(x) - A(y)| \approx |A(y)| \approx |y|^d.$$

Remark 3.2 We will fix M to be the number as in Lemma 3.1 throughout this paper.

Proof. That $|A(x) - A(y)| \leq C|x - y|(|x|^{d-1} + |y|^{d-1})$ is trivial for any polynomial. For (1), note that

$$\begin{aligned} \frac{x^d - y^d}{x - y} &= \sum_{j=0}^{d-1} x^{d-j-1}y^j \\ &= \frac{1}{2}x^{d-1} + \frac{1}{2}y^{d-1} + \frac{1}{2}(x^2 + 2xy + y^2) \sum_{j=1}^{(d-1)/2} x^{d-(2j+1)}y^{2(j-1)} \\ &\geq \frac{1}{2}(x^{d-1} + y^{d-1}) \end{aligned}$$

since d is odd. Therefore we have

$$|A(x) - A(y)| \geq \frac{1}{2}|x - y|(|x|^{d-1} + |y|^{d-1}).$$

Let $A(x) = \sum_{j=1}^d a_j x^j$. Assume that $a_d = 1$ without loss of generality. Then, we have

$$\begin{aligned} |A(x) - A(y)| &\geq |x^d - y^d| - \sum_{j=1}^{d-1} |a_j| |x^j - y^j| \\ &\geq \frac{1}{2}|x - y|(|x|^{d-1} + |y|^{d-1}) - C|x - y|(|x|^{d-2} + |y|^{d-2}) \\ &\geq C|x - y|(|x|^{d-1} + |y|^{d-1}). \end{aligned}$$

For (3), we let $A(x) = \sum_{n=1}^r a_n x^{2n}$ with $2r = d$. Observe that

$$\begin{aligned} |x^{2n} - y^{2n}| &= |x - y||x + y||x^{2n-2} + x^{2n-4}y^2 + \dots + x^2y^{2n-4} + y^{2n-2}| \\ &\approx |x - y||x + y|(|x|^{2n-2} + |y|^{2n-2}) \end{aligned}$$

by the same reason as before. Therefore, there exist constants C_1 and C_2 such that

$$\begin{aligned} |A(x) - A(y)| &= \left| \sum_{n=1}^r a_n(x^{2n} - y^{2n}) \right| \\ &\geq |x - y||x + y|[C_1(|x|^{2n-2} + |y|^{2n-2}) - C_2(|x|^{2n-4} + |y|^{2n-4})] \\ &\geq C|x - y||x + y|(|x|^{2n-2} + |y|^{2n-2}) \end{aligned}$$

as long as $|x| + |y|$ is large. It is easy to show that

$$|A(x) - A(y)| \leq C|x - y||x + y|(|x|^{2n-2} + |y|^{2n-2})$$

and hence we obtain (3). (2) and (4) are trivial. This completes the proof.

Remark 3.2. The polynomial $A(x) = x^4 - x^3$ is not included in (3.1). The estimate (3) in Lemma 3.1 is actually false for $A(x) = x^4 - x^3$. It would be interesting to see whether the Cauchy transform on the curve $y = x^4 - x^3$ is L^2 -bounded or not.

4. L^2 boundedness

Throughout this paper \mathcal{C}_A denotes the Cauchy transform on the curve $y = A(x)$. This section is devoted to the proof of Main Theorem:

MAIN THEOREM. *Let $A(x)$ be a polynomial of the form*

$$(4.1.1) \quad \begin{cases} A(x) \text{ is any polynomial if } p \text{ is an odd integer} \\ A(x) = \sum_{j=1}^n a_j x^{2j} \text{ if } p = 2n \text{ is an even integer.} \end{cases}$$

Then, the Cauchy transform \mathcal{C}_A is bounded on L^p for any $1 < p < \infty$.

By the classical theory of singular integral operators, it suffices to prove when $p = 2$. Recall that the integral kernel $K(x, y)$ of \mathcal{C}_A is given by

$$(4.1.2) \quad K(x, y) = \frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))}.$$

The kernel $K(x, y)$ does not satisfy the standard estimates. If $|y|$ is large and if x and y are close, then the estimate $K(x, y) \leq C|x - y|^{-1}$ does not hold. We overcome this obstacle by decomposing the kernel into two standard kernels by introducing an appropriate cut-off function.

Let $\tilde{\phi}$ be a C^∞ smooth function such that

$$(4.1.3) \quad \begin{cases} \tilde{\phi}(x) = 1 & \text{if } |x| < \frac{1}{2} \\ \tilde{\phi}(x) = 0 & \text{if } |x| \geq \frac{4}{5} \\ \|\tilde{\phi}'\|_{L^\infty} \leq 10 \end{cases}$$

and we let

$$(4.1.4) \quad \phi(x, y) = \tilde{\phi}\left(\frac{x - y}{1 + |x|}\right).$$

Let $K(x, y)$ be the Cauchy kernel as given in (4.1.1). We define K_1 and K_2 by $K_1(x, y) = K(x, y)\phi(x, y)$ and $K_2(x, y) = K(x, y)(1 - \phi(x, y))$. We denote by \mathcal{C}_1 and \mathcal{C}_2 the integral operators defined by K_1 and K_2 respectively. We show that both \mathcal{C}_1 and \mathcal{C}_2 are bounded on L^2 in the following subsections by using $T(1)$ -theorem. Each subsection corresponds to a condition of $T(1)$ -Theorem. Since $\mathcal{C}_A = \mathcal{C}_1 + \mathcal{C}_2$, Main Theorem follows. We now recall the Weak boundedness property and $T(1)$ -Theorem of David and Journé:

$T(1)$ -THEOREM (David and Journé). *Let T be the integral operator defined by*

$$\langle Tf, g \rangle = \int_{\mathbf{R}} \int_{\mathbf{R}} K(x, y) f(x) g(y) \, dx dy$$

for any bounded functions with compact supports f and g such that $\text{supp}(f) \cap \text{supp}(g) = \emptyset$. Suppose that an integral kernel $K(x, y)$ satisfies

(1) *Standard Estimates:*

$$|K(x, y)| \leq C \frac{1}{|x - y|}$$

$$|\nabla_x K(x, y)| + |\nabla_y K(x, y)| \leq C \frac{1}{|x - y|^2}$$

for all $x \neq y \in \mathbf{R}$.

(2) *Weak Boundedness Property:* there exist constants N and C such that for any pair of functions φ and ψ in $C_0^\infty(\mathbf{R})$ satisfying $\varphi(x) = \psi(x) = 0$ if $|x| > 1$ and $\|\varphi\|_{C^N} \leq 1$ and $\|\psi\|_{C^N} \leq 1$, for any $x \in \mathbf{R}$ and $t > 0$,

$$|\langle T\varphi^{x,t}, \psi^{x,t} \rangle| \leq Ct$$

where $\varphi^{x,t}(y) = \varphi\left(\frac{x + y}{t}\right)$.

(3) $T1 \in BMO$.

(4) $T^*1 \in BMO$.

Then T can be extended as an operator bounded on $L^2(\mathbf{R})$.

For notational convenience we put, throughout this paper,

$$(4.1.5) \quad Q(x, y) = \frac{A(x) - A(y)}{x - y}.$$

4.1. Standard estimates

In this subsection we show that the decomposed kernels K_1 and K_2 satisfy the standard estimates. We remark that if A is a polynomial of odd degree, there is no need of decomposing the kernel, namely, the Cauchy kernel $K(x, y)$ itself satisfy the standard estimates. However, we decompose the kernel even in this case since we want to deal with all the polynomials one time.

PROPOSITION 4.1.1. $K_1(x, y)$ satisfies the standard estimates: there exists a positive constant C such that

$$(4.1.6) \quad |K_1(x, y)| \leq C \frac{1}{|x - y|}$$

$$(4.1.7) \quad |\nabla_x K_1(x, y)| + |\nabla_y K_1(x, y)| \leq C \frac{1}{|x - y|^2}.$$

Proof. Note that $K_1(x, y) = 0$ if $|x - y| > \frac{4}{5}(1 + |x|)$. Note also that $\nabla_{x,y}\phi(x, y) \neq 0$ only if $\frac{1}{2}(1 + |x|) \leq |x - y| \leq \frac{4}{5}(1 + |x|)$. Hence

$$(4.1.8) \quad |\nabla_{x,y}\phi(x, y)| \leq C \frac{1}{|x - y|}.$$

Let M be the number in Lemma 3.1. If $|x| > 5M$ and if $|x - y| \leq \frac{4}{5}(1 + |x|)$, then $|y| > M$, x and y have the same signs, and $|x| \approx |y|$. It then follows from Lemma 3.1 that

$$|Q(x, y)| \approx |x|^{d-1} \approx |A'(y)|$$

and

$$|\nabla_{x,y}Q(x, y)| \leq \frac{(|A'(x)| + |A'(y)|)|x - y| + |A(x) - A(y)|}{|x - y|^2} \leq C \frac{|y|^{d-1}}{|x - y|}.$$

Therefore,

$$\begin{aligned} \left| \nabla_{x,y} \left(\frac{1 + iA'(y)}{1 + iQ(x, y)} \right) \right| &\leq \frac{|A'(y)| (1 + |Q(x, y)|^2)^{1/2}}{1 + |Q(x, y)|^2} \\ &\quad + \frac{(1 + |A'(y)|^2)^{1/2} |\nabla_{x,y}(Q(x, y))|}{1 + |Q(x, y)|^2} \end{aligned}$$

$$\leq C \frac{1}{|x - y|}.$$

Combining all these estimates we have

$$|K_1(x, y)| = \left| \frac{1}{|x - y|} \frac{1 + iA'(y)}{1 + iQ(x, y)} \phi(x, y) \right| \leq C \frac{1}{|x - y|}$$

and

$$|\nabla_{x,y} K_1(x, y)| = \left| \nabla_{x,y} \left(\frac{1}{|x - y|} \frac{1 + iA'(y)}{1 + iQ(x, y)} \phi(x, y) \right) \right| \leq C \frac{1}{|x - y|^2}$$

provided that $|x| > 5M$.

On the other hand, if $|x| \leq 5M$ and $|x - y| \leq \frac{4}{5}(1 + |x|)$, then $|y| \leq 10M$ and hence $|A'(y)|$ is bounded. So, it is easy to derive the estimates (4.1.6) and (4.1.7) by using (4.1.8). This completes the proof.

PROPOSITION 4.1.2. $K_2(x, y)$ satisfies the following estimates: there exists a constant C such that

$$(4.1.9) \quad |K_2(x, y)| \leq C \frac{1}{|x + y|}$$

$$(4.1.10) \quad |\nabla_x K_2(x, y)| + |\nabla_y K_2(x, y)| \leq C \frac{1}{|x + y|^2}.$$

Moreover, if either $|x| \leq M$ or $|y| \leq M$ where M is the number in Lemma 3.1, then $|K_2(x, y)| + |\nabla_{x,y} K_2(x, y)|$ is bounded.

Remark 3.5. We note that (4.1.9) and (4.1.10) are not standard estimates. But $K_2(x, -y)$ satisfies the standard estimates and we can apply $T(1)$ -Theorem to $K_2(x, -y)$.

Proof. Since $K_2(x, y) = 0$ if $|x - y| \leq 1/2(|x| + 1)$, we assume that $|x - y| > 1/2(|x| + 1)$. Then, by the triangular inequality, we have

$$(4.1.11) \quad 1 + |x| + |y| \leq 4|x - y|.$$

Let M be the number in Lemma 3.1. We first deal with the easier case. Let A be a polynomial of odd degree. If either $|x| > M$ or $|y| > M$, then by Lemma 3.1 (1) and (4.1.8),

$$\begin{aligned}
 (4.1.12) \quad |K_2(x, y)| &\leq \left| \frac{1 + iA'(y)}{A(x) - A(y)} \right| \\
 &\leq C \frac{|y|^{d-1} + 1}{|x - y| (|x|^{d-1} + |y|^{d-1})} \\
 &\leq C \frac{1}{|x - y|} \leq C \frac{1}{|x + y| + 1}.
 \end{aligned}$$

If $|x| \leq M$ and $|y| \leq M$, then it is easy to get $|K_2(x, y)| \leq C$. (4.1.10) can be proved in the same way.

We now suppose that $A(x) = \sum_{k=1}^n a_k x^{2k}$. Recall that $|x - y| > 1/2(1 + |x|)$. If $|y| > M$, then by Lemma 3.1 (2) and (3) and (4.1.11),

$$\begin{aligned}
 (4.1.13) \quad |K_2(x, y)| &\leq \left| \frac{1 + iA'(y)}{A(x) - A(y)} \right| \\
 &\leq C \frac{|y|^{d-1} + 1}{|x - y| |x + y| (|x|^{d-2} + |y|^{d-2})} \leq C \frac{1}{|x + y|}.
 \end{aligned}$$

If $|y| < M$, then since $|x - y| > 1/2(1 + |x|)$, we have

$$\begin{aligned}
 (4.1.14) \quad |K_2(x, y)| &\leq \left| \frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))} \right| \\
 &\leq C \frac{M^{d-1}}{|x - y|} \leq C \frac{1}{|x + y| + 1}.
 \end{aligned}$$

In order to derive (4.1.10), we first observe as in the proof of Proposition 4.1.1 that $\nabla_{x,y}\phi(x, y) \neq 0$ only if $1/2(1 + |x|) \leq |x - y| \leq 4/5(1 + |x|)$ and hence

$$|\nabla_{x,y}\phi(x, y)| \leq C \frac{1}{1 + |x|} \leq C \frac{1}{|x + y| + 1}.$$

If $|y| > M$, then by Lemma 3.1 and (4.1.11)

$$\begin{aligned}
 (4.1.15) \quad &\left| \nabla_{x,y} \left(\frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))} \right) \right| \\
 &\leq C \frac{(1 + |A(x)|) |y|^{d-1} + |y|^{2d-2}}{|x - y|^2 |x + y|^2 (|x|^{d-2} + |y|^{d-2})^2} \leq C \frac{1}{|x + y|^2}.
 \end{aligned}$$

Hence

$$(4.1.16) \quad |\nabla_{x,y}K_2(x, y)| = \left| \nabla_{x,y} \left(\frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))} (1 - \phi(x, y)) \right) \right|$$

$$\leq C \frac{1}{|x + y|^2}.$$

If $|y| \leq M$, $|A'(y)|$ is bounded and it is easy to derive

$$(4.1.17) \quad |\nabla_{x,y}K_2(x, y)| \leq C \frac{1}{|x - y|^2 + 1} \leq C \frac{1}{|x + y|^2}.$$

Combining estimates (4.1.12)-(4.1.16), we can derive (4.1.9) and (4.1.10). One also can see that if either $|x| \leq M$ or $|y| \leq M$, then $|K_2(x, y)| + |\nabla_{x,y}K_2(x, y)|$ is bounded. This completes the proof.

4.2. Weak boundedness

We now show that the operators \mathcal{C}_1 and \mathcal{C}_2 satisfy the weak boundedness property. We first show that \mathcal{C}_A itself is weakly bounded and then show how the weak boundedness of \mathcal{C}_1 and \mathcal{C}_2 follows.

PROPOSITION 4.2.1 (Weak Boundedness). *Let $A(x)$ be a polynomial of the form (4.1.1). There exists a constant $C > 0$ such that*

$$|\langle \mathcal{C}_A \phi_1^{u,t}, \phi_2^{u,t} \rangle| \leq Ct$$

for any $u \in \mathbf{R}$, for any $t > 0$, and for any $\phi_i \in C_0^\infty$ supported in $\{x \in \mathbf{R} : |x| \leq 1\}$ and $\|\phi_i\|_{C^1} \leq 1$. Here $\phi^{u,t}(x) = \phi\left(\frac{x + u}{t}\right)$.

Proof. Let $u = tv$ and let $\phi^v(x) = \phi(x + v)$. By a change of variables, we have

$$\begin{aligned} \langle \mathcal{C}_A \phi_1^{u,t}, \phi_2^{u,t} \rangle &= \lim_{\epsilon \rightarrow 0} \int \int_{|x-y| > \epsilon} K(x, y) \phi_1^{u,t}(x) \phi_2^{u,t}(y) dx dy \\ &= t \lim_{\epsilon \rightarrow 0} \int \int_{|x-y| > \epsilon} \frac{1}{x - y} \left(\frac{1 + iA'(ty)}{1 + iQ(tx, ty)} \right) \phi_1^v(x) \phi_2^v(y) dx dy. \end{aligned}$$

We then write

$$\begin{aligned} (4.2.1) \quad \langle \mathcal{C}_A \phi_1^{u,t}, \phi_2^{u,t} \rangle &= t \lim_{\epsilon \rightarrow 0} \int \int_{|x-y| > \epsilon} \frac{1}{x - y} \phi_1^v(x) \phi_2^v(y) dx dy \\ &\quad - it \lim_{\epsilon \rightarrow 0} \int \int_{|x-y| > \epsilon} \frac{tP(tx, ty)}{1 + iQ(tx, ty)} \phi_1^v(x) \phi_2^v(y) dx dy \\ &:= t(I_1(v) - iI_2(v, t)) \end{aligned}$$

where we put

$$(4.2.2) \quad P(x, y) = \frac{A'(y) - Q(x, y)}{y - x}.$$

Hence it suffices to show that I_1 and I_2 are bounded uniformly in v and t . Note that $I_1(v) = \langle H\phi_1^v, \phi_2^v \rangle$ where H is the Hilbert transform and hence I_1 is bounded. So, it remains to show that I_2 is bounded. By using the polar coordinates, we have

$$(4.2.3) \quad I_2(v, t) = \int_0^\infty \int_{-\pi}^\pi \frac{tr P(tr, \theta)}{1 + iQ(tr, \theta)} \phi(v, r, \theta) d\theta dr$$

where

$$Q(r, \theta) = Q(r \cos \theta, r \sin \theta)$$

$$P(r, \theta) = P(r \cos \theta, r \sin \theta)$$

$$\phi(v, r, \theta) = \phi_1^v(r \cos \theta) \phi_2^v(r \sin \theta).$$

Note that $\phi_1^v(x) \phi_2^v(y) \neq 0$ only if $|x + v| \leq 1$ and $|y + v| \leq 1$. Therefore, the set $\{r \in [0, \infty) : \phi(v, r, \theta) \neq 0\}$ is included in the interval $[|v| - 2, |v| + 2]$. In particular, $|\{r \in [0, \infty) : \phi(v, r, \theta) \neq 0\}| \leq 4$. Therefore, in order to prove that I_2 is bounded, it suffices to show that

$$(4.2.4) \quad F(v, s, r) := \int_{-\pi}^\pi \frac{sP(s, \theta)}{1 + iQ(s, \theta)} \phi(v, r, \theta) d\theta$$

is bounded.

Let $A(x) = \sum_{j=1}^d a_j x^j$ be a polynomial of the form (4.1.1). If we put

$$Q_j(x, y) = \frac{x^j - y^j}{x - y} = \sum_{k=1}^j x^{j-k} y^{k-1} \quad \text{for } j \geq 1,$$

then

$$Q(x, y) = \frac{A(x) - A(y)}{x - y} = \sum_{j=1}^d a_j Q_j(x, y).$$

Since

$$P(x, y) = \frac{A'(y) - Q(x, y)}{y - x} = \sum_{j=2}^d a_j \left(\sum_{k=1}^{j-1} y^{k-1} Q_{j-k}(x, y) \right),$$

one can see that

$$(4.2.5) \quad |P(s, \theta)| \leq Cs^{d-2} \quad \text{and} \quad \left| \frac{\partial P(s, \theta)}{\partial \theta} \right| \leq Cs^{d-2}.$$

Let $Q_j(\theta) = Q_j(\cos \theta, \sin \theta)$. Then, by the homogeneity, we have

$$(4.2.6) \quad F(v, s, r) = \int_{-\pi}^{\pi} \frac{sP(s, \theta)}{1 + i \sum_{j=1}^d s^{j-1} a_j Q_j(\theta)} \phi(v, r, \theta) d\theta.$$

Now suppose that d is an odd integer. Then,

$$Q_d(x, y) = 2^{-1}[x^{d-1} + x^{d-3}(x+y)^2 + \cdots + y^{d-3}(x+y)^2 + y^{d-1}]$$

and hence we have

$$(4.2.7) \quad Q_d(\theta) \geq 2^{-1} [(\cos \theta)^{d-1} + (\sin \theta)^{d-1}] \geq 2^{-d}.$$

Since $|s^{d-1} Q_d(\theta)| \geq 1/2 |\sum_{j=1}^{d-1} s^{j-1} a_j Q_j(\theta)|$ for all large s by (4.2.7), it follows from (4.2.5) that

$$\left| \frac{sP(s, \theta)}{1 + i \sum_{j=1}^d s^{j-1} a_j Q_j(\theta)} \right| \leq C.$$

It also follows that $F(v, s, r)$ is bounded.

We now deal with the case when d is even. Let $2n = d$ and $A(x) = \sum_{j=1}^n a_j x^{2j}$. Then, we have

$$(4.2.8) \quad F(v, s, r) = \int_{-\pi}^{\pi} \frac{sP(s, \theta)}{1 + i \sum_{j=1}^n s^{2j-1} a_j Q_{2j}(\theta)} \phi(v, r, \theta) d\theta.$$

Using the identity $\sin \theta + \cos \theta = \sqrt{2} \sin(\theta + \frac{\pi}{4})$, we can write Q_{2j} as

$$(4.2.9) \quad Q_{2j}(\theta) = \sqrt{2} \sin\left(\theta + \frac{\pi}{4}\right) \sum_{l=1}^j (\cos \theta)^{2(j-l)} (\sin \theta)^{2(l-1)}.$$

Let us put

$$(4.2.10) \quad q_j(\theta) = \sum_{l=1}^j (\cos \theta)^{2(j-l)} (\sin \theta)^{2(l-1)} \quad \text{for } j = 1, 2, \dots, n.$$

Observe that

$$(4.2.11) \quad q_j\left(-\theta - \frac{\pi}{4}\right) = q_j\left(\theta - \frac{\pi}{4}\right) \quad \text{and} \quad q_j(\theta) \geq 2^{-j}.$$

We now begin to estimate F in (4.2.8). We split the integral in (4.2.8) as

$$\begin{aligned}
 F(v, s, r) &= \int_{-\pi}^{\pi} \frac{sP(s, \theta)}{1 + i \left(\sqrt{2} \sin \left(\theta + \frac{\pi}{4} \right) \sum_{j=1}^n s^{2j-1} a_j q_j(\theta) \right)} \phi(v, r, \theta) d\theta \\
 &= \int_{-\pi}^{-\frac{\pi}{2}} + \int_{-\frac{\pi}{2}}^0 + \int_0^{\frac{\pi}{2}} + \int_{\frac{\pi}{2}}^{\pi} \cdots d\theta = I_1 + I_2 + I_3 + I_4.
 \end{aligned}$$

Since $|\sqrt{2} \sin \left(\theta + \frac{\pi}{4} \right) q_n(\theta)| \geq 2^{-p}$ if $-\pi \leq \theta \leq -\pi/2$ or $0 \leq \theta \leq \pi/2$, I_1 and I_3 are bounded by the same reason as above. We now estimate I_2 . By translating by $\frac{\pi}{4}$, we have

$$(4.2.12) \quad I_2 = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{sP\left(s, \theta - \frac{\pi}{4}\right)}{1 + i \sqrt{2} \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)} \phi\left(v, r, \theta - \frac{\pi}{4}\right) d\theta.$$

Let I and II be the real part and the imaginary part of I_2 respectively. Since $q_j\left(-\theta - \frac{\pi}{4}\right) = q_j\left(\theta - \frac{\pi}{4}\right)$ for any j , we have

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\sqrt{2} \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)}{1 + \left(\sqrt{2} \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)\right)^2} d\theta = 0.$$

Therefore, by (4.2.5) and (4.2.9), we have

$$\begin{aligned}
 |II| &= \left| \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\sqrt{2} \sin \theta \sum_{j=1}^d s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)}{1 + \left(\sqrt{2} \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)\right)^2} \right. \\
 &\quad \times \left. \left(P\left(s, \theta - \frac{\pi}{4}\right) \phi\left(v, r, \theta - \frac{\pi}{4}\right) - P\left(-\frac{\pi}{4}\right) \phi\left(v, r, -\frac{\pi}{4}\right) \right) d\theta \right| \\
 &\leq C \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{\left| \sqrt{2} s \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right) \right|}{1 + \left(\sqrt{2} \sin \theta \sum_{j=1}^d s^{2j-1} a_j q_j\left(\theta - \frac{\pi}{4}\right)\right)^2} s^{d-2} |\theta| d\theta.
 \end{aligned}$$

Since $q_n(\theta) \geq 2^{-n}$, we have

$$|II| \leq C \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{s^{2d-2} |\theta \sin \theta|}{1 + [s^{d-1} \sin \theta]^2} d\theta \leq C.$$

Finally, for $I = \Re I_2$, we use the fact $d(\theta) > 2^{-p}$ to have

$$\begin{aligned}
 |I| &= \left| \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{1 + \left(\sqrt{2} \sin \theta \sum_{j=1}^n s^{2j-1} a_j q_j \left(\theta - \frac{\pi}{4} \right) \right)^2} \right. \\
 &\quad \left. \times P\left(s, \theta - \frac{\pi}{4} \right) \phi\left(v, r, \theta - \frac{\pi}{4} \right) d\theta \right| \\
 &\leq C \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{s^{2d-1}}{1 + [s^{d-1} \sin(\theta)]^2} d\theta \leq C.
 \end{aligned}$$

This proves that I_2 is bounded. I_2 can be proved to be bounded in a similar way.

PROPOSITION 4.2.2. \mathcal{C}_1 and \mathcal{C}_2 satisfy the weak boundedness property.

Proof. Proofs are similar to the proof of Proposition 4.2.1. As in (4.2.1), we have

$$\begin{aligned}
 \langle \mathcal{C}_1 \phi_1^{u,t}, \phi_2^{v,t} \rangle &= t \lim_{\varepsilon \rightarrow 0} \int \int_{|x-y|>\varepsilon} \frac{1}{x-y} \phi(tx, ty) \phi_1^v(x) \phi_2^v(y) dx dy \\
 &\quad - it \lim_{\varepsilon \rightarrow 0} \int \int_{|x-y|>\varepsilon} \frac{tP(tx, ty)}{1 + iQ(tx, ty)} \phi(tx, ty) \phi_1^v(x) \phi_2^v(y) dx dy \\
 &:= t(I_1(t, v) - iI_2(t, v)).
 \end{aligned}$$

Here ϕ is the cut-off function defined in (4.1.4). For I_2 , (4.2.2) can be changed as

$$I_2 = \int_0^\infty \int_{-\pi}^\pi \frac{trP(tr, \theta)}{1 + iQ(tr, \theta)} \phi(t, v, r, \theta) d\theta dr$$

where

$$\phi(t, r, \theta, v) = \phi(tr \cos \theta, tr \sin \theta) \phi_1^v(r \cos \theta, r \sin \theta) \phi_1^v(r \cos \theta, r \sin \theta).$$

Again since the set $\{r \in [0, \infty) : \phi(t, v, r, \theta) \neq 0\}$ is included in the interval $[|v| - 2, |v| + 2]$, it suffices to show that

$$F(t, r, s) := \int_{-\pi}^\pi \frac{trP(tr, \theta)}{1 + iQ(tr, \theta)} \phi(t, v, r, \theta) d\theta$$

is bounded. Now we can repeat the same argument as in Proposition 4.2.1 to show that I_2 is bounded.

I_1 looks almost like a truncated Hilbert transform. However, unlike the usual truncation, the size of truncation in I_1 varies depending on x . So, we include the proof of the boundedness of I_1 even if it follows a standard argument. Note that, since $\phi(tx, ty) = 1$ if $|x - y| \leq 1/2(t^{-1} + |x|)$ and $\phi_1^v(x)\phi_2^v(y) = 0$ if $|x - y|$

≥ 2 , $\phi(tx, ty)\phi_1^v(x)\phi_2^v(y) = \phi_1^v(x)\phi_2^v(y)$ if either $|x| > 4$ or $t \leq \frac{1}{4}$. Note also that, if $|v| > 8$, then $\phi_1^v(x) \neq 0$ only if $|x| \geq 7$. Hence if either $|v| > 8$ or $t \leq 1/4$, then $\phi(tx, ty)\phi_1^v(x)\phi_2^v(y) = \phi_1^v(x)\phi_2^v(y)$ and hence $I_1(t, v) = \langle H\phi_1^v, \phi_2^v \rangle$ where H is Hilbert transform. Therefore, I_1 is bounded.

Suppose now that $|v| \leq 8$ and $t \geq 1/4$. Then

$$\begin{aligned} I_1(t, v) &= \frac{1}{2} \int \int \frac{1}{x-y} (\phi(tx, ty)\phi_1^v(x)\phi_2^v(y) - \phi(ty, tx)\phi_1^v(y)\phi_2^v(x)) \, dx dy \\ &= \frac{1}{2} \int \int \frac{1}{x-y} (\phi(tx, ty) - \phi(ty, tx))\phi_1^v(x)\phi_2^v(y) \, dx dy \\ &\quad + \frac{1}{2} \int \int \frac{1}{x-y} \phi(tx, ty)(\phi_1^v(x)\phi_2^v(y) - \phi_1^v(y)\phi_2^v(x)) \, dx dy. \end{aligned}$$

But from (4.1.3) and (4.1.4), we obtain

$$|\phi(tx, ty) - \phi(ty, tx)| \leq 10|x-y|(h(t, x, y) + h(t, y, x))$$

where

$$h(t, x, y) = \begin{cases} \frac{1}{t+|x|} & \text{for } \frac{1}{2}\left(\frac{1}{t} + |x|\right) \leq |x-y| \leq \frac{4}{5}\left(\frac{1}{t} + |x|\right), \\ 0 & \text{otherwise.} \end{cases}$$

Since $\phi_1^v(x)\phi_2^v(y) = 0 = \phi_1^v(y)\phi_2^v(x)$ for $|x| + |y| > 20$,

$$\begin{aligned} |I_1(t, v)| &\leq C \int \int_{|x|+|y| \leq 20} (h(t, x, y) + h(t, y, x)) \, dx dy \\ &\quad + \int \int_{|x|+|y| \leq 20} \phi(ty, tx) (\|\phi_1^v\|_{C^1} + \|\phi_2^v\|_{C^1}) \, dx dy \\ &\leq C. \end{aligned}$$

This completes the proof.

4.3. Estimates for $\mathcal{E}_1 1$

We now show that $\mathcal{E}_1 1, \mathcal{E}_1^* 1 \in \text{BMO}$.

PROPOSITION 4.3.1. $\mathcal{E}_1 1 \in \text{BMO}$.

Proof. Since $\phi(x, y) = 0$ if $|x-y| > \frac{4}{5}(|x|+1)$ and 1 if $|x-y|$

$< \frac{1}{2}(|x| + 1)$, we can divide $\mathcal{C}_1 1(x)$ as follows;

$$\begin{aligned} \mathcal{C}_1 1(x) &= \text{p.v.} \int_{-\infty}^{\infty} K_1(x, y) dy \\ &= \text{p.v.} \int_{|x-y| \leq \frac{1}{2}} \frac{1}{x-y} \left(\frac{1+iA'(y)}{1+iQ(x,y)} - 1 \right) \\ &\quad + \int_{\frac{1}{2} \leq |x-y| \leq \frac{1}{2}(|x+1|)} \frac{1}{x-y} \frac{1+iA'(y)}{1+iQ(x,y)} dy \\ &\quad + \int_{\frac{1}{2}(|x+1|) \leq (|x-y|) \leq \frac{4}{5}(|x+1|)} \frac{1}{x-y} \frac{1+iA'(y)}{1+iQ(x,y)} \phi(x, y) dy \\ &:= I_1(x) + I_2(x) + I_3(x). \end{aligned}$$

Let

$$\Psi(x, y) = \frac{1+iA'(y)}{1+iQ(x,y)} - 1 = \frac{1+iA'(y)}{1+iQ(x,y)} - \frac{1+iA'(y)}{1+iA'(y)}.$$

Then, by the mean value theorem and Lemma 3.1 (1) and (3), we have

$$|\Psi(x, y)| \leq |x-y| \sup_{|x-y| < 1} \left| \frac{\partial \Psi}{\partial y}(x, y) \right| \leq C|x-y|.$$

Therefore, $I_1(x)$ is bounded. Let M be the large number given in Lemma 3.1. If $|x| \leq M$ and if $|x-y| \leq \frac{4}{5}(|x| + 1)$, then, $|y| \leq 2M$. Hence, I_2 and I_3 are bounded if $|x| \leq M$.

We now handle the case when $|x| > M$. Assume that $x > M$ without loss of generality. If $\frac{1}{2}(x+1) \leq |x-y| \leq \frac{4}{5}(1+x)$ and if $|x| \geq M$, then by Lemma 3.1

$$\left| \frac{1}{x-y} \left(\frac{1+iA'(y)}{1+iQ(x,y)} \right) \right| \leq C|x|^{-1}$$

and therefore I_3 is bounded for $x > M$. Hence we only need to show $I_2 \in \text{BMO}$. We decompose I_2 as

$$\begin{aligned} (4.3.1) \quad I_2(x) &= \int_{1 \leq |x-y| \leq \frac{1}{2}(x+1)} \frac{1}{x-y} \left(\frac{1+iA'(y)}{1+iQ(x,y)} - \frac{A'(y)}{Q(x,y)} \right) dy \\ &\quad + \int_{1 \leq |x-y| \leq \frac{1}{2}(x+1)} \frac{A'(y)}{A(x) - A(y)} dy \end{aligned}$$

$$:= F(x) + G(x).$$

If $d = 1$, clearly $I_2 \in \text{BMO}$. Assume that $d > 1$. Then, by Lemma 3.1, we have

$$\left| \frac{1 + iA'(y)}{1 + iQ(x, y)} - \frac{A'(y)}{Q(x, y)} \right| \leq C \frac{|Q(x, y) - A'(y)|}{|Q(x, y)|^2} \leq Cx^{-d+1}$$

if $x > M$ and $1 \leq |x - y| \leq \frac{1}{2}(x + 1)$. Hence $F(x)$ is bounded for $x \geq M$. So, in order to prove that $I_2 \in \text{BMO}$, it suffices to show

$$(4.3.2) \quad |G'(x)| < Cx^{-1} \quad \text{for sufficiently large } x$$

by Lemma 2.1. To prove (4.3.2), note that

$$\begin{aligned} G'(x) &= \int_{1 \leq |x-y| \leq \frac{1}{2}(x+1)} \left(\frac{\partial}{\partial y} + \frac{\partial}{\partial x} \right) \left(\frac{A'(y)}{A(x) - A(y)} \right) dy + J(x) \\ &= \int_{1 \leq |x-y| \leq \frac{1}{2}(x+1)} \frac{A''(y)[A(x) - A(y)] - A'(y)[A'(x) - A'(y)]}{[A(x) - A(y)]^2} dy + J(x) \end{aligned}$$

where

$$J(x) = \frac{1}{2} \frac{A'\left(\frac{1}{2}x - \frac{1}{2}\right)}{A(x) - A\left(\frac{1}{2}x - \frac{1}{2}\right)} + \frac{1}{2} \frac{A'\left(\frac{3}{2}x + \frac{1}{2}\right)}{A(x) - A\left(\frac{3}{2}x + \frac{1}{2}\right)}.$$

Clearly $J(x)$ is bounded for $x > M$. If $1 \leq |x - y| \leq \frac{1}{2}(x + 1)$ and $x \geq M$, then

$$\begin{aligned} |A(x) - A(y)| &\geq C|x|^{d-1}|y - x| \\ A''(y)[A(x) - A(y)] &= A''(y)[A'(y)(x - y) + (x - y)^2 O(|x|^{d-2})] \\ A'(y)[A'(x) - A'(y)] &= A'(y)[A''(y)(x - y) + (x - y)^2 O(|x|^{d-2})] \end{aligned}$$

and therefore

$$|A''(y)[A(x) - A(y)] - A'(y)[A'(x) - A'(y)]| \leq C|x - y|^2|x|^{d-2}.$$

Hence for $x > M$,

$$|G'(x)| \leq C \int_{1 \leq |x-y| \leq \frac{1}{2}x} \frac{x^{2d-4}}{x^{2d-2}} dy \leq Cx^{-1}.$$

This completes the proof.

PROPOSITION 4.3.2. $\mathcal{C}_1^* \mathbf{1} \in BMO$.

Proof. Proposition 4.3.2 can be proved in the same way as Proposition 4.3.1. We include a sketch of the proof for reader’s sake. As in the proof of Proposition 4.3.1, we divide $\mathcal{C}_1^* \mathbf{1}$ as follows;

$$\begin{aligned} \mathcal{C}_1^* \mathbf{1}(x) &= \text{p.v.} \int_{-\infty}^{\infty} K_1(y, x) dy \\ &= \text{p.v.} \int_{|x-y| < \frac{1}{2}} \frac{1}{x-y} \left(\frac{1+iA'(x)}{1+iQ(x,y)} - 1 \right) \\ &\quad + \int_{\frac{1}{2} \leq |x-y| \leq \frac{1}{2}(|y|+1)} \frac{1}{x-y} \left(\frac{1+iA'(x)}{1+iQ(x,y)} \right) dy \\ &\quad + \int_{\frac{1}{2}(|y|+1) \leq |x-y| \leq \frac{4}{5}(|y|+1)} \frac{1}{x-y} \left(\frac{1+iA'(x)}{1+iQ(x,y)} \right) \phi(y,x) dy \\ &:= I_1(x) + I_2(x) + I_3(x). \end{aligned}$$

Then, by the same reasons as in the proof of Proposition 4.3.1, I_1 is bounded and I_2 and I_3 are bounded if $|x| \leq M$.

Assume that $|x| \geq M$ and that $x > 0$. Then, by Lemma 3.1, we have

$$\begin{aligned} |I_3(x)| &\leq \int_{\frac{1}{2}(|y|+1) \leq |x-y| \leq \frac{4}{5}(|y|+1)} \left| \frac{1}{x-y} \right| \left| \frac{1+iA'(x)}{1+iQ(x,y)} \right| dy \\ &\leq Cx^{d-1} \int_{\frac{1}{2}x \leq |y|} \frac{1}{|y|^d} dy \leq C. \end{aligned}$$

In order to prove that $I_2 \in BMO$, it suffices to show that

$$G_*(x) = \int_{\frac{1}{2} \leq |x-y| \leq \frac{1}{2}(x+1)} \left(\frac{A'(x)}{A(x) - A(y)} \right) dy$$

satisfies the estimates $|G'_*(x)| < Cx^{-1}$ for $x \geq M$ as in the proof of Proposition 4.3.1. We then note that

$$G'_*(x) = \int_{1 \leq |x-y| \leq \frac{1}{4}x} \left(\frac{\partial}{\partial y} + \frac{\partial}{\partial x} \right) \left(\frac{A'(x)}{A(x) - A(y)} \right) dy + J_*(x)$$

where

$$J_*(x) = \frac{2}{3} \frac{A'(x)}{A(x) - A\left(\frac{2}{3}x - \frac{1}{3}\right)} + \frac{A'(x)}{A(x) - A(2x + 1)}.$$

One can see as in the proof of Proposition 4.3.1 that if $1 \leq |x - y| \leq \frac{1}{2}(|y| + 1)$ and $x \geq M$, then

$$\left| \left(\frac{\partial}{\partial y} + \frac{\partial}{\partial x} \right) \left(\frac{A'(x)}{A(x) - A(y)} \right) \right| \leq Cx^{-2}$$

and therefore $G'_*(x) = O(x^{-1})$ for $x > M$. This completes the proof.

4.4 Estimates for $\mathcal{E}_2\mathbf{1}$

In this subsection, we finally show that $\mathcal{E}_2\mathbf{1}, \mathcal{E}_2^*\mathbf{1} \in \text{BMO}$.

PROPOSITION 4.4.1 $\mathcal{E}_2\mathbf{1} \in \text{BMO}$.

Proof. For given x , we let $Q = [-2|x|, 2|x|]$. Then, $\mathcal{E}_2\chi_Q(0)$ is finite. So, $\mathcal{E}_2\mathbf{1}(x)$ is understood to be

$$\mathcal{E}_2\mathbf{1}(x) = \mathcal{E}_2\chi_Q(x) - \mathcal{E}_2\chi_Q(0) + \int_{Q^c} [K_2(x, y) - K_2(0, y)] dy$$

where χ_Q is the characteristic function for Q . Then, by Proposition 4.1.2

$$\begin{aligned} \int_{Q^c} [K_2(x, y) - K_2(0, y)] dy &\leq \int_{|y|>2|x|} \sup_{|\xi| \leq |x|} |\nabla_x K_2(\xi, y)| |x| dy \\ &\leq C|x| \int_{|y|>2|x|} \frac{1}{|y|^2} dy \leq C. \end{aligned}$$

So, it remains to show that $\mathcal{E}_2\chi_Q(x)$ and $\mathcal{E}_2\chi_Q(0)$ belong to BMO. If $|x| < 2M$, then both $\mathcal{E}_2\chi_Q(x)$ and $\mathcal{E}_2\chi_Q(0)$ are bounded. Suppose that $|x| \geq 2M$. Then,

$$\mathcal{E}_2\chi_Q(0) = \int_{1/2 < |y| < 4/5} K_2(0, y) dy + \int_{4/5 \leq |y| \leq 2|x|} \frac{1 + iA'(y)}{y + iA(y)} dy.$$

The first integral in the right hand side is bounded and the second one is $\log(2|x| + iA(|x|)) + C$ which belongs to BMO by Lemma 2.1. Finally we show that $\mathcal{E}_2\chi_Q(x) \in \text{BMO}$.

$$\begin{aligned} \mathcal{E}_2\chi_Q(x) &= \int_{\substack{|y| \leq 2|x| \\ 1/2(1+|x|) < |x-y| < 4/5(1+|x|)}} K_2(x, y) dy + \int_{\substack{|y| \leq 2|x| \\ 4/5(1+|x|) < |x-y|}} K_2(x, y) dy \\ &= I_1(x) + I_2(x). \end{aligned}$$

If $|y| \leq 2|x|$ and $1/2(1 + |x|) < |x - y| < 4/5(1 + |x|)$, then $|K_2(x, y)|$

$\leq C|x + y|^{-1} \leq C|x|^{-1}$ and hence

$$|I_1(x)| \leq C \int_{1/2|x| \leq |y| \leq 9/5|x|} |K_2(x, y)| dy \leq C.$$

If $4/5(1 + |x|) \leq |x - y|$, then $\phi(x, y) = 0$ and hence

$$\begin{aligned} I_2(x) &= \int_{-2|x|}^{1/5|x|-4/5} + \int_{9/5|x|+4/5}^{2|x|} \frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))} dy \\ &= \log((x + 2|x|) + i(A(x) - A(-2|x|))) \\ &\quad - \log((x - 1/5|x| + 4/5) + i(A(x) - A(1/5|x| - 4/5))) \\ &\quad - \log((x - 2|x|) + i(A(x) - A(2|x|))) \\ &\quad + \log((x - 9/5|x| - 4/5) + i(A(x) - A(9/5|x| - 4/5))) \in \text{BMO}. \end{aligned}$$

This completes the proof.

PROPOSITION 4.4.2. $\mathcal{E}_2^*1 \in \text{BMO}$.

Proof. We may assume $d = \text{deg}A \geq 2$. Recall that

$$\mathcal{E}_2^*1(x) = \int_{-\infty}^{\infty} \frac{1 + iA'(y)}{(x - y) + i(A(x) - A(y))} (1 - \phi(y, x)) dy.$$

If $|x| < 2M$, then

$$\begin{aligned} |\mathcal{E}_2^*1(x)| &\leq C \int_{|x-y|>1/2} \frac{1}{|x - y| + |A(x) - A(y)|} dy \\ &\leq C \int_{-\infty}^{\infty} \frac{1}{1 + |y|^d} dy < C. \end{aligned}$$

We now suppose that $|x| \geq 2M$ and assume that $x > 0$ without loss of generality. We then split \mathcal{E}_2^*1 as

$$\begin{aligned} \mathcal{E}_2^*1(x) &= \int_{|y|>2x} + \int_{-2x \leq y \leq -M} + \int_{-M < y \leq 2x} \overline{K_2(y, x)} dy \\ &= I_1(x) + I_2(x) + I_3(x). \end{aligned}$$

Estimates of I_1 and I_3 are easier parts. In fact, by Lemma 3.1,

$$|I_1(x)| \leq C|A'(x)| \int_{|y|>2x} \frac{1}{|y|^{d-1}} dy < C,$$

and

$$\begin{aligned}
 |I_3(x)| &\leq C \int_{\substack{-M \leq y \leq M \\ 1/2(1+|y|) < |x-y|}} \frac{|A'(x)|}{|A(x)|} dy \\
 &\quad + C \int_{\substack{M \leq y \leq 2x \\ 1/2(1+|y|) < |x-y|}} \frac{|A'(x)|}{|x-y|[1+|x+y|(x^{d-2}+|y|^{d-2})]} dy \\
 &\leq \frac{C}{x} + Cx^{d-1} \int_{\frac{1}{10}x}^{2x} \frac{1}{y^d} dy \leq C.
 \end{aligned}$$

For $I_2(x)$, we let

$$f(x) = A'(x) \int_{-2x}^{-M} \frac{1}{(x-y) + i(A(x) - A(y))} (1 - \phi(y, x)) dy.$$

Note that if $-2x \leq y \leq -M$ and $x > 2M$, then $4/5(1 + |y|) < |x - y|$ and hence $\phi(y, x) = 0$. If $A(x)$ is of odd degree, then it is easy to see that $f(x)$ is bounded. Therefore, we assume that A is of even degree. We use Lemma 2.1. Note that

$$\begin{aligned}
 f'(x) &= A''(x) \int_{-2x}^{-M} \frac{1}{(x-y) + i(A(x) - A(y))} dy \\
 &\quad + A'(x) \int_{-2x}^{-M} \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial y} \right) \left(\frac{1}{(x-y) + i(A(x) - A(y))} \right) dy + E(x)
 \end{aligned}$$

where

$$E(x) = \frac{A'(x)}{(x-M) - i(A(x) - A(M))} - \frac{A'(x)}{3x - i(A(x) - A(-2x))}.$$

It then follows that

$$\begin{aligned}
 f'(x) &= \int_{-2x}^{-M} \frac{A''(x)(x-y) - 2A'(x)}{[(x-y) + i(A(x) - A(y))]^2} dy \\
 &\quad + i \int_{-2x}^{-M} \frac{A''(x)(A(y) - A(x)) + A'(x)(A'(y) + A'(x))}{[(x-y) + i(A(x) - A(y))]^2} dy + E(x) \\
 &= J_1(x) + iJ_2(x) + E(x).
 \end{aligned}$$

Since $|A''(x)(x-y) - 2A'(x)| \leq C(x^{d-2}|x-y| + x^{d-1})$ by Lemma 3.1 (2) and $|x-y| \approx |x|$ if $-2x < y < -M$, we have

$$\begin{aligned}
 |J_1(x)| &\leq C \int_{-2x}^{-M} \frac{x^{d-2} |x-y| + x^{d-1}}{|x-y|^2 [1 + |x+y|(x^{d-2} + |y|^{d-2})]^2} dy \\
 &\leq Cx^{d-3} \int_{-2x}^{-M} \frac{1}{1 + |x+y|^2 x^{2(d-2)}} dy \leq Cx^{-1}.
 \end{aligned}$$

We now estimate $J_2(x)$. Since A is even, we have

$$\begin{aligned}
 &A''(x)(A(y) - A(x)) + A'(x)(A'(y) + A'(x)) \\
 &= A''(-x)(A(y) - A(-x)) - A'(-x)(A'(y) - A'(-x)) \\
 &= (x+y)^2 \sum_{j=2}^d \frac{1}{j!} [A^{(j)}(-x)A^{(j)}(-x) - A'(-x)A^{(j+1)}(-x)](x+y)^{j-2}
 \end{aligned}$$

and hence

$$\begin{aligned}
 &|A''(x)[A(y) - A(x)] + A'(x)[A'(y) + A'(x)]| \\
 &\leq C|x+y|^2 [|x+y|^{d-2} + x^{2(d-2)}].
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 |J_2(x)| &\leq C \int_{-2x}^{-M} \frac{|x+y|^2 (|x+y|^{d-2} + x^{2(d-2)})}{|x|^2 [1 + |x+y|(|x|^{d-2} + |y|^{d-2})]^2} dy \\
 &\leq \frac{C}{x^2} \int_{-x}^{-M+x} \frac{t^2(t^{d-2} + |x|^{2(d-2)})}{(1 + |t||x|^{d-2})^2} dt \leq x^{-1}.
 \end{aligned}$$

It is easy to see that $|E(x)| \leq C|x|^{-1}$. In conclusion, we have $|f'(x)| \leq C|x|^{-1}$ if $|x| \geq 2M$. By Lemma 2.1, $f \in \text{BMO}$. It follows that $I_2 \in \text{BMO}$. This completes the proof.

5. Non- L^2 -boundedness

In this section, we give two examples of A for which \mathcal{C}_A are not L^2 -bounded. The first example of A has two many zeros while the derivative of the second A grows too fast relatively to A itself.

THEOREM 5.1. *Let $A'(x) = x \sin x$. Then, \mathcal{C}_A is not bounded on L^2 .*

Proof. For each positive integer n , we let f_n be the characteristic function on $[2n\pi + \pi/4, 2n\pi + 3\pi/4]$. Then, $\|f_n\|_2 = \pi/2$ for each n . Note that

$$|A(x) - A(y)| = |-x \cos x + y \cos y + \sin x - \sin y| \leq 2(|x - y| + |y| + 1).$$

If $n \geq 2, j \geq 2$, and if $2n\pi + \pi/4 \leq y \leq 2n\pi + 3\pi/4$ and $2(n + j)\pi \leq x \leq 2(n + j + 1)\pi$, then

$$|A(x) - A(y)| \leq 2(|x - y| + |y| + 1) \leq \frac{1}{2} y \sin y (x - y).$$

It then follows that

$$\begin{aligned} |\mathcal{C}_A f_n(x)| &\geq |\Im \mathcal{C}_A f_n(x)| \\ &= \left| \int_{2n\pi+\pi/4}^{2n\pi+3\pi/4} \frac{y \sin y (x - y) + (A(x) - A(y))}{(x - y)^2 + (A(x) - A(y))^2} dy \right| \\ &\geq \int_{2n\pi+\pi/4}^{2n\pi+3\pi/4} \frac{y \sin y |x - y| - |A(x) - A(y)|}{(x - y)^2 + (A(x) - A(y))^2} dy \\ &\geq \frac{1}{10} \int_{2n\pi+\pi/4}^{2n\pi+3\pi/4} \frac{y \sin y |x - y|}{(|x - y| + |y| + 1)^2} dy \geq C \frac{nj}{(n + j)^2} \end{aligned}$$

for some constant C . Therefore,

$$\|\mathcal{C}_A f_n\|_2^2 \geq \sum_{j=2}^{\infty} \int_{2(n+j)\pi}^{2(n+j+1)\pi} |\mathcal{C}_A f_n(x)|^2 dx \geq C \sum_{j=2}^{\infty} \frac{n^2 j^2}{(n + j)^4}.$$

Since

$$\sum_{j=2}^{\infty} \frac{j^2}{(n + j)^4} \geq \frac{C}{n},$$

we have $\|\mathcal{C}_A f_n\|_2 \geq C \sqrt{n} \|f_n\|_2$ for each $n \geq 2$. This completes the proof.

THEOREM 5.2. *Let $A(x) = \exp(x^2)$. Then, \mathcal{C}_A is not bounded on L^2 .*

Proof. For each positive integer n , we let f_n be the characteristic function on $[n, n + 1]$. Then, $\|f_n\|_2 = 1$ for each n . If $x \in [0, 1]$, we have

$$|\mathcal{C}_A f_n(x)| \geq C \int_n^{n+1} \frac{y \exp(y^2) |A(x) - A(y)|}{(x - y)^2 + (A(x) - A(y))^2} dy \geq C$$

for some constant C independent of n . So $\|\mathcal{C}_A f_n\|_2 \geq Cn \|f_n\|_2$ for each n . This completes the proof.

REFERENCES

- [Cal1] A. P. Calderón, Cauchy integrals on Lipschitz curves and related operators, Proc. Nat. Acad. Sci. USA, **74** (1977), 1324–1327.
- [Cal2] —, Commutators, Singular integrals on Lipschitz curves and applications, Proceedings of ICM (Helsinki, 1978), Acad. Sci. Fennica, Helsinki (1980), 85–96.
- [Chr] M. Chris, Lectures on Singular Integral Operators, CBMS 77, Amer. Math. Soc., 1990.
- [C.D.M] R. R. Coifman, G. David, Y. Meyer, La solution des conjectures de Calderón, Advances in Math., **48** (1983), 144–148.
- [C.J.S] R. R. Coifman, P. Jones, and S. Semmes, Two elementary proofs of the L^2 -boundedness of Cauchy integrals on Lipschitz curves, J. Amer. Math. Soc., **2** (1989), 553–564.
- [C.M.M] R. R. Coifman, A. McIntosh, Y. Meyer, L'intégrale de Cauchy définit un opérateur bornée sur L^2 pour courbes lipschitziennes, Ann. of Math., **116** (1982), 361–387.
- [C.M] R. R. Coifman and Y. Meyer, Au delà des opérateurs pseudo-différentiels, Astérisque 57, Société Mathématique de France, 1978.
- [D.J] G. David and J.-L. Journé, A boundedness criterion for generalized Calderón-Zygmund operators, Ann. of Math., **120** (1984), 371–397.
- [Jou] J.-L. Journé, Calderón-Zygmund operators, pseudo-differential operators, and the Cauchy integral of Calderón, Lecture Notes in Math., **994**, Springer Verlag, New York, 1983.
- [Mur] T. Murai, A real variable method for the Cauchy transform, and analytic capacity, Lecture Note in Math., **1307**, Springer-Verlag, New York, 1988.
- [St1] E. Stein, Singular Integrals and Differentiability Properties of Functions, Princeton Univ. Press, Princeton, 1970.
- [St2] —, Beijing Lectures in harmonic analysis, Princeton Univ. Press Princeton, 1986.
- [Tor] A. Torchinsky, Real-variable methods in Harmonic Analysis, Academic Press, 1986.

H. Kang
Department of Mathematics
Soong Sil University,
Sangdo-Dong, Dongjak-Gu
Seoul, 156-743
Korea

J. K. Seo
GARC
Seoul National University
Shinlim-Dong, Kwanak-Gu
Seoul, 135-110
Korea

Current address of J. K. Seo
Department of Mathematics
POSTECH
Pohang, Korea