

Conclusions: Cluster B personality disorder, admission for heroin detoxification, and pre-admission binge-pattern substance use are factors associated with voluntary treatment discontinuation.

Disclosure of Interest: None Declared

Child and Adolescent Psychiatry

EPP0086

Associations between cerebellar development and autistic traits during adolescence: a population-based cohort study

C. Morimoto^{1*} and S. Koike²

¹Faculty of Allied Health Science, Niigata University of Rehabilitation, Niigata and ²Department of Neuropsychiatry, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.312

Introduction: Brain maturation is associated with adolescent socio-cognitive development. The lateral posterior region of the cerebellum plays a critical role in higher cognitive processes, and deviations of this region are associated with autism-related behaviors. Hence, it is plausible that developmental changes in this region of the cerebellum during adolescence are different along a variation in autistic traits. Additionally, its difference may be moderated by parental age at birth and weight growth during infancy, which have effects on brain development.

Objectives: The aim of this study was two folds: (1) to test whether cerebellar development during adolescence is different along a variation in autistic traits (2) to test whether parental age at birth and weight growth during infancy moderate the results of (1).

Methods: Longitudinal study was conducted over a 6-year period with 256, 230 and 187 participants ranging from 10.5 to 17.6 years, observing adolescent development respectively at 2-year time periods. We undertook a detailed investigation into differences in the lateral posterior region of the cerebellum volume. The 50-item Autism-Spectrum Quotient (AQ) was rated by primary parents. Weight growth and parental age were evaluated using maternal and child health handbook records. A multiple regression analysis was performed to examine whether AQ subscales, sex, and their interactions affected cerebellar development. Moderation analysis assessed whether parental age and weight growth moderated associations between cerebellar development and autistic traits. All participants provided written informed consent, and the study was approved by the Ethics Committee (No.10069).

Results: Interactions between sex and attention switching and sex and attention to detail were significantly associated with cerebellar development in the bilateral gray matter (GM) and white matter (WM) of Crus I and Crus II (Fig1, 2). Simple slope analyses showed that the slopes of cerebellar development were significant for girls ($p_{FDR} < 0.001$). Although no significant interaction was found between them, the main effect of attention to detail was significantly associated with cerebellar development in WM of VIIB ($p_{FDR} = 0.006$). Further, moderation analysis found that the association between the cerebellar development and autistic traits were significantly moderated by maternal age; the magnitude of its effect was significant for high maternal age in boys ($p_{FDR} = 0.036$, Fig3).

Paternal age, early (0-9 months) and late weight growth (4-18 months) also moderated associations between them, however, no significance remained after FDR controlling.

Image:




Figure.1

Image 2:




Figure.2

Image 3:




Figure.3

Conclusions: There are significant associations between cerebellar development during adolescence and autistic traits, and its pattern of association can be moderated by parental ages at birth and weight growth during infancy in a cerebellar region- and sex-specific manner.

Disclosure of Interest: None Declared

EPP0087

Changes in Parental Reflective Functioning before and after a postpartum depression group therapy

L. Erkoreka^{1,2,3*}, Z. Alonso⁴ and L. Pérez Cabeza⁵

¹Galdakao-Usansolo University Hospital, Galdakao; ²University of the Basque Country UPV/EHU, Leioa; ³Biocruces Bizkaia Health Research Institute, Barakaldo; ⁴Bizkaia Mental Health Network, Getxo and ⁵Bizkaia Mental Health Network, Bilbao, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.313

Introduction: Parental Reflective Functioning (PRF) refers to parents' ability to view their children's and their own behavior by considering internal mental states, such as thoughts, desires, and intentions. Depression has been described as compromising reflective functioning in female samples, whereas other studies have not detected differences in RF between depressed and non-depressed mothers.

Objectives: We aim to study whether a group intervention focused on postpartum depression, which we have already observed to cause significant changes in the mother-child bond and the severity of depressive, also improves parental reflective functioning.

Methods: To that end, we analyzed pre-post data from two different groups (N=12), composed of mothers who had been clinically diagnosed with postpartum depression. They received the 6-week Mothers & Babies Program© and completed the Parental Bonding

Questionnaire (PBQ), the Edinburgh Postnatal Depression Scale (EPDS) and the Parental Reflective Functioning Questionnaire (PRFQ) before and after group therapy. Pre-post data from the PRFQ were analyzed using the repeated measures t-test. The correlation between changes in the three questionnaires was also analyzed using Pearson's correlation test.

Results: Significant changes were observed in the Pre-Mentalization Modes (pre=2.37±.457, post=2.03±.520, t=2.0206, p=0.027) and Certainty About Mental States (pre=2.87±1.141, post=3.68±.908, t=-2.814, p=0.010) subscales of the PRFQ, with no significant changes in the Interest and Curiosity subscale (t=-.516, p=0.309). A significant correlation was also observed between pre-post change in EPDS scores and pre-post change in the Certainty About Mental States subscale of the PRFQ (r=-.640, p<.05), while no significant correlations were observed with the rest of the PRFQ subscales, nor with the PBQ.

Conclusions: A brief cognitive-behavioral group therapy developed specifically to treat postpartum depression improves pre-post scores on the Pre-Mentalization (lower post- than pre- score) and Certainty About Mental States (higher post- than pre- score) subscales of the PRFQ. Although a control group is needed to determine the actual effect of the intervention, as time could also play a role in the observed changes, this is an encouraging result. Moreover, the improvement obtained in Certainty About Mental States is inversely correlated with the pre-post changes observed in the EPDS, meaning that the greater the improvement in depression, the greater the improvement in the aforementioned subscale of the PRFQ. A larger sample is needed to assess a hypothetical mediating effect of depression in the observed change.

Disclosure of Interest: None Declared

EPP0088

The Effect of Depression on Smartphone Addiction: The Medicating Effects of Interpersonal Problem in Korean Adolescents

Y. Seo*, Y. Nam and J. Lee

Department of Psychiatry, Yonsei University Wonju College of Medicine, Wonju, Korea, Republic Of

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.314

Introduction: Problematic smartphone use is twice as common among teenagers as it is among adults. Smartphone addiction is associated with anxiety, depression, attention deficit disorder, impulsivity, and sleep problems, among other issues.

Objectives: To assess whether interpersonal relationship problems mediate the relationship between smartphone addiction and depression among adolescents (ages 12-17) currently enrolled in middle and high school.

Methods: A cross-sectional study was conducted among 653 middle and high school students living in Wonju, South Korea between September 1 and November 30, 2019. Depression was measured by Center for Epidemiologic Studies Depression Scale (CES-D). In order to evaluate smartphone addiction, the Smartphone Addiction Scale Short Form Version (SAS-SV) was used. To examine interpersonal problems, the Korea Inventory of Interpersonal Problems