



Winter Meeting, 6–7 December 2016, Diet, Nutrition and Mental Health and Wellbeing

## Malnutrition in hospitalised older adults: A multi-centre observational study of prevalence, associations and outcomes

Emma O' Shea<sup>1</sup>, Steven Trawley<sup>2</sup>, Edmund Manning<sup>1</sup>, Aoife Barrett<sup>1</sup>, Vanessa Browne<sup>1</sup> and Suzanne Timmons<sup>1</sup>

<sup>1</sup>Centre for Gerontology and Rehabilitation, School of Medicine, University College Cork, Cork, Ireland and

<sup>2</sup>Centre for Health and Social Research, Australian Catholic University, Melbourne, Australia

Malnutrition is common in older adults, and is associated with high healthcare costs and adverse outcomes, particularly in hospital settings<sup>(1,2)</sup>. The prevalence and correlates of malnutrition in hospitalised older adults are currently not clear; much of the existing research in this area is limited methodologically; studies are typically based on small samples and/or narrow populations<sup>(3,4)</sup>, exclude people with dementia<sup>(3,4)</sup>, are uni-centre<sup>(5)</sup>, and/or use tools not designed for use with older adults<sup>(5)</sup>. The present study addresses this gap, investigating the prevalence, correlates and outcomes of malnutrition in older adults on admission to hospital.

In total, 606 (70+ years) older adults were included in a prospective cohort study across six hospitals in the Republic of Ireland. All elective and acute admissions to any speciality were eligible. Day-case admissions and those moribund on admission were excluded. All participants were clinically assessed for dementia on admission (see Timmons et al.<sup>(6)</sup>). Socio-demographic and clinical data, including nutritional status (Mini-Nutritional Assessment – short form<sup>(7)</sup>), was collected within 36 hours of admission. Outcome data was collected prospectively on length of stay, in-hospital mortality and institutionalisation.

The mean age was 79.7; 51 % were female; 29 % were elective admissions; 67 % were admitted to a medical specialty. Nutrition scores were available for 602/606; 37 % had a 'normal' status, 45 % were 'at-risk', and 18 % were 'malnourished'. Malnutrition was more common in females, acute admissions, older patients and those who were widowed/ separated. Dementia, functional dependency, comorbidity and frailty independently predicted a) malnutrition and b) being at-risk of malnutrition ( $p < .001$ ). Malnutrition was also associated with an increased length of stay ( $p < .001$ ), institutionalisation ( $p < 0.001$ ) and in-hospital mortality ( $p < .001$ ).

These findings support the prioritisation of nutritional screening in clinical practice and public health policy, for all  $\geq 70$  on admission to hospital, and in particular for people with dementia, increased functional dependency and/or multi-morbidity, and those who are frail.

1. Rice N, Normand C (2012) The cost associated with disease-related malnutrition in Ireland. *Public Health Nutr* 15(10): 1966.
2. Rasheed S, Woods RT (2013) Malnutrition and associated clinical outcomes in hospitalized patients aged 60 and older: An observational study in rural Wales. *J Nutr Gerontol Geriatr* 32(1): 71–80.
3. Chen CC, Bai YY, Huang GH *et al.* (2007) Revisiting the concept of malnutrition in older people. *J Clin Nurs* 16(11): 2015–26.
4. Vanderwee K, Clays E, Bocquaert I *et al.* (2011) Malnutrition and nutritional care practices in hospital wards for older people. *J Adv Nurs* 67(4): 736–46.
5. Eide HD, Halvorsen K, Almendingen K (2015) Barriers to nutritional care for the undernourished hospitalised elderly: Perspectives of nurses. *J Clin Nurs* 24(5–6): 696–706.
6. Timmons S, Manning E, Barrett A *et al.* (2015) Dementia in older people admitted to hospital: A regional multi-hospital observational study of prevalence, associations and case recognition. *Age Ageing* 44(6): 993.
7. Kaiser M, Bauer JM, Ramsch C *et al.* (2009) Validation of the mini nutritional assessment short-form (MNA®-SF): A practical tool for identification of nutritional status. *J Nutr Health Aging* 13(9): 782–8.