

# Spaces of Whitney Functions on Cantor-Type Sets

Bora Arslan, Alexander P. Goncharov and Mefharet Kocatepe

*Abstract.* We introduce the concept of logarithmic dimension of a compact set. In terms of this magnitude, the extension property and the diametral dimension of spaces  $\mathcal{E}(K)$  can be described for Cantor-type compact sets.

## 1 Introduction

We introduce a concept of logarithmic dimension as the following generalization of the Hausdorff dimension: take the function  $\psi(r) = \frac{1}{\log \frac{1}{r}}$ ,  $r > 0$ , corresponding to the logarithmic measure; then for any compact set  $K \subset \mathbb{R}$  there exists a critical value  $\lambda_0 = \lambda_0(K) \in [0, \infty]$  (called the logarithmic dimension of  $K$ ) such that for  $\lambda < \lambda_0$  the  $\psi^\lambda$ -measure of  $K$  is  $\infty$ , for  $\lambda > \lambda_0$  it is zero.

The aim of this article is to show that for the class  $\mathcal{E}(K)$  of Whitney functions defined on generalized Cantor sets the logarithmic dimension is highly suitable for the investigation of the following two problems.

The problem of geometric characterization of the extension property of  $K$  (that is the existence of a continuous linear extension operator  $L: \mathcal{E}(K) \rightarrow C^\infty(\mathbb{R}^n)$ ) was posed by Mityagin [11, Section 8.5]. Some particular results were given by Stein [16], Bierstone [2], Pawłucki and Pleśniak [13] and others. In [17] Tidten suggested to consider perfect sets of class  $(\alpha)$  (see [18], [6] for definition) and proved that the condition  $K \in (1)$  is sufficient for the extension property of  $K$  whereas “ $K \in (\alpha)$  for some  $\alpha$ ” is necessary. In particular he showed that the classical Cantor set  $C$  (clearly  $\lambda_0(C) = \infty$ ) has the extension property. Here we show that the generalized Cantor set  $K$  has the extension property if  $\lambda_0(K) > 1$  and it has not if  $\lambda_0(K) < 1$ . Examples of perfect sets of finite class without extension property can be given easily (compare this with [6]).

On the other hand, the diametral dimension of the space  $\mathcal{E}(K)$  in our case can be described in terms of the logarithmic dimension of  $K$  (Chapter 4) and what is more, we have complete (up to the case of equal logarithmic dimensions) isomorphic classification of given spaces  $\mathcal{E}(K)$  (Theorem 3). Examples of continua of pairwise nonisomorphic spaces of considered type appear directly (compare with [8]). It should be noted that the diametral dimension can not be applied to distinguish nuclear spaces of type  $C^\infty(\bar{\Omega})$  or  $\mathcal{E}(K)$  with  $\overset{\circ}{K} \neq \emptyset$ . In fact, all these spaces contain a subspace which is isomorphic to the space  $s$  of rapidly decreasing sequences and thus their diametral dimension  $\Gamma(X)$  is not larger than  $\Gamma(s)$  (see for example [11, Proposition 7]).

Received by the editors March 6, 2001.  
AMS subject classification: 46E10, 31A15, 46A04.  
©Canadian Mathematical Society 2002.

On the other hand, the space  $s$  has the smallest diametral dimension in the class of nuclear spaces and we get  $\Gamma(X) = \Gamma(s)$ .

## 2 Logarithmic Dimension

We consider the following generalization of the Cantor ternary set. Let  $(l_n)_1^\infty$  be a sequence of positive numbers and  $(N_n)_1^\infty$  be a sequence of integers,  $N_n \geq 2$  for all  $n$ . Then  $K = K((l_n), (N_n)) = \bigcap_{n=0}^\infty E_n$ , where  $E_0 = I_{0,1} = [0, 1]$  and  $E_n, n \geq 1$ , is a union of  $N_1 N_2 \cdots N_n$  disjoint closed intervals  $I_{n,k}$  of length  $l_n$  and  $E_{n+1}$  is obtained by replacing each interval by  $N_{n+1}$  disjoint subintervals  $I_{n+1,j}$  of length  $l_{n+1}$  with  $N_{n+1} - 1$  gaps of length  $h_{n+1}$ . The intervals  $I_{n,k}$  that make up the set  $E_n$  are called *basic intervals*. The set  $K$  is well-defined if for all  $n$  we have  $l_{n-1} > N_n l_n$  with  $l_0 = 1$ . Then  $h_n = \frac{l_{n-1} - N_n l_n}{N_n - 1}$ . We will restrict ourselves to the case  $l_n \leq h_n$ , since otherwise  $K$  is uniformly perfect and has the extension property.

Let  $\alpha_1 = 1$  and for  $n \geq 2$  let  $\alpha_n$  satisfy  $l_n = l_{n-1}^{\alpha_n}$ . For  $n \geq 2$  set  $\lambda_n = \frac{\log N_n}{\log \alpha_n}$ . We will analyze two regular cases:  $N_n = N$  for all  $n$  and  $N_n \nearrow \infty$  as  $n \rightarrow \infty$ . The corresponding compact set of finite type will be denoted by  $K_N$ ;  $K_\infty$  stands for the infinite case. If in particular  $\alpha_n = \alpha, n \geq 2$ , then we will write  $K_N^{(\alpha)}$  and  $K_\infty^{(\alpha)}$ .

Let us introduce a parameter which can be applied in classifying such rarefied sets as  $K_N^{(\alpha)}$ . Let  $\psi(r) = \frac{1}{\log \frac{1}{r}}, r > 0$ . Here and subsequently,  $\log$  denotes the natural logarithm,  $[x]$  denotes the greatest integer in  $x$ . For  $0 < \lambda < \infty$  consider the  $\psi^\lambda$ -measure of a compact set  $K$  (see e.g. [12, V.6.2] or [3], [15]): let  $m_\epsilon(K, \psi^\lambda) = \inf \sum \psi^\lambda(r_i)$  where the greatest lower bound is taken over all covers  $\bigcup B_i$  of  $K$  by balls  $B_i$  with  $\text{diam } B_i = r_i \leq \epsilon$ ; then  $m(K, \psi^\lambda) = \lim_{\epsilon \rightarrow 0} m_\epsilon(K, \psi^\lambda)$ . As in the definition of the Hausdorff dimension (see e.g. [15, 10.1], [3, 1.2]) we see that there exists a critical value  $\lambda_0 = \lambda_0(K), 0 \leq \lambda_0 \leq \infty$ , such that  $m(K, \psi^\lambda) = \infty$  for  $\lambda < \lambda_0$  and  $m(K, \psi^\lambda) = 0$  for  $\lambda > \lambda_0$ . Since the function  $\psi$  corresponds to the logarithmic measure we will say that the value  $\lambda_0$  is the *logarithmic dimension* of  $K$ .

**Proposition 1** *Suppose that for  $K = K((l_n), (N_n))$  the limit  $\lambda_0 = \lim_n \lambda_n$  exists in the set of extended real numbers. Then  $\lambda_0$  is the logarithmic dimension of  $K$ . In particular,  $\lambda_0(K_N^{(\alpha)}) = \frac{\log N}{\log \alpha}$ .*

**Proof** For  $\lambda > \lambda_0$  consider  $m_\epsilon(K, \psi^\lambda)$ . Notice that  $\alpha_n \rightarrow 1$ . In fact, otherwise we would have  $\lambda_0 = \infty$ . For  $\epsilon = l_n$  we have the covering of  $K$  by  $N_1 N_2 \cdots N_n$  intervals  $I_{n,k}$  of length  $\epsilon$ . Thus

$$m_\epsilon(K, \psi^\lambda) \leq N_1 N_2 \cdots N_n \log^{-\lambda} \left( \frac{1}{l_n} \right).$$

Since  $l_n = l_1^{\alpha_1 \alpha_2 \cdots \alpha_n}$  and  $N_n = \alpha_n^{\lambda_n}$  we get

$$m_\epsilon(K, \psi^\lambda) \leq \alpha_1^{\lambda_1 - \lambda} \cdots \alpha_n^{\lambda_n - \lambda} \log^{-\lambda} \left( \frac{1}{l_1} \right),$$

which tends to 0 as  $n \rightarrow \infty$  since  $\lambda > \lim \lambda_n$  and  $\alpha_n \rightarrow 1$ . Thus,  $m(K, \psi^\lambda) = 0$ .

Suppose now that  $\lambda < \lambda_0$ . Then we can modify the arguments in [3, 2.3]. For  $\epsilon > 0$  there exists a finite covering  $\bigcup_{i=1}^M U_i$  of  $K$  by open intervals  $U_i$ ,  $\text{diam } U_i = r_i < 2\epsilon$  such that  $\sum \psi^\lambda(r_i) \leq 2m_\epsilon(K, \psi^\lambda)$ . For each  $r_i$  fix  $n = n(i) \in \mathbb{N}$  with  $l_n \leq r_i < l_{n-1}$ . Let  $n_0 = \min_{i \leq M} n(i)$ ,  $n_1 = \max_{i \leq M} n(i)$ . To simplify calculations we set  $l_1 = 1/e$ . Then

$$\psi^\lambda(r_i) \geq \psi^\lambda(l_n) \geq (\alpha_1 \cdots \alpha_n)^{-\lambda}.$$

Let  $\epsilon$  be so small that  $\lambda_n > \lambda$  for  $n \geq n_0$ . Then

$$\begin{aligned} (\alpha_1 \cdots \alpha_n)^\lambda &= (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \cdot (\alpha_{n_0}^\lambda \cdots \alpha_n^\lambda) \\ &\leq (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \cdot (\alpha_{n_0}^{\lambda_{n_0}} \cdots \alpha_n^{\lambda_n}) \\ &= (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \cdot N_{n_0} \cdots N_n. \end{aligned}$$

We decompose the sum  $\sum \psi^\lambda(r_i)$  into two parts. Let  $\sum'$  be the sum over all  $i$  such that  $l_n \leq r_i < \frac{l_{n-1}}{N_n}$ , and  $\sum''$  be the sum over the remaining  $i$ 's. Since  $\frac{l_{n-1}}{N_n} < l_n + h_n$ , for any  $i$  in the sum  $\sum'$ , the interval  $U_i$  can intersect at most two basic intervals of  $E_n$ . By construction, it can intersect at most  $2N_{n+1}$  basic intervals of  $E_{n+1}; \dots; 2N_{n+1} \cdots N_{n_1}$  basic intervals of  $E_{n_1}$ . Then

$$\begin{aligned} 2N_{n+1} \cdots N_{n_1} &\leq 2N_{n+1} \cdots N_{n_1} \cdot (\alpha_1 \cdots \alpha_n)^\lambda \cdot \psi^\lambda(r_i) \\ &\leq 2N_{n_0} \cdots N_{n_1} \cdot (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \cdot \psi^\lambda(r_i). \end{aligned}$$

For  $i$  in the second sum  $\sum''$ , fix  $j$ ,  $j = 1, 2, \dots, N_n - 1$ , such that  $\frac{j}{N_n} l_{n-1} \leq r_i < \frac{j+1}{N_n} l_{n-1}$ . Then the interval  $U_i$  can intersect at most  $j + 2$  basic intervals of  $E_n$  and thus  $(j + 2)N_{n+1} \cdots N_{n_1}$  basic intervals of  $E_{n_1}$ . Here

$$\psi^\lambda(r_i) \geq \psi^\lambda\left(\frac{j}{N_n} l_{n-1}\right) \geq \left(\alpha_1 \cdots \alpha_{n-1} + \log \frac{N_n}{j}\right)^{-\lambda}.$$

If  $\log \frac{N_n}{j} \geq \alpha_1 \cdots \alpha_{n-1}$ , then  $1 \leq 2^\lambda \log^\lambda\left(\frac{N_n}{j}\right) \psi^\lambda(r_i) \leq C'_\lambda \frac{N_n}{j} \psi^\lambda(r_i)$ . Therefore

$$\begin{aligned} (j + 2)N_{n+1} \cdots N_{n_1} &\leq C''_\lambda N_n N_{n+1} \cdots N_{n_1} \psi^\lambda(r_i) \\ &\leq C''_\lambda N_{n_0} \cdots N_{n_1} \cdot (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \psi^\lambda(r_i). \end{aligned}$$

On the other hand, if  $\log \frac{N_n}{j} < \alpha_1 \cdots \alpha_{n-1}$ , then  $1 \leq 2^\lambda (\alpha_1 \cdots \alpha_{n-1})^\lambda \psi^\lambda(r_i)$ , therefore

$$\begin{aligned} (j + 2)N_{n+1} \cdots N_{n_1} &\leq (N_n + 1)N_{n+1} \cdots N_{n_1} (\alpha_1 \cdots \alpha_{n-1})^\lambda 2^\lambda \psi^\lambda(r_i) \\ &\leq 2^{\lambda+1} N_{n_0} \cdots N_{n_1} (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \psi^\lambda(r_i). \end{aligned}$$

Thus any interval  $U_i$ ,  $i \leq M$  can intersect at most

$$C_\lambda N_{n_0} \cdots N_{n_1} (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \psi^\lambda(r_i)$$

basic intervals of  $E_{n_1}$ . Here  $C_\lambda = \max\{C'_\lambda, 2^{\lambda+1}\}$ . Since the covering  $\bigcup U_i$  intersects all basic intervals of  $E_{n_1}$ , we have

$$N_1 \cdots N_{n_1} \leq C_\lambda N_{n_0} \cdots N_{n_1} (\alpha_1 \cdots \alpha_{n_0-1})^\lambda \sum \psi^\lambda(r_i)$$

and so

$$\sum \psi^\lambda(r_i) \geq C_\lambda^{-1} N_1 \alpha_2^{\lambda_2-\lambda} \cdots \alpha_{n_0-1}^{\lambda_{n_0-1}-\lambda}.$$

This bound implies that the sum of type  $\sum \psi^\lambda(r_i)$  must be arbitrarily large for small enough  $\epsilon$ , that is  $m(K, \psi^\lambda) = \infty$ .

In fact, for  $\alpha_n \rightarrow 1$  it is obvious as  $\lambda_n > \lambda$  for large  $n$ . Clearly,  $\epsilon \rightarrow 0$  gives  $n_0 \rightarrow \infty$ . If  $\alpha_n \rightarrow 1$ , then  $\alpha_n^{\lambda_n-\lambda} = \frac{N_n}{\alpha_n^\lambda} \geq \frac{2}{\alpha_n^\lambda} \rightarrow 2$  as  $n \rightarrow \infty$  and the result follows. ■

The value  $\lambda_0 = 1$  is critical in Potential Theory: if  $\lambda_0(K) < 1$ , then the logarithmic measure of  $K$  is 0 and the set  $K$  is exceptional. We show that this bound is crucial as well for the extension property of Cantor-type sets.

### 3 Extension Property

Let  $\mathcal{E}(K)$  denote the space of Whitney functions on a perfect compact set  $K$  with the topology defined by the norms

$$\|f\|_q = |f|_q + \sup\{|(R_y^q f)^{(j)}(x)| \cdot |x-y|^{j-q} : x, y \in K, x \neq y, j \leq q\}, \quad q = 0, 1, \dots,$$

where  $|f|_q = \sup\{|f^{(j)}(x)| : x \in K, j \leq q\}$  and  $R_y^q f(x) = f(x) - T_y^q f(x)$  is the Taylor remainder. Each function  $f \in \mathcal{E}(K)$  is extendable to a  $C^\infty$ -function on the line. If there exists a linear continuous extension operator  $L: \mathcal{E}(K) \rightarrow C^\infty(\mathbb{R})$ , then we say that the compact set  $K$  has the *extension property*. In [17] Tidten showed that the extension property of  $K$  and the property DN of the space  $\mathcal{E}(K)$  are equivalent.

A Fréchet space  $X$  with a fundamental system of seminorms  $(\|\cdot\|_q)$  is said to have the property DN [21] (see also the class  $D_1$  in [23]) if

$$\exists p \forall q \exists r, C > 0 : \|\cdot\|_q \leq t \|\cdot\|_p + \frac{C}{t} \|\cdot\|_r, \quad t > 0.$$

Here  $p, q, r \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$ .

**Proposition 2** For  $X = \mathcal{E}(K)$ , the following statements are equivalent to DN:

- (a)  $\exists p \exists R > 0 \forall q \exists r, C : \|\cdot\|_q \leq t^{R \cdot q} \|\cdot\|_p + \frac{C}{t^q} \|\cdot\|_r, t > 0;$
- (b)  $\exists p \forall \epsilon > 0 \forall q \exists r, C : \|\cdot\|_q^{1+\epsilon} \leq C \|\cdot\|_p \|\cdot\|_r^\epsilon.$

**Proof** In [10, Lemma 29.10] and in [5] it was shown that DN is equivalent to the conditions (a), (b) with  $\|\cdot\|_q$  instead of  $|\cdot|_q$ . Frerick [4] and Tidten [19] proved that we can replace  $\|\cdot\|_q$  by  $|\cdot|_q$ .

We first generalize Theorems 2 and 3 in [6].

**Theorem 1** If  $\underline{\lim} \alpha_n > N$ , then  $K_N$  does not have the extension property. If  $\overline{\lim} \alpha_n < N$ , then  $K_N$  has the extension property.

**Proof** Let  $\beta$  be such that  $\liminf \alpha_n > \beta > N$  and find  $i_0$  such that  $\alpha_i > \beta, i \geq i_0$ . Fix  $0 < \epsilon < \frac{\beta-N}{2(N-1)}$  and  $M \in \mathbb{N}$  such that  $M \geq \frac{2\beta}{\beta-N}$ . We want to show

$$\forall p \exists \epsilon \exists q : \forall r > q \exists (f_n) \in \mathcal{E}(K_N) : \frac{\|f_n\|_p \cdot \|f_n\|_r^\epsilon}{\|f_n\|_q^{1+\epsilon}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

For arbitrary  $p \in \mathbb{N}$  let  $q = Mp$ . For any  $r > q$  take  $s \in \mathbb{N}$  with  $N^s \geq \frac{r}{q} > N^{s-1}$ . Fix a natural number  $n, n \geq s + i_0$  and consider first  $N^s$  intervals of  $E_n$ . Let  $c_j$  denote the midpoint of  $I_{n,j}, j = 1, 2, \dots, N^s$ . Set  $f_n(x) = g^q(x)$  where  $g(x) = \prod_{j=1}^{N^s} (x - c_j)$  for  $x \in K_N \cap [0, l_{n-s}]$  and  $g(x) = 0$  otherwise on  $K_N$ . Let us estimate the norms of  $f_n$ .

Fix  $k \in \mathbb{N}, k \leq p$  and  $x \in \bigcup_{j=1}^{N^s} I_{n,j}$ . By Lemma 1 in [6], we have

$$(1) \quad |f_n^{(k)}(x)| \leq C_{p,r} |g(x)|^{q-k}$$

where  $C_{p,r} = \frac{(N^s \cdot q)!}{(N^s \cdot q - k)!} < (N \cdot r)^p$  does not depend on  $n$ .

By the structure of  $K_N$  we have  $|g(x)| < l_n \tau^{N-1}$  where  $\tau = l_{n-1} l_{n-2} \dots l_{n-s}^{N^{s-1}}$ . Thus

$$(2) \quad |f_n|_p \leq C_{p,r} (l_n \tau^{N-1})^{q-p}.$$

Next we estimate

$$A_p := \frac{|(R_x^p f_n)^{(k)}(y)|}{|x - y|^{p-k}}, \quad k \leq p, x \neq y, x, y \in K_N.$$

If  $|x - y| < h_n$ , then  $x, y$  belong to the same  $I_{n,j}$  for some  $j$ . Hence applying the Lagrangian form for Taylor's remainder we find  $\xi \in I_{n,j}$  such that

$$(R_x^p f_n)^{(k)}(y) = [f^{(p)}(\xi) - f^{(p)}(x)] \frac{(y - x)^{p-k}}{(p - k)!}$$

and so  $A_p \leq 2C_{p,r} (l_n \tau^{N-1})^{q-p}$ .

If  $|x - y| \geq h_n$ , then by (1), (2)

$$\begin{aligned} A_p &\leq |f_n^{(k)}(y)| \cdot |x - y|^{k-p} + \sum_{i=k}^p |f_n^{(i)}(x)| \cdot \frac{|x - y|^{i-p}}{(i - k)!} \\ &\leq C_{p,r} (l_n \cdot \tau^{N-1})^{q-p} \left[ \left( \frac{l_n \cdot \tau^{N-1}}{h_n} \right)^{p-k} + \sum_{i=k}^p \frac{1}{(i - k)!} \left( \frac{l_n \cdot \tau^{N-1}}{h_n} \right)^{p-i} \right]. \end{aligned}$$

Since  $l_n \cdot \tau^{N-1} < h_n$  we get

$$A_p \leq C_{p,r} (1 + e) (l_n \cdot \tau^{N-1})^{q-p} \quad \text{and} \quad \|f\|_p \leq C_{p,r} (2 + e) (l_n \cdot \tau^{N-1})^{q-p}.$$

Clearly

$$|f_n|_q \geq |f^{(q)}(c_1)| = q! |g'(c_1)|^q;$$

$$|g'(c_1)| = \prod_{j=2}^{N^s} (c_j - c_1) > \left(\frac{l_{n-1}}{N}\right)^{N-1} \cdot \left(\frac{l_{n-2}}{N}\right)^{N^2-N} \cdots \left(\frac{l_{n-s}}{N}\right)^{N^s-N^{s-1}}$$

$$= \tau^{N-1} \cdot N^{1-N^s}.$$

Thus,

$$|f_n|_q \geq q! N^{-r \cdot N} \cdot \tau^{(N-1)q}.$$

Analogously,  $\|f_n\|_r \leq C_r$ .

Finally, we conclude for some  $D$  independent of  $n$  that

$$\frac{\|f_n\|_p \|f_n\|_r^\epsilon}{|f_n|_q^{1+\epsilon}} \leq D l_n^{q-p} \tau^{-(N-1)(\epsilon q+p)}$$

$$= D l_{n-s}^{(\alpha_n \cdots \alpha_{n-s+1})(q-p)} [l_{n-s}^{\alpha_{n-1} \cdots \alpha_{n-s+1}} l_{n-s}^{(\alpha_{n-2} \cdots \alpha_{n-s+1})N} \cdots l_{n-s}^{N^{s-1}}]^{-(N-1)(\epsilon q+p)}.$$

Let us now show that

$$w := (\alpha_n \cdots \alpha_{n-s+1})(q-p)$$

$$- (N-1)(\epsilon q+p)[(\alpha_{n-1} \cdots \alpha_{n-s+1}) + N(\alpha_{n-2} \cdots \alpha_{n-s+1}) + \cdots + N^{s-1}]$$

which is the exponent of  $l_{n-s}$ , is positive and bounded away from zero. Indeed for the expression in the square brackets we have

$$[\cdots] \leq \alpha_{n-1} \cdots \alpha_{n-s+1} \cdot \frac{\beta}{\beta-N} \leq \alpha_n \cdots \alpha_{n-s+1} \cdot \frac{1}{\beta-N} =: w_1,$$

therefore

$$w \geq w_1 \cdot [(\beta-N)(q-p) - (N-1)(\epsilon q+p)]$$

$$= w_1 \cdot p \cdot \{M[\beta-N-\epsilon(N-1)] - \beta+1\} \geq w_1 \cdot p$$

due to the choice of  $M$  and  $\epsilon$ .

So  $w \geq \beta^s \frac{p}{\beta-N}$  from which it follows that

$$\lim_{n \rightarrow \infty} \frac{\|f_n\|_p \|f_n\|_r^\epsilon}{|f_n|_q^{1+\epsilon}} = 0.$$

Therefore the condition (b) is not fulfilled and the compact set  $K_N$  does not have the extension property.

The second statement of the theorem can be proved quite similarly to Theorem 3 in [6], so we omit it. ■

**Corollary** For a compact set  $K_N$ , let the limit  $\alpha = \lim \alpha_n$  exist and be not equal to  $N$ . Then  $K_N$  has the extension property if and only if  $\lambda_0(K_N) > 1$ .

**Remark** The set  $K_N^{(\alpha)}$ ,  $\alpha > N$ , gives us an example of a perfect set of finite class without extension property (for the definition and details see [18], [6]).

We now turn to the case  $N_n \nearrow \infty$ . The compact set  $K_\infty$  loses its Cantor-type set nature in the following sense: topologically the space  $\mathcal{E}(K_\infty)$  is closer to the space  $\mathcal{E}(K)$  with  $K$  having the form of a sequence of intervals tending to a point than to  $\mathcal{E}(K_N)$ . Compare the next statement with Theorem 2 in [7].

**Theorem 2** *The set  $K_\infty$  has the extension property if and only if there exists a constant  $M$  such that*

$$l_n \geq h_n^M, \quad \forall n.$$

**Proof** Assume the space  $\mathcal{E}(K_\infty)$  has the property DN. Let us fix  $p$  in the condition (b) of Proposition 2. For  $\epsilon = 1$  and  $q = p + 1$  we find  $r$  and  $C$  such that (b) is fulfilled.

Defining

$$f_n(x) = \begin{cases} \frac{x^q}{q!} & \text{if } x \in K \cap [0, l_n] \\ 0 & \text{otherwise} \end{cases}$$

we obtain the estimates

$$|f_n|_q \geq 1, \quad \|f_n\|_p \leq 4l_n, \quad \|f_n\|_r \leq 4h_n^{q-r}.$$

Now, Proposition 2 (b) gives  $1 \leq 16C \cdot l_n \cdot h_n^{q-r}$ , from which the necessity follows.

In order to prove the sufficiency we use a simplified version of Lemma 2 in [7] (see also [8]):

**Lemma** *Let  $K \subset \mathbb{R}$  be a compact set containing  $r + 1$  distinct points  $x_0, x_1, \dots, x_r$  such that for some  $h$  and a constant  $N$*

$$h \leq |x_i - x_j| \leq N \cdot h, \quad i, j = 0, 1, \dots, r, \quad i \neq j.$$

*Then for all  $k \leq r$  and  $f \in \mathcal{E}(K)$ ,*

$$|f^{(k)}(x_0)| \leq C \cdot h^{-k} |f|_0 + C \cdot h^{r-k} \|f\|_r,$$

*where  $C$  depends only on  $r$  and  $N$ .*

We will show that

$$(3) \quad \exists p \forall q \exists r, C : |\cdot|_q \leq Ct^{Mq} \cdot \|\cdot\|_p + \frac{C}{t^q} \|\cdot\|_r, \quad t > 0,$$

which is equivalent to (a) and hence to the property DN. The constant  $M$  here is the same as the one in the statement of the theorem.

Let  $p = 0$ . Given  $q$ , let  $r = 2q$ . Let  $n_0$  be such that  $\frac{N_n}{2} \geq r$  for  $n \geq n_0$ . Let  $t_0 = 1/l_{n_0-1}$ . Given  $t \geq t_0$ , find  $n \geq n_0$  such that

$$l_n < \frac{1}{t} \leq l_{n-1}.$$

Let  $x_0 \in K$ . Then  $x_0 \in I_{n,j_0}$  where without loss of generality we may assume that  $1 \leq j_0 \leq N_n$ . Let  $a = a_{n,r+1}$  be the left endpoint of  $I_{n,r+1}$ . Suppose that  $j_0 \leq \frac{N_n}{2}$  and consider the interval  $[x_0, x_0 + \frac{1}{2t}]$ . (If  $j_0 > \frac{N_n}{2}$ , then similar arguments apply to  $[x_0 - \frac{1}{2t}, x_0]$ .)

**Case 1:**  $\frac{1}{2t} \geq a$ . Let  $J$  be such that the interval  $[x_0, x_0 + \frac{1}{2t}]$  intersects  $J + 1$  basic intervals of the set  $E_n$ . Then  $J \geq r$  due to the choice of  $t$  and  $\frac{1}{2}(l_n + h_n) < \frac{1}{2t} < 2J(l_n + h_n)$ , as is easy to check. Then  $S := [J/r] \geq 1$ , the points  $x_i = x_0 + iS(l_n + h_n)$ ,  $i = 1, \dots, r$ , belong to  $K$  and for  $i \neq j$ ,  $|x_i - x_j| \geq S(l_n + h_n) > \frac{1}{8rt}$  since  $[\frac{1}{r}] > \frac{1}{2r}$  for  $J \geq r$ . On the other hand,  $|x_i - x_j| \leq rS(l_n + h_n) < \frac{1}{t}$ , so by the Lemma

$$|f^{(k)}(x_0)| \leq C \cdot t^k |f|_0 + C \cdot t^{k-r} \|f\|_r.$$

**Case 2:**  $\frac{1}{2t} < a$ . Now we choose the points  $x_i$ ,  $i = 1, \dots, r$  in the interval  $I_{n,j_0}$  in a similar way such that  $\frac{h_n}{4r} \leq |x_i - x_j| \leq \frac{h_n}{2}$  for  $i \neq j$ . Since  $a = r(l_n + h_n) \leq 2rh_n \leq 2r \cdot l_n^{\frac{1}{M}}$ , by the condition, we can apply the Lemma with  $h = \frac{h_n}{4r} > (4r)^{-M-1} \cdot t^{-M}$  and  $h < (4rt)^{-1}$ :

$$|f^{(k)}(x_0)| \leq C \cdot t^{kM} |f|_0 + C \cdot t^{k-r} \|f\|_r.$$

We have this estimate for any  $x_0 \in K$  and  $k \leq q$ , thus (3) is proved. ■

We observe that the logarithmic dimension is not related to the extension property of compact sets  $K_\infty$  of infinite type as seen by the following proposition and example. But in the finite case we can use it as well for isomorphic classification of corresponding spaces.

**Proposition 3** *If  $K_\infty$  has the extension property, then  $\lambda_0(K_\infty) = \infty$ .*

**Proof** If  $\overline{\lim} \alpha_n < \infty$ , then clearly  $\lim_n \lambda_n = \infty$ . So we consider the case of  $\overline{\lim} \alpha_n = \infty$ . By the previous theorem there is  $M \geq 1$  such that  $l_n \geq h_n^M$  for all  $n$ . Let  $I = \{n \in \mathbb{N} : \alpha_n \geq 2M\}$ ,  $J = \mathbb{N} \setminus I$ . Then  $\lim_{n \in J} \lambda_n = \infty$ . For  $n \in I$ ,

$$\begin{aligned} l_n^{1/\alpha_n} &= l_{n-1} \leq 2N_n h_n \leq 2N_n l_n^{1/M} \Rightarrow \frac{1}{2} l_n^{\frac{1}{\alpha_n} - \frac{1}{M}} \leq N_n \Rightarrow \log N_n \\ &\geq \frac{1}{2} \left( \frac{1}{M} - \frac{1}{\alpha_n} \right) \log \left( \frac{1}{l_n} \right) \geq \frac{1}{4M} \alpha_n \cdots \alpha_2 \log \left( \frac{1}{l_1} \right) \Rightarrow \lambda_n \\ &= \frac{\log N_n}{\log \alpha_n} \geq \frac{\log(l_1^{-1})}{4M} \cdot \frac{\alpha_n \cdots \alpha_2}{\log \alpha_n} \rightarrow \infty \text{ as } n \rightarrow \infty. \end{aligned}$$

But the converse of the above proposition is not true as the following example shows:

**Example 1** Let  $l_n = \exp(-n!)$  and  $N_n = [n^{\log n} + 1]$ . Then  $\alpha_n = n$  and  $\lambda_n \geq \log n$ . Thus  $\lambda_0 = \infty$ . But

$$h_n > \frac{l_{n-1}}{e \cdot (N_n - 1)} \geq \frac{1}{n^{\log n} e^{(n-1)!+1}} > \frac{1}{e^{\epsilon n!}} = l_n^\epsilon, \quad n \geq n_0$$

for every  $\epsilon > 0$  which means that  $K_\infty$  does not have the extension property.



### 4 Diametral Dimension of $\mathcal{E}(K)$

Approximative and diametral dimensions, introduced by Kolmogorov [9], Pełczyński [14] and Bessaga, Pełczyński and Rolewicz [1], were the first linear topological invariants applicable to isomorphic classification of nonnormed Fréchet spaces. We follow the notation of [11].

Let  $X$  be a Fréchet space with a fundamental system of neighborhoods  $(U_q)$ , let  $d_n(U_q, U_p)$  denote the  $n$ -th Kolmogorov diameter of  $U_q$  with respect to  $U_p$ . Then

$$\Gamma(X) = \{(\gamma_n)_{n=0}^\infty : \forall p \exists q : \gamma_n \cdot d_n(U_q, U_p) \rightarrow 0 \text{ as } n \rightarrow \infty\}.$$

We will consider the counting function corresponding to the diametral dimension

$$\beta(t) = \beta(U_p, U_q, t) = \min\{\dim L : t \cdot U_q \subset U_p + L\}, \quad t > 0.$$

It is easy to see that  $\beta(t) = |\{n : d_n(U_q, U_p) > \frac{1}{t}\}|$ , where  $|Z|$  denotes the cardinality of the set  $Z$ . If  $X$  is a Schwartz space and  $p, q$  are sufficiently apart, then the function  $\beta$  takes finite values. Clearly, the diametral dimension can be characterized in terms of  $\beta$  in the following way.

**Proposition 4**  $(\gamma_n) \in \Gamma(X) \Leftrightarrow \forall p \exists q : \forall C \exists n_0 : \beta(U_p, U_q, C\gamma_n) \leq n \text{ for } n \geq n_0.$

But we will directly compare asymptotic behavior of counting functions of isomorphic spaces. The proof of the following proposition is straightforward.

**Proposition 5** *If Fréchet spaces  $X$  and  $Y$  are isomorphic, then*

$$\forall p_1 \exists p \forall q \exists q_1, \epsilon > 0 : \beta_Y(V_{p_1}, V_{q_1}, \epsilon t) \leq \beta_X(U_p, U_q, t), \quad t > 0.$$

An analogous condition holds after interchanging  $\beta_X$  and  $\beta_Y$ . Here  $(V_k)_0^\infty$  is a fundamental system of neighborhoods of  $Y$ .

**Theorem 3** *Let  $X = \mathcal{E}(K)$  with  $K = K((l_n), (N_n))$ , let  $p$  and  $q, p < q$  be fixed natural numbers. If  $t \leq \frac{1}{5}l_n^{p-q}$ , then  $\beta(U_p, U_q, t) \leq (q + 1)N_1 \cdots N_n$ . If  $t \geq 5(q - p)!l_n^{p-q}$ , then  $\beta(U_p, U_q, t) \geq N_1 \cdots N_n$ .*

**Proof: Upper bound of  $\beta$**  If for some subspace  $L$  we have  $t \cdot U_q \subset U_p + L$ , then  $\beta(t) \leq \dim L$ . Let us fix  $n$  such that  $5t \leq l_n^{p-q}$ . Set  $M = N_1 \cdots N_n$ . In the union  $E_n = \bigcup_{k=1}^M I_{n,k}$  let  $I_{n,k} = [a_k, b_k]$  briefly. For  $k = 1, 2, \dots, M$  and  $j = 0, 1, \dots, q$ , let  $e_{k,j}(x) = \frac{(x-a_k)^j}{j!}$  if  $x \in K \cap I_{n,k}$  and  $e_{k,j}(x) = 0$  otherwise on  $K$ . We take  $L = \text{Span}(e_{k,j})_{k=1, j=0}^{M,q}$ . Then  $\dim L = (q + 1)M$  and it is enough to show that for any function  $f$  with  $\|f\|_q \leq t$  there exists a function  $g \in L$  such that  $\|f - g\|_p \leq 1$ . Given  $f \in tU_q$  let  $g = \sum_{k=1}^M \sum_{j=0}^q f^{(j)}(a_k) \cdot e_{k,j}$ . Clearly, if  $x \in I_{n,k}$ , then  $(f - g)(x) = R_{a_k}^q f(x)$ . Since  $|x - a_k| \leq l_n$  we get

$$(4) \quad |(f - g)^{(i)}(x)| \leq \|f\|_q \cdot |x - a_k|^{q-i} \leq t \cdot l_n^{q-i}, \quad i \leq p.$$

Let now  $A_p = \frac{|(R_y^p(f-g))^{(i)}(x)|}{|x-y|^{p-i}}$ ,  $x, y \in K, x \neq y, i \leq p$ .

If  $x, y \in I_{n,k}$  for some  $k$ , then as in [7, p. 569] we can use the following representation

$$R_y^p(f-g)(x) = R_y^q f(x) + \sum_{m=p+1}^q (R_{a_k}^q f)^{(m)}(y) \cdot \frac{(x-y)^m}{m!}.$$

Therefore

$$A_p \leq \|f\|_q \cdot |x-y|^{q-i} \cdot |x-y|^{i-p} + \sum_{m=p+1}^q \|f\|_q \cdot |y-a_k|^{q-m} \cdot \frac{|x-y|^{m-i}}{(m-i)!} \cdot |x-y|^{i-p}.$$

Here  $|x-y| \leq l_n$ , hence  $A_p \leq t \cdot l_n^{q-p} (1 + \sum \frac{1}{(m-i)!}) < (e+1)t \cdot l_n^{q-p}$ .

On the other hand, if the points  $x, y$  are situated on different basic intervals of  $E_n$ , then  $|x-y| \geq h_n \geq l_n$  by assumption. For this case

$$A_p \leq |(f-g)^{(i)}(x)| \cdot |x-y|^{i-p} + \sum_{m=i}^p \frac{|(f-g)^{(m)}(y)|}{(m-i)!} |x-y|^{m-i+i-p}.$$

From (4) it follows that  $A_p \leq t \cdot l_n^{q-p} + t \cdot l_n^{q-p} \sum_{m=i}^p \frac{1}{(m-i)!}$ . Thus,  $\|f-g\|_p \leq t \cdot l_n^{q-p} (2+e) < 1$ , by condition and this establishes the upper bound of  $\beta$ .

**Lower bound of  $\beta$**  In order to find a lower estimate for Kolmogorov diameters we use the Tikhomirov theorem [20] (see also [11, Proposition 6]): if  $d \cdot U_p \cap L \subset U_q$  with  $\dim L = n+1$ , then  $d_n(U_q, U_p) \geq d$ .

Therefore,  $\beta(t) \geq \dim L$  if  $U_p \cap L \subset (1-\epsilon_0)t \cdot U_q$  with some  $\epsilon_0 > 0$ . Let us take  $L = \text{Span}(e_{k,q})_{k=1}^M$  and fix  $f = \sum_{k=1}^M C_k \cdot e_{k,q} \in L \cap U_p$ . Since  $1 \geq \|f\|_p \geq |f^{(p)}(b_k)| \geq |C_k| \frac{l_n^{p-p}}{(q-p)!}$ , we have  $|C_k| \leq (q-p)! l_n^{p-q}$  for all  $k$ . Let  $x \in I_{n,k}$ . Then  $|f^{(i)}(x)| \leq |C_k| \frac{l_n^{q-i}}{(q-i)!}, i \leq q$ , hence  $|f|_q \leq (q-p)! l_n^{p-q}$ .

If  $x, y \in I_{n,k}$  then  $R_y^q f(x) = 0$ . Otherwise,  $|x-y| \geq h_n \geq l_n$  and arguing as before, we obtain

$$\begin{aligned} |(R_y^q f)^{(i)}(x)| \cdot |x-y|^{i-q} &\leq |f^{(i)}(x)| \cdot h_n^{i-q} + \sum_{m=i}^q \frac{f^{(m)}(y)}{(m-i)!} h_n^{m-q} \\ &\leq \frac{|C_k|}{(q-i)!} + \sum_{m=i}^q \frac{|C_{k_1}|}{(m-i)!} \leq (q-p)! l_n^{p-q} (1+e). \end{aligned}$$

If we take  $\epsilon_0$  with  $2+e < 5(1-\epsilon_0)$ , then  $\|f\|_q \leq (q-p)! l_n^{p-q} (2+e) < t \cdot (1-\epsilon_0)$  and  $\beta(t) \geq \dim L = M$ . ■

Now we can easily find the diametral dimension of  $\mathcal{E}(K)$  for concrete compact set  $K$ . In particular for classical Cantor set we have  $\beta(U_p, U_q, t) \sim t^{\frac{\log 2}{(q-p) \log 3}}$ , that is the

diametral dimension of  $\mathcal{E}(K)$  is the same as  $\Gamma(s)$ . Here and subsequently,  $F \sim G$  means that for some  $C, t_0$  we have

$$\frac{1}{C}F\left(\frac{t}{C}\right) \leq G(t) \leq C \cdot F(Ct), \quad t > t_0.$$

The most interesting case we have is for  $K = K_N^{(\alpha)}$ . Let  $\lambda_0$  denote as before the logarithmic dimension of  $K$ .

**Corollary 1** *Let  $X = \mathcal{E}(K_N^{(\alpha)})$ . Then  $\beta(U_p, U_q, t) \sim \log^{\lambda_0} t, t \rightarrow \infty$ .*

**Corollary 2**  $\Gamma(\mathcal{E}(K_N^{(\alpha)})) = \{(\gamma_n) : \gamma_n \cdot \exp(-n^{\frac{1}{\lambda_0}}) \rightarrow 0 \text{ as } n \rightarrow \infty\}$ .

**Corollary 3** *If spaces of the type  $\mathcal{E}(K_N^{(\alpha)})$  are isomorphic, then the corresponding compact sets have the same logarithmic dimension.*

**Corollary 4** *If  $\alpha < N$ , then the space  $\mathcal{E}(K_N^{(\alpha)})$  is isomorphic to a complemented subspace of  $s$ , but is not isomorphic to  $s$ .*

**Proof** Corollary 1 is the result of a simple computation. Corollary 2 follows from Proposition 4, and Corollary 3 does so from Proposition 5.

By definition,  $\mathcal{E}(K_N^{(\alpha)})$  is a quotient space of  $s$ . If  $\alpha < N$ , then according to Theorem 1, the space  $\mathcal{E}(K_N^{(\alpha)})$  has the property DN. Thus, due to Vogt’s characterization (see, e.g. [22]) it is a complemented subspace of  $s$ . But since  $\Gamma(\mathcal{E}(K_N^{(\alpha)})) \neq \Gamma(s)$ , they are not isomorphic.

## 5 Isomorphic Classification

First we shall generalize Corollary 3 to Theorem 3 for compact sets of finite type.

Let  $K_N = K((l_n), N), K_M = K((L_m), M)$ . Without loss of generality we can take  $l_1 = L_1 = 1/e$ . Then  $l_n = \exp(-\alpha_1 \cdots \alpha_n), L_m = \exp(-\alpha'_1 \cdots \alpha'_m)$ .

**Theorem 4** *Let  $X = \mathcal{E}(K_N), Y = \mathcal{E}(K_M)$ . Suppose that the corresponding limits  $\lambda_X = \lim_{n \rightarrow \infty} \frac{\log N}{\log \alpha_n}, \lambda_Y = \lim_{m \rightarrow \infty} \frac{\log M}{\log \alpha'_m}$  exist in the set of extended real numbers and are not equal. Then the spaces  $X$  and  $Y$  are not isomorphic.*

**Proof** To be definite, assume that  $\lambda_X < \lambda_Y$ . If  $\lambda_X = 0$ , then  $\alpha_n \rightarrow \infty$  and  $\alpha'_n \not\rightarrow \infty$ . By Theorem 1 in [8], it follows that the space  $Y$  has the property DN (for definition see e.g. [10]) whereas  $X$  does not. So they are not isomorphic.

Consider now the case  $0 < \lambda_X < \lambda_Y < \infty$ . Suppose, contrary to our claim, that  $X \simeq Y$ . Then by Proposition 4, for  $p_1 = 0$  there exists  $p \in \mathbb{N}$  such that for  $q = p + 1$  one can find  $q_1 \in \mathbb{N}, \epsilon > 0$  with

$$\beta_Y(V_0, V_{q_1}, \epsilon t) \leq \beta_X(U_p, U_{p+1}, t), \quad t > 0.$$

Assume  $n$  and  $m$  are large enough and are such that

$$(5) \quad 5 \cdot q_1! L_m^{-q_1} \leq \frac{\epsilon}{5} t_n^{-1}.$$

Then by taking  $t = \frac{1}{5} t_n^{-1}$  and using Theorem 3, we obtain

$$M^m \leq \beta_Y(V_0, V_{q_1}, \epsilon t) \leq \beta_X(U_p, U_{p+1}, t) \leq (p + 2)N^n.$$

Set  $\alpha = \lim \alpha_n$ ,  $\alpha' = \lim \alpha'_n$ . Clearly,  $1 < \alpha, \alpha' < \infty$ . Take  $\rho > 0$  with  $\lambda_X = (1 - 2\rho)\lambda_Y$  and  $\delta > 0$  such that

$$\frac{\log(\alpha - \delta)}{\log(\alpha' + \delta)} \cdot \frac{\log \alpha'}{\log \alpha} > 1 - \rho.$$

For this  $\delta$  one can take  $n_0, m_0$  such that  $|\alpha_n - \alpha| < \delta$ ,  $\log \frac{25q_1!}{\epsilon} \leq \frac{\alpha_1 \cdots \alpha_n}{2}$  for  $n \geq n_0$  and  $|\alpha'_m - \alpha'| < \delta$ ,  $m \geq m_0$ .

Let now

$$C_1 = \frac{1}{2q_1} \frac{\alpha_1 \cdots \alpha_{n_0}}{\alpha'_1 \cdots \alpha'_{m_0}} \frac{(\alpha' + \delta)^{m_0}}{(\alpha - \delta)^{n_0}}, \quad C_2 = \frac{\log C_1}{\log(\alpha' + \delta)},$$

$$m = m(n) = \left\lceil n \cdot \frac{\log(\alpha - \delta)}{\log(\alpha' + \delta)} + C_2 \right\rceil.$$

Then

$$(\alpha' + \delta)^m \leq C_1(\alpha - \delta)^n,$$

and so for  $n \geq n_0, m \geq m_0$  we get

$$(6) \quad \log \frac{25 \cdot q_1!}{\epsilon} + q_1 \cdot \alpha'_1 \cdots \alpha'_m \leq \alpha_1 \cdots \alpha_n.$$

This is equivalent to (5) and so we have

$$M^m \leq (p + 2)N^n \quad \text{or} \quad m \cdot \lambda_Y \cdot \log \alpha' \leq \log(p + 2) + n \cdot \lambda_X \cdot \log \alpha.$$

But this contradicts the choice of  $\delta$  and  $m(n)$ , as it is easy to check. Thus the spaces  $X, Y$  are not isomorphic.

It remains to consider the case  $\lambda'_m \rightarrow \lambda_Y = \infty, 0 < \lambda_X < \infty$ . Since  $(\alpha'_m)^{\lambda'_m} = M$  for all  $m$ , we see that there exists  $m_0$  such that  $\alpha'_m < 1 + \frac{\log 2M}{\lambda'_m}$  for  $m \geq m_0$ . Let  $\delta < \alpha - 1$  and for  $n \geq n_0$  let  $\alpha_n \geq \alpha - \delta$ . As before the inequality (6) implies the boundedness of  $\frac{m}{n}$ .

If now  $\alpha'_1 \cdots \alpha'_m \rightarrow \infty$  as  $m \rightarrow \infty$ , then we have (6) for some  $n_1$  and all  $m$ , a contradiction. Otherwise let  $C_3 = \frac{2q_1 \alpha'_1 \cdots \alpha'_{m_0}}{\alpha_1 \cdots \alpha_{n_0}} \cdot (\alpha - \delta)^{n_0}$ . For (6), it is enough to have

$$(7) \quad C_3 \cdot \left( 1 + \frac{\log 2M}{\lambda'_m} \right)^{m-m_0} \leq (\alpha - \delta)^n.$$

If now the sequence  $(m/\lambda'_m)$  is bounded, then (7) is valid for an arbitrary  $m$  and some fixed  $n$  which is a contradiction. If  $\lim_{m \rightarrow \infty} \frac{m}{\lambda'_m} = \infty$ , then we take large  $n, m$  with  $n - 1 < \frac{2 \log 2M}{\log(\alpha - \delta)} \cdot \frac{m - m_0}{\lambda'_m} \leq n$ . Here we have (7) and therefore (6) for some subsequences  $(m_j), (n_j)$  but the ratio  $\frac{m_j}{n_j}$  together with  $\lambda'_{m_j}$  are unbounded. This contradiction completes the proof. ■

The situation with compact sets of infinite type is much more complicated. By similar arguments we can distinguish now the spaces corresponding to different types. Let  $K_\infty = K((L_m), (M_m)), M_m \nearrow \infty, L_1 = 1/e$  and hence as before  $L_m = \exp(-\alpha'_1 \cdots \alpha'_m)$ . Set  $H_m = \frac{L_{m-1} - M_m \cdot L_m}{M_m - 1}$ .

**Theorem 5** *Let  $X = \mathcal{E}(K_N^{(\alpha)}), Y = \mathcal{E}(K_\infty)$  and assume the limit  $\lambda_Y = \lim_{m \rightarrow \infty} \frac{\log M_m}{\log \alpha'_m}$  exists in the set of extended real numbers. Then the spaces  $X$  and  $Y$  are not isomorphic.*

**Proof** We may assume that there exists a constant  $C_1 \geq 1$ , such that  $L_m \geq H_m^{C_1}, \forall m$ . Otherwise by Theorem 3 in [8] the space  $Y$  does not have the property DN and the result follows. Then we get

$$L_{m-1} < 2M_m \cdot L_m^{\frac{1}{C_1}}$$

and

$$(8) \quad 2M_m \geq \exp\left(\alpha'_1 \cdots \alpha'_{m-1} \left(\frac{\alpha'_m}{C_1} - 1\right)\right).$$

Suppose that the spaces  $X, Y$  are isomorphic. Arguing as above, we see that if  $m$  and  $n$  satisfy

$$(9) \quad \log \frac{25 \cdot q_1!}{\epsilon} + q_1 \cdot \alpha'_1 \cdots \alpha'_m \leq \alpha^n$$

then

$$(10) \quad M_1 \cdots M_m \leq (p + 2)N^n.$$

Consider the case  $\lambda_Y < \infty$ . Here  $\alpha'_m \rightarrow \infty$ , with  $m \rightarrow \infty$  and  $M_m < (\alpha'_m)^{2\lambda_Y}$  for large enough  $m$ . This contradicts (8).

Let now  $\lambda_Y = \infty$ . If  $\alpha'_m \rightarrow 1$ , then we take  $m = n$  in (9) and obtain a contradiction in (10) as  $M_m \nearrow \infty$ . It suffices to consider the case  $\lambda_Y = \infty, \alpha'_m \not\rightarrow 1$ . For large  $m$  let us take  $n = C_2 + \frac{\log(\alpha'_1 \cdots \alpha'_m)}{\log \alpha}$ , where  $C_2$  is fixed such that  $n \in \mathbb{N}$  satisfies the condition (9).

From (10) we deduce that

$$(\alpha'_1)^{\lambda'_1} \cdots (\alpha'_m)^{\lambda'_m} \leq (p + 2)N^{C_2} \cdot (\alpha'_1 \cdots \alpha'_m)^{\frac{\log N}{\log \alpha}},$$

which is impossible as  $\lambda'_m \rightarrow \lambda_Y = \infty, \alpha'_m \not\rightarrow 1$ . ■

**Question** Does the space  $\mathcal{E}(K_\infty)$  have a basis?

## References

- [1] C. Bessaga, A. Pełczyński and S. Rolewicz, *On Diametral approximative dimension and linear homogeneity of  $F$ -spaces*. Bull. Acad. Pol. Sci. **9**(1961), 677–683.
- [2] E. Bierstone, *Extension of Whitney-Fields from Subanalytic Sets*. Invent. Math. **46**(1978), 277–300.
- [3] K. Falconer, *Fractal Geometry, Mathematical Foundations and Applications*. John Wiley & Sons, 1990.
- [4] L. Frerick, *Extension operators for spaces of infinitely differentiable Whitney functions*. Habilitation thesis, 2001.
- [5] A. Goncharov, *A compact set without Markov's property but with an extension operator for  $C^\infty$ -functions*. Studia Math. (1) **119**(1996), 27–35.
- [6] ———, *Perfect sets of finite class without the extension property*. Studia Math. (2) **126**(1997), 161–170.
- [7] A. Goncharov and M. Kocatepe, *Isomorphic classification of the spaces of Whitney functions*. Michigan Math. J. **44**(1997), 555–577.
- [8] ———, *A continuum of pairwise nonisomorphic spaces of Whitney functions on Cantor-type sets*. Linear Topol. Spaces Complex Anal. **3**(1997), 57–64.
- [9] A. N. Kolmogorov, *On the linear dimension of topological vector spaces*. Dokl. Akad. Nauk SSSR **120**(1958), 239–341 (Russian).
- [10] R. Meise and D. Vogt, *Introduction to functional analysis*. Clarendon Press, Oxford, 1997.
- [11] B. S. Mityagin, *Approximate dimension and bases in nuclear spaces*. Russian Math. Surveys (4) **16**(1961), 59–127 (English translation).
- [12] R. Nevanlinna, *Analytic Functions*. Springer-Verlag, Berlin-New York, 1970.
- [13] W. Pawłucki and W. Pleśniak, *Extension of  $C^\infty$  functions from sets with polynomial cusps*. Studia Math. **88**(1988), 279–287.
- [14] A. Pełczyński, *On the approximation of  $S$ -spaces by finite dimensional spaces*. Bull. Acad. Polon. Sci. **5**(1957), 879–881.
- [15] Ch. Pomerenke, *Boundary Behavior of Conformal Maps*. Springer-Verlag, Berlin-Heidelberg, 1992.
- [16] E. M. Stein, *Singular integrals and differentiability properties of functions*. Princeton Univ. Press, 1970.
- [17] M. Tidten, *Fortsetzungen von  $C^\infty$ -Funktionen, welche auf einer abgeschlossenen Menge in  $\mathbb{R}$  definiert sind*. Manuscripta Math. **27**(1979), 291–312.
- [18] ———, *Kriterien für die Existenz von Ausdehnungoperatoren zu  $\mathcal{E}(K)$  für kompakte Teilmengen  $K$  von  $\mathbb{R}$* . Arch. Math. **40**(1983), 73–81.
- [19] ———, *A geometric characterization for the property (DN) of  $\mathcal{E}(K)$  for arbitrary compact subsets  $K$  of  $\mathbb{R}$* . Arch. Math. **77**(2001), 247–252.
- [20] V. M. Tikhomirov, *On  $n$ -th diameters of compact sets*. Dokl. Akad. Nauk SSSR (4) **130**(1960), 734–737.
- [21] D. Vogt, *Charakterisierung der Unterräume von  $(s)$* . Math. Z. **155**(1977), 109–117.
- [22] D. Vogt and M. J. Wagner, *Charakterisierung der Quotienträume von  $s$  und eine Vermutung von Martineau*. Studia Math. **67**(1980), 225–240.
- [23] V. P. Zahariuta, *Some linear topological invariants and isomorphisms of tensor products of scale's centers*. Izv. Severo-Kavkaz. Nauchn. Centra Vyssh. Shkoly **4**(1974), 62–64 (in Russian).

Bilkent University  
 Department of Mathematics  
 06533 Bilkent  
 Ankara  
 Turkey  
 email: barslan@firstlinux.net  
 goncha@fen.bilkent.edu.tr  
 kocatepe@fen.bilkent.edu.tr