

Our findings showed a significant increase in VAS score after COVID infection (3.82 vs 1.69; $p < 0.001$). Sixty-eight (56.2%) participants reported spontaneously fatigability after the infection and 52 (43%) reported spontaneously a deterioration in memory capacity either with or without previous memory dysfunction. Fatigability was statistically associated to cognitive impairment (55.9% vs 26.4%; $P = 0.02$). In addition, fatigability and cognitive impairment were statistically associated with pain ($P = 0.001$ and $P = 0.022$ respectively).

There was no significant association of fatigability nor cognitive impairment with the gender of the survivors.

Conclusions: The clinician should keep in mind to screen for possible somatic or psychological distress, in particular pain, fatigability and cognitive impairment even after resolution of the COVID infection, in order to guarantee a better quality of life.

Disclosure of Interest: None Declared

EPP0675

Impact of the COVID-19 pandemic on the mental health of hospital employees: single center experience

M. Kristic¹, V. Kuresic¹, I. Miskulin^{1*}, J. Dumic², L. Dumic¹ and M. Miskulin¹

¹Department of Public Health, Faculty of Medicine Osijek and

²Department of Dental Medicine, Faculty of Dental Medicine and Health Osijek, Osijek, Croatia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.766

Introduction: Hospital employees are at high risk of developing mental health issues during the coronavirus (COVID-19) pandemic. Indeed, several studies have shown increased rates of anxiety, depression, stress, and other mental health issues but existing studies show inconsistencies, and each country has some local specificities.

Objectives: This study aimed to investigate the influence of the COVID-19 pandemic on various aspects of the mental health of hospital employees (health workers and non-health workers) from Croatia.

Methods: This cross-sectional questionnaire study was conducted from February to April 2023 period. A validated, anonymous questionnaire that contained questions regarding demographic data, as well as the Pittsburgh Sleep Quality Index (PSQI), the Zung Self-Rating Anxiety Scale, and the Zung Self-Rating Depression Scale was self-administered to a convenient sample of hospital employees from one general hospital in northwestern Croatia.

Results: The study sample included 360 subjects with a median age of 42 years (interquartile range 35-50), 24.7% males, and 75.3% females. According to the PSQI, 21.1% of subjects presented sleep disturbances. According to the Zung Self-Rating Anxiety Scale, there were 39.4% of subjects with anxiety while according to the Zung Self-Rating Depression Scale, there were 6.4% of subjects with depression. Sleep disturbances were more frequent among subjects who considered their socioeconomic status as under average ($p = 0.040$), and among health workers in comparison to non-health workers employed in hospital ($p = 0.040$). Anxiety was more frequent among females ($p = 0.010$), and subjects with lower levels of education (only elementary school) ($p = 0.040$). Depression was more frequent among females ($p = 0.030$).

Conclusions: The COVID-19 pandemic has a significant negative influence on the mental health of hospital employees where health workers in comparison to non-health workers, females, subjects with lower levels of education, and subjects who considered their socioeconomic status as under average are more prone to the development of investigated mental health issues. The development of appropriate supportive programs that enhance the mental health of all hospital employees during pandemics is needed to address mental health issues in this vulnerable population.

Disclosure of Interest: None Declared

Post-Traumatic Stress Disorder

EPP0678

Understanding the Traumatic Impact of Serious Chronic Illness

M. Theodoratou^{1*} and H. Diamanti¹

¹Social Sciences, Hellenic Open University, Patras, Greece

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.767

Introduction: The diagnosis of a severe chronic illness represents a deeply impactful traumatic event, frequently giving rise to initial adverse consequences that can manifest as post-traumatic stress. The duration and characteristics of these effects exhibit considerable variation among individuals.

Objectives: This study aims to explore the levels of post-traumatic stress, post-traumatic growth, and psychosocial adaptation among individuals coping with chronic diseases.

Methods: This cross-sectional study involved 92 participants with chronic illnesses, recruited through convenience and snowball sampling. Data collection utilized an online questionnaire that included both demographic questions to provide a comprehensive understanding of participants' experiences, as well as psychometric scales for measuring post-traumatic stress, post-traumatic growth, and psychosocial adaptation.

Instruments used :

1. PTSD Checklist for DSM-5 (PCL-5)
2. Posttraumatic Growth Inventory (PTGI) and Tedeschi and Calhoun Posttraumatic Growth Inventory (TCGI).
3. Psychosocial Adjustment to Illness Scale (PAIS).

Analysis included descriptive statistics and inductive analysis using SPSS ($p < 0.05$). Ethical considerations were observed, with informed consent and data confidentiality.

Results: The study revealed the presence of low to moderate levels of post-traumatic stress ($M = 2.45$), moderate levels of post-traumatic growth ($M = 2.90$), and moderate levels of psychosocial adaptation in various aspects of participants' lives, including work ($M = 2.36$), sexuality ($M = 2.11$), sociability ($M = 2.28$), relationships with partners and family members ($M = 1.92$), and perception of their health ($M = 1.94$). Furthermore, the overall psychosocial situation of the participants was found to range from low to moderate ($M = 2.48$). Notably, individuals with fewer chronic illnesses tended to experience lower levels of post-traumatic stress and exhibited less adaptation in their work. Additionally, higher levels of post-traumatic growth were observed in women and patients with higher educational backgrounds. The analysis revealed a positive and statistically