

phagocytic cell types contain Mtb is critical. **METHODS/STUDY POPULATION:** To determine the impact T cells have on different phagocyte cell populations' host defense mechanisms, groups of wild-type and T cell deficient TCRA-/- mice were infected with an Mtb strain expressing fluorescent mScarlet protein. At four weeks post-infection, a time when T cell help contributes to control of Mtb, lungs were homogenized and cells sorted based on detection of mScarlet, indicating Mtb-infected cells. Cell suspensions from each mouse background were underwent single-cell RNA sequencing analysis to reveal the heterogenous cellular transcriptional response of different phagocyte populations. **RESULTS/ANTICIPATED RESULTS:** We found that Mtb-infected phagocytes from wild-type and TCRA-/- mouse lungs contain the same dominant cell phenotypic clusters, but these have different patterns of gene expression. Without T cells, phagocytes are prone to a more inflammatory phenotype. **DISCUSSION/SIGNIFICANCE:** This will translate fundamental biological data to test the hypothesis that Mtb encounters different environmental stresses exerted by different phagocytic cell types. This work could reveal host intracellular niches that enable bacterial persistence and elucidate new pathways that could be targeted for traditional antibiotic therapies for TB.

493

Prevalence and Clinical Presentation of Chronic Neck Pain in Individuals with Generalized Joint Hypermobility

Rebecca Abbott, Paula Ludewig, Victor Barocas, Arin M Ellingson
University of Minnesota

OBJECTIVES/GOALS: Evidence suggests that individuals with generalized joint hypermobility (GJH), or excessive joint range of motion, are at higher risk of developing chronic neck pain. The objective of this study is to determine the prevalence and clinical presentation of chronic neck pain in GJH and investigate its associations with other measures of spine health. **METHODS/STUDY POPULATION:** Data was collected at the Driven to Discover Research Facility at the 2022 Minnesota State Fair. Individuals 18 years and older were invited to participate. All enrolled participants completed Phase 1, which included: the Beighton Score (measure of GJH), the 5-Point-Questionnaire (self-report survey for current or historical GJH), and a custom self-report survey for demographics and musculoskeletal pain. A subset of participants was also asked to complete Phase 2 of the study. Phase 2 consisted of additional self-report surveys (Neck Disability Index (NDI) and PROMIS-10 Global) and the following physical measures: neck range of motion in all planes, neck strength in flexion-extension and lateral bending, and grip strength. **RESULTS/ANTICIPATED RESULTS:** A total of 559 participants were enrolled in the study. All participants completed Phase 1, and 285 of those individuals completed Phase 2. Those with a Beighton Score ≥ 4 were categorized as having GJH. The overall prevalence of GJH was 23.8% for females and 9.1% for males. Consistent with previous studies, multiple linear regression analysis ($R^2=0.20$, $F(2,552) = 69.37$, p **DISCUSSION/SIGNIFICANCE:** This is one of the largest studies investigating GJH, pain, and physical measures of neck function in the general population. The results highlight the higher prevalence of chronic neck pain in those with GJH and will form the basis for a subsequent study to identify mechanisms and potential therapeutic targets for individuals with GJH and chronic pain.

495

Radon and Fracking Exposures and Lymphoma Risk in a Canine Model of non-Hodgkin Lymphoma*

Ashleigh Tindle, Lauren Trepanier
University of Wisconsin-Madison

OBJECTIVES/GOALS: The objective of this study was to determine whether residential radon and proximity to horizontal oil and drilling (fracking) are risk factors for the development of multicentric lymphoma in pet dogs, a spontaneous, immunocompetent model for non-occupational risk for NHL in humans. **METHODS/STUDY POPULATION:** Two case-control populations of dogs with multicentric lymphoma were utilized, with a focus on two dog breeds at high risk for lymphoma. Control dogs were matched for age, breed, and sex. Home addresses were collected for 54 Golden retrievers with lymphoma and 108 Golden retriever controls, and for 56 boxer dogs with lymphoma and 84 unaffected boxer controls. Counties of residence were matched to radon zones and percentage of home radon tests that exceeded the actionable level of 4 pCi/L, available by county through the EPA and the CDC National Environmental Public Health Tracking Network from 2008 to 2017. Locations of horizontal oil and gas wells were obtained from the Enverus Database, and distances from dog homes to the closest well, and well density by county, were calculated for each case and control. **RESULTS/ANTICIPATED RESULTS:** We found no significant differences in radon zones, county level radon measurements, or residential proximity to active fracking wells between dogs with lymphoma and unaffected controls in either the Golden Retriever or boxer populations. **DISCUSSION/SIGNIFICANCE:** Canine multicentric lymphoma resembles human NHL and is a valuable model of non-occupational environmental risk for NHL in people. Although we did not find geographic associations between radon and fracking wells, follow-up studies will measure household radon, as well as household air, water, and dog urine for potentially genotoxic chemicals.

498

Structural Determination of the CqsR CACHE Domain and its Autoinducer*[†]

Andrew Guarnaccia¹, Wai-Leung Ng², Anjali Steenhaut², Sandra Olenic², Lark J. Perez³, Matthew B. Neiditch¹

¹Department of Microbiology, Biochemistry, and Molecular Genetics, New Jersey Medical School, Rutgers, ²Department of Molecular Biology and Microbiology, Tufts University School of Medicine, ³Department of Chemistry & Biochemistry, Rowan University

OBJECTIVES/GOALS: Our goal is to determine the structure of the CACHE domain of the *Vibrio cholerae* quorum sensing receptor CqsR as well as its autoinducer (AI). We are performing X-ray crystallography on the protein in its apo form, with the fractions containing the AI, and with known ligand ethanolamine (ETA). **METHODS/STUDY POPULATION:** We have transformed BL21(DE3) *E. coli* cells with a pTB146 vector to contain the gene for the CqsR CACHE domain. We grow these cells to high optical density and induce protein expression, at which point we harvest them and purify the protein. This entails lysing the cells, separating the protein with Ni-NTA resin, cleaving our protein tag, and column chromatography. With purified protein, high-throughput screens are set up to find crystallization conditions of apo CqsR, CqsR-ETA, and CqsR-AI. We then determine conditions that best lead

[†]Wai-Leung Ng's name has been corrected. Additionally, middle initials for two authors have been added and the affiliations have been corrected. A corrigendum detailing these changes has been published (doi:10.1017/cts.2023.551).

to crystal formation and optimize them. Crystals are then diffracted with X-rays, process the data, and determine the structure of protein and AI. RESULTS/ANTICIPATED RESULTS: We anticipate finding the structure of the CqsR CACHE domain to a high resolution in addition to the identity of its autoinducer. Previous results found that the structure is homologous to another *V. cholerae* chemoreceptor, Mlp37, and we expect the results from this project to confirm this. In addition, we know that the autoinducer weighs approximately 62 daltons, the same as the known ligand, ethanolamine. Given that CACHE domains bind specifically to their ligands, we anticipate that the autoinducer will be structurally similar to ethanolamine. DISCUSSION/SIGNIFICANCE: The results will reveal the structure of the CqsR CACHE domain and its autoinducer. This knowledge will better allow researchers to treat cholera, as both autoinducer identity and receptor conformational changes will be uncovered, allowing for drug development to inhibit cell growth.

500

The Aging Exposome: Characterizing Bidirectional Effects of Exposures and Aging

Ram Gouripeddi, Caden Stewart, Julio Facelli
University of Utah

OBJECTIVES/GOALS: The objective of this study is to synthetically generate and use records of exposure, and so that we can understand the effects of exposure on aging and vice-versa. METHODS/STUDY POPULATION: Quantifying bidirectional effects of environment and aging requires time series of data from all contributing exposures which can span endogenous processes within the body, biological responses of adaptation to environment, and socio-behavioral factors. Gaps in measured data may need to be filled with computationally modeled data. Essentially, the challenge in generating aging exposome is the absence of readily available records for individuals over the course of their life. Instead, these would need to be assimilated from historic person reported data (e.g. residential location, durations, behaviors) along with publically available data. This could lead to potential gaps and uncertainties that would need inform on how the exposomic records can be used for aging research. RESULTS/ANTICIPATED RESULTS: We present a pragmatic approach to generation of longitudinal exposomic and aging records as required for different study archetypes. Such records can then be used to understand the bidirectional effects of exposures and aging. DISCUSSION/SIGNIFICANCE: Effects of a lifetime of environmental and lifestyle exposures on aging or age-associated diseases are not well understood. Characterizing differential, additive and intense sporadic multi-agent exposures require advanced big data and artificial intelligence methods.

502

The Prospective Geriatric Patient Reported Outcomes (GERI-PRO) Study: Understanding Post-Traumatic Injury Recovery from the Patient's Perspective

Mira Ghneim, Deborah Stein

R Adams Cowley Shock Trauma Center. The University of Maryland School of Medicine

OBJECTIVES/GOALS: Patient reported outcomes (PROs) provide unique insight to the patients experience with their healthcare related quality of life QoL. This study aims to 1. Characterize geriatric

trauma patients'(GTPs) perceived QoL, at time of injury vs. 3- and 6-months post-injury. 2. Introduce and validate a PROs tool, known as the Five Favorite Activities. METHODS/STUDY POPULATION: This is a prospective cohort study of older adults (≥ 65) presenting to our trauma center with mild traumatic brain injury and/or mild spine, thoracic or extremity fractures. Participants will be asked to complete the NIH-validated Patient-Reported Outcome Measure Information System (PROMIS)-29, PROMIS Cognitive and Functional Abilities, Life-Space Levels and Five Favorite Activities assessment (a list of the five favorite overall and daily activities) tools. Cognitive function will be measured using Montreal Cognitive Assessment tool. Physical function will be evaluated using the Activity Measure for Post-Acute Care 6-click tool. Patients will be contacted at 3- and 6- months post discharge and asked to complete the assessment tools listed above to evaluate changes in QoL during the recovery process. RESULTS/ANTICIPATED RESULTS: We hypothesize that geriatric trauma patients will experience a decline in QoL, physical and cognitive function post-injury. This decline will be associated with a decrease in return to the ability to participate in their pre-injury Five Favorite Activities. DISCUSSION/SIGNIFICANCE: First, this study is one of the first to evaluate PROMs in GTPs. Second, the Five Favorite Activities PROM, will provide a unique, direct and individualized characterization of what GTPs find important to their recovery post injury compared to the current generic PROMs. This information can be utilized in the future to align goal of care with expectations

504

Topical adenosine treatment inhibits inflammation and mucus production in viral acute rhinosinusitis

Xiaoyang Hua¹, Kody A. Waldstein², Maria Ganama³, Steven M. Varga⁴, Stephen Tilley⁵, Xiaoyang Hua³

¹Department of Otolaryngology-Head and Neck Surgery, University of Iowa ²Interdisciplinary Graduate Program in Immunology, University of Iowa, Iowa City, IA 52242, USA ³Department of Otolaryngology-Head and Neck Surgery, University of Iowa, Iowa City, IA 52242, USA ⁴Department of Pathology, University of Iowa, Iowa City, IA 52242, USA ⁵Department of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC 27514

OBJECTIVES/GOALS: Viral acute rhinosinusitis (ARS), a.k.a, the common cold, affects millions every year. The symptoms caused by viral ARS dramatically affect the general well-being and functional levels of patients, causing work and school absence, and antibiotic abuse. In this study, we examined the therapeutic potential of topical adenosine in viral ARS METHODS/STUDY POPULATION: Rhinosinusitis was induced in WT and adenosine receptor (AR) knockout mice by respiratory syncytial virus (RSV) infection in the upper airways. Mice were subjected to adenosine or vehicle control within the sinuses. Adenosine receptor expression, inflammatory cytokine expression, and histologic mucus and inflammation score was assessed. The effect of endogenous adenosine accumulation within the sino-nasal tract was assessed in adenosine deaminase knockout (ADA^{-/-}) mice. RESULTS/ANTICIPATED RESULTS: Topical administration of adenosine significantly inhibited the expression of pro-inflammatory cytokines, mucus production, and cell damage in the nose of mice with viral ARS, without prolonging virus clearance. This inhibitory effect was primarily mediated by the A2A adenosine receptor (AR). We also examined and compared the