

The Summer Meeting of the Nutrition Society was held at the University of Glasgow on 29 June – 2 July 1999

## Plenary Lecture

### Factors affecting newborn bone mineral content: *in utero* effects on newborn bone mineralization

Ran Namgung<sup>1</sup> and Reginald C. Tsang<sup>2\*</sup>

<sup>1</sup>Yonsei University College of Medicine, Department of Pediatrics, 134 Shinochon-Dong, Sudaemoon-Ku, Seoul 120-752, Korea

<sup>2</sup>University of Cincinnati Children's Hospital Medical Centre, Division of Neonatology, 231 Bethesda Avenue, Cincinnati, OH 45267-0541, USA

Several factors have been found recently to have a significant impact on newborn bone mineral content (BMC) and developing fetal bone. Recently we showed that maternal vitamin D deficiency may affect fetal bone mineralization. Korean winter-born newborn infants had extremely low serum 25-hydroxyvitamin D (25-OHD), high serum cross-linked carboxy-terminal telopeptide of type I collagen (ICTP; a bone resorption marker), and markedly lower (8 %) total body BMC than summer-born newborn infants. Infant total body BMC was positively correlated with cord serum 25-OHD and inversely correlated with ICTP, which was also negatively correlated with vitamin D status. In three separate studies on North American neonates we found markedly lower (8–12 %) BMC in summer newborn infants compared with winter newborn infants, the opposite of the findings for Korean neonates. The major reason for the conflicting BMC results might be the markedly different maternal vitamin D status of the North American and Korean subjects. Recently, we found evidence of decreased bone formation rates in infants who were small-for-gestational age (SGA) compared with infants who were appropriate-for-gestational age; we reported reduced BMC, cord serum osteocalcin (a marker of bone formation) and 1,25-dihydroxyvitamin D (the active metabolite of vitamin D), but no alterations in indices of fetal bone collagen metabolism. In theory, reduced utero-placental blood flow in SGA infants may result in reduced transplacental mineral supply and reduced fetal bone formation. Infants of diabetic mothers (IDM) have low BMC at birth, and infant BMC correlated inversely with poor control of diabetes in the mother, specifically first trimester maternal mean capillary blood glucose concentration, implying that factors early in pregnancy might have an effect on fetal BMC. The low BMC in IDM may be related to the decreased transplacental mineral transfer. Cord serum ICTP concentrations were higher in IDM than in control subjects, implying increased intrauterine bone resorption. BMC is consistently increased with increasing body weight and length in infants. Race and gender differences in BMC appear in early life, but not at birth. Ethanol consumption and smoking by the mother during pregnancy affect fetal skeletal development.

**Bone mineral content: Fetal development: Vitamin D: Birth weight:  
Maternal diabetes**

Fetal bone mineralization is affected by season of birth (Namgung *et al.*, 1992, 1993, 1994, 1998), low weight relative to gestation (Minton *et al.* 1983; Petersen *et al.* 1989; Pohlandt & Mathers, 1989; Namgung *et al.* 1993) and diabetes in the mother (Mimouni *et al.* 1988). In theory,

fetal bone mineralization is determined by placental mineral transfer, fetal bone mineral accretion and fetal bone resorption. Indices of fetal bone turnover appear to be affected by gestation (immaturity) and compromised fetal growth (Mora *et al.* 1997; Harrast & Kalkwarf, 1998; Ogueh

**Abbreviations:** AGA, appropriate-for-gestational age; BMC, bone mineral content; BMD, bone mineral density; ICTP, cross-linked carboxy-terminal telopeptide of type I collagen; IDM, infants of diabetic mothers; 25-OHD, 25-hydroxyvitamin D; 1,25-(OH)<sub>2</sub>D, 1,25-dihydroxyvitamin D; PICP, carboxy-terminal propeptide of type I collagen; PTH, parathyroid hormone; SGA, small-for-gestational age.

\*Corresponding author: Professor R. C. Tsang, fax +1 513 558 7770, email tsangrc@email.uc.edu

*et al.* 1998). Since effects on fetal bone are often probably primarily brought about through the mother, any disturbance in maternal mineral metabolism (i.e. maternal nutrient deficiency or diseases) or conditions that affect placental mineral transfer could be factors affecting fetal bone mineralization. For example, we have shown recently that maternal vitamin D deficiency during pregnancy has a major impact on fetal bone mineralization (Namgung *et al.* 1998). In the present review factors associated with alterations in neonatal bone mineralization are summarized.


### Season

Fetal bone metabolism, in theory, can be influenced by season through changes in vitamin D metabolism. Vitamin D is essential for mineral homeostasis and adequate bone mineralization. A lack of vitamin D during pregnancy may adversely affect maternal mineral metabolism, which in turn could affect the fetus, since normal fetal mineral accretion is thought to be derived from the maternal skeleton (Comar, 1956). Maternal vitamin D status may influence placental Ca transfer (Durand *et al.* 1983*b*); in rats, maternal administration of the synthetic analogue 1- $\alpha$  hydroxycholecalciferol (a highly bioactive form of vitamin D) resulted in increased fetal Ca content (Durand *et al.* 1983*a*). In the UK higher newborn cord serum parathyroid hormone (PTH) levels have been found in vitamin D-deficient Asian Indians than in white subjects, which is thought to be related to a reduction in placental Ca transport, leading to high fetal PTH levels (Okonofua *et al.* 1987).

Seasonal variations in maternal vitamin D status and their effects on the fetus have been reported. Earlier studies have shown seasonal variations in serum 25-hydroxyvitamin D (25-OHD; the major vitamin D metabolite in blood) concentrations in both pregnant women and their newborn infants, with low values in winter (Kuroda *et al.* 1981; Verity *et al.* 1981). Severe vitamin D deficiency in pregnancy can lead to intrauterine growth retardation (Brooke *et al.* 1980), adverse effects on the appearance of neonatal ossification centres (Specker *et al.* 1992) and congenital rickets (Moncrieff & Fadahunsi, 1974), confirming the potential impact of maternal vitamin D deficiency on fetal bone development.

The effect of maternal vitamin D deficiency on fetal bone mineralization (BMC) and metabolism (bone turnover markers) has not been studied previously. We suggested that seasonally-related changes in maternal vitamin D metabolism may affect fetal bone metabolism (Namgung *et al.* 1992, 1993, 1994, 1998). In vitamin D deficiency bone mineralization is low and bone resorption is thought to be high (Ooms *et al.* 1995; Scharla *et al.* 1996; Peacock *et al.* 1998). Thus, we suggested that BMC would be lower and bone turnover markers higher in winter-born infants than in summer-born infants, possibly related to increased bone turnover. We tested our hypothesis in four separate studies; one in Korean neonates (Namgung *et al.* 1998) and three in American neonates (Namgung *et al.* 1992, 1993, 1994).


In our recent study Korean winter newborn infants had markedly lower ( $P=0.0001$ ) serum 25-OHD levels (97 % below normal,  $< 11$  ng/ml), significantly higher ( $P=0.0002$ ) concentrations of cross-linked carboxy-terminal telopeptide



**Fig. 1.** Total body bone mineral content (TBBMC; g) of Korean newborn infants in winter and summer in a population without routine vitamin D supplementation. Mean value for TBBMC winter-born newborn infants was significantly lower than that for summer-born newborn infants: ( $*P=0.0002$ ). (Reproduced with permission from Namgung *et al.* 1998.)

of type I collagen (ICTP; a bone resorption marker), and markedly lower (8 %;  $P=0.0002$ ) total body BMC than summer newborn infants (Fig. 1). Infant total body BMC was positively correlated with cord serum 25-OHD ( $r=0.24$ ,  $P=0.047$ ) and negatively correlated with ICTP ( $r=-0.33$ ,  $P=0.008$ ), which was also negatively correlated with vitamin D status ( $r=-0.39$ ,  $P=0.001$ ; Fig. 2; Namgung *et al.* 1998). The finding of low total body BMC in Korean winter newborn infants is presumably related to poor materno-fetal vitamin D status, and is consistent with bone mineral alterations (high bone resorption and low BMC) related to seasonally-induced changes in vitamin D status. The seasonal effect (8 % differential) on BMC is markedly greater than the seasonal difference (less than 2 %) in BMC or bone mineral density (BMD) of adults (Aitken *et al.* 1973; Krølner 1983; Bergstralh *et al.* 1990). In Korean winter newborn infants cord serum 1,25-dihydroxyvitamin D (1,25-(OH) $_2$ D; the most bioactive form of vitamin D) concentrations were lower than those of summer newborn infants, and was correlated with 25-OHD. Production of 1,25-(OH) $_2$ D is substrate (25-OHD)-dependent, and serum 1,25-(OH) $_2$ D may be correlated with 25-OHD in vitamin D deficiency (Bouillon *et al.* 1987). Low 1,25-(OH) $_2$ D levels in Korean winter newborn infants may reflect decreased synthesis in the fetal kidney associated with limited substrate (25-OHD) availability, and theoretically may be a factor affecting placental Ca transfer and fetal bone mineralization. Cord serum Ca concentration was significantly lower in winter than in summer, but cord serum P, Mg, and PTH concentrations were similar.

However, in three separate studies carried out with North American newborn infants we found markedly lower (8–12 %) BMC in summer newborn infants compared with winter newborn infants from Cincinnati, OH, USA, both for appropriate-for-gestational-age (AGA) infants (Fig. 3) and for small-for-gestational-age (SGA) infants (Fig. 4; Namgung *et al.* 1992, 1993, 1994), which is the opposite of the findings for Korean neonates, and for adults (Aitken




**Fig. 2.** (A) Relationship between total body bone mineral content (TBBMC) and cord serum 25-hydroxyvitamin D (25-OHD; ng/ml).  $TBBMC = 87.256 + 0.425 (25-OHD)$ ;  $r = 0.24$ ,  $P = 0.047$ . (B) Relationship between TBBMC and serum cross-linked carboxy-terminal telopeptide of type I collagen (ICTP; a bone resorption marker; ng/ml).  $TBBMC = 103.74 - 0.149 (ICTP)$ ;  $r = -0.33$ ,  $P = 0.008$ . (C) Relationship between serum ICTP and 25-OHD.  $ICTP = 98.3 - 1.5 (25-OHD)$ ;  $r = -0.39$ ,  $P = 0.001$ . (Reproduced with permission from Namgung *et al.* 1998.)

*et al.* 1973; Krølner *et al.* 1983; Bergstralh *et al.* 1990). American summer-born newborn infants had higher cord serum osteocalcin (a bone turnover marker) and 1,25-(OH)<sub>2</sub>D concentrations than American winter-born newborn infants. Osteocalcin is the most abundant non-collagenous bone matrix protein, synthesized by osteoblasts, and reflects osteoblastic activity. Theoretically, the high serum osteocalcin concentrations in summer newborn infants may be an adaptive response to the lower BMC of these infants, possibly secondary to the raised plasma 1,25-(OH)<sub>2</sub>D concentrations. Cord serum Ca concentration was lower in summer than in winter, but P, Mg, ionized Ca, PTH and 25-OHD concentrations were similar.


Based on these findings in North American and Korean neonates, we suggested that reduced bone mass in North American summer-born newborn infants is related to reduced maternal vitamin D status in the preceding winter, i.e. 6 months before summer birth. We speculated that sunshine deprivation in winter alters vitamin D metabolism

in the mother and fetus during an important period of early fetal skeletal development, resulting in low BMC at birth 6 months later.

The major reason for this discrepancy (conflicting BMC results in American neonates *v.* Korean neonates) might be the markedly different vitamin D status in North America compared with Korea. During the first trimester of pregnancy most women in the USA have not received much prenatal care and are less likely to be taking multivitamin pills that contain vitamin D. Thus, the mother is relatively 'unprotected' (i.e. not given vitamin D supplements), and seasonal changes in vitamin D status can occur. We suggest that these changes in seasonal vitamin D status result in first-trimester bone alterations in winter that are evident 6 months later as reduced bone mass at birth in summer compared with winter. After the first trimester most women receive vitamin D supplements. At the time of delivery in North America there are no differences in serum vitamin D concentrations according to season, presumably because of vitamin D supplementation (Namgung *et al.* 1994). In



**Fig. 3.** Bone mineral content (BMC; mg/mm) of newborn infants in winter and summer by (A) race (□, white; ▨, black) and (B) gender (□, male; ▨, female) in a population where vitamin D fortification is practiced (Cincinnati, OH, USA). Values are least square means adjusted for birth weight with their standard errors represented by vertical bars for nos. of newborn infants shown in parentheses. Summer-born infants had significantly lower BMC than winter-born infants ( $P=0.009$ ). Race and gender differences in BMC corresponded to birth weight differences; after adjusting for birth weight, race and gender, differences were  $P=0.09$  and  $P=0.17$  respectively. Seasonal differences in BMC remained significant after adjusting for race and gender effects ( $P=0.002$  and  $P=0.001$  respectively). (Reproduced with permission from Namgung *et al.* 1994.)



**Fig. 4.** Bone mineral content (BMC; mg/mm) in newborn infants for small-for-gestational-age (SGA) v. appropriate-for-gestational-age (AGA) infants by season of birth (□, winter; ▨, spring; ■, summer) in Cincinnati, OH, USA. The BMC was significantly lower for SGA than for AGA infants ( $P=0.0001$ ), and differed by season of birth; spring or summer BMC values were lower than winter values in both SGA and AGA infants ( $P=0.003$ ). The BMC differences between groups remained significant after adjusting for season ( $P=0.0001$ ). Values are means with their standard errors represented by vertical bars for nos. of infants shown in parentheses. (Reproduced with permission from Namgung *et al.* 1993.)

contrast, Koreans are often vitamin D deficient throughout pregnancy, and there is little vitamin D supplementation in practice. Korean winter-born neonates had cord serum 25-OHD concentrations in the vitamin D-deficient range (4.3 (SD 3) ng/ml). Of winter-born neonates, 97 % have subnormal vitamin D levels, and thus significant negative effects on fetal bone in winter might be expected. The period of greatest vulnerability of the fetus to altered maternal vitamin D status, if any, is unclear. However, from our studies it could happen either in the early and/or late trimester of pregnancy, depending on the maternal vitamin D status.

In adults the BMC of forearm and BMD of lumbar spine (Aitken *et al.* 1973; Krølner *et al.* 1983; Bergstralh *et al.* 1990) differ according to the season, and are related to seasonal variations in vitamin D status (Dawson-Hughes *et al.* 1991; Sherman *et al.* 1992; Ooms *et al.* 1995). These seasonal differences are, however, much lower (< 2 %) than those (8–12 %) we have found in newborn infants (Namgung *et al.* 1992, 1993, 1994, 1998). Suboptimal vitamin D status in the elderly is associated with high serum PTH concentrations (Orwoll & Meier, 1986; Lukert *et al.* 1987; Dawson-Hughes *et al.* 1991; Ooms *et al.* 1995), osteopenia (Orwoll & Meier, 1986; Fonsca *et al.* 1988), and low BMD of radius or hip (Sherman *et al.* 1992; Ooms *et al.* 1995). In elderly women (>60 years) in North America (Peacock *et al.* 1998) and in post-menopausal women in Germany (Scharla *et al.* 1996) serum 25-OHD levels were negatively correlated with urinary deoxypyridinoline excretion (an index of bone resorption), and positively related to BMD of the proximal femur, consistent with our findings in Korean neonates. Biochemical and hormonal markers of bone metabolism vary significantly ( $P<0.05$ ) according to the season, consistent with an acceleration of bone turnover and possible subclinical vitamin D deficiency in winter (Woitge *et al.* 1998). In young women a reduction in serum 25-OHD was associated with decreased intestinal Ca and P absorption during wintertime, but was not associated with the changes in bone turnover markers (serum carboxy-terminal propeptide of type I procollagen (PICP) and urinary deoxypyridinoline; Zittermann *et al.* 1998).

### Small-for-gestational age

Low weight relative to gestation (SGA or intrauterine growth retardation) could be a factor influencing fetal bone metabolism, possibly through alterations in placental mineral transfer. In a previous study from our group using *in situ* perfused placentas of intrauterine growth-retarded rat fetuses (modified Wigglesworth model in the rat; Wigglesworth, 1964; Kollee *et al.* 1979), materno-fetal Ca transfer was reduced across the placentas of growth-retarded



*v.* control fetuses, and the mean total Ca content of growth-retarded fetuses was significantly lower than that of control fetuses (Mughal *et al.* 1989). Presumably the reduced placental blood supply that causes experimental fetal growth retardation also causes reduced placental Ca transport proportional to the reduction in body size. In an earlier study from our group (Minton *et al.* 1983) we found that the BMC of the distal radius (measured using single-photon absorptiometry) was significantly lower ( $P < 0.01$ ) and serum PTH concentrations were higher ( $P < 0.01$ ) for term SGA infants than for term AGA infants at birth. We suggested that the SGA fetus responds to possible *in utero* Ca deficiency by increasing the secretion of PTH.

In a recent study (Namgung *et al.* 1993) we found evidence of decreased bone formation, low BMC and low serum osteocalcin concentrations, and low serum 1,25-(OH)<sub>2</sub>D concentrations in term SGA infants compared with term AGA infants. There were no differences in cord serum Ca, P, Mg, 25-OHD and PTH concentrations between SGA and AGA infants; BMC was inversely correlated with cord serum PTH ( $r = -0.228$ ,  $P = 0.006$ ). We suggested that low 1,25-(OH)<sub>2</sub>D in SGA infants might be related to a reduction in fetal-placental production of 1,25-(OH)<sub>2</sub>D, possibly due to reduced utero-placental blood flow in SGA infants, resulting also in low serum osteocalcin (an index of bone formation) and low BMC. Fetal serum PTH values may be relatively increased because of a reduction in placental mineral supply. The finding of low cord serum osteocalcin in SGA *v.* AGA infants supports the concept that decreased fetal bone formation may lead to reduced BMC in SGA infants.

Chunga Vega *et al.* (1996) have recently examined whether low cord serum Zn concentrations in SGA infants are associated with fetal growth retardation and bone mineral alteration, related to low 1,25-(OH)<sub>2</sub>D. They reported that BMD, measured by dual-energy X-ray absorptiometry, was lower (by 20 %) in term SGA *v.* AGA infants, and linearly related to growth and nutritional measures (weight, length, head circumference, mid-arm circumference and surface area). Cord serum Zn concentrations were in the normal range and similar for SGA and AGA infants. Cord serum PTH, 25-OHD, 1,25-(OH)<sub>2</sub>D, osteocalcin, total Ca and inorganic P concentrations were not different between groups. They speculated that low BMD in SGA infants could be caused by growth and nutritional deficiencies, possibly related to a reduction in transplacental nutrient substrate supply to the fetus.

We questioned whether alterations in bone collagen type I metabolism could be a cause of low BMC in SGA infants. In theory, conditions that affect fetal growth can lead to changes in bone collagen type I metabolism. Type I collagen, the major collagen product synthesized by bone cells, is the only collagen type found in mineralized bone (Burgeson, 1988). Serum PICP and ICTP are markers of bone collagen type I biosynthesis and degradation respectively (Parfitt *et al.* 1987; Eriksen *et al.* 1993).

Thus, we tested the hypothesis that serum PICP would be low and serum ICTP would be high in term SGA infants compared with term AGA infants, based on the *in vitro* data of a regulatory role of insulin-like growth factor in bone collagen type I metabolism. In comparisons

of SGA infants (Thieriot-Prevost *et al.* 1988; Lassarre *et al.* 1991) and growth-retarded animals with controls (Bernstein *et al.* 1991; Unterman *et al.* 1993), serum insulin-like growth factor-I concentrations were reduced by approximately 30–50 %; insulin-like growth factor-I stimulates type I collagen synthesis and decreases collagen degradation. In SGA *v.* AGA infants serum markers of bone collagen type I synthesis (PICP) and degradation (ICTP) were not different; serum ICTP was correlated with osteocalcin ( $r = 0.596$ ,  $P = 0.007$ ) and with PICP ( $r = 0.663$ ,  $P = 0.002$ ) in SGA infants but not in AGA infants. These findings are not consistent with the theory that altered fetal bone collagen type I formation or breakdown causes the low BMC of SGA infants. Thus, we concluded that the low BMC of SGA *v.* AGA infants may be predominantly related to low mineral supply rather than defective regulation of bone collagen type I metabolism (Namgung *et al.* 1996).

In contrast to our findings in term SGA infants, a recent report from our centre showed that amniotic fluid PICP concentrations were low in SGA infants compared with controls (AGA infants;  $P = 0.07$ ), both PICP and ICTP were inversely associated with gestational age ( $P < 0.0001$ ), and amniotic fluid concentrations of PICP increased exponentially in relation to infant birth weight ( $P = 0.008$ ). These findings are consistent with a high bone turnover early in fetal life, and a reduction in bone formation when fetal growth is compromised (Harrast & Kalkwarf, 1998). These findings are also consistent with the concept of decreased bone formation in SGA infants beginning *in utero*, and add to our finding of low cord serum osteocalcin in term SGA *v.* AGA infants. An alternative explanation is that gestational age (and thus maturity) may affect clearance of bone markers in blood, and thus affect blood concentrations. It is also possible that fetal growth retardation *per se* has an effect on maturation of clearance pathways.

### Infants of diabetic mothers

Maternal diabetes could be a factor influencing fetal bone metabolism, through alterations in maternal mineral metabolism or limited mineral availability in a pregnancy where the mother is diabetic. In a diabetic pregnant rat model fetuses of diabetic rats have been reported to have decreased mineralization, as measured by the number of ossification centres (delay in bone maturation) and total ash weight. Ca and P content per fetus, and Ca and P content per unit fetal body weight (Verhaeghe *et al.* 1986, 1988, 1999; Demignon & Robut-Bonneton, 1988; Husain *et al.* 1994).

The pathogenesis of fetal hypomineralization in a pregnancy where the mother is diabetic remains uncertain. Maternal-fetal Ca transport is decreased in perfused placentas from diabetic rats; placental unidirectional materno-fetal flux of Ca is reduced (by 61 % of the control value) in untreated diabetic rats compared with control or insulin-treated diabetic rats. Fetal Ca content was also significantly lower in the untreated diabetic rats compared with the other two groups. Since placental calbindin 9K:cyclophilin mRNA was 11- to 12-fold lower in the untreated diabetic rats, it appears that untreated maternal diabetes mellitus reduces fetal Ca accretion through an effect on the expression of placental transport components


involved in the materno-fetal transfer of Ca (Husain *et al.* 1994).

Using spontaneously-diabetic rats fed on a 10 g Ca/kg diet (a standard rat chow; Trouw, Ghent, Belgium), Verhaeghe *et al.* (1999) recently showed that fetuses of diabetic rats had no mineralization defect (as measured by Ca content per unit body weight, although the mean Ca content was lower in fetuses of diabetic rats *v.* control), caused by disturbed maternal duodenal Ca absorption or transplacental Ca transport. However, they did report a delay in bone maturation. This finding is not consistent with previous findings from the same authors using diabetic BB rats (Verhaeghe *et al.* 1988). The difference between these reports might be related to differences in litter size, since impairment in mineralization was most pronounced in the large litters in the earlier study (Verhaeghe *et al.* 1988).

Decreased BMC has been reported in diabetic patients (Levine *et al.* 1976), and might be related to insulin deficiency, since insulin therapy improves BMC in juvenile diabetes. Insulin has been shown to stimulate proliferation of osteoblast-like cells *in vitro*, and to increase histomorphometric indices of bone formation *in vivo* (Cornish *et al.* 1996). Neonatal hypocalcaemia and neonatal hypomagnesaemia occur frequently in infants born of mothers with insulin-dependent diabetes; the incidence and severity of hypocalcaemia are directly related to the severity of maternal diabetes, and related to the urinary Mg losses in diabetes (Tsang *et al.* 1972).

In an earlier study from our group (Mimouni *et al.* 1988), we suggested that infants of diabetic mothers (IDM) may have decreased BMC at birth, associated with poor control of diabetes during pregnancy. Since elevation of 1,25-(OH)<sub>2</sub>D concentrations has been reported in IDM (Steichen *et al.* 1981), raised 1,25-(OH)<sub>2</sub>D may be an adaptive response to the possible poor transplacental mineral supply. BMC was measured at birth (by single-photon absorptiometry) in forty-five infants born of diabetic mothers (class B-RT; 38.2 (1SE 0.2) weeks; body weight percentile 77.4 %), and in fifty-five normal term newborn infants (38.1 (1SE 0.4) weeks; body weight percentile 53.2 %) born of non-diabetic mothers. The birth weight percentiles were significantly higher ( $P < 0.01$ ) in IDM *v.* control infants. BMC at birth was significantly decreased ( $P < 0.001$ ) in IDM compared with control infants. BMC was inversely correlated with first trimester maternal mean blood glucose concentration ( $r = -0.664$ ,  $P < 0.01$ ; Fig. 5), implying that factors early in pregnancy might have an effect on fetal BMC (Mimouni *et al.* 1988). The low BMC in IDM (despite greater body weight) may be related to the decreased mineral transfer across the placenta in early pregnancy, possibly secondary to possible decreased mineral availability in a pregnancy where the mother is diabetic.

In contrast to the earlier findings in IDM, the changes in whole-body BMC in IDM, recently measured using dual-energy X-ray absorptiometry, has been reported as variable. Lapillonne *et al.* (1997b) recently reported that in IDM, the whole-body BMC Z score when adjusted for weight of each infant at the time of dual-energy X-ray absorptiometry measurement (whole-body BMC Z score/W, where W is weight) was positive (1.3 (SD 0.9)) and significantly



**Fig. 5.** Neonatal bone mineral content *v.* mean maternal first trimester postprandial blood glucose concentration for infants and their diabetic mothers. There was a significant inverse correlation between bone mineral content and maternal blood glucose concentration ( $r = -0.66$ ,  $P < 0.01$ ). (Reproduced with permission from Mimouni *et al.* 1998.)

different from zero ( $P < 0.0001$ ), but was not influenced either by *in utero* growth (AGA or large-for-gestational age) or by the type of diabetes mellitus of the mother. The reference values for term infants used by Lapillonne *et al.* (1997b) were from different studies (Lapillonne *et al.* 1997a). The difference in these findings might be related to variations in maternal diabetes control, variability in the population studied and lack of an adequate control population. Recent reports on outcomes of pregnancies where the mother was diabetic have documented improved control of the diabetes compared with earlier studies (Mehta *et al.* 1998). The inverse relationship between infant BMC and maternal blood glucose concentration reported by Mimouni *et al.* (1988) supports this conclusion, since if maternal blood glucose control improves with time, infant BMC values in IDM should rise also.

The mechanism for BMC changes in IDM is not clear. A study from our group (Demarini *et al.* 1995) found evidence of increased intrauterine bone resorption in term infants born of mothers with insulin-dependent diabetes (class B-RT). Cord serum ICTP concentrations (a marker of bone resorption) were higher in IDM than in control subjects, supporting the concept of increased osteoclastic activity in IDM. IDM had significantly higher birth weight than the controls. Serum PICP concentration (a marker of bone formation) was not different between groups.

Recently, our centre reported that diabetes (insulin-dependent and gestational) in the mother was not associated with alterations in amniotic fluid PICP and ICTP concentrations; there was no relationship between either of the bone markers in amniotic fluid and maternal diabetes (Harrast & Kalkwarf, 1998). The physiological significance of chemical substances found in amniotic fluid is often unclear, and make comparisons with cord blood studies difficult. A recent report indicates that urinary excretion of N-terminal telopeptide of type I collagen (a bone resorption marker) during the first 48 h in seventeen IDM were comparable with those of control subjects (Mora *et al.*

1997). This finding could also be due to improved metabolic control of diabetes during pregnancy; all mothers in the study had values for blood glycosylated haemoglobin of <8 % total haemoglobin. It appears that osteopenia in IDM and factors promoting bone resorption may be ameliorated with improved diabetes management in pregnancy.

#### **Other factors: anthropometric measures, race, gender and maternal ethanol consumption and smoking**

BMC is consistently found to increase with increasing body weight and length in infants and children (Li *et al.* 1989; Rupich *et al.* 1996). Race and gender differences in BMC appear in early life, but probably not at birth. In one study a higher total body bone mass was observed in male infants than in female infants aged 1–18 months, even after controlling for weight, length and race (Rupich *et al.* 1996). Race differences appear by 1 year of age, with black infants having greater BMC than white infants (Li *et al.* 1989). No gender or race differences in BMC were observed in term infants at birth (Steichen *et al.* 1976; Namgung *et al.* 1993). Both decreased urinary Ca excretion and increased intestinal Ca absorption have been found in older black children compared with white children, possibly providing a mechanism for race differences (Abrams *et al.* 1995); no data are available in infants.

Maternal ethanol consumption can have deleterious effects on both adult and developing bone. Pregnant rats fed on a liquid diet with ethanol showed low Ca content of maternal bone, low maternal blood ionized Ca concentration and elevated PTH compared with rats fed on diets without ethanol (pair-fed control). Mean fetal body weight and fetal skeletal ossification were reduced in the ethanol-fed rats compared with controls, but no group differences were found in fetal Ca content (Keiver *et al.* 1996). Chronic alcohol consumption in young actively-growing (4-week-old) rats during the age of bone development reduced bone volume and trabecular number in cancellous bone. The rate of bone formation is reduced in alcohol-fed animals, indicating a reduced osteoblastic activity (Sampson *et al.* 1996; Sampson, 1998).

The effects of ethanol on fetal bone development do not appear to be related to alterations in fetal rat serum levels of calcitropic hormones, as shown by the finding of no differences in fetal rat serum levels of PTH and calcitonin between ethanol-fed and control groups (Keiver *et al.* 1997). There might be interactive effects of ethanol intake and maternal nutritional status, since increasing the protein content of the alcohol-containing diet from 180 to 250 g/kg significantly increased ossification of fetal bone in rat (Weinberg *et al.* 1990). Comparable studies in human infants have not been carried out. The effect of alcohol on bone may be related to a more direct effect of alcohol on bone cells (Sampson *et al.* 1998).

Smoking by the mother is detrimental to the developing fetus. In rats there is a dose-related retardation in embryonic growth, and also a reduction in the number of skeletal ossification centres (developmental delay; Seller & Bnait, 1995). There is evidence of a long-term negative association between maternal smoking during pregnancy and both growth and bone mass in children born at term (Jones *et al.*

1999). Smoking by the mother during pregnancy is associated with deficits in growth in 8-year-old children (lower weight and height), and disproportionate deficit in bone mass; children whose mothers smoked during pregnancy had lower size-adjusted bone mass at the lumbar spine and femoral neck but not total body. This association was only present for children born at term. Mothers who smoked during pregnancy also had lower placental weight, and further adjustment for placental weight led to non-significant results for smoking in both growth and bone variables, suggesting that these associations may be mediated through placental size and function.

Tobacco smoking affects the fetoplacental unit; carboxyhaemoglobinaemia and chronic hypoxaemia, vasoconstriction of uteroplacental circulation and intermediary metabolism disturbance are the aetiopathogenetic base of intrauterine fetal growth retardation, i.e. the fetal smoking syndrome (Habek, 1998). Oxidant stress from increased free radicals produced by smoking may have some role in the adverse effects on the skeleton of subjects who smoke, possibly related to alterations in bone resorption (Melhus *et al.* 1999).

In full-term newborn infants of mothers who smoked during pregnancy, smoking had a clear dose-dependent negative effect on all anthropometric measures in infants (Zaren *et al.* 1996). Newborn infants born of non-smoking but exposed mothers (passive smoking) or mothers who were light smokers (less than ten cigarettes per d) showed a significant reduction in fat mass and most anthropometric measurements (birth weight, crown–heel length and skinfolds). Intrauterine growth seems to be negatively influenced not only by active smoking, but also by passive and light active smoking (Luciano *et al.* 1998). The possible effect on fetal bone development has not been examined in this situation.

#### **References**

- Abrams SA, O'Brien KO, Liang LK & Stuff JE (1995) Differences in calcium absorption and kinetics between black and white girls. *Journal of Bone Mineral Research* **10**, 829–833.
- Aitken JM, Anderson JB & Horton PW (1973) Seasonal variations in bone mineral content after menopause. *Nature* **241**, 59–60.
- Bergstralh EJ, Sinaki M, Offord KP, Wahner HW & Melton LJ (1990) Effect of season on physical activity score, back extensor muscle strength, and lumbar bone mineral density. *Journal of Bone Mineral Research* **5**, 371–377.
- Bernstein IM, DeSouza M & Copeland KC (1991) Insulin-like growth factor I in substrate-deprived, growth-retarded fetal rats. *Pediatric Research* **30**, 154–157.
- Bouillon RA, Auwerx JD, Lissens WD & Pelemans WK (1987) Vitamin D status in the elderly: seasonal substrate deficiency causes 1,25-dihydroxycholecalciferol deficiency. *American Journal of Clinical Nutrition* **45**, 755–763.
- Brooke OG, Brown IRF, Bone CDM, Carter ND, Cleeve HJW, Maxwell JD, Robinson VP & Winder SM (1980) Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *British Medical Journal* **280**, 751–754.
- Burgeson RE (1988) New collagen, new concepts. *Annual Review of Cell Biology* **4**, 551–557.
- Chunga Vega F, Gomez de Tejada MJ, Gonzalez Hachero J, Perez Cano R & Coronel Rodriguez C (1996) Low bone mineral density in small for gestational age infants: correlation with cord



- blood zinc concentrations. *Archives of Disease in Childhood* **75**, F126–F129.
- Comar CL (1956) Radiocalcium studies in pregnancy. *Annals of the New York Academy of Sciences* **64**, 281–298.
- Cornish J, Callon KE & Reid IR (1996) Insulin increases histomorphometric indices of bone formation in vivo. *Calcified Tissue International* **59**, 492–495.
- Dawson-Hughes B, Dallal Ge, Krall EA, Harris S, Sokoll LJ & Falconer G (1991) Effect of vitamin D supplementation on wintertime and overall bone loss in healthy postmenopausal women. *Annals of Internal Medicine* **115**, 505–512.
- Demarini S, Specker BL, Sierra RI, Miodovnik M & Tsang RC (1995) Evidence of increased intrauterine bone resorption in term infants of mothers with insulin-dependent diabetes. *Journal of Pediatrics* **126**, 796–798.
- Demignon J & Robut-Bonneton C (1988) Effects of experimental diabetes on the vitamin D metabolism of pregnant rats and their fetuses. *Calcified Tissue International* **42**, 127–135.
- Durand D, Barlet JP & Braithwaite GD (1983a) The influence of 1,25-dihydroxycholecalciferol on the mineral content of foetal guinea-pigs. *Reproduction Nutrition Development* **23**, 235–244.
- Durand D, Braithwaite GD & Barlet JP (1983b) The effect of 1- $\alpha$ -hydroxycholecalciferol on the placental transfer of calcium and phosphate in sheep. *British Journal of Nutrition* **49**, 475–480.
- Eriksen EF, Charles P, Melsen F, Mosekilde L, Risteli L & Risteli J (1993) Serum markers of type I collagen formation and degradation in metabolic bone disease: correlation with bone histomorphometry. *Journal of Bone Mineral Research* **8**, 127–132.
- Fonsca V, Agnew JE, Nag D & Dandona P (1988) Bone density and cortical thickness in nutritional vitamin D deficiency: effect of secondary hyperparathyroidism. *Annals of Clinical Biochemistry* **25**, 271–274.
- Habek D (1998) Effect of smoking on the fetoplacental unit. *Lijec Vjesn* **120**, 215–219.
- Harrast SD & Kalkwarf HJ (1998) Effects of gestational age, maternal diabetes, and intrauterine growth retardation on markers of fetal bone turnover in amniotic fluid. *Calcified Tissue International* **62**, 205–208.
- Husain SM, Birdsey TJ, Glazier JD, Mughal MZ, Garland HO & Sibley CP (1994) Effect of diabetes mellitus on maternofetal flux of calcium and magnesium and calbindin9K mRNA expression in rat placenta. *Pediatric Research* **35**, 376–381.
- Jones G, Riley M & Dwyer T (1999) Maternal smoking during pregnancy, growth, and bone mass in prepubertal children. *Journal of Bone Mineral Research* **14**, 146–151.
- Keiver K, Ellis L, Anzarut A & Weinberg J (1997) Effect of prenatal ethanol exposure on fetal calcium metabolism. *Alcoholism Clinical and Experimental Research* **21**, 1612–1618.
- Keiver K, Herbert L & Weinberg J (1996) Effect of maternal ethanol consumption on maternal and fetal calcium metabolism. *Alcoholism Clinical and Experimental Research* **20**, 1305–1312.
- Kollee L, Mannens L, Trijbels J, Veerkamp J, Jassen A & Haard-Hustings H (1979) Experimental intrauterine growth retardation in the rat; evaluation of the Wigglesworth model. *Early Human Development* **3**, 295–300.
- Krølner B (1983) Seasonal variation of lumbar spine bone mineral content in normal women. *Calcified Tissue International* **35**, 145–147.
- Kuroda E, Okano T & Mizuno N (1981) Plasma levels of 25-hydroxyvitamin D<sub>2</sub> and 25-hydroxyvitamin D<sub>3</sub> in maternal, cord and neonatal blood. *Journal of Nutritional Science and Vitaminology* **27**, 55–65.
- Lapillonne A, Brillon P, Claris O, Chatelain PG, Delmas PD & Salle BL (1997a) Body composition in appropriate and in small for gestational age infants. *Acta Paediatrica* **86**, 196–200.
- Lapillonne A, Guerion S, Braillon P, Claris O, Delmas PD & Salle BL (1997b) Diabetes during pregnancy does not alter whole body bone mineral content in infants. *Journal of Clinical Endocrinology and Metabolism* **82**, 3993–3997.
- Lassarre C, Hardouin S, Daffos F, Forestier F, Frankenne F & Binoux M (1991) Serum insulin-like growth factors and insulin-like growth factor binding proteins in the human fetus. Relationships with growth in normal subjects and in subjects with intrauterine growth retardation. *Pediatric Research* **29**, 219–225.
- Levine ME, Boisseau VC & Aviolo LV (1976) Effects of diabetes mellitus on bone mass in juvenile and adult onset diabetes. *New England Journal of Medicine* **294**, 241–245.
- Li JY, Specker BL, Ho ML & Tsang RC (1989) Bone mineral content in black and white children 1 to 6 years of age. *American Journal of Diseases in Children* **143**, 1346–1349.
- Luciano A, Bolognani M, Biondani P, Ghizzi C, Zoppi G & Signori E (1998) The influence of maternal passive and light active smoking on intrauterine growth and body composition of the newborn. *European Journal of Clinical Nutrition* **52**, 760–763.
- Lukert BP, Carey M, McCarty B, Tiemann S, Goodnight L, Helm M, Hassanein R, Stevenson C, Stoskopf M & Doolan L (1987) Influence of nutritional factors on calcium regulating hormones and bone loss. *Calcified Tissue International* **40**, 119–125.
- Mehta KC, Kalkwarf HJ, Mimouni F, Khoury J & Tsang RC (1998) Randomized trial of magnesium administration to prevent hypocalcemia in infants of diabetic mothers. *Journal of Perinatology* **18**, 352–356.
- Melhus H, Michalsson K, Holmberg L, Wolk A & Ljunghall S (1999) Smoking, antioxidant vitamins, and the risk of hip fracture. *Journal of Bone Mineral Research* **14**, 129–135.
- Mimouni F, Steichen JJ, Tsang RC, Hertzberg V & Miodovnik M (1988) Decreased bone mineral content in infants of diabetic mothers. *American Journal of Perinatology* **5**, 339–343.
- Minton SD, Steichen JJ & Tsang RC (1983) Decreased bone mineral content in small for gestational age infants compared with appropriate for gestational age infants: normal serum 25-hydroxyvitamin D and decreasing parathyroid hormone. *Pediatrics* **71**, 383–388.
- Moncrieff M & Fadahunsi TO (1974) Congenital rickets due to maternal vitamin D deficiency. *Archives of Disease in Childhood* **49**, 810–811.
- Mora S, Prinster C, Bellini A, Weber G, Proverbio MC, Puziovio M, Bianchi C & Chiumello G (1997) Bone turnover in neonates: Changes of urinary excretion rate of collagen type I cross-linked peptides during the first days of life and influence of gestational age. *Bone* **20**, 563–566.
- Mughal MZ, Ross R & Tsang RC (1989) Clearance of calcium across in situ perfused placentas in intrauterine growth-retarded rat fetuses. *Pediatric Research* **24**, 420–422.
- Namgung R, Mimouni F, Campaigne BN, Ho ML & Tsang RC (1992) Low bone mineral content in summer-born compared with winter-born infants. *Journal of Pediatric Gastroenterology and Nutrition* **15**, 285–288.
- Namgung R, Tsang RC, Lee C, Han DG, Ho ML & Sierra RI (1998) Low total body bone mineral content and high bone resorption in Korean winter-born versus summer-born newborn infants. *Journal of Pediatrics* **132**, 421–425.
- Namgung R, Tsang RC, Sierra RI & Ho ML (1996) Normal serum indices of bone collagen biosynthesis and degradation in small for gestational age infants. *Journal of Pediatric Gastroenterology and Nutrition* **23**, 224–228.
- Namgung R, Tsang RC, Specker BL, Sierra RI & Ho ML (1993) Reduced serum osteocalcin and 1,25-dihydroxyvitamin D concentrations and low bone mineral content in small for



- gestational age infants: Evidence of decreased bone formation rates. *Journal of Pediatrics* **122**, 269–275.
- Namung R, Tsang RC, Specker BL, Sierra RI & Ho ML (1994) Low bone mineral content and high serum osteocalcin and 1,25(OH)<sub>2</sub> vitamin D in summer- versus winter-born newborn infants: An early fetal effect? *Journal of Pediatric Gastroenterology and Nutrition* **19**, 220–227.
- Ogeuh O, Khastgir G, Studd J, Jones J, Alaghband-Zadeh J & Johnson MR (1998) The relationship of fetal serum markers of bone metabolism to gestational age. *Early Human Development* **29**, 109–112.
- Okonofua F, Menon RK, Houlder S, Thomas M, Robinson D, O'Brien S & Dandona P (1987) Calcium, vitamin D and parathyroid hormone relationships in pregnant Caucasian and Asian women and their neonates. *Annals of Clinical Biochemistry* **24**, 22–28.
- Ooms ME, Lips P, Roos JC, van der Vijgh WJF, Popp-Snijders C, Bezemer PD & Bouter LM (1995) Vitamin D status and sex hormone binding globulin: determinants of bone turnover and bone mineral density in elderly women. *Journal of Bone Mineral Research* **10**, 1177–1184.
- Orwoll ES & Meier DE (1986) Alterations in calcium, vitamin D, and parathyroid hormone physiology in normal men with aging: relationship to the development of senile osteopenia. *Journal of Clinical Endocrinology and Metabolism* **63**, 1262–1269.
- Parfitt AM, Simon LS, Villanueva AR & Krane SM (1987) Procollagen type I carboxyterminal extension peptide in serum as a marker of collagen biosynthesis in bone. Correlation with iliac bone formation rates and comparison with total alkaline phosphatase. *Journal of Bone Mineral Research* **2**, 427–436.
- Peacock M, Liu G, Carey M, Ambrosius W, Turner CH, Hui S & Johnston CC Jr (1998) Bone mass and structure at the hip in men and women over the age of 60 years. *Osteoporosis International* **8**, 231–239.
- Petersen F, Gotfredsen A & Knudsen FU (1989) Total body bone mineral in light-for-gestational-age infants and appropriate-for-gestational-age infants. *Acta Paediatrica Scandinavica* **78**, 347–350.
- Pohlandt F & Mathers N (1989) Bone mineral content of appropriate and light for gestational age preterm and term newborn infants. *Acta Paediatrica Scandinavica* **78**, 835–839.
- Rupich RC, Specker BL, Lieuw-A-Fam N & Ho M (1996) Gender and race differences in bone mass during infancy. *Calcified Tissue International* **58**, 395–397.
- Sampson HW (1998) Effect of alcohol consumption on adult and aged bone: a histomorphometric study of the rat animal model. *Alcoholism Clinical and Experimental Research* **22**, 2029–2034.
- Sampson HW, Hebert VA, Booe HL & Champney TH (1998) Effect of alcohol consumption on adult and aged bone: composition, morphology, and hormone levels of a rat animal model. *Alcoholism Clinical and Experimental Research* **22**, 1746–1753.
- Sampson HW, Perks N, Champney TH & DeFee B II (1996) Alcohol consumption inhibits bone growth and development in young actively growing rats. *Alcoholism Clinical and Experimental Research* **20**, 1375–1384.
- Scharla SH, Scheidt-Nave C, Leidig G, Woitge H, Wuster C, Seibel MJ & Ziegler R (1996) Lower serum 25-hydroxyvitamin D is associated with increased bone resorption markers and lower bone density at the proximal femur in normal females: a population-based study. *Experimental and Clinical Endocrinology and Diabetes* **104**, 289–292.
- Seller MJ & Bnait KS (1995) Effects of tobacco smoke inhalation on the developing mouse embryo and fetus. *Reproductive Toxicology* **9**, 449–459.
- Sherman SS, Tobin JD, Hollis BW, Gundberg CM, Roy TA & Plato CC (1992) Biochemical parameters associated with low bone density in healthy men and women. *Journal of Bone Mineral Research* **7**, 1123–1130.
- Specker BL, Ho ML, Oestreich A, Yin TA, Shui QM, Chen XC & Tsang RC (1992) Prospective study of vitamin D supplementation and rickets in China. *Journal of Pediatrics* **120**, 733–739.
- Steichen JJ, Kaplan B, Edwards N & Tsang RC (1976) Bone mineral content in full-term infants measured by direct photon absorptiometry. *American Journal of Roentgenology* **126**, 1283–1285.
- Steichen JJ, Tsang RC & Ho ML (1981) Perinatal magnesium, calcium and 1,25-dihydroxyvitamin D in relation to prospective randomized management of maternal diabetes. *Pediatric Research* **15**, 683A.
- Thieriot-Prevost G, Boccara JF, Francoual C, Badoual J & Job JC (1988) Serum insulin-like growth factor I and serum growth-promoting activity during the first postnatal year in infants with intrauterine growth retardation. *Pediatric Research* **24**, 380–382.
- Tsang RC, Kleinman L, Sutherland JM & Light IJ (1972) Hypocalcemia in infants of diabetic mothers: studies in Ca, P and Mg metabolism and in parathormone responsiveness. *Journal of Pediatrics* **80**, 384–395.
- Unterman TG, Simmons RC, Glick RP & Ogata ES (1993) Circulating levels of insulin, insulin-like growth factor-I (IGF-I), IGF-II, and IGF-binding proteins in the small for gestational age fetal rat. *Endocrinology* **132**, 327–336.
- Verhaeghe J, Bouillon R, Lissens W, Visser WJ & Van Assche FA (1988) Diabetes and low Ca-P diets have opposite effects on adult and fetal bone and mineral metabolism. *American Journal of Physiology* **254**, E496–E504.
- Verhaeghe J, Bouillon R, Nyomba BL, Lissens W & Van Assche FA (1986) Vitamin D and bone mineral homeostasis during pregnancy in the diabetic BB rat. *Endocrinology* **118**, 1019–1025.
- Verhaeghe J, Van Bree R, Van Herck E, Rummens K, Vercruysse L, Bouillon R & Pijnenborg R (1999) Pathogenesis of fetal hypomineralization in diabetic rats: evidence for delayed bone maturation. *Pediatric Research* **45**, 209–217.
- Verity CM, Burman D, Beadle PC, Holton JB & Morris A (1981) Seasonal changes in perinatal vitamin D metabolism: maternal and cord blood biochemistry in normal pregnancies. *Archives of Disease in Childhood* **56**, 943–948.
- Weinberg J, D'Alquen G & Bezio S (1990) Interactive effects of ethanol intake and maternal nutritional status on skeletal development of fetal rats. *Alcohol* **7**, 383–388.
- Wigglesworth J (1964) Experimental growth retardation in the foetal rat. *Journal of Pathology and Bacteriology* **88**, 1–13.
- Woitge HW, Scheidt-Nave C, Kissling C, Leidig-Bruckner G, Meyer K, Grauer A, Scharla SH, Ziegler R & Seibel MJ (1998) Seasonal variation of biochemical indexes of bone turnover: results of a population-based study. *Journal of Clinical Endocrinology and Metabolism* **83**, 68–75.
- Zaren B, Lindmark G & Gebre-Medhin M (1996) Maternal smoking and body composition of the newborn. *Acta Paediatrica* **85**, 213–219.
- Zittermann A, Scheld K & Stehle P (1998) Seasonal variations in vitamin D status and calcium absorption do not influence bone turnover in young women. *European Journal of Clinical Nutrition* **52**, 501–506.