

Circulation and colonisation of *Blastocystis* subtypes in schoolchildren of various ethnicities in rural northern Thailand[‡]

Original Paper

Cite this article: McCain A, Grunec L, Popluechai S, Tsaousis AD, Gentekaki E (2023). Circulation and colonisation of *Blastocystis* subtypes in schoolchildren of various ethnicities in rural northern Thailand. *Epidemiology and Infection*, **151**, e77, 1–12 <https://doi.org/10.1017/S0950268823000596>

Received: 14 December 2022

Revised: 12 April 2023

Accepted: 13 April 2023

Keywords:

Blastocystis; BMI; children; colonisation; One Health; subtyping

Corresponding author:


Anastasios D. Tsaousis;

Email: A.Tsaousis@kent.ac.uk

Eleni Gentekaki;

Email: gentekaki.eleni@mfu.ac.th

[‡]This article has been updated since its original publication. A correction notice has been published detailing the changes can be found here: <https://doi.org/10.1017/S095026882300078X>

Abby McCain¹, Lucsme Grunec², Siam Popluechai^{1,2},
Anastasios D. Tsaousis³  and Eleni Gentekaki^{1,2}

¹School of Science, Mae Fah Luang University, Chiang Rai, Thailand; ²Gut Microbiome Research Group, Mae Fah Luang University, Chiang Rai, Thailand and ³Laboratory of Molecular and Evolutionary Parasitology, RAPID Group, School of Biosciences, University of Kent, Canterbury, UK

Abstract

Blastocystis is a protist of controversial pathogenicity inhabiting the gut of humans and other animals. Despite a century of intense study, understanding of the epidemiology of *Blastocystis* remains fragmentary. Here, we aimed to explore its prevalence, stability of colonisation and association with various factors in a rural elementary school in northern Thailand. One hundred and forty faecal samples were collected from 104 children at two time points (tp) 105 days apart. For tp2, samples were also obtained from 15 animals residing on campus and seven water locations. Prevalence in children was 67% at tp1 and 89% at tp2, 63% in chickens, 86% in pigs, and 57% in water. Ten STs were identified, two of which were shared between humans and animals, one between animals and water, and three between humans and water. Eighteen children (out of 36) carried the same ST over both time points, indicating stable colonisation. Presence of *Blastocystis* (or ST) was not associated with body mass index, ethnicity, birth delivery mode, or milk source as an infant. This study advances understanding of *Blastocystis* prevalence in an understudied age group, the role of the environment in transmission, and the ability of specific STs to stably colonise children.

Introduction

Blastocystis is a unicellular eukaryote inhabiting the gastrointestinal tract of vertebrates [1–3] and invertebrates [4, 5]. The organism comprises the most common protist identified in human stool samples [6]. The genetic heterogeneity of *Blastocystis* is extremely high based on the small subunit ribosomal RNA (SSU rRNA) gene [7–14]. To date, 34 subtypes (STs; ST1–ST17, ST21, and ST23–ST38) have been identified in avian and mammalian hosts. Of these, 14 have been recorded in humans (ST1–ST10, ST12, ST14, ST16, and ST23) [13, 15–20], with ST1–ST3 accounting for the majority of human carriage [21, 22]. The remaining STs mostly colonise non-human endothermic hosts [1, 6].

The pathogenicity of *Blastocystis* is a topic of ongoing debate. Its presence in the gut was initially linked to gastrointestinal symptoms, primarily due to a lack of information regarding carriage in healthy populations. The recent burst of studies reporting *Blastocystis* in individuals with no gastrointestinal symptoms has brought forth the hypothesis that the organism is a common member of the human gut microbiota [23, 24]. To that end, low prevalence of *Blastocystis* has been negatively associated with irritable bowel syndrome (IBS) [25]. However, positive associations between IBS patients and the presence of *Blastocystis* have also been observed. These conflicting results reinforce its controversial pathogenicity status [26–31]. Links between *Blastocystis* STs and/or strains and pathogenicity have also been proposed, but not conclusively shown [32–34].

Blastocystis has a worldwide distribution, having been found in both industrialised and non-industrialised countries [35–39]. ST3 is the most commonly distributed globally, followed by ST1 and ST2 [21, 22]. Several factors have been looked at in association with the presence or absence of *Blastocystis*. For instance, prevalence in rural communities is typically higher than that in urban areas [40, 41]. The age of the host seems to also play a role in colonisation, with carriage being generally higher in children [42–45].

The prevalence of *Blastocystis* in children varies, ranging from 4% [46] to 100% [36]. The majority of studies in this age group are microscopy-based; hence, subtyping information is relatively sparse [46–48]. In these studies, ST1–ST3 were the most commonly detected, following the global trend [49–52]. ST6 and ST7 have also been identified, most prominently in low- and middle-income countries (LMICs), but with lower carriage [44, 53–55]. Recently, the first occurrence of ST10, ST14, and ST16 was reported in children [16, 17]. The presence of *Blastocystis* in children was linked to sanitary habits, water source/treatment, type of housing,

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

and socioeconomic status of parents [48, 56, 57]. Most subtyping studies based on children have originated in South America [17, 54, 58], with only comparatively few from Asia. Given these gaps, studies addressing both the presence and STs of *Blastocystis* in children are needed worldwide.

In this study, we collected faecal samples from children of different ethnicities with no gastrointestinal symptoms in grades 1–6 attending a rural school in the Chiang Rai province, northern Thailand. The prevalence and genetic diversity of *Blastocystis* were determined. Information on body mass index (BMI), ethnicity, age, birth delivery mode, and milk source as an infant was also collected and assessed for association with *Blastocystis* presence. Faecal samples from animals raised on the school grounds and environmental samples were also examined to assess the circulation of the organism.

Methods

Ethics

The Ethics committee of Mae Fah Luang University approved the collection of human faecal samples (Ethics Registry: EC19359-11). The process, conditions, and ethical rules are in compliance with the Declaration of Helsinki. Data were strictly anonymised with each sample assigned an individual barcode. The parents and/or legal guardians of all participants provided a signed informed consent form.

Study area and sample collection

The present study took place in a rural area in the northern part of Thailand (Figure 1). Students enrolled in grades 1–6 (6–14 years old) at an elementary school in the Mae Fah Luang district of Chiang Rai province were recruited for the study. Recruitment of students took place via voluntary participation. The principal, staff, and parents and/or legal guardians were invited to attend a meeting, during which the objectives of the study were conveyed in detail in

the local language. Proper sterile technique in the collection of stool samples was also emphasised. Time was set aside for discussion, during which both children and adults were given the opportunity to ask questions and/or clarifications about any part of the project. A total of 104 volunteers participated. The participants were Akha ($n = 29$), Burmese ($n = 1$), Chinese ($n = 25$), Lahu ($n = 5$), Thai ($n = 19$), and Thai Yai ($n = 25$).

Samples were collected in November 2019 [time point 1 (tp1)] and February 2020 [time point 2 (tp2)] over the course of several days each time. Prior to collection, students were provided with a sterile container labelled with a unique ID code, ensuring confidentiality. On the morning of collection, one sample from each child was obtained at their homes and in some cases at the school. Samples were immediately stored at 4°C. Containers containing stool were collected from the students upon arrival to school and placed on ice until transport to the laboratory, where they were stored at –80°C. Information on ethnicity, age, sex, childbirth delivery mode (normal birth vs. Caesarean section), and mode of milk feeding (breast milk, formula, and mix of both) as an infant was recorded from questionnaires distributed to students (Supplementary Table S1). The questionnaires were filled in by the parents with the help of the school principal. Body weight and height information was also obtained at this time and used to calculate BMI. The BMI was converted into gender-specific z-scores for BMI-for-age according to BMI cut-offs for children (5–19 years) set by the World Health Organization. z-scores for BMI-for-age were classified into five groups: severely underweight (< -3 SD; $n = 1$), underweight (≥ -3 SD to < -2 SD; $n = 4$), healthy (≥ -2 SD to ≤ 1 SD; $n = 65$), overweight (OV; > 1 SD to ≤ 2 SD; $n = 16$), and obese (OB; > 2 SD; $n = 17$). None of the students had a history of gastrointestinal disease, were presented with diarrhoea episodes, or had taken probiotic supplements one month prior to either collection. Diarrhoea was defined as loose, watery stool three or more times per day. None had been treated with antibiotics two months prior to either sample collection. Students eat lunch at the school canteen and are involved in taking care of the school gardens.




Figure 1. Left panel: Map of Thailand. Middle panel: Close-up of Chiang Rai Province. Right panel: Details of the school grounds. Red dots indicate collection sites.

At tp2, faecal samples were also obtained from chickens ($n = 8$) and pigs ($n = 7$). A large dog visits the school grounds regularly; however, it was not possible to collect its stool. The chickens roam freely on school grounds and have been present for several years (Supplementary Figure S1). Chicken stool samples were collected in various areas of the school. The pigs were kept in different cement enclosures based on age (Supplementary Figure S2). Pre-weaned piglets were housed in two different enclosures, totalling over a dozen altogether. Adult pigs were housed in two enclosures (three in one enclosure and two in another). Two faecal samples from each pig enclosure were collected, excluding one adult enclosure where only one sample was obtained. Care was taken to collect the stool at the moment of defecation, and none of the animals were diarrheic. The animal stool was immediately inoculated into tubes containing liquid yeast serum growth media (LYSGM) media [59, 60] and incubated at 37°C for up to 3 days before further analysis.

Water samples ($n = 7$) were obtained at tp2 from seven different locations at the elementary school. Of those, five were collected from different water sources on campus, all flowing from two different origins. Water is supplied from two identical water treatment systems (Supplementary Figure S3). Each system is composed of four cylindrical cement tanks. All tap water available on campus runs from these two water treatment systems via plastic pipes. Samples were collected from taps outside of the bathroom, a water collection container inside the bathroom, a tap in the food preparation area, directly from a water tank, and a tap outside of the classroom (Supplementary Figure S4). One water sample was collected from a water container from which students drink daily, whereas another was obtained from the creek that runs along the campus (Supplementary Figure S5). The creek is approximately 3 metres wide and had a slow steady flow at the site of collection. During the time of collection, the water level was very low (a few centimetres) and the rock bed was above water in the middle of the creek. The collected water was slightly cloudy, as a small amount of sediment was present. The creek is accessible to the chickens roaming the school grounds. Approximately 1.5–3 ml of water was collected using sterile, disposable pipettes. The water was immediately inoculated into LYSGM media to a maximum volume of 15 ml and tubes were incubated at 37°C for up to 3 days.

DNA extraction

Total genomic DNA was extracted from 200 to 400 mg of human faeces using the QIAamp DNA stool mini kit (QIAGEN Inc., Hilden, Germany) according to the manufacturer's specifications. DNA was extracted for all 104 faecal samples from tp2, whereas for tp1, 36 samples were randomly selected. DNA from animal and environmental samples was extracted prior to the

first passage of culture using 200–400 ml of sediment from each sample. For these samples, the G-spin™ Total DNA Extraction Mini Kit (iNtRON, Seongnam, Korea) was used. The isolated DNA from all the samples was kept at –20°C until analysed.

Molecular detection and subtyping of *Blastocystis* sp.

Polymerase chain reaction

PCR amplification of the SSU rRNA gene was performed using a nested PCR reaction. In the first reaction, the broadly specific oligonucleotide primers RD5 (5'-GGAAGCTTATCTGGTTGATCCTGCCAGTA-3') and RD3 (5'-GGGATCCTGATCCTCCG-CAGGTTACCTAC-3') were used [61]. The thermocycling profile was as follows: denaturing at 94°C for 3 min; 35 cycles of 100 s at 94°C, 100 s at 65°C, 100 s at 72°C, and a final extension of 10 min at 72°C. In the second reaction, the product from the first PCR reaction was used as the template along with the barcoding primers BsRD5F (5'-ATCTGGTTGATCCTGCCAGT-3') and BhrDR9R (5'-GAGCTT TTTAACTGCAACAACG-3') [62]. The thermocycling profile was as follows: denaturing at 94°C for 3 min; 35 cycles of 1 min at 94°C, 1 min at 60°C, 90 s at 72°C, and a final extension of 10 min at 72°C. The final reaction amplified an approximately 600 bp region that is typically used for the subtyping of *Blastocystis* [62]. The PCR amplicons of the target size were purified using AccuPrep® Gel Purification Kit (Bioneer Corporation, Daejeon, Republic of Korea) according to the manufacturer's specifications. Purified products were sent for sequencing to Bionics Co., Ltd., Seoul, South Korea. qPCR was performed on all DNA samples that were negative by conventional PCR following a previously developed protocol [15, 63]. Amplifications were performed using the primers BL18SPPF1 (5'-AGTAGTCATACGCTCGTCTCAAA-3') and BL18SR2PP (5'-TCTTCGTTACCCGTTA CTGC-3'), generating a 320 bp product. All PCR and qPCR products were sequenced unidirectionally using the reverse PCR (BhrDR9) and qPCR (BL18SR2PP) primers, respectively.

Cloning

Samples with unclear chromatograms were cloned using pLUG-Prime® TA-cloning Vector Kit II (iNtRON, Seongnam, Korea). A total of 32 samples were cloned. Of these, 25 were qPCR products, whereas 7 were PCR products. Thirty-one of the samples were from humans (3 from tp1 and 28 from tp2), and one was from a water source. With regard to PCR products, two colonies per transformation were screened in four cases, and one colony in the remaining two. For the qPCR products, two colonies per transformation were screened in 13 cases, and one colony in the remaining 15. In total, 47 clones were screened. Forty-five of the clones were from children's samples, whereas two were from a water sample.

Table 1. Prevalence and subtype distribution in children, animal, and water samples

Source	Prevalence (%)	ST1	ST2	ST3	ST5	ST6	ST7	ST15	ST26	UNK/mix
Human	93/104 (89)	7	19	42	1	–	14	–	1	5/2
Chicken	5/8 (63)	–	–	–	–	1	4	–	–	–
Pig	6/7 (86)	–	–	–	5	–	–	1	–	–
Water	4/7 (57)	2	–	1	–	–	–	–	–	0/1
	Total	9	19	43	6	1	18	1	1	5/3

Phylogeny and sequence analysis

Raw data chromatograms were edited, and ambiguous quality bases were removed from the 5' and 3' ends using the software AliView [64]. The sequences were used as queries to perform a BLAST search to check for contamination. A dataset was assembled containing reference sequences from all STs, including reptile and insect lineages. Sequences were aligned using MAFFT version 7 [65]. Ambiguous positions were removed using trimAl version 1.3 (<http://phylemon.bioinfo.cipf.es/>) [66]. One hundred and seventy-five taxa were present in the alignment, and 1,392 positions were left after trimming. A maximum likelihood phylogeny was inferred using RAxML version 8 on XSEDE available on the CIPRES gateway (<http://www.phylo.org/portal2/home.action>) [67]. The estimates of the genetic distance between sequences that were identified as the same ST and derived from children and water or children and animal were analysed using Kimura two-parameter model embedded in MEGAX [68–70].

Statistical analysis

Fisher's exact test with Monte Carlo simulation was used to determine the associations of BMI *z*-score, ethnicity, delivery mode, and milk source variables with the presence of *Blastocystis*. The associations of these variables with the prevalence of *Blastocystis* STs were also determined using the same test. One sample was excluded from the BMI *z*-score analysis, as weight and height information was not collected. Multiple correspondence analysis (MCA) was performed to explore the relationships between variables and the prevalence of *Blastocystis* STs using FactoMineR version 2.4 [71]. The confidence ellipses around the categories (variables) represented in the MCA plot were plotted using Factoextra version 1.0.7 [72]. Contingency tables were visualised using the function balloon plot (R gplots package version 3.1.1) [73]. All the analyses were performed in R software version 4.0.3 [74].

Results

Blastocystis prevalence in faecal and environmental samples

For tp1, 12 human faecal samples were positive for *Blastocystis* using nested PCR and 12 additional ones using qPCR (67%, 24/36). For tp2, 27 human faecal samples were positive for *Blastocystis* using nested PCR. Of these, five were false positives (no significant match by BLAST or plant species). The remaining 82 samples were subsequently screened by qPCR, and 71 of them were positive. Thus, the overall prevalence of *Blastocystis* in children at tp2 was 89% (93/104) (Table 1). Sanger sequencing was used to sequence all the samples that were positive by either PCR or qPCR. The overall prevalence of *Blastocystis* in pigs and chickens was 86% (6/7) and 63% (5/8), respectively. Of the seven water samples, four were positive for *Blastocystis*, giving an overall prevalence of 57%. All the sequences generated in this study were submitted to GenBank under accession numbers QQ571480–QQ571626.

Subtyping of *Blastocystis*

A total of 25 sequences (21 PCR/qPCR products + 4 clones) were subtyped at tp1, and in total five STs were identified: ST1 (2/25,

Table 2. Samples collected at two time points, 105 days apart and the subtypes present

Sample	tp1	tp2
MA203	ST1/ST3	ST3
MA204	–	ST7
MA205	ST2	ST2
MA210	ST3	ST3
MA217	ST3	ST3
MA227	ST3	ST3
MA235	–	ST3
MA236	ST10	ST2
MA241	ST3	ST3
MA246	–	ST7
MA256	–	ST3
MA270	–	ST7
MA272	ST3	ST1
MA280	ST3	ST3
MA282	–	ST7
MA286	ST1	ST1
MA291	ST2	ST2
MA296	ST3	ST3
MA300	–	ST3
MA302	–	ST3
MA303	–	ST7
MA307	ST3	ST2
MA309	ST3	ST3
MA311	ST3	ST3
MA318	–	ST7
MA320	ST3	ST3
MA323	–	ST2
MA325	ST3	ST3
MA327	ST2	ST2
MA331	ST1	ST1
MA332	ST10	ST7
MA334	ST3	ST3
MA336	–	ST7
MA341	ST3	ST1
MA344	ST3	ST3
MA348	ST23	ST7

Note: Dash indicates a negative result.
Abbreviation: tp, time point.

8%), ST2 (3/25, 12%), ST3 (15/25, 60%), ST10 (2/25, 8%), and ST23 (1/25, 4%). A case of ST1 and ST3 co-occurrence was also noted. At tp2, a total of eight STs were identified: ST1–ST3, ST5–ST7, ST15, and ST26 (Table 1). Six out of the eight STs identified at tp2 were found in children: ST3 (*n* = 43), ST2 (*n* = 19), ST7 (*n* = 14), ST1 (*n* = 8), ST5 (*n* = 1), and ST26

($n = 1$). There were four co-occurrences in the children's samples: ST3/UNK ($n = 1$), ST7/UNK ($n = 1$), and ST1/ST3 ($n = 2$). Five sequences were designated as unknown. Although it was possible to discern that they belonged to *Blastocystis* due to their short length, an ST could not be confidently assigned. Chickens had ST7 (4/5, 80%) and ST6 (1/5, 20%); pigs had ST5 (5/6, 83%) and ST15 (1/6, 17%). ST1 (2/4, 50%) and ST3 (1/4, 25%) along with one co-occurrence of ST1/ST7 were identified in water samples.

Stability of *Blastocystis* colonisation

Of the 36 children for which both time points are available, 12 were negative at tp1 (Table 2). All 12 tp1 negatives were positive at tp2 as follows: one with ST2, four with ST3, and seven with ST7. In six instances, a different ST was detected in the two time points. Eighteen children carried the same ST at both time points as follows: ST1 ($n = 2$), ST2 ($n = 3$), and ST3 ($n = 13$). In seven cases, the sequences between the two time points were identical: ST1 ($n = 1$), ST2 ($n = 2$), and ST3 ($n = 4$). In the remaining 11 cases, the sequences differed between 0.18% and 2.97% (Supplementary Table S2).

Circulation of *Blastocystis* subtypes

This analysis involved only sequences from tp2. ST sharing was noted among all three categories herein: humans, animals, and water (Figure 2). ST5 and ST7 were shared between humans and animals, ST1, ST3, and ST7 between humans and water, and ST7 between animals and water. ST7 was shared by all three. The ST5 sequences from humans and pigs were not identical. Similarly, ST1 sequences from water and children differed. ST3 sequences from the water creek, and 16 children were identical. ST7 sequences from two children samples and a water sample obtained from the tap outside the school bathroom were also identical.

Phylogenetic analysis

For the phylogenetic analysis, 23 sequences were chosen to represent the different STs present in the sample cohort (Figure 3). ST15 found in the pig sample grouped with previously published ST15 sequences and together with ST28 and sequences from ectothermic hosts placed at the base of the tree. Newly generated sequences placed within clades consisting of known STs. Sequences identified

as ST10 and ST23 were grouped with similar sequences from our previous One Health study [15].

Association of *Blastocystis* with BMI, ethnicity, birth delivery mode, and milk source

For this analysis, only samples from tp2 were included. The BMI data of two participants were excluded. In the case of one participant, the data were not sufficient to calculate BMI, whereas another was underweight. As there was only one participant in the underweight category, it was not included in the BMI analysis calculations. A graph of contingency tables was constructed, and Fisher's exact tests were performed.

At the *Blastocystis* level, there was no significant dependence between its presence and BMI z -score, ethnicity, birth delivery mode, or milk source groups ($p > 0.05$) (Figure 4). At the ST level, there were also no significant associations between their prevalence and any of the variables ($p > 0.05$) (Figure 5). The four most abundant STs (ST1–ST3 and ST7) were found across all groups, with two exceptions: ST1 and ST7 were absent in the Thai Yai and Thai ethnicities, respectively.

MCA explained 13.8% and 13.1% of individual variability in Dim1 and Dim2, respectively (Figure 6). In Dim1, Lahu ethnicity highly contributed to the dimension (coordinate = 0.80, $p < 0.001$). Breastfeeding (coordinate = 0.42, $p < 0.0001$) and natural birth (coordinate = 0.14, $p = 0.03$) showed a contrasting profile to formula feeding (coordinate = -0.42 , $p < 0.0001$) and caesarean (coordinate = -14 , $p = 0.03$), respectively. In Dim 2, the variation of observations was mostly characterized by ethnicity ($R^2 = 0.51$, $p < 0.0001$), BMI z -score ($R^2 = 0.37$, $p < 0.0001$), and *Blastocystis* STs ($R^2 = 0.34$, $p < 0.0001$), respectively. According to the MCA plots of the observations and categories (Figure 6), there were two density zones clustering around the caesarean, obese, and healthy groups as well as around the natural birth, Akha ethnicity, breastfeeding, and ST2 groups, indicating that the majority of individuals in this study shared a similar pattern among these two clustered groups of variables.

Discussion

This study took place at an elementary school located in a rural area of northern Thailand. The overall prevalence of *Blastocystis* was




Figure 2. *Blastocystis* subtypes (STs) present in humans, animals, and water in a rural elementary school in northern Thailand. STs on arrows indicate overlap. Shaded STs indicate the presence in all sources considered in this study.




Figure 3. Maximum likelihood phylogeny inferred from 175 *Blastocystis* sequences and 1,392 sites of the SSU rRNA gene. Newly generated sequences are in bold font. Numerical values indicate bootstrap support. Only values above 70 are shown.

89%, the highest rate reported in children in Thailand to date. Previous studies on this age group in Thailand focused on children from childcare centres and orphanages located in rural or low

socioeconomic areas with overall prevalence ranging from 4.8% to 31.9% [50, 57, 75–78]. The variable sensitivity of the methods used for detection across studies likely accounts for the different




Figure 4. Balloon plots of contingency tables of the relationship between variables (rows) and the presence of *Blastocystis* (columns) in 102 samples. Values represent the frequencies of the presence/absence of *Blastocystis*. (a) Body mass index z-score. (b) Ethnicity. (c) Birth delivery mode. (d) Milk source.

rates of prevalence. Regardless, the high prevalence rate herein matches previous works on children residing in LMICs in Asia, Africa, and South America [16, 43–45, 55, 79]. The highest carriage ever observed was 100% in children from the Senegal River basin [36].

Identification of *Blastocystis* in children has been primarily microscopy-based with only a limited number of studies focusing on subtyping. In this study, ST3 was the most dominant with a prevalence of 45% followed by ST2 (20%) and ST7 (15%). This pattern does not follow the global trend, where the top three STs are ST1–ST3 [7, 16, 22, 24]. Nonetheless, differences in ST distribution based on geographic location have been observed. For example, in Indonesia and at the China–Myanmar border, ST1, ST3, and ST4 were identified in children, the latter being an ST most commonly observed in European countries [43, 45]. Here, a total of eight STs were identified in children: ST1–ST3, ST5, ST7, ST10, ST23, and ST26. This high ST diversity in terms of number matches previous research on children living in rural areas of South American and African countries [16, 17, 53–55]. In contrast, subtyping work in this age range from southeast Asia has identified remarkably less ST diversity [43, 45, 46, 50, 57, 76]. ST10 and ST23 were previously identified in adults from the same province, but from different districts, suggesting that these are usual, but low-frequency STs in the area. In this study, we also report the first occurrence of ST26 in humans, which has so far been identified only in artiodactyls in the United States, China, and Spain [9, 80, 81]. Even though the sequence is relatively short,

both the blast result and the robust phylogenetic placement confirm this identification. The increasing occurrence of these ‘non-human’ STs challenges our previous understanding of the presence of *Blastocystis* in humans.

To examine the stability of *Blastocystis* colonisation, we sampled 36 children at two time points 105 days apart. Only three studies have so far explored temporal colonisation, all of which took place in Europe [82–85]. Two of these involved adults, whereas one focused on infants and toddlers. Nevertheless, this is the first study focusing on children (ages 6–14) from an Asian country having a high number of positive samples at both time points. For 18 children, the same STs were detected at both sampling points. These included ST1–ST3, matching the previous temporal colonisation studies, where the same STs along with ST4 and ST8 were found [82–85]. In the rest of the cases herein, the STs were either lost or switched. Notably, even though ST7 was present in seven individuals at the second time point, it was not detected in any of them at the first. This raises the interesting possibility of whether ST7 can stably colonise the human gut or if it is merely a passenger. Alternatively, the observed losses and/or switches could be due to the cycling of STs in the host or double colonisation, whereby only the dominant ST is amplified. Regardless, our data point towards ST1–ST3 being stable colonisers in these children.

In our previous One Health study of *Blastocystis*, we identified similar STs circulating between humans, animals, and the environment and suggested expanding this to additional communities [15]. Hence, here, we collected human and animal stool and




Figure 5. Balloon plots of contingency tables of the relationship between variables (rows) and the prevalence of *Blastocystis* subtypes (STs; columns) in 81 samples. Values are the frequencies of *Blastocystis* STs. (a) Body mass index z-score. (b) Ethnicity. (c) Birth delivery mode. (d) Milk source.

environmental samples to explore the circulation of *Blastocystis* and its STs. We found shared STs between hosts and the environment. The zoonotic and waterborne transmission of the various STs has been suggested multiple times [2, 86–92]. Taking this a step further, our genetic distance comparisons identified identical ST3 and ST7 sequences only between children and water. This hints at the waterborne transmission of these STs. A possible explanation for their environmental persistence is a higher resistance to a water environment [93, 94]. Future studies focusing on *Blastocystis* circulation should consider the sequences beyond the ST level.

Fisher's exact test was used to determine the association between the occurrence of *Blastocystis* and certain variables. The results indicated that there was no association between the presence of *Blastocystis* and BMI z-score, ethnicity, birth delivery mode, and milk source. This is likely due to the high prevalence of the organism across all categories in this cohort. To date, there have not been *Blastocystis* studies in children that evaluate any of these factors. BMI has been considered in past investigations involving adults [95, 96]. The presence of *Blastocystis* has been observed to have a strong negative correlation with BMI; specifically, higher prevalence is found in lean individuals [35, 97, 98]. Data from these studies have not shown *Blastocystis* and BMI to be ST-specific [35]. Our statistical analysis suggested that lean children were more likely to be colonised with ST1, whereas obese children were more likely to be

colonised with ST3. Further investigations are necessary for the overall microbiome and metabolome [99] level to disentangle relationships between STs and BMI.

This study combines a One Health approach in a rural school community focused on a microbial eukaryote of controversial pathogenicity. Aside from the high prevalence and stability of *Blastocystis* ST1–ST3 colonisation in children, this study raises a few important questions: which STs are colonisers or passengers and what are the determining factors? Do anthropometric factors and ethnicity play a role? What is the environmental contribution in transmission? Are all these factors associated with colonisation at the ST level? Studies of larger cohorts at the national and international levels are urgently needed to explore these questions and broaden our understanding of this enigmatic organism.

Supplementary material. The supplementary material for this article can be found at <http://doi.org/10.1017/S0950268823000596>.

Data availability statement. Molecular data generated in this project have been submitted to GenBank and are publicly available.

Acknowledgements. We thank all the volunteers and their parents/legal guardians for their enthusiastic participation. We are grateful to Mr. Supachoke Piyasant for his ceaseless support of this project. We wish to thank Nantinee Moontan, Channarong Wanthunjai, and Vasana Jinatham for their assistance in sample collection. We would also like to thank Daisy Shaw from the Tsaousis Lab for proofreading the manuscript.




Figure 6. Multiple correspondence analysis (MCA) plots of the association between variables and the prevalence of *Blastocystis* subtypes ($n = 81$). (a) MCA plot displaying the observations and the categories. Density (grey) curves indicate the zones that are highly concentrated. (b) MCA factor map with 95% confidence ellipses surrounding the variables used in this study.

Author contribution. Conceptualization: E.G.; Formal analysis: A.M., L.G.; Funding acquisition: E.G.; Investigation: A.M.; Methodology: A.M., L.G., S.P.; Project administration: E.G.; Resources: S.P., A.D.T., E.G.; Supervision: S.P., A.D.T., E.G.; Visualisation: A.M.; Writing – original draft: A.M.; Writing – review and editing: A.D.T., E.G.

Financial support. This research was supported by the Thailand Research Fund (Grant No. RSA6080048) awarded to E.G. The corresponding authors would also like to acknowledge the COST Action network grant (CA21105 – *Blastocystis* under One Health).

Competing interest. The authors declare none.

References

- [1] Alfellani MA, Taner-Mulla D, Jacob AS, Imeede CA, Yoshikawa H, Stensvold CR and Clark CG (2013) Genetic diversity of *Blastocystis* in livestock and zoo animals. *Protist* **164**, 497–509. <https://doi.org/10.1016/j.protis.2013.05.003>
- [2] Cian A, el Safadi D, Osman M, Moriniere R, Gantois N, Benamrouz-Vanneste S, Delgado-Viscogliosi P, Guyot K, Li LL, Monchy S, Noël C, Poirier P, Nourrisson C, Wawrzyniak I, Delbac F, Bosc S, Chabé M, Petit T, Certad G and Viscogliosi E (2017) Molecular epidemiology of *Blastocystis* sp. in various animal groups from two French zoos and evaluation of potential zoonotic risk. *PLoS One* **12**, 1–29.
- [3] Lhotská Z, Jirků M, Hložková O, Brožová K, Jirsová D, Stensvold CR, Kolísko M and Jirků Pomajbíková K (2020) A study on the prevalence and subtype diversity of the intestinal protist *Blastocystis* sp. in a gut-healthy human population in the Czech Republic. *Frontiers in Cellular & Infection Microbiology* **10**, 544335. <https://doi.org/10.3389/fcimb.2020.544335/full>
- [4] Yoshikawa H, Wu Z, Howe J, Hashimoto T, Geok-Choo N and Tan KSW (2007) Ultrastructural and phylogenetic studies on *Blastocystis* isolates from cockroaches. *Journal of Eukaryotic Microbiology* **54**, 33–37. <https://doi.org/10.1111/j.1550-7408.2006.00141.x>
- [5] Zaki M, Zaman V and Sheikh N A (1996) Resistance of *Blastocystis hominis* cysts to chlorine. *Journal of Pakistan Medical Association* **46**, 178–179
- [6] Clark CG, van der Giezen M, Alfellani MA and Stensvold CR (2013) Recent developments in *Blastocystis* research. *Advances in Parasitology* **82**, 1–32. <https://doi.org/10.1016/B978-0-12-407706-5.00001-0>
- [7] Alfellani MA, Jacob AS, Perea NO, Kreczek RC, Taner-Mulla D, Verweij JJ, Levecke B, Tannich E, Clark CG and Stensvold CR (2013) Diversity

- and distribution of *Blastocystis* sp. subtypes in non-human primates. *Parasitology* **140**, 966–971.
- [8] Higuera A, Herrera G, Jimenez P, García-Corredor D, Pulido-Medellín M, Bulla-Castañeda DM, Pinilla JC, Moreno-Pérez DA, Maloney JG, Santín M and Ramírez JD (2021) Identification of multiple *Blastocystis* subtypes in domestic animals from Colombia using amplicon-based next generation sequencing. *Frontiers in Veterinary Science* **8**, 732129. <https://doi.org/10.3389/fvets.2021.732129>
- [9] Maloney JG, da Cunha MJR, Molokin A, Cury MC and Santin M (2021) Next-generation sequencing reveals wide genetic diversity of *Blastocystis* subtypes in chickens including potentially zoonotic subtypes. *Parasitology Research* **120**, 2219–2231. <https://doi.org/10.1007/s00436-021-07170-3>
- [10] Maloney JG, Molokin A, da Cunha MJR, Cury MC and Santin M (2020) *Blastocystis* subtype distribution in domestic and captive wild bird species from Brazil using next generation amplicon sequencing. *Parasite Epidemiology & Control* **9**, e00138. Available at <https://www.sciencedirect.com/science/article/pii/S2405673120300076>.
- [11] Maloney JG, Molokin A and Santin M (2020) Use of Oxford Nanopore MinION to generate full-length sequences of the *Blastocystis* small subunit (SSU) rRNA gene. *Parasites & Vectors* **13**, 595. <https://doi.org/10.1186/s13071-020-04484-6>
- [12] Stensvold CR (2012) Thinking *Blastocystis* out of the box. *Trends in Parasitology* **28**, 305. <https://doi.org/10.1016/j.pt.2012.05.004>
- [13] Stensvold CR and Clark CG (2020) Pre-empting Pandora's Box: *Blastocystis* subtypes revisited. *Trends in Parasitology* **36**, 229–232. <https://doi.org/10.1016/j.pt.2019.12.009>
- [14] Zhao GH, Hu XF, Liu TL, Hu RS, Yu ZQ, Yang WB, Wu YL, Yu SK and Song JK (2017) Molecular characterization of *Blastocystis* sp. in captive wild animals in Qinling Mountains. *Parasitology Research* **116**, 2327–2333. <https://doi.org/10.1007/s00436-017-5506-y>
- [15] Jinatham V, Maxamhud S, Popluechai S, Tsaousis AD and Gentekaki E (2021) *Blastocystis* one health approach in a rural community of northern Thailand: Prevalence, subtypes and novel transmission routes. *Frontiers in Microbiology* **12**, 746340. Available at <https://europepmc.org/articles/PMC8696170>.
- [16] Khaled S, Gantois N, Ly AT, Senghor S, Even G, Dautel E, Dejager R, Sawant M, Baydoun M, Benamrouz-Vanneste S, Chabé M, Ndiaye S, Schacht AM, Certad G, Riveau G and Viscogliosi E (2020) Prevalence and subtype distribution of *Blastocystis* sp. in Senegalese school children. *Microorganisms* **8**, 1–17. <https://doi.org/10.3390/microorganisms8091408>
- [17] Osorio-Pulgarin MI, Higuera A, Beltran-Álzate JC, Sánchez-Jiménez M and Ramírez JD (2021) Epidemiological and molecular characterization of *Blastocystis* infection in children attending daycare centers in Medellín, Colombia. *Biology (Basel)* **10**, 669. Available at <https://www.mdpi.com/2079-7737/10/7/669>.
- [18] Ramírez JD, Sánchez A, Hernández C, Flórez C, Bernal MC, Giraldo JC, Reyes P, López MC, García L, Cooper PJ, Vicuña Y, Mongi F and Casero RD (2016) Geographic distribution of human *Blastocystis* subtypes in South America. *Infection Genetics & Evolution* **41**, 32–35. Available at <https://www.sciencedirect.com/science/article/pii/S1567134816300867>.
- [19] Baek S, Maloney JG, Molokin A, George NS, Cortés Vecino JA and Santin M (2022) Diversity of *Blastocystis* subtypes in horses in Colombia and identification of two new subtypes. *Microorganisms* **10**, 1693.
- [20] Maloney JG, Molokin A, Seguí R, Maravilla P, Martínez-Hernández F and Villalobos G, Tsaousis AD, Gentekaki E, Muñoz-Antolí C, Klisiowicz DR, Oishi CY, Toledo R, Esteban JG, Köster PC, de Lucio A, Dashti A, Bailo B, Calero-Bernal R, González-Barrio D, Carmena D and Santín M (2023) Identification and molecular characterization of four new *Blastocystis* subtypes designated ST35–ST38. *Microorganisms* **11**, 46.
- [21] Alfellani MA, Stensvold CR, Vidal-Lapiedra A, Onuoha ESU, Fagbenro-Beyioku AF and Clark CG (2013) Variable geographic distribution of *Blastocystis* subtypes and its potential implications. *Acta Tropica* **126**, 11–18. Available at <https://linkinghub.elsevier.com/retrieve/pii/S0001706X12003993>.
- [22] Stensvold CR and Clark CG (2016) Current status of *Blastocystis*: A personal view. *Parasitology International* **65**, 763–771. Available at <https://linkinghub.elsevier.com/retrieve/pii/S1383576916301544>.
- [23] Stensvold CR and van der Giezen M (2018) Associations between gut microbiota and common luminal intestinal parasites. *Trends in Parasitology* **34**, 369–377. <https://doi.org/10.1016/j.pt.2018.02.004>
- [24] Tito RY, Chaffron S, Caenepeel C, Lima-Mendez G, Wang J, Vieira-Silva S, Falony G, Hildebrand F, Darzi Y, Rymenans L, Verspecht C, Bork P, Vermeire S, Joossens M and Raes J (2019) Population-level analysis of *Blastocystis* subtype prevalence and variation in the human gut microbiota. *Gut* **68**, 1180–1189. <https://doi.org/10.1136/gutjnl-2018-316106>
- [25] Krogsgaard LR, Engsbro AL, Stensvold CR, Nielsen HV and Bytzer P (2015) The prevalence of intestinal parasites is not greater among individuals with irritable bowel syndrome: A population-based case-control study. *Clinical Gastroenterology & Hepatology* **13**, 507–513.e2. <https://doi.org/10.1016/j.cgh.2014.07.065>
- [26] Jimenez-Gonzalez DE, Martinez-Flores WA, Reyes-Gordillo J, Ramirez-Miranda ME, Arroyo-Escalante S, Romero-Valdivinos M, Stark D, Souza-Saldivar V, Martinez-Hernandez F, Flisser A, Olivio-Diaz A and Maravilla P (2012) *Blastocystis* infection is associated with irritable bowel syndrome in a Mexican patient population. *Parasitology Research* **110**, 1269–1275. <https://doi.org/10.1007/s00436-011-2626-7>
- [27] Poirier P, Wawrzyniak I, Vivarès CP, Delbac F and el Alaoui H (2012) New insights into *Blastocystis* spp.: A potential link with irritable bowel syndrome. *PLoS Pathogens* **8**, e1002545. <https://doi.org/10.1371/journal.ppat.1002545>
- [28] Dogruman-al F, Simsek Z, Boorum K, Ekici E, Sahin M, Tuncer C, Kustimur S and Altinbas A (2010) Comparison of methods for detection of *Blastocystis* infection in routinely submitted stool samples, and also in IBS/IBD patients in Ankara, Turkey. *PLoS One* **5**, e15484. <https://doi.org/10.1371/journal.pone.0015484>
- [29] Giacometti A, Cirioni O, Fiorentini A, Fortuna M and Scalise G (1999) Irritable bowel syndrome in patients with *Blastocystis hominis* infection. *European Journal of Clinical Microbiology & Infectious Diseases* **18**, 436–439. <https://doi.org/10.1007/s100960050314>
- [30] Yakob J, Jafri W, Beg MA, Abbas Z, Naz S, Islam M and Khan R (2010) *Blastocystis hominis* and *Dientamoeba fragilis* in patients fulfilling irritable bowel syndrome criteria. *Parasitology Research* **107**, 679–684. <https://doi.org/10.1007/s00436-010-1918-7>
- [31] Yakob J, Jafri W, Beg MA, Abbas Z, Naz S, Islam M and Khan R (2010) Irritable bowel syndrome: Is it associated with genotypes of *Blastocystis hominis*. *Parasitology Research* **106**, 1033–1038 <https://doi.org/10.1007/s00436-010-1761-x>
- [32] Mirza H and Tan KSW (2009) *Blastocystis* exhibits inter- and intra-subtype variation in cysteine protease activity. *Parasitology Research* **104**, 355–361 <https://doi.org/10.1007/s00436-008-1203-1>
- [33] Scanlan PD (2012) *Blastocystis*: Past pitfalls and future perspectives. *Trends in Parasitology* **28**, 327–334. <https://doi.org/10.1016/j.pt.2012.05.001>
- [34] Tan KSW, Mirza H, Teo JDW, Wu B and MacAry PA (2010) Current views on the clinical relevance of *Blastocystis* spp. *Current Infectious Disease Reports* **12**, 28–35. <https://doi.org/10.1007/s11908-009-0073-8>
- [35] Beghini F, Pasolli E, Truong TD, Putignani L, Cacciò SM and Segata N (2017) Large-scale comparative metagenomics of *Blastocystis*, a common member of the human gut microbiome. *The ISME Journal* **11**, 2848–2863. <https://doi.org/10.1038/ismej.2017.139>
- [36] el Safadi D, Gaayeb L, Meloni D, Cian A, Poirier P, Wawrzyniak I, Delbac F, Dabboussi F, Delhaes L, Seck M, Hamze M, Riveau G and Viscogliosi E (2014) Children of Senegal River Basin show the highest prevalence of *Blastocystis* sp. ever observed worldwide. *BMC Infectious Diseases* **14**, 1–11. <https://doi.org/10.1186/1471-2334-14-164>
- [37] Jiménez PA, Jaimes JE and Ramírez JD (2019) A summary of *Blastocystis* subtypes in North and South America. *Parasites & Vectors* **12**, 376. <https://doi.org/10.1186/s13071-019-3641-2>
- [38] Roberts T, Stark D, Harkness J and Ellis J (2013) Subtype distribution of *Blastocystis* isolates from a variety of animals from New South Wales, Australia. *Veterinary Parasitology* **196**, 85–89. Available at <https://linkinghub.elsevier.com/retrieve/pii/S0304401713000368>.
- [39] Yowang A, Tsaousis AD, Chumphonsuk T, Thongsin N, Kullawong N, Popluechai S and Gentekaki E (2018) High diversity of *Blastocystis* subtypes isolated from asymptomatic adults living in Chiang Rai,

- Thailand. *Infection Genetics & Evolution* **65**, 270–275. <https://doi.org/10.1016/j.meegid.2018.08.010>
- [40] El Fatni C, Olmo F, El Fatni H, Romero D and Rosales MJ (2014) First genotyping of *Giardia duodenalis* and prevalence of enteroparasites in children from Tetouan (Morocco). *Parasite* **21**, 48. <https://doi.org/10.1051/parasite/2014049>
- [41] Nithyamathi K, Chandramathi S and Kumar S (2016) Predominance of *Blastocystis* sp. infection among school children in Peninsular Malaysia. *PLoS One* **11**, 1–14. <https://doi.org/10.1371/journal.pone.0136709>
- [42] de Boer MD, Schuurs TA, Vermeer M, Ruijs GJHM, van der Zanden AGM, Weel JF and Bruijnesteijn van Coppenraet LES (2020) Distribution and relevance of *Dientamoeba fragilis* and *Blastocystis* species in gastroenteritis: Results from a case–control study. *European Journal of Clinical Microbiology & Infectious Diseases* **39**, 197–203. <https://doi.org/10.1007/s10096-019-03710-z>
- [43] Gong B, Liu X, Wu Y, Xu N, Xu M, Yang F, Tong L, Zhou K, Cao J, Liu A and Shen Y (2019) Prevalence and subtype distribution of *Blastocystis* in ethnic minority groups on both sides of the China–Myanmar border, and assessment of risk factors. *Parasite* **26**. <https://doi.org/10.1051/parasite/2019046>
- [44] Efunshile AM, Nelson JA, Stensvold CR and Poulsen CS (2016) Epidemiological aspects of *Blastocystis* colonisation in children in Ilero, Nigeria. *American Journal of Tropical Medicine & Hygiene* **95**, 175–179. <https://doi.org/10.4269/ajtmh.16-0074>
- [45] Sari IP, Benung MR, Wahdini S and Kurniawan A (2018) Diagnosis and identification of *Blastocystis* subtypes in primary school children in Jakarta. *Journal of Tropical Pediatrics* **64**, 208–214. <https://doi.org/10.1093/tropej/fmx051>
- [46] Liao CW, Chiu KC, Chiang IC, Cheng PC, Chuang TW, Kuo JH, Tu YH and Fan CK (2017) Prevalence and risk factors for intestinal parasitic infection in schoolchildren in Battambang, Cambodia. *American Journal of Tropical Medicine & Hygiene* **96**, 583–588. <https://doi.org/10.4269/ajtmh.16-0681>
- [47] Diarthini NLPE, Swastika IK, Ariwati L, Isyaputri R, Fitri N MY, Hidajati S and Basuki S (2018) *Blastocystis* and other intestinal parasites infections in elementary school children in Dukuh Village, Karangasem District, Bali. *Indonesian Journal of Tropical & Infectious Disease* **7**, 57. <https://doi.org/10.20473/ijtid.v7i3.7323>
- [48] Rebolla MF, Silva EM, Gomes JF, Falcão AX, Rebolla MV and Franco RM (2016) High prevalence of *Blastocystis* spp. infection in children and staff members attending public urban schools in São Paulo State, Brazil. *Revista do Instituto de Medicina Tropical de Sao Paulo* **58**, 31. <https://doi.org/10.1590/S1678-9946201658031>
- [49] Perea M, Vásquez V, Pineda V, Samudio F, Calzada JE and Saldaña A (2020) Prevalence and subtype distribution of *Blastocystis* sp. infecting children from a rural community in Panama. *Parasite Epidemiology & Control* **9**, e00139. <https://doi.org/10.1016/j.parepi.2020.e00139>
- [50] Leelayoova S, Mungthin M, Aunpad R, Naaglor T, Rangsin R and Pipatsatitpong D (2015) Prevalence and risk factors for *Blastocystis* infection among children and caregivers in a child care center, Bangkok, Thailand. *American Journal of Tropical Medicine & Hygiene* **93**, 310–315. <https://doi.org/10.4269/ajtmh.14-0492>
- [51] Qi M, Wei Z, Zhang Y, Zhang Q, Li J, Zhang L and Wang R (2020) Genetic diversity of *Blastocystis* in kindergarten children in southern Xinjiang, China. *Parasites & Vectors* **13**, 15. <https://doi.org/10.1186/s13071-020-3890-0>
- [52] Yoshikawa H, Wu Z, Pandey K, Pandey BD, Sherchand JB, Yanagi T and Kanbara H (2009) Molecular characterization of *Blastocystis* isolates from children and rhesus monkeys in Kathmandu, Nepal. *Veterinary Parasitology* **160**, 295–300. Available at <https://linkinghub.elsevier.com/retrieve/pii/S0304401708006742>.
- [53] Dacal E, Saugar JM, de Lucio A, Hernández-de-Mingo M, Robinson E, Köster PC, Aznar-Ruiz-de-Alegria ML, Espasa M, Ninda A, Gandasegui J, Sulleiro E, Moreno M, Salvador F, Molina I, Rodríguez E and Carmena D (2018) Prevalence and molecular characterization of *Strongyloides stercoralis*, *Giardia duodenalis*, *Cryptosporidium* spp., and *Blastocystis* spp. isolates in school children in Cubal, Western Angola. *Parasites & Vectors* **11**, 1–18. <https://doi.org/10.1186/s13071-018-2640-z>
- [54] Ramírez JD, Flórez C, Olivera M, Bernal MC and Giraldo JC (2017) *Blastocystis* subtyping and its association with intestinal parasites in children from different geographical regions of Colombia. *PLoS One* **12**, 1–13. <https://doi.org/10.1371/journal.pone.0172586>
- [55] Sánchez A, Muñoz M, Gómez N, Tabares J, Segura L, Salazar Á, Restrepo C, Ruiz M, Reyes P, Qian Y, Xiao L, López MC and Ramírez JD (2017) Molecular epidemiology of *Giardia*, *Blastocystis* and *Cryptosporidium* among indigenous children from the Colombian Amazon basin. *Frontiers in Microbiology* **8**, 1–14. <https://doi.org/10.3389/fmicb.2017.00248>
- [56] Assavapongpaiboon B, Bunkasem U, Sanprasert V and Nuchprayoon S (2018) A cross-sectional study on intestinal parasitic infections in children in suburban public primary schools, Saraburi, the central region of Thailand. *American Journal of Tropical Medicine & Hygiene* **98**, 763–767. <https://doi.org/10.4269/ajtmh.17-0240>
- [57] Naaglor T, Siripattanapipong S, Mungthin M, Thathaisong U, Taamasri P, Leelayoova S and Piyaraj P (2008) Drinking water: A possible source of *Blastocystis* spp. subtype 1 infection in schoolchildren of a rural community in central Thailand. *American Journal of Tropical Medicine & Hygiene* **79**, 401–406.
- [58] Villamizar X, Higuera A, Herrera G, Vasquez-A LR, Buitron L, Muñoz LM, Gonzalez-C FE, Lopez MC and Giraldo JC, Ramírez JD (2019) Molecular and descriptive epidemiology of intestinal protozoan parasites of children and their pets in Cauca, Colombia: A cross-sectional study. *BMC Infectious Diseases* **19**, 190. <https://doi.org/10.1186/s12879-019-3810-0>
- [59] Diamond LS (1987) A new liquid medium for xenic cultivation of *Eatamoeba histolytica* and other lumen-dwelling protozoa. *Journal of Parasitology* **68**, 958–959.
- [60] Diamond LS (1957) The Establishment of various trichomonads of animals and man in axenic cultures. *Journal of Parasitology* **43**, 488–490. Available at <http://www.jstor.org/stable/3274682>.
- [61] Clark CG (1997) Extensive genetic diversity in *Blastocystis hominis*. *Molecular & Biochemical Parasitology* **87**, 79–83. [https://doi.org/10.1016/s0166-6851\(97\)00046-7](https://doi.org/10.1016/s0166-6851(97)00046-7)
- [62] Scicluna SM, Tawari B and Clark CG (2006) DNA barcoding of *Blastocystis*. *Protist* **157**, 77–85. <https://doi.org/10.1016/j.protis.2005.12.001>
- [63] Poirier P, Wawrzyniak I, Albert A, el Alaoui H, Delbac F and Livrelli V (2011) Development and evaluation of a real-time PCR assay for detection and quantification of *Blastocystis* parasites in human stool samples: Prospective study of patients with hematological malignancies. *Journal of Clinical Microbiology* **49**, 975–983. <https://doi.org/10.1128/JCM.01392-10>
- [64] Larsson A (2014) AliView: A fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**, 3276–3278. <https://doi.org/10.1093/bioinformatics/btu531>
- [65] Katoh K and Toh H (2010) Parallelization of the MAFFT multiple sequence alignment program. *Bioinformatics* **26**, 1899–1900. <https://doi.org/10.1093/bioinformatics/btq224>
- [66] Capella-Gutiérrez S, Silla-Martínez JM and Gabaldón T (2009) trimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- [67] Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *2010 Gateway Computing Environments Workshop (GCE)*. New York: Cold Spring Harbor Laboratory Press, pp. 1–8.
- [68] Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* **16**, 111–120. <https://doi.org/10.1007/BF01731581>
- [69] Stecher G, Tamura K and Kumar S (2020) Molecular evolutionary genetics analysis (MEGA) for macOS. *Molecular Biology & Evolution* **37**, 1237–1239. Available at <https://academic.oup.com/mbe/article/37/4/1237/5697095>.
- [70] Tamura K, Stecher G and Kumar S (2021) MEGA 11: Molecular evolutionary genetics analysis version 11. *Molecular Biology & Evolution* **38**, 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- [71] Lê S, Josse J and Husson F (2008) FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software* **25**, 1–18. Available at <https://www.jstatsoft.org/index.php/jss/article/view/v025i01>.

- [72] Kassambara A and Mundt F (2020) Factoextra: Extract and visualise the results of multivariate data analyses. R Package Version 1.0.7. Available at <https://CRAN.R-project.org/package=factoextra>.
- [73] Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw A, Lumley T, Maechler M, Magnusson A, Moeller S, Schwartz M and Venables B (2020) gplots: Various R programming tools for plotting data. R package version 3.1.0.
- [74] R Core Team (2020) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at <https://www.r-project.org/>.
- [75] Pipatsatitpong D, Rangsin R, Leelayoova S, Naaglor T and Mungthin M (2012) Incidence and risk factors of *Blastocystis* infection in an orphanage in Bangkok, Thailand. *Parasites & Vectors* 5, 37. <https://doi.org/10.1186/1756-3305-5-37>
- [76] Leelayoova S, Pipatsatitpong D, Siripattanapipong S, Naaglor T, Mungthin M, Thathaisong U and Tan-ariya P (2013) Identification of *Blastocystis* subtype 1 variants in the home for girls, Bangkok, Thailand. *American Journal of Tropical Medicine & Hygiene* 88, 352–358. <https://doi.org/10.4269/ajtmh.2012.12-0237>
- [77] Yaicharoen R, Ngrenngarmert W, Wongjindanon N, Sripochang S and Kiatfuengfoo R (2006) Infection of *Blastocystis hominis* in primary schoolchildren from Nakhon Pathom province, Thailand. *Tropical Biomedicine* 23, 117–122.
- [78] Ruang-areerate T, Suwannahitatorn P, Sirirungreung A, Thita T, Naaglor T, Sitthichot N, Hempatawee N, Piyaraj P, Rangsin R, Taamasri P, Leelayoova S and Mungthin M (2019) Prevalence and distribution of *Blastocystis* infection among children from four primary schools after water treatment in rural Thailand. *The Southeast Asian Journal of Tropical Medicine & Public Health* 50, 217–225.
- [79] Oliveira-Arbex AP, David ÉB and Guimaraes S (2018) *Blastocystis* genetic diversity among children of low-income daycare center in Southeastern Brazil. *Infection Genetics & Evolution* 57, 59–63. <https://doi.org/10.1016/j.meegid.2017.11.005>
- [80] Abarca N, Santín M, Ortega S, Maloney JG, George NS, Molokin A, Cardona GA, Dashti A, Köster PC, Bailo B, Hernández-de-Mingo M, Muadica AS, Calero-Bernal R, Carmena D and González-Barrio D (2021) Molecular detection and characterization of *Blastocystis* sp. and *Enterocytozoon bieneusi* in cattle in northern Spain. *Veterinary Science* 8, 191. <https://doi.org/10.3390/vetsci8090191>
- [81] Wang X, Xue NY, Qin LT, Liu YY, Wang HX, Zhao Q, Ni HB and Lyu C (2021) Molecular characterization of *Blastocystis* from beef cattle in Northeastern China. *Vector-Borne & Zoonotic Diseases* 21, 955–960. <https://doi.org/10.1089/vbz.2021.0056>
- [82] Forsell J, Bengtsson-Palme J, Angelin M, Johansson A, Evengård B and Granlund M (2017) The relation between *Blastocystis* and the intestinal microbiota in Swedish travellers. *BMC Microbiology* 17, 1–9. <https://doi.org/10.1186/s12866-017-1139-7>
- [83] Scanlan PD, Hill CJ, Ross RP, Ryan CA, Stanton C and Cotter PD (2018) The intestinal protist *Blastocystis* is not a common member of the healthy infant gut microbiota in a Westernized country (Ireland). *Parasitology* 145, 1274–1278. <https://doi.org/10.1017/S0031182018000033>
- [84] Scanlan PD, Stensvold CR, Rajilić-Stojanović M, Heilig HGHJ, de Vos WM, O'Toole PW and Cotter PD (2014) The microbial eukaryote *Blastocystis* is a prevalent and diverse member of the healthy human gut microbiota. *FEMS Microbiology Ecology* 90, 326–330. <https://doi.org/10.1111/1574-6941.12396>
- [85] Hernández-Castro C, Dashti A, Vusirikala A, Balasegaram S, Köster PC, Bailo B, Imaña E, López A, Llorente MT, González-Barrio D, Sánchez S and Carmena D (2023) Prevalence and temporal dynamics of *Cryptosporidium* spp., *Giardia duodenalis*, and *Blastocystis* sp. among toddlers attending day-care centres in Spain. A prospective molecular-based longitudinal study. *European Journal of Pediatrics* 182, 213–223. <https://doi.org/10.1007/s00431-022-04662-x>
- [86] Greige S, el Safadi D, Bécu N, Gantois N, Pereira B, Chabé M, Benamrouz-Vanneste S, Certad G, el Hage R, Chemaly M, Hamze M and Viscogliosi E (2018) Prevalence and subtype distribution of *Blastocystis* sp. isolates from poultry in Lebanon and evidence of zoonotic potential. *Parasites & Vectors* 11, 1–10. <https://doi.org/10.1186/s13071-018-2975-5>
- [87] Udonsom R, Prasertbun R, Mahittikorn A, Mori H, Changbunjong T, Komalamisra C, Pintong AR, Sukthana Y and Popruk S (2018) *Blastocystis* infection and subtype distribution in humans, cattle, goats, and pigs in central and western Thailand. *Infection Genetics & Evolution* 65, 107–111. Available at <https://linkinghub.elsevier.com/retrieve/pii/S1567134818304805>.
- [88] Parkar U, Traub RJ, Vitali S, Elliot A, Levecke B, Robertson I, Geurden T, Steele J, Drake B and Thompson RCA (2010) Molecular characterization of *Blastocystis* isolates from zoo animals and their animal-keepers. *Veterinary Parasitology* 169, 8–17. <https://doi.org/10.1016/j.vetpar.2009.12.032>
- [89] Wang W, Owen H, Traub RJ, Cuttler L, Inpankaew T and Bielefeldt-Ohmann H (2014) Molecular epidemiology of *Blastocystis* in pigs and their in-contact humans in Southeast Queensland, Australia, and Cambodia. *Veterinary Parasitology* 203, 264–269. <https://doi.org/10.1016/j.vetpar.2014.04.006>
- [90] Yan Y, Su S, Ye J, Lai X, Lai R, Liao H, Chen G, Zhang R, Hou Z and Luo X (2007) *Blastocystis* sp. subtype 5: A possibly zoonotic genotype. *Parasitology Research* 101, 1527–1532. <https://doi.org/10.1007/s00436-007-0672-y>
- [91] Jinatham V, Nonebudsri C, Wandee T, Popluechai S, Tsaousis AD and Gentekaki E (2022) *Blastocystis* in tap water of a community in northern Thailand. *Parasitology International* 91, 102624. <https://doi.org/10.1016/j.parint.2022.102624>
- [92] Adamska M (2022) First report of *Blastocystis* sp. subtypes in natural water bodies in north-western Poland: A one-year monitoring. *International Journal of Environmental Health Research* 32, 862–869. <https://doi.org/10.1080/09603123.2020.1803804>
- [93] Martín-Escolano R, Ng GC, Tan KSW, Stensvold CR, Gentekaki E and Tsaousis AD (2023) Resistance of *Blastocystis* to chlorine and hydrogen peroxide. *Parasitology Research* 122, 167–176. <https://doi.org/10.1007/s00436-022-07713-2>
- [94] Tsaousis AD, Hamblin KA, Elliott CR, Young L, Rosell-Hidalgo A, Gourlay CW, Moore AL and van der Giezen M (2018) The human gut coloniser *Blastocystis* respire using complex II and alternative oxidase to buffer transient oxygen fluctuations in the gut. *Frontiers in Cellular & Infection Microbiology* 8, 371. <https://doi.org/10.3389/fcimb.2018.00371>
- [95] Andersen LOB and Stensvold CR (2016) *Blastocystis* in health and disease: Are we moving from a clinical to a public health perspective? *Journal of Clinical Microbiology* 54, 524–528. <https://doi.org/10.1128/JCM.02520-15>
- [96] Mirjalali H, Latifi A, Taghipour A, Yadegar A, Hatami B, Sadeghi A, Ehsani MJ and Zali MR (2020) Association between *Blastocystis* and body mass index in healthy subjects; a theoretical pilot study. *Journal of Diabetes & Metabolic Disorders* 19, 129–134. <https://doi.org/10.1007/s40200-019-00483-2>
- [97] Andersen LOB, Bonde I, Nielsen HB and Stensvold CR (2015) A retrospective metagenomics approach to studying *Blastocystis*. *FEMS Microbiology Ecology* 91, 1–9. <https://doi.org/10.1093/femsec/fiv072>
- [98] Asnicar F, Berry SE, Valdes AM, Nguyen LH, Piccinno G, Drew DA, Leeming E, Gibson R, le Roy C, Khatib HA, Francis L, Mazidi M, Mompeo O, Valles-Colomer M, Tett A, Beghini F, Dubois L, Bazzani D, Thomas AM, Mirzayi C, Khleborodova A, Oh S, Hine R, Bonnett C, Capdevila J, Danzanvilliers S, Giordano F, Geistlinger L, Waldron L, Davies R, Hadjigeorgiou G, Wolf J, Ordovás JM, Gardner C, Franks PW, Chan AT, Huttenhower C, Spector TD and Segata N (2021) Microbiome connections with host metabolism and habitual diet from 1,098 deeply phenotyped individuals. *Nature Medicine* 27, 321–332. <https://doi.org/10.1038/s41591-020-01183-8>
- [99] Betts EL, Newton JM, Thompson GS, Sarzhanov F, Jinatham V, Kim MJ, Popluechai S, Dogruman-al F, Won EJ, Gentekaki E and Tsaousis AD (2021) Metabolic fluctuations in the human stool obtained from *Blastocystis* carriers and non-carriers. *Metabolites* 11, 883. <https://doi.org/10.3390/metabo11120883>