

GLACIER OUTBURST FLOODS AT MOUNT RAINIER, WASHINGTON STATE, U.S.A.

by

Carolyn L. Driedger

(U.S. Geological Survey, Pacific Northwest District, Suite 600, 1201 Pacific Avenue,
Tacoma, WA 98402, U.S.A.)

and

Andrew G. Fountain

(U.S. Geological Survey, P.O. Box 25046, MS-412 Denver, CO 80225, U.S.A.)

ABSTRACT

During the twentieth century, glacial outburst floods have been the most destructive natural events on Mount Rainier, a stratovolcano in the Cascade Range in Washington State, U.S.A. In the period between 1930 and 1980 numerous floods were reported from five glaciers on the mountain, most of which flowed from Nisqually, Kautz, or South Tahoma Glaciers on its southern flank. Such floods threaten lives and property because they occur without warning and quickly mobilize the loose volcanic debris into debris flows.

A monitoring program was begun in 1987 which was designed to measure the dimensions and timing of outburst floods, but this has been unsuccessful because no floods have yet occurred on the monitored streams. Four floods did burst from South Tahoma Glacier that was unmonitored, but in spite of this they have been useful in providing evidence of flood storage and release mechanisms. All flood volumes were found to be of approximately similar orders of magnitude, of $1 \times 10^5 \text{ m}^3$ of water, indicating that all floods probably had similar mechanisms for storage and release of water. Hydraulic pressure considerations indicate that such a large volume of flood water would be stored at the bed of the glacier rather than in isolated englacial cavities. The stepped bedrock terrain provides an ideal setting for the formation of subglacial cavities capable of storing the volumes of flood water noted.

INTRODUCTION

Glacier outburst floods, jökulhlaups, are sudden discharges from water bodies dammed within or at the margins of glaciers. At Mount Rainier, the largest dormant volcano in the central Cascade Range of Washington State, U.S.A., one or more glacier outburst floods may occur in any given year, although in some years none occur. Glacial streams on Mount Rainier erode through loose volcanic debris and sudden increases in discharge such as those that occur in outburst floods commonly mobilize this eroded material as debris flows. In the geological record for Mount Rainier, and similarly for other Cascade volcanoes such as Mount Shasta, California, U.S.A., stratigraphic evidence of debris flows (Osterkamp and others, 1985) indicates that these flows are seldom associated with eruptive activity (Crandell, 1971). The outburst floods at Mount Rainier apparently originate from below the glacier surface and no ice-dammed lakes have been observed on the mountain.

This paper presents the results of an analysis of the historical flood records for the Mount Rainier National Park, and theorizes about the source, storage, and release of outburst water.

THE SETTING

Mount Rainier rises to 4400 m a.s.l. and is a strato-volcano composed of overlapping layers of lava and tephra deposits capped by glaciers (Fig. 1). It last erupted in the mid-nineteenth century (Crandell, 1971), and some low-level geothermal activity persists. Twenty-five glaciers exist on Mount Rainier, with a total area of 92 km^2 of snow and ice cover (Driedger and Kennard, 1986). The morphological features of five of Mount Rainier's flood-producing glaciers, Nisqually, Kautz, South Tahoma, Carbon, and Winthrop, are shown in Table I. Five major rivers have their origins in glaciers on Mount Rainier, Kautz Creek, Tahoma Creek, and the Nisqually River drain the major outburst-producing glaciers, and each has a summer discharge of approximately




Fig. 1. Glaciers and permanent snowfields cover about 92 km^2 of Mount Rainier. The three glaciers producing most floods, South Tahoma, Kautz, and Nisqually, are located on the southern flank of the mountain. Some floods have emanated from Winthrop Glacier, and outburst flood activity is also suspected on South and North Mowich Glaciers, and Carbon Glacier.

TABLE I. CHARACTERISTICS OF OUTBURST-FLOOD-PRODUCING GLACIERS, MOUNT RAINIER, WASHINGTON, U.S.A.

Glacier	Surface area (km ²)	Mean surface slope	Surface type	No. of floods in record
Nisqually	4.6	25°	alternating ice falls and basins	9
South Tahoma	2.8	23°	alternating ice falls and basins	12
Kautz-Success	1.8	29°	alternating ice falls and basins	5
Carbon	11.2	18°	steep accumulation area, lesser slope in ablation area	1
Winthrop	9.1	21°	steep accumulation area, lesser slope in ablation area	2

$1 \times 3 \text{ m}^3 \text{ s}^{-1}$. The three glaciers with the highest historically noted recurrence of floods, Nisqually, Kautz, and South Tacoma Glaciers, drain into the Nisqually River. Nisqually and Kautz Glaciers are situated on the southern flank of Mount Rainier, and both flow from near to the summit crater across a series of basins and cliffs. South Tahoma Glacier is located on the south-western flank of Mount Rainier and discharges water into Tahoma Creek. The glacier is bordered by ridges to the north and south, and also by a 1000 m high headwall. The most obvious feature of the glacier is a 1 km long stretch of stagnating ice that was partially disconnected from the glacier body by an earlier flood and now conceals the margin of the currently active terminus.

APPRAISAL OF RECORDS OF GLACIER OUTBURST FLOODS

Compilation of data from outburst events helps in identifying which glaciers are likely to be susceptible to outburst floods, and in determining their regional hazard potential (Post and Mayo, 1971; Young, 1980). The work of Haerberli (1983) correlates the morphological characteristics of glaciers, such as area and slope, with frequency and type of outburst. He has found that, in Switzerland, approximately two-thirds of all floods are caused by lake outbursts and the remaining one-third caused by outbursts from unobservable water pockets. The water-pocket floods have tended to originate from those glaciers with steeper-than-average slopes, which are usually greater than 15°. Frequency-distribution data indicate that both water-pocket and lake-outburst floods most usually occur in July and August.

The available information about past floods at Mount Rainier has been reported by Richardson (1968), by park personnel, and by others. Some of this information is based on intermittent observations of events by untrained observers. Many small floods have probably gone unrecorded, but Table II is thought to represent a reasonably complete record of the major events of this century. Richardson (1968) determined that floods most often occurred during the period of the year from August to October, with a frequency of about one every 3 to 10 years. Hodge (1972) noted some additional events to those which Richardson had reported, and suggested that Nisqually Glacier has generated floods once every 2 years. Recently, even frequency has increased to nine observed events between 1985 and 1987, but many other outburst floods have probably gone unnoticed because of their small volumes or remote locations. During the 87 years since the establishment of Mount Rainier National Park, floods and resulting debris flows have destroyed four highway bridges, several sections of highway, camping grounds and picnic areas, and numerous trail bridges (Richardson, 1968). The

transformation of an outburst flood into a debris flow increases its destructive potential. The large number of visitors, the increased number of flood events, and the recent destruction of visitors facilities by floods, have all drawn attention to the hazard.

Records for all glaciers on the mountain show that many of the floods occurred in a cluster in time but that, when averaged over the decades, the years 1967 and 1987 had a frequency exceeding one flood per year. However, from 1985 to 1987, the frequency of events averaged three floods per year. In all, there have been nine floods with peak discharges about $1 \times 10^2 \text{ m}^3 \text{ s}^{-1}$, 14 with peak values $1 \times 10^3 \text{ m}^3 \text{ s}^{-1}$, and one with a value of $1 \times 10^4 \text{ m}^3 \text{ s}^{-1}$. 60% of the outbursts reported produced debris flows, and 74% either damaged or destroyed Park facilities. 40% of the outburst floods occurred during or immediately after rainfall, and these were some of the largest reported. Several outburst floods were reported to have occurred in surges with periodicities ranging from several minutes to several hours. No precursory stream-flow events have been recorded. It was noted that many floods peaked during the afternoon or evening. The lack of evidence for localized melting indicates that intermittent local area geothermal activity is not responsible for recent flood formation. All floods reported, except for that occurring during 1987 on Winthrop Glacier, appeared to have originated beneath the glacier surface.

A field program was begun in 1987 in order to monitor both glacial outburst floods and possible precursory Sydrologic events occurring within the Park. Measurements of stream-gauge height, electrical conductivity, turbidity, and of meteorological variables were made at two sites each lying both on Kautz Creek and the Nisqually River. Gauge height and conductivity data were also collected from the non-glacierized Paradise River so as to provide a reference set for background conditions. In the summer of 1987 there were no outburst floods from the monitored basins, although four were released from the unmonitored South Tahoma Glacier. Several people witnessed flood events at South Tahoma Creek, and reported similar accounts of floods on 29 June, 28 August, 31 August, and 26 September 1987. The material of the flow was described as being brownish black in colour, with a viscosity similar to that of wet concrete, and the flow carried boulders of up to several meters in diameter. Furthermore, the flood wave was accompanied by strong winds and boulders of up to about 400 mm in diameter that were thrown above the railings of the suspension bridge which stands about 20 m above the stream bed. Down-valley, each of the four floods formed boulder levees along their margins of flow, and rerouted the original stream beds. They inundated a picnic area and a parking lot, deposited material ranging in depth from a few tenths of a meter to several meters of sediment, and buried signs and picnic tables.

All four floods at Tahoma Creek in 1987 occurred

TABLE II. GLACIER-OUTBURST FLOOD CHRONOLOGY

Nisqually Glacier						
Date	Type of event	Estimate of peak discharge ($\text{m}^3 \text{s}^{-1}$)	Weather	Effects of outburst	Remarks	References
October 1926	Debris flow	>100–200	Occurred during first heavy rain at end of summer	Destroyed Nisqually River road bridge	Outburst flood supplemented by rainfall	Richardson, 1968, p. 79
14 October 1932	Debris flow	>100–200	No rain from 25 September to 9 October. Rain 10, 14 October = 220 mm	Nisqually River road bridge destroyed	Debris flow 7 m high and 50 m wide. Landslide on lateral moraine resulted from outburst	Richardson, 1968, p. 79
24–25 October 1934	Debris flow	>100–200	20–25 October, rain = 252 mm	Moderate damage caused to bridge and motorized equipment	Water flooded in three or four distinct surges	Richardson, 1968, p. 80
1–2 October 1947	Water flood	>100–200	1500 mm rain at Paradise on 2 October 1947	High-water surge damaged motorized equipment; V-shaped gorge eroded into glacier	Outburst flood supplemented by rainfall	Richardson, 1968 p. 81
25 October 1955	Debris flow	2000	96 mm rain in previous 32 h	Concrete bridge removed; some damage at Longmire visitor facilities	Five or six surges of water occurred within 45 min. Ice and boulders in flow. Velocity estimated at 6.1 m s^{-1} ; 30% water by volume	Richardson, 1968 p. 82
2 June 1968	Water flood	<83	112 mm rain in 24 h	Damaged recorder	Peak discharge three times that of normal daily discharge	Hodge, 1972, p. 319
4 July 1970	Water flood	85	Dry	Stream-gauging station undermined	Two peaks observed on hydrograph about 2 h apart	Hodge, 1972, p. 315
14 July 1972	Water flood	140–170	(6 d period) 63 mm rain	Shelter and recorder on gauging station at Glacier Bridge destroyed	Probably included englacial waters, though there is no report of sudden release	Richardson, personal communication
22 June 1986	Water flood	>1.7	40 mm rain in previous month	None observed	Flood originated at base of active ice	Authors
South Tahoma Glacier						
29 August 1967	Water flood	3 (11 km downstream of glacier)	Exceptionally warm and dry summer. No rain in 2 months	Footbridge 1.9 km below glacier destroyed	Water emerged at ice fall at about 2 300 m a.s.l. Occurred at 08.40 h	Richardson, 1968, p. 83
31 August 1967	Debris flow	680	Exceptionally warm and dry summer. No rain in 2 months	Destroyed part of camping ground. Channel eroded in glacier; About $38 \times 10^3 \text{ m}^3$ of material eroded	Break-out at 2 300 m a.s.l. Flood had consistency of wet concrete, occurred during evening hours. 50% water flow. Flood dissipated 10.4 km below glacier	Richardson, 1968 p. 83
15 September 1967	Debris flow	<680	Exceptionally warm and dry summer. No rain in 2 months	No damage noted	Moved down-valley as far as the camping ground; noted as a <i>small lahar</i>	Crandell, 1971, p. 60
Summer 1968	Water floods	<680		No damage noted	... "Floods not associated with rainfall also moved down the valley from time to time during the summer of 1968." ...	Crandell, 1971, p. 60
21 August 1970	Water flood	1000	Dry weather, total rainfall <10 mm for month	Wonderland Trail Bridge destroyed	Flood inundated picnic area. Cline's slides show mudline about 2 m high below trail bridge	Crandell, 1971; and Cline USGS, Tacoma, WA (personal communication)
10 August 1971	Water flood	1000	Zero rain in month previous to flood	Damaged trees		Mount Rainier National Park collection
July–August 1979	Water flood	1000	Dry weather with brief periods of light rain	Water covered parts of the trail-head parking lot	Muddy water splashed on >10 m high cable railings of the suspension bridge	Gene Casey, National Park Service, Mount Rainier
August 1981	Water flood	1000	<250 mm rain for month	Low trail bridge was destroyed	2–3 m high wall of water seen by park personnel	C. Harvey via G. Casey, National Park Service, Mount Rainier
26 October 1986	Debris flow	1000	Dry weather in month preceded by 58 mm rain on day of flood and day previous	Trail bridge; parking areas; part of picnic area. Creek re-routed towards west	Debris flow dissipated 6 km down-valley of glacier	K. Scott, USGS, Vancouver, WA
29 June 1987	Debris flow	(approximately) 1000	18 mm rain during June	Rocks thrown over trail bridge 20 m above stream bed; Deposited 1+ m of mud in picnic area. Levees constructed 3–4 m over stream-water level; Destroyed picnic area	Concrete-like mass carried boulders, occurred 14.30 h; flood-wave of debris preceded by rush of wind; sound like aircraft	J. Fielding and M. Starkey, park visitors

Date	Type of event	Estimate of peak discharge (m ³ s ⁻¹)	Weather	Effects of outburst	Remarks	References
28 August 1987	Debris flow	1000	Unseasonably dry weather. 8 mm rain during August	Stream re-routed within existing channel	Occurred 17.00 h	Authors
31 August 1987	Debris flow	1000	Unseasonably dry weather. 8 mm rain during August	1 m aggradation of river bed at picnic area	Occurred during evening hours	Authors
26 September 1987	Debris flow	1000	Below normal rainfall, 41 mm rain in September before flood date	Re-routed stream bed towards west, small percentage flowed over highway; destroyed signs and outhouse	Velocity >2 m s ⁻¹ through parking lot; occurred 17.30 h	Authors
Kautz Glacier						
2 October 1947	Debris flow	Average daily discharge 280 peak discharge was "many times larger"	Abnormally warm summer. 149 mm of rain in 24 h during first day of flood	Eroded 20 m deep channel in glacier; covered large area of forest, highway and bridge with debris. Destroyed 1.5 km segment of glacier	Outburst flood triggered by "an abnormally intense downpour of rain." 40% of flood was water. Total <i>lahar</i> flood volume 38 × 10 ⁶ m ³ (Grater, 1947, 1948)	Richardson, 1968, p. 83
23 August 1961	"Surge of muddy water"	Discharge <1947 flood	Flood occurred during long dry	Cut stream bed 20–30 mm below its previous levee and destroyed two trail bridges	Debris flow was preceded by water flow over its top	Richardson, 1968, p. 83
September 1975	Debris flow	No estimate	Dry weather with only 3 mm rain in September	Debris flowed off moraine towards Van Trump Park.	Estimated volume of debris flow 3000 m ³	D.R. Cline, personal slides (USGS, Tacoma, WA)
20 July 1985	Debris flow	Similar to flood of 1986	Dry weather with <10 mm in month previous to flood	Destroyed trail bridge	Debris depth at trail bridge 1 m+	W.G. Sikonia (USGS, Tacoma, WA)
3–6 September 1986	Debris flow	84–114	Dry weather with 17 mm rain in month previous to flood	Destroyed trail bridge	Debris depth at trail bridge 1–2 m	K. Scott, (USGS Vancouver, WA)

during late afternoon or evening. Geomorphological evidence allowed an estimate of peak discharge for each of the four floods to be made; all were of the same order of magnitude, namely about 1 × 10³ m³ s⁻¹. The flood of 26 September mobilized debris up-stream, scouring the upper canyon to a depth of 2 m and depositing sediment farther down-stream to a depth of 1 m or more. Mean velocity of the surface flow was calculated as being 2–3 m s⁻¹. Mud lines on the lee side of trees were used to measure draw-down in a picnic area 7 km down-stream of the glacier.

THEORY OF SOURCE, STORAGE, AND RELEASE OF WATER

Because neither geothermal activity nor glacier-dammed lakes have been observed it is conjectured that water is stored within the glacier. Several possibilities are now examined. Water-filled crevasses are common in glaciers, and are often found in marginal shear zones. In order to account for flood volumes of 1 × 10⁵ m³, the value estimated from events in the summer of 1987 at South Tahoma Glacier, the numbers and volumes of crevasse reservoirs must be estimated. If a crevasse depth of 10 m, a surface width of 5 m, and a triangular cross-section are assumed, then 40 crevasses, each 100 m long, would be required to provide the estimated flood volume, the volume per unit length of a typical crevasse being 25 m³. No water-filled crevasses have been observed, making it improbable that the flood water originated at the glacier surface; the same problem of space-accounting applies to connected englacial voids.

The manner in which water is most probably ponded is in subglacial cavities. To account for the large volumes of flood water involved, subglacial cavities may be either small and numerous, or few and very large. Such cavities would be able to provide a sudden flux if water is held under high pressure and thereby relatively isolated from the low-pressure, subglacial hydraulic system. It is thought more likely that a few large cavities would remain isolated than

that many small ones would do so.

Given the rough terrain of Mount Rainier, and observations on the bedrock of unglaciated parts of the mountains, cavities could form in the lee of bedrock steps. Detailed descriptions of the mechanics of cavity formation under a variety of conditions have been given by Lliboutry (1968), Iken (1981), and Kamb (1987). If one quadrant of an elliptical profile in a cavity is assumed (Lliboutry, 1978), then volume per unit length (*V*) is given by

$$V = \frac{\pi}{4} hw \tag{1}$$

where *h* is the rock-step height and *w* is the cavity width. Breaks of slope occur in the subglacial terrain as the result of numerous overlapping lava flows; on Mount Rainier there are several steps each of roughly 30 m in height. In order to calculate potential water volume we assume a maximum cavity length of 10 m, for which the volume per unit width is 235 m³ m⁻¹. If this cavity extends along one-half of the distance across a glacier 1 km wide, as would be the case in South Tahoma Glacier where we believe the flood originated, the total cavity volume would be 1.5 × 10⁵ m³. Although this view is a gross simplification, it does provide the basis for a theory in which large cavities are feasible. A single cavity could possibly be solely responsible for the observed floods, although the presence of several smaller hydraulically linked cavities is more likely.

In this situation of relatively isolated cavities, the approach of Walder (1986) seems most applicable. The length, *L*, of a cavity in the ice-flow direction is the product of the time required for the ice to close an opening created by a step of height *R*, and the sliding speed of the glacier, so that

$$L = \frac{U_s R}{U_c} \tag{2}$$

where *U_c* is the creep-closure rate, and *U_s* is the sliding speed (Walder's equation (6)). Substituting bore-hole

closure calculation values (Nye, 1953) for creep-closure rate, ignoring any enlargement caused by heat generated from the viscous dissipation of energy from flowing water, and assuming an elliptical profile of the cavity, Equation (2) becomes

$$S = \frac{R}{4\pi} U_s \left(\frac{nA}{P_e} \right)^n \quad (3)$$

where S is the cross-sectional area, n and A are the exponent and coefficient of the ice-flow law, and P_e is effective pressure in the cavity (Walder's Equation (9), without the melting term). The size of the cavity is linearly related to step and to glacier sliding speed. Haerberli (1983) suggested that water-pocket floods are most frequently associated with steep glaciers, and this suggestion can be explained by proposing that the implied greater sliding speed creates larger cavities than would result on more gently sloping glaciers. The most important term in Equation (3) is the effective pressure, which has an inverse power relationship with cavity size. Unfortunately this is the term about which least direct information is available. Larger cavities will decrease the basal shear stress of a glacier and this will result in increased glacier speed (Iken, 1981), a fact which explains the conjecture (Hodge, 1972) that surface speed would increase prior to flood events.

The stated relationships now permit a qualitative hypothesis of flood generation from subglacial cavities to be offered. Glacier ice separates from its bed when it encounters a step in the bedrock and creates a cavity. If water flows into this cavity through a connection to a surface crevasse, or from a debris layer, the cavity will enlarge to new dimensions determined by a combination of step size, sliding speed, ice-overburden pressure and water pressure in the cavity, as indicated in Equation (3). If water pressure is high, as is suggested by studies of subglacial debris layers (Clarke, 1987), the cavity may become extremely large. The cavity will continue to enlarge itself until a connection has been established with a low-pressure hydraulic system, such as a conduit (Röthlisberger, 1972). Once the connection is made, water is forced from the cavity by the pressure difference between the cavity and conduit, and the flood proceeds as it would do from an ice-dammed lake. Subsequent increased water flux enlarges the conduit using the heat produced by viscous dissipation of energy in the flowing water. Large conduits allow greater water flux, resulting in more heat dissipation, until the cavity reservoir is nearly emptied. When water flux drops rapidly, the pressure of the overlying ice squeezes the cavity connection closed and ends the flood.

There are several implications arising from such a theory of subglacial cavity flood genesis. Hydrographs of stream flow will probably not show precursors of an impending flood, since the flood occurrence depends upon the sudden establishment of a subglacial hydraulic connection. A calculated glacier-water balance, similar to that of Tangborn and others (1975), would probably be useful as a means of indicating the magnitude of any sizeable water body already stored in a subglacial cavity. This model could also help to quantify the processes producing the observed higher frequency of floods in mid-to-late summer and leading to the storage of water during spring already estimated by Tangborn and others (1975).

The theory of flood genesis from subglacial cavities is also able to explain why a series of outburst events sometimes occurs on Mount Rainier. Once a flood has ended the cavities collapse from the pressure of the ice-overburden pressure and the seal at the cavity exit reforms. Water may continue to drain into the cavity if a connection is maintained either with the surface or with a debris layer on the bedrock steps. When pressure is sufficiently increased the cavity will again become enlarged, setting the stage for another event. As can be seen from Table II, floods do not necessarily occur in a cyclic manner. Their clustered or singular occurrence is dependent on melt-water flux, subglacial conditions, and the presence of cavity connections. Because of multiple possibilities, an accurate numerical model of flood occurrence would be difficult to construct.

CONCLUSION

Outburst floods on Mount Rainier are dangerous because they occur without warning and trigger debris flows. South Tahoma, Kautz, and Nisqually Glaciers generate the majority of the observed floods on the mountain, although it is recognized that small floods can occur in remote regions throughout the Park and that in any location they may go unnoticed. Glaciers on the south-western slopes of the mountain exhibit a greater flood frequency than those on the north-eastern side, probably because the nature of the stepped bedrock bench at 2100 m a.s.l. on the south-western side makes it more susceptible to cavity development, and also because the greater exposure to storms and more intense solar radiation on the south-western side of the mountain allows a greater water flux through Kautz, Nisqually, and South Tahoma Glaciers. The source of flood water is apparently within, or at the beds of, the glaciers and its release may be subject to the occurrence of appropriate changes in cavity size. This theory reasonably explains why no precursory hydrological events have been seen to precede an outburst occurrence and also why floods sometimes develop in series, perhaps with recurring cavity enlargements. Future modelling of glacier hydrology, and particularly of the subglacial drainage system, is a first step towards estimating flood timing and magnitude.

REFERENCES

- Clarke, G. 1987. Subglacial till: a physical framework for its properties and processes. *J. Geophys. Res.*, **92**(B9), 9023-9036.
- Crandell, D.R. 1971. Postglacial lahars from Mt. Rainier volcano, Washington. *U.S. Geol. Surv. Prof. Pap.* 677.
- Driedger, C.L. and P. Kennard. 1986. Ice volumes on the Cascade volcanoes: Mount Rainier, Mount Hood, Three Sisters, and Mount Shasta. *U.S. Geol. Surv. Prof. Pap.* 1365.
- Haerberli, W. 1983. Frequency and characteristics of glacier floods in the Swiss Alps. *Ann. Glaciol.*, **4**, 85-90.
- Hodge, S. Unpublished. The movement and basal conditions of the Nisqually Glacier, Mount Rainier. (Ph.D. thesis, University of Washington, 1972.)
- Iken, A. 1981. The effect of the subglacial water pressure on the sliding velocity of a glacier in an idealized numerical model. *J. Glaciol.*, **27**(97), 407-421.
- Kamb, B. 1987. Glacier surge mechanism based on linked cavity configuration of the basal water conduit system. *J. Geophys. Res.*, **92**(B9), 9083-9100.
- Llibouty, L. 1968. General theory of subglacial cavitation and sliding of temperate glaciers. *J. Glaciol.*, **7**(49), 21-58.
- Llibouty, L. 1978. Glissement d'un glacier sur un plan parsemé d'obstacles hémisphériques. *Ann. Geophys.*, **34**(2), 147-162.
- Nye, J. 1953. The flow law of ice from measurements in glacier tunnels, laboratory experiments and the Jungfraufirn borehole experiment. *Proc. R. Soc. London, Ser. A*, **219**(1139), 477-489.
- Osterkamp, W., C. Hupp, and J. Blodgett. 1985. Magnitude and frequency of debris flows and areas of hazard on Mount Shasta, northern California. *U.S. Geol. Surv. Open-File Rep.* 85-425.
- Post, A. and L. Mayo. 1971. Glacier dammed lakes and outburst floods in Alaska. *U.S. Geol. Surv. Hydrol. Invest. Atlas* HA-455.
- Richardson, D. 1968. Glacier outburst floods in the Pacific northwest. *U.S. Geol. Surv. Prof. Pap.* 600-D, D79-D86.
- Röthlisberger, H. 1972. Water pressure in intra- and subglacial channels. *J. Glaciol.*, **11**(62), 177-203.
- Tangborn, W.V., R.M. Krimmel, and M.F. Meier. 1975. A comparison of glacier mass-balance by glaciological, hydrological and mapping methods, South Cascade Glacier, Washington. *International Association of Hydrological Sciences Publication* 104 (*General Assembly of Moscow 1971 - Snow and Ice*), 185-196.
- Walder, J.S. 1986. Hydraulics of subglacial cavities. *J. Glaciol.*, **32**(112), 439-445.
- Young, G. 1980. Monitoring glacier outburst floods. *Nord. Hydrol.*, **11**(5), 285-300.