

# Bias in genomic predictions for populations under selection

Z. G. VITEZICA<sup>1\*</sup>, I. AGUILAR<sup>2</sup>, I. MISZTAL<sup>3</sup> AND A. LEGARRA<sup>4</sup>

<sup>1</sup> Université de Toulouse, UMR 1289 TANDEM, INRA/INPT-ENSAT/ENVT, F-31326 Castanet-Tolosan, France

<sup>2</sup> Instituto Nacional de Investigación Agropecuaria Las Brujas, Canelones 90200, Uruguay

<sup>3</sup> Department of Animal and Dairy Science, University of Georgia, Athens, Georgia 30602, USA

<sup>4</sup> INRA, UR 631 SAGA, F-31326 Castanet-Tolosan, France

(Received 27 January 2011; revised 27 March 2011; first published online 18 July 2011)

## Summary

Prediction of genetic merit or disease risk using genetic marker information is becoming a common practice for selection of livestock and plant species. For the successful application of genome-wide marker-assisted selection (GWMAS), genomic predictions should be accurate and unbiased. The effect of selection on bias and accuracy of genomic predictions was studied in two simulated animal populations under weak or strong selection and with several heritabilities. Prediction of genetic values was by best-linear unbiased prediction (BLUP) using data either from relatives summarized in pseudodata for genotyped individuals (multiple-step method) or using all available data jointly (single-step method). The single-step method combined genomic- and pedigree-based relationship matrices. Predictions by the multiple-step method were biased. Predictions by a single-step method were less biased and more accurate but under strong selection were less accurate. When genomic relationships were shifted by a constant, the single-step method was unbiased and the most accurate. The value of that constant, which adjusts for non-random selection of genotyped individuals, can be derived analytically.

## 1. Introduction

Selection of economically important quantitative traits in animals and plants is traditionally based on phenotypic records of an individual and its relatives. High-density panels of single-nucleotide polymorphisms (SNP) have recently been used to map genes of domestic animal species and select desirable livestock (Goddard & Hayes, 2009). The genetic merit of an individual can be estimated by genome-wide marker-assisted selection (GWMAS), also known as genomic selection (Meuwissen *et al.*, 2001). Such selection is expected to improve the precision of genetic merit predictions because some markers from a dense SNP panel will be in linkage disequilibrium with quantitative trait loci (QTLs) (Hayes *et al.*, 2009). Genomic selection will lead to faster genetic gain than that achieved with traditional selection methods based on pedigree and phenotypic data only.

Genome-wide evaluation methods use statistical tools to combine phenotypes with high-density

marker data to predict the genetic merit of individuals with complex traits. No agreement exists currently on which genome-wide evaluation method is the most appropriate (Daetwyler *et al.*, 2010; Hill, 2010). Most developments in genome-wide evaluation methods published to date assume that all animals have been genotyped. However, it is rarely the case, and most often non-genotyped close family members exist with phenotypic information (Garrick *et al.*, 2009). Ignoring this information results in less accurate predictions and possible bias because of selection (VanRaden *et al.*, 2009*a, b*). Unbiased predictions are of paramount importance in selection for accurate estimates of the genetic trend and also comparison of animals across generations (Henderson *et al.*, 1959). The acceptance and wide use of genomic predictions will largely depend on correct statistical modelling and ease of breeder application. Therefore, a genetic evaluation method that produces accurate and unbiased predictions is critically important.

Combining pedigree, phenotypic and marker information to calculate genomic predictions is a challenge. The number of genotyped individuals is extremely small compared with the total number of

\* Corresponding author: UMR 1289 TANDEM, ENSAT, Avenue de l'Agrobiopole, Postal Box 32607, 31326 Auzeville Tolosane, France. E-mail: zulma.vitezica@ensat.fr

individuals. Some genotyped individuals (as in dairy cattle) do not have a phenotype of their own. Thus, most proposed methods currently are based on multiple-step procedures (VanRaden *et al.*, 2009a). First, phenotypic data from relatives are summarized to create pseudodata for genotyped individuals (Garrick *et al.*, 2009); then in the second step, genomic predictions are computed by a genome-wide method from pseudodata and marker information. For example, in dairy cattle, pseudodata for males are measures of (precorrected) daughter performance called daughter yield deviations (DYD). The use of DYD may involve several problems (VanRaden *et al.*, 2009a,b): information loss of animals with few progeny, which leads to accuracy loss; heterogeneity caused by different amounts of information in the original dataset; and bias (caused by selection). For other species (e.g. sheep or swine) or traits (e.g. maternal traits), pseudodata are more difficult to estimate or even define. Thus, multiple-step methods of computing genomic predictions are not only complicated but likely sub-optimal for GWMAS.

Joint use of pedigree, phenotypic and genomic information should theoretically solve such problems. A single-step method based on a linear mixed model and a pedigree relationship matrix augmented with genomic information has been developed recently (Legarra *et al.*, 2009; Aguilar *et al.*, 2010; Christensen & Lund, 2010). This method combines pedigree and all available phenotypes and genotypes, needs no creation of pseudodata, and it has been applied to population sizes in the millions (Aguilar *et al.*, 2010). Further, the method is general with straightforward extension to other models or species.

In traditional pedigree-based evaluation, all information about selection decisions is included in phenotypes and the relationship matrix, and no bias exists from the selection (Sorensen & Kennedy, 1984; Im *et al.*, 1989). However, how selection is accounted for in GWMAS procedures is unclear, although this is becoming a serious concern (Aguilar *et al.*, 2010; Mäntysaari *et al.*, 2010; Chen *et al.*, 2011). Models for GWMAS implicitly assume an unselected genotyped population (Hayes *et al.*, 2009). However, in practice, genotyped individuals are highly selected, and GWMAS models do not take this selection into account. This is in contrast to classical procedures in which models refer to a base unselected population. Thus, GWMAS models appear to be unable to consider past selection based on pedigree and phenotypes, which might cause bias as well as accuracy loss.

The objective of this study was to investigate how unbiased genetic values can be predicted by GWMAS. The effects of selection and genome-wide evaluation method (single- or multiple-step) on bias and prediction accuracy were examined. The effect of trait heritability was also investigated.

## 2. Materials and methods

### (i) Theory

The single-step genomic prediction approach (Legarra *et al.*, 2009; Aguilar *et al.*, 2010; Christensen & Lund, 2010) is based on the model  $\mathbf{y} = \mathbf{X}\mathbf{b} + \mathbf{Z}\mathbf{u} + \mathbf{e}$ , where  $\mathbf{y}$  is the phenotype vector,  $\mathbf{X}$  and  $\mathbf{Z}$  are incidence matrices,  $\mathbf{b}$  denotes fixed effects,  $\mathbf{e}$  is the residual and  $p(\mathbf{u}) \sim N(\mathbf{0}, \mathbf{H}\sigma_u^2)$  involves the genetic effect for non-genotyped ( $\mathbf{u}_1$ ) and genotyped ( $\mathbf{u}_2$ ) individuals and the genetic variance  $\sigma_u^2$ . Here  $\mathbf{H}^{-1}$  is derived as in Legarra *et al.*, (2009) and Christensen & Lund (2010):

$$\mathbf{H}^{-1} = \begin{bmatrix} \mathbf{A}^{11} & \mathbf{A}^{12} \\ \mathbf{A}^{21} & \mathbf{A}^{22} + \mathbf{G}^{-1} - \mathbf{A}_{22}^{-1} \end{bmatrix}, \quad (1)$$

where  $\mathbf{G}$  is a genomic relationship matrix and  $\mathbf{A} = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{bmatrix}$  and  $\mathbf{A}^{-1} = \begin{bmatrix} \mathbf{A}^{11} & \mathbf{A}^{12} \\ \mathbf{A}^{21} & \mathbf{A}^{22} \end{bmatrix}$  are the pedigree-based relationship matrix and its inverse partitioned into non-genotyped and genotyped individuals, respectively. Creation of  $\mathbf{H}$  (and  $\mathbf{H}^{-1}$ ) can be seen as a projection of genetic merit (or marker genotypes) from genotyped to non-genotyped individuals using pedigree relationships (Legarra *et al.*, 2009; Christensen & Lund, 2010).

The pedigree relationship matrix  $\mathbf{A}$  implies that the mean genetic value of the base population is 0. Also, according to VanRaden (2008) and Hayes *et al.* (2009), the genomic relationship matrix  $\mathbf{G}$  automatically sets the mean genetic value of the genotyped population to 0 if raw means in the genotyped population are used to estimate current allele frequencies. That is not the case if frequencies for the base population are used, but accurate estimates of those frequencies are difficult to obtain.

If selection occurs, the mean genetic value of genotyped individuals ( $\mathbf{u}_2$ ) on the scale of the whole population (i.e. relative to the base population) may have a value different from 0, say  $\mu$ . This would be particularly true if genotyped individuals were elite individuals or in the last generations of selection. Let us assume that  $\mu$  is a random variable; for instance, in (conceptual) repetitions of the selection process,  $\mu$  will vary due to drift, in other words, because of finite sampling of replacement animals. Thus,  $p(\mu) \sim N(0, \alpha\sigma_u^2)$ ; and,  $p(\mathbf{u}_2|\mu) \sim N(\mathbf{1}\mu, \mathbf{G}\sigma_u^2)$ , where  $\mathbf{1}$  is a vector of ones. Equivalently,  $p(\mathbf{u}_2|\alpha) \sim N(\mathbf{0}, (\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha)\sigma_u^2)$ .

As in Legarra *et al.* (2009), the distribution of genetic values of non-genotyped individuals conditioned on genetic values of genotyped individuals (using multivariate normality) is  $p(\mathbf{u}_1|\mathbf{u}_2) \sim N(\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{u}_2, (\mathbf{A}^{11})^{-1}\sigma_u^2)$ , where  $(\mathbf{A}^{11})^{-1} = \mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}$ . Thus,  $p(\mathbf{u}_1|\alpha) \sim N(\mathbf{0}, (\mathbf{A}^{11})^{-1}\sigma_u^2 + \mathbf{A}_{12}\mathbf{A}_{22}^{-1}(\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha)\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\sigma_u^2)$  and  $p(\mathbf{u}) \sim N(\mathbf{0}, \mathbf{H}^\dagger\sigma_u^2)$ , where  $\mathbf{H}^\dagger$  is equivalent to  $\mathbf{H}$  with  $\mathbf{G}$  substituted for

$\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha$ . The mixed model equations are

$$\begin{bmatrix} \mathbf{X}'\mathbf{X} & \mathbf{X}'\mathbf{Z} \\ \mathbf{Z}'\mathbf{X} & \mathbf{Z}'\mathbf{Z} + \mathbf{H}^{\dagger-1}\lambda \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ \mathbf{u} \end{bmatrix} = \begin{bmatrix} \mathbf{X}'\mathbf{y} \\ \mathbf{Z}'\mathbf{y} \end{bmatrix},$$

where

$$\mathbf{H}^{\dagger-1} = \begin{bmatrix} \mathbf{A}^{11} & \mathbf{A}^{12} \\ \mathbf{A}^{21} & \mathbf{A}^{22} + (\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha)^{-1} - \mathbf{A}_{22}^{-1} \end{bmatrix}. \tag{2}$$

An equivalent model for unbiased genomic predictions based on a genetic group model (Quaas, 1988) is shown in the appendix.

To determine why  $\mu$  can be presumed as a random variable and obtain the value of  $\alpha$ , traditional best-linear unbiased prediction (BLUP) will be assumed to be unbiased and able to account properly for selection and drift (Sorensen & Kennedy, 1984). We suggest that the mean value  $\mu$  of genetic effects of genotyped individuals  $\mathbf{u}_2$  can be expressed as  $\mu = \frac{1}{n}\mathbf{1}'\mathbf{u}_2$ , where  $n$  is the number of individuals. Since  $\mu$  is a function of random variables, it is a random variable in itself. The variable  $\mu$  can be estimated from either pedigree ( $\mu_p$ ) or single-step ( $\mu_s$ ) procedures. If genetic prediction based on pedigree (and phenotypic data) is unbiased and accounts for selection and drift, the distribution of  $\mu_p$  accounts properly for bias, selection and drift as well. The prior distributions for genetic values of  $\mathbf{u}_2$  are  $\mathbf{u}_{2p} \sim N(\mathbf{0}, \mathbf{A}_{22}\sigma_u^2)$  and  $\mathbf{u}_{2s} \sim N(\mathbf{0}, (\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha)\sigma_u^2)$ ; thus, the distribution of  $\mu$  is  $\mu_p \sim N(\mathbf{0}, \frac{1}{n^2}\mathbf{1}'\mathbf{A}_{22}\mathbf{1}\sigma_u^2)$  and

$$\begin{aligned} \mu_s &\sim N\left(\mathbf{0}, \frac{1}{n^2}\mathbf{1}'(\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha)\mathbf{1}\sigma_u^2\right) \\ &= N\left(\mathbf{0}, \frac{1}{n^2}\mathbf{1}'\left(\mathbf{G} + \begin{bmatrix} \alpha & \cdots & \alpha \\ \vdots & & \vdots \\ \alpha & \cdots & \alpha \end{bmatrix}\right)\mathbf{1}\sigma_u^2\right). \end{aligned}$$

Since the  $\mathbf{1}\mathbf{1}'$  are simply summations,

$$\text{var}(\mu_p) = \sigma_u^2 \frac{1}{n^2} \sum_i \sum_j \mathbf{A}_{22(i,j)} \tag{3}$$

and

$$\text{var}(\mu_s) = \sigma_u^2 \left( \alpha + \frac{1}{n^2} \sum_i \sum_j \mathbf{G}_{i,j} \right). \tag{4}$$

In order to construct a model with features similar to pedigree-based BLUP, we equate the two variances in eqns (3) and (4) and this gives

$$\alpha = \frac{1}{n^2} \left( \sum_i \sum_j \mathbf{A}_{22(i,j)} - \sum_i \sum_j \mathbf{G}_{i,j} \right). \tag{5}$$

Thus,  $\alpha$  is simply the difference between means for  $\mathbf{A}_{22}$  and  $\mathbf{G}$ . The  $\alpha$  accounts for the fact that genotyped animals in  $\mathbf{u}_2$  are more related through pedigree

(in reference to the base population), which is correctly considered in  $\mathbf{A}_{22}$ . The genomic relationship matrix  $\mathbf{G}$  does not correctly reflect this fact, especially if current allele frequencies are used, which sets the genomic base as the genotyped individuals (Oliehoek *et al.*, 2006; Van Raden, 2008). For  $\mathbf{G}$  to be correct, base allele frequencies would be required. In practice, current allele frequencies are used because base allele frequencies are difficult to estimate. For a population of unrelated individuals where  $\mathbf{A}_{22} = \mathbf{I}$ ,  $\alpha = 0$ .

From Wright's  $F_{ST}$ , another interpretation of  $\alpha$  is also possible. The  $F_{ST}$  can be defined as the mean relationship between gametes in a recent population with respect to an older base population (Cockerham, 1969, 1973; Powell *et al.*, 2010). Then  $\mathbf{A}_{22}$  involves relationships of genotyped individuals with reference to the base population, and  $\mathbf{G}$  corresponds to relationships within the current population. Consequently,  $\alpha$  is equal to twice  $F_{ST}$ ; the factor of two is needed because  $F_{ST}$  is referred to as co-ancestries, which are half individual additive relationships. The correction suggested by Powell *et al.* (2010) is  $F_{Qid} = F_{new} + (1 - F_{new}) F_{ST}$ , which is equivalent to  $\mathbf{G}^S = (1 - \frac{1}{2}\alpha)\mathbf{G} + \mathbf{1}\mathbf{1}'\alpha$  and similar to our suggestion.

(ii) Simulations

To evaluate the effectiveness of the proposed approach for genomic prediction, two selection scenarios with different heritabilities were simulated. The simulator QMSim (Sargolzaei & Schenkel, 2009) was used to generate historical (undergoing drift and mutation) and recent (undergoing selection) population structures. In total, 10 chromosomes of equal length (100 cM) were simulated. Biallelic markers (10 000) were distributed at random along the chromosomes with equal frequency in the first generation of the historical population. Potentially, 250 QTLs affect the phenotype; QTLs allele effects were sampled from a Gamma distribution with a shape parameter of 0.4. The mutation rate of the markers (recurrent mutation process) and QTL was assumed to be  $2.5 \times 10^{-5}$  per locus per generation (Solberg *et al.*, 2008).

First, a base population of 200 males and 2600 females was generated by mutation and drift over 100 generations ( $t$ ) in a historical population with an effective population size of 100 ( $t = 1-95$ ) and gradually expanded to 3000 offspring ( $t = 100$ ). Then, 10 generations ( $t = 101-110$ ) of selection for a sex-limited trait with a phenotypic variance of 1 were simulated. Three heritabilities (0.05, 0.30 and 0.50) were used to examine the effect of heritability on genomic predictions. In each generation, 200 males were mated to 2600 females to produce 2600 offspring following random phenotype ( $P_Y$ ) or positive assortative (matings among best males and females based on estimated breeding values (EBVs);  $P_{EBV}$ ) designs. For the

Table 1. Mean  $\alpha$  for different heritabilities under  $P_Y$  or  $P_{EBV}$  selection

Selection design	Heritability		
	0.05	0.30	0.50
$P_Y$	0.0045	0.0046	0.0047
$P_{EBV}$	0.0834	0.0703	0.0615

$\alpha$  is the difference between pedigree-based and genome-based relationships for genotyped animals; selection was based either on random phenotype ( $P_Y$ ) or assortative mating using EBV ( $P_{EBV}$ ).

next generation, 80 males and 520 females were selected based on  $P_Y$  or  $P_{EBV}$ . For  $P_{EBV}$ , EBVs were computed in each generation with data accumulated so far by a BLUP procedure that includes phenotypes and pedigree, which mimics recent selection procedures for livestock populations. At the end of each simulation, true genetic values (TBVs) and pedigree information were available for all ten generations (28 800 individuals in pedigree), phenotypes were available for all generations except for the last one (13 100 records). All males (840 sires of females with records) were genotyped as well as 260 individuals in generation 110, which were potential candidates for selection. No fixed effects were simulated. For each scenario, 20 replicates were run.

(iii) Prediction methods

The 260 genotyped selection candidates in the last generation were evaluated using genomic information. Genetic merit of the selection candidates was estimated using five methods based on BLUP. The first method was a mixed model based on the pedigree relationship matrix and phenotypes ( $BLUP_{PED}$ ). The second method was a two-step procedure, in which DYD were computed using a regular method based on pedigree and phenotypes and then used for genomic prediction ( $BLUP_{DYD}$ ). For  $BLUP_{DYD}$ , 840 males were included with DYD information from 14 ( $\pm 10$ ) daughters on average. The DYD were weighted by their accuracy. For the third method, the full dataset (pedigree, phenotypes and genotypes) was directly used in a single-step procedure that used  $G$  ( $BLUP_{1STEP}$ ). The fourth method ( $BLUP_\alpha$ ) is also a single-step procedure with the correction of genetic differences among genotyped and non-genotyped individuals simply by using  $H^{\dagger-1}$ , which uses  $G^\dagger = G + 11'\alpha$  instead of  $G$ . The correction of  $G$  proposed by Powell *et al.* (2010) was also implemented using  $G^S = (1 - \frac{1}{2}\alpha)G + 11'\alpha$  and tested in a single-step procedure ( $BLUP_{FST}$ ).

All single-step methods ( $BLUP_{1STEP}$ ,  $BLUP_\alpha$  and  $BLUP_{FST}$ ) and  $BLUP_{DYD}$  used  $G = ZZ'/2\sum p_i(1-p_i)$ ,




Fig. 1. Bias (dotted curve with solid circles), coefficient of regression of true on EBV ( $b$ ; solid curve with squares), and squared correlation between true and EBVs ( $R^2$ ; dashed curve with circles) as functions of the correction factor  $\alpha$  (difference between pedigree-based and genome-based relationships for genotyped animals) for a given replicate with heritability of 0.30 under assortative mating selection based on EBV.

where  $z_i$  was coded as  $-p_i$  or  $1-p_i$  for the first or second allele, respectively, and  $p_i$  is the allele frequency of the second allele (VanRaden, 2008) computed as raw mean from all available genotypes.

Prediction quality was checked for all 260 selection candidates (validation data). Bias was measured as the difference between predicted and simulated breeding values of the candidates. Regression of TBV on EBV was used as a measure of the inflation of the prediction method, where a regression coefficient of one denotes no inflation. Accuracy of evaluation methods was computed as the square of the correlation between TBV and EBV. In addition, prediction error variance (PEV), which is a measure of candidate prediction error, and mean-squared error (MSE), which is a measure of overall fit of the model to the data, were computed. Results were the mean of the 20 replicates of each scenario.

3. Results

(i) Correction factor

Table 1 shows mean  $\alpha$  for the three heritabilities under  $P_Y$  or  $P_{EBV}$ . The variance  $\alpha$  is higher for selection predominantly on relatives (e.g. low heritability traits under  $P_{EBV}$ ). Bias, inflation and accuracy of predictions as a function of  $\alpha$  for a given replicate are presented in Fig. 1. In the example, the theoretical value of  $\alpha$  according to eqn (5) was 0.07, which was  $\alpha$  where bias was close to zero but not the smallest. Optimally,

Table 2. Means (SDs) of breeding values from different heritabilities and prediction methods for selection candidates under  $P_Y$  and  $P_{EBV}$

Heritability	Prediction method	$P_Y$	$P_{EBV}$
0.05	True value	0.09 (0.02)	0.49 (0.08)
	BLUP <sub>PED</sub>	0.09 (0.01)	0.52 (0.07)
	BLUP <sub>DYD</sub>	0.02 (0.01)	0.16 (0.03)
	BLUP <sub>1STEP</sub>	0.05 (0.01)	0.24 (0.05)
	BLUP <sub>α</sub>	0.08 (0.01)	0.52 (0.08)
	BLUP <sub>FST</sub>	0.08 (0.01)	0.52 (0.07)
0.30	True value	0.53 (0.03)	2.01 (0.15)
	BLUP <sub>PED</sub>	0.54 (0.03)	2.05 (0.14)
	BLUP <sub>DYD</sub>	0.17 (0.02)	0.61 (0.05)
	BLUP <sub>1STEP</sub>	0.29 (0.02)	1.41 (0.17)
	BLUP <sub>α</sub>	0.52 (0.04)	2.10 (0.15)
	BLUP <sub>FST</sub>	0.52 (0.04)	2.10 (0.15)
0.50	True value	0.90 (0.07)	2.92 (0.23)
	BLUP <sub>PED</sub>	0.90 (0.06)	2.96 (0.21)
	BLUP <sub>DYD</sub>	0.31 (0.03)	0.88 (0.89)
	BLUP <sub>1STEP</sub>	0.49 (0.03)	2.16 (0.29)
	BLUP <sub>α</sub>	0.90 (0.07)	3.00 (0.23)
	BLUP <sub>FST</sub>	0.89 (0.07)	3.01 (0.24)

Prediction methods were based on BLUP: a mixed model based on the pedigree relationship matrix and phenotypes (BLUP<sub>PED</sub>), a two-step procedure in which DYD were computed from a regular method based on pedigree and phenotypes and then used for genomic prediction (BLUP<sub>DYD</sub>), a single-step procedure that used the genomic relationship matrix and the full dataset of pedigree, phenotypes and genotypes (BLUP<sub>1STEP</sub>), a single-step procedure with genetic differences among genotyped and non-genotyped individuals corrected by considering the difference between pedigree-based and genome-based relationships for genotyped animals (BLUP<sub>α</sub>) and a single-step procedure that corrected the genomic relationship matrix as proposed by Powell *et al.* (2010) (BLUP<sub>FST</sub>). Selection was based either on random phenotype ( $P_Y$ ) or assortative mating using EBV ( $P_{EBV}$ ).

high accuracy ( $R^2$ ) and low inflation ( $b$ ) should be balanced to avoid large bias increases.

(ii) Bias

Table 2 shows TBV and EBV means from the five prediction methods under  $P_Y$  or  $P_{EBV}$  for the selection candidates in the last generation. Mean TBV of the last generation was 3.2–5.4 times larger for  $P_{EBV}$  than for  $P_Y$ . As expected, even with selection and different heritabilities, BLUP<sub>PED</sub> predicted mean TBV correctly. The BLUP<sub>DYD</sub> and BLUP<sub>1STEP</sub> methods underestimated mean TBV in the last generation, but BLUP<sub>α</sub> and BLUP<sub>FST</sub> were unbiased.

(iii) Inflation

The degree of inflation from the prediction methods is indicated by the coefficient of regression of TBV

Table 3. Coefficients (SDs) for regression of true on EBV for different heritabilities and prediction methods under  $P_Y$  and  $P_{EBV}$

Heritability	Prediction method	$P_Y$	$P_{EBV}$
0.05	BLUP <sub>PED</sub>	0.94 (0.25)	0.86 (0.22)
	BLUP <sub>DYD</sub>	1.02 (0.13)	0.84 (0.14)
	BLUP <sub>1STEP</sub>	1.01 (0.11)	0.66 (0.24)
	BLUP <sub>α</sub>	1.02 (0.11)	0.66 (0.27)
	BLUP <sub>FST</sub>	1.02 (0.11)	0.69 (0.24)
	0.30	BLUP <sub>PED</sub>	1.00 (0.01)
BLUP <sub>DYD</sub>		0.97 (0.07)	0.89 (0.08)
BLUP <sub>1STEP</sub>		0.98 (0.07)	0.86 (0.11)
BLUP <sub>α</sub>		0.97 (0.07)	0.87 (0.09)
BLUP <sub>FST</sub>		0.97 (0.07)	0.88 (0.08)
0.50		BLUP <sub>PED</sub>	0.96 (0.07)
	BLUP <sub>DYD</sub>	0.96 (0.07)	0.89 (0.05)
	BLUP <sub>1STEP</sub>	0.99 (0.05)	0.99 (0.05)
	BLUP <sub>α</sub>	0.97 (0.05)	0.90 (0.05)
	BLUP <sub>FST</sub>	0.97 (0.05)	0.90 (0.05)

Prediction methods were based on BLUP: a mixed model based on the pedigree relationship matrix and phenotypes (BLUP<sub>PED</sub>), a two-step procedure in which DYD were computed from a regular method based on pedigree and phenotypes and then used for genomic prediction (BLUP<sub>DYD</sub>), a single-step procedure that used the genomic relationship matrix and the full dataset of pedigree, phenotypes and genotypes (BLUP<sub>1STEP</sub>), a single-step procedure with genetic differences among genotyped and non-genotyped individuals corrected by considering the difference between pedigree-based and genome-based relationships for genotyped animals (BLUP<sub>α</sub>), and a single-step procedure that corrected the genomic relationship matrix as proposed by Powell *et al.* (2010) (BLUP<sub>FST</sub>). Selection was based either on random phenotype ( $P_Y$ ) or assortative mating using EBV ( $P_{EBV}$ ).

on EBV (Table 3). The optimal prediction method of genetic merit of young individuals would have a regression coefficient close to 1. For each scenario, the differences between the methods were small, and the approaches achieved very similar inflation. A strong selection, under  $P_{EBV}$ , increased inflation with values lower than 1. According to Kennedy *et al.* (1988) relationship matrix accounts for selection, drift and non-random mating, but it does not account for a wrong definition of the base population or a finite number of loci. Potentially 250 QTLs affected the phenotype, and so this discards the second reason. Ideally, the base population should be infinite; this is indeed not the case, and thus the first assumption is violated. Consider animals 1 and 2 (e.g. father and son). If they are related, the distribution of the genetic value of 2 can be expressed as conditioned on the genetic value of 1. Thus, knowledge of the EBV of 1 would decrease uncertainty (variance) of the EBV of 2. However, if this relationship is unknown, the conditional distribution cannot be written. Thus,

Table 4. Squared correlations between true and EBVs (SDs) for different heritabilities and prediction methods under  $P_Y$  and  $P_{EBV}$

Heritability	Prediction method	$P_Y$	$P_{EBV}$
0.05	BLUP <sub>PED</sub>	7 (4)	10 (4)
	BLUP <sub>DYD</sub>	23 (5)	28 (7)
	BLUP <sub>ISTEP</sub>	29 (5)	25 (7)
	BLUP <sub><math>\alpha</math></sub>	29 (5)	27 (8)
	BLUP <sub><math>F_{ST}</math></sub>	29 (5)	27 (7)
0.30	BLUP <sub>PED</sub>	20 (4)	23 (6)
	BLUP <sub>DYD</sub>	49 (5)	56 (6)
	BLUP <sub>ISTEP</sub>	54 (4)	47 (6)
	BLUP <sub><math>\alpha</math></sub>	55 (5)	60 (5)
	BLUP <sub><math>F_{ST}</math></sub>	55 (5)	60 (5)
0.50	BLUP <sub>PED</sub>	21 (4)	30 (6)
	BLUP <sub>DYD</sub>	56 (5)	64 (6)
	BLUP <sub>ISTEP</sub>	61 (5)	49 (7)
	BLUP <sub><math>\alpha</math></sub>	61 (5)	67 (5)
	BLUP <sub><math>F_{ST}</math></sub>	61 (5)	67 (5)

Squared correlations between true and EBVs were expressed as percentages. Prediction methods were based on BLUP: a mixed model based on the pedigree relationship matrix and phenotypes (BLUP<sub>PED</sub>), a two-step procedure in which DYD were computed from a regular method based on pedigree and phenotypes and then used for genomic prediction (BLUP<sub>DYD</sub>), a single-step procedure that used the genomic relationship matrix and the full dataset of pedigree, phenotypes and genotypes (BLUP<sub>ISTEP</sub>), a single-step procedure with genetic differences among genotyped and non-genotyped individuals corrected by considering the difference between pedigree-based and genome-based relationships for genotyped animals (BLUP <sub>$\alpha$</sub> ) and a single-step procedure that corrected the genomic relationship matrix as proposed by Powell *et al.* (2010) (BLUP <sub>$F_{ST}$</sub> ). Selection was based either on random phenotype ( $P_Y$ ) or assortative mating using EBV ( $P_{EBV}$ ).

not knowing relationships will increase the variance of EBVs and cause inflation; this has not been verified.

#### (iv) Accuracy

Squared correlations between TBV and EBV (i.e. reliability) by heritability and prediction method are shown in Table 4. Squared correlations are presented in Fig. 2 for all replicates using a heritability of 0.30 under  $P_{EBV}$ . Compared with BLUP<sub>PED</sub>, all genomic prediction methods increased accuracy by about 37 percentage units and 22 percentage units under assortative mating based on EBV and mass selection, respectively.

With low heritability under  $P_{EBV}$ , accuracy from BLUP <sub>$\alpha$</sub>  and BLUP <sub>$F_{ST}$</sub>  was as good as from BLUP<sub>DYD</sub> (Table 4). When these accuracies are comparable, the single-step procedures with correction (BLUP <sub>$\alpha$</sub>  and BLUP <sub>$F_{ST}$</sub> ) have an advantage over BLUP<sub>DYD</sub> because they provide a unified framework that eliminates all




Fig. 2. Accuracy (squared correlations between true and EBV) of BLUP methods for estimating breeding value of selection candidates across 20 replicates with heritability of 0.30 under assortative mating selection based on EBV. Prediction methods were a mixed model based on the pedigree relationship matrix and phenotypes (BLUP<sub>PED</sub>; solid line with triangles), a two-step procedure in which DYD were computed from a regular method based on pedigree and phenotypes and then used for genomic prediction (BLUP<sub>DYD</sub>; dashed line with solid circles), a single-step procedure with genetic differences among genotyped and non-genotyped individuals corrected by considering the difference between pedigree-based and genome-based relationships for genotyped animals (BLUP <sub>$\alpha$</sub> ; solid line with squares) and a single-step procedure that corrected the genomic relationship matrix as proposed by Powell *et al.* (2010) (BLUP <sub>$F_{ST}$</sub> ) (dashed line with x markers).

the assumptions applied in multiple-step methods. With medium or high heritabilities under  $P_{EBV}$ , BLUP <sub>$\alpha$</sub>  and BLUP <sub>$F_{ST}$</sub>  have the highest accuracies (Table 4; Fig. 2). The lower accuracy of BLUP<sub>DYD</sub> compared with BLUP <sub>$\alpha$</sub>  and BLUP <sub>$F_{ST}$</sub>  results from ignoring parent average in BLUP<sub>DYD</sub>. Including parent average in predictions increases the accuracy in multiple-step methods, but it is complicated (VanRaden *et al.*, 2009a). Parent averages are automatically included in genomic predictions with single-step methods.

#### (v) Other comparison measures

In addition to assessing the quality of genetic evaluations through bias and accuracy from the difference and correlation between TBV and EBV, PEV and MSE were also considered (Table 5). All genomic methods had the lowest PEV under  $P_Y$  for all heritabilities. Under  $P_{EBV}$ , PEV also were lowest for genomic methods except with low heritability. Compared with MSE for BLUP<sub>DYD</sub> predictions, MSE for BLUP <sub>$\alpha$</sub>  were 18–66% lower under  $P_Y$  and 60–96%

Table 5. PEVs (SDs) and MES (SDs) for different heritabilities and prediction methods under  $P_Y$  and  $P_{EBV}$ 

Heritability	Prediction method	$P_Y$		$P_{EBV}$	
		PEV	MSE	PEV	MSE
0.05	BLUP <sub>PED</sub>	0.047 (0.002)	0.047 (0.002)	0.045 (0.003)	0.046 (0.003)
	BLUP <sub>DYD</sub>	0.038 (0.002)	0.044 (0.004)	0.037 (0.004)	0.156 (0.040)
	BLUP <sub>1STEP</sub>	0.036 (0.003)	0.037 (0.003)	0.049 (0.019)	0.106 (0.027)
	BLUP <sub><math>\alpha</math></sub>	0.035 (0.002)	0.036 (0.002)	0.059 (0.057)	0.062 (0.057)
	BLUP <sub>F<sub>ST</sub></sub>	0.036 (0.003)	0.036 (0.002)	0.047 (0.021)	0.049 (0.021)
0.30	BLUP <sub>PED</sub>	0.241 (0.011)	0.242 (0.012)	0.230 (0.021)	0.234 (0.021)
	BLUP <sub>DYD</sub>	0.153 (0.017)	0.289 (0.028)	0.134 (0.019)	2.088 (0.304)
	BLUP <sub>1STEP</sub>	0.137 (0.014)	0.198 (0.020)	0.166 (0.025)	0.531 (0.106)
	BLUP <sub><math>\alpha</math></sub>	0.137 (0.015)	0.138 (0.014)	0.127 (0.020)	0.138 (0.024)
	BLUP <sub>F<sub>ST</sub></sub>	0.137 (0.015)	0.138 (0.014)	0.126 (0.019)	0.136 (0.023)
0.50	BLUP <sub>PED</sub>	0.393 (0.020)	0.395 (0.019)	0.349 (0.031)	0.352 (0.032)
	BLUP <sub>DYD</sub>	0.222 (0.028)	0.582 (0.069)	0.185 (0.032)	4.384 (0.615)
	BLUP <sub>1STEP</sub>	0.198 (0.023)	0.369 (0.051)	0.257 (0.035)	0.854 (0.200)
	BLUP <sub><math>\alpha</math></sub>	0.195 (0.024)	0.196 (0.024)	0.170 (0.028)	0.179 (0.028)
	BLUP <sub>F<sub>ST</sub></sub>	0.195 (0.024)	0.196 (0.024)	0.169 (0.028)	0.177 (0.027)

PEV and MSE are shown. Prediction methods were based on BLUP: a mixed model based on the pedigree relationship matrix and phenotypes (BLUP<sub>PED</sub>), a two-step procedure in which DYD were computed from a regular method based on pedigree and phenotypes and then used for genomic prediction (BLUP<sub>DYD</sub>), a single-step procedure that used the genomic relationship matrix and the full dataset of pedigree, phenotypes and genotypes (BLUP<sub>1STEP</sub>), a single-step procedure with genetic differences among genotyped and non-genotyped individuals corrected by considering the difference between pedigree-based and genome-based relationships for genotyped animals (BLUP <sub>$\alpha$</sub> ), and a single-step procedure that corrected the genomic relationship matrix as proposed by Powell *et al.* (2010) (BLUP<sub>F<sub>ST</sub></sub>). Selection was based either on random phenotype ( $P_Y$ ) or assortative mating using EBV ( $P_{EBV}$ ).

lower under  $P_{EBV}$ , with differences increasing with heritability under both selection designs. The greatly decreased MSE for all scenarios indicated that the multiple-step BLUP<sub>DYD</sub> was less accurate. Considering the potential genetic gain from selection, BLUP <sub>$\alpha$</sub>  and BLUP<sub>F<sub>ST</sub></sub> were the most advantageous prediction methods.

#### 4. Discussion

Comparison of the four genomic prediction methods at various heritabilities under two selection intensities demonstrated that a single-step method with correction (either BLUP <sub>$\alpha$</sub>  or BLUP<sub>F<sub>ST</sub></sub>) was a preferred method of accounting for bias in genomic predictions. Furthermore, two possible  $G$  for prediction of genetic merit were tested. Genomic preselection of animals often is included in dairy cattle breeding programmes now. Thus, bias becomes an important concern with the multiple-step method, because computation of DYD assumes random selection of the Mendelian sampling, which is clearly violated (Patry & Ducrocq, 2011). Because single-step methods do not create pseudodata and are unbiased, they may be an attractive genomic evaluation method.

Bias differences among the methods may be explained by how the different methods account for genotypes of highly selected individuals (i.e. males). Classical theory for modelling covariance among

individuals assumes no selection. Thus, covariances among TBVs of selected individuals are no longer described well by any (genomic or pedigree-based) relationship matrix, unless all records used in the selection are accounted for (as in BLUP<sub>PED</sub>) (Sorensen & Kennedy, 1984; Kennedy *et al.*, 1988). Although all records are used in BLUP<sub>1STEP</sub>, only genotyped (and mostly selected) individuals are included in  $G$ . Since genetic values of non-genotyped animals are, a priori, conditioned on genetic values of genotyped animals (Legarra *et al.*, 2009), bias is alleviated but not fully corrected. In a single-step method with correction, such as BLUP <sub>$\alpha$</sub> , bias is eliminated by referencing genomic and pedigree-based relationship matrices to the base population. With correction, BLUP <sub>$\alpha$</sub>  (or BLUP<sub>F<sub>ST</sub></sub>) is the most accurate for GWMAS.

A pertinent question is which evaluation criterion should be used to maximize genetic gain in selection schemes. According to classical selection theory, best prediction is ideal for selection (Henderson, 1973; Fernando & Gianola, 1986). The best predictor minimizes MSE, and is unbiased and not inflated ( $b = 1$ ) by construction (Henderson, 1973). If we were using the best predictor, accuracy, PEV or MSE should provide the same ranking of methods. The best predictor requires knowing the true model generating the data. This is not the case here because true QTLs were not included in the model for genetic evaluation.

Instead, we used a linear model assuming multivariate normality.

In practice, the relevance of the criterion used depends on the selection schemes. If the parents of the next generation come from (and only from) genotyped selection candidates, then accuracy is the criterion to maximize, because selection candidates share a common mean (i.e. they belonged to the same generation) and thus bias is not a concern.

If for different candidates there are different amounts of information (e.g. comparing progeny-tested males vs. newborn animals), PEV or MSE have to be considered. For example, in the presence of bias, genetic gain is over (or under) estimated. Thus, newborns are thought to be better than what they are. In this case, MSE is possibly the criterion of choice, because it also includes bias. For instance, Roehe & Kennedy (1993), who showed that a wrong model resulted in an artificial overestimation of genetic trend, which raised the estimated merit of young selection candidates. Similarly, inflation ( $b < 1$ , which is included in PEV or MSE but not in accuracy) results in exaggeration (both over and under) of estimated genetic merit of newborns with respect to progeny-tested animals.

However, accuracy is currently used to assess genomic prediction methods in cross-validation studies (e.g. VanRaden *et al.*, 2009a), although bias is becoming an increasing concern (e.g. Luan *et al.*, 2009; VanRaden *et al.*, 2009b; Mäntysaari *et al.*, 2010). Consideration of bias, accuracy and inflation, possibly through MSE, is strongly recommended for the comparison of future genomic selection strategies.

### 5. Conclusion

Overall, a single-step genomic prediction method with corrected  $\mathbf{G}$  ( $\text{BLUP}_\alpha$  or  $\text{BLUP}_{F_{ST}}$ ) was unbiased, similarly inflated and more accurate than other

prediction approach is promising for animal/plant breeding.

We are grateful to Paul VanRaden, Daniel Gianola, Daniel Sorensen, Shogo Tsuruta and Ching-Yi Chen for their helpful and constructive comments. This work was partly financed by Holstein Association USA, Smithfield Premium Genetics, the National Beef Cattle Evaluation Consortium and AFRI grants 2009-65205-05665 and 2010-65205-20366 from the USDA NIFA Animal Genome Program (ZV, IA, IM) as well as Apisgene and ANR projects AMASGEN and SheepSNPQTL and POCTEFA project Genomia (AL). The project also was partly supported by the Genopole Toulouse Midi-Pyrénées bioinformatics platform.

### Appendix

#### Genetic group model for unbiased genomic predictions

From the model of unbiased genomic predictions presented in section (i), we have that

$$p(\mathbf{u}_1|\mu) \sim N(\mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{1}\mu, (\mathbf{A}^{11})^{-1}\sigma_u^2 + \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{G}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\sigma_u^2),$$

$$p(\mathbf{u}_2|\mu) \sim N(\mathbf{1}\mu, \mathbf{G}\sigma_u^2),$$

and  $p(\mathbf{u}|\mu) \sim N(\mathbf{Q}\mu, \mathbf{H}\sigma_u^2)$ , where  $\mathbf{H}$  is according to eqn (1) and  $\mathbf{Q}$  is  $\mathbf{Q} = \begin{bmatrix} \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{1} \\ \mathbf{1} \end{bmatrix}$ .

The setting is like a genetic group model (Quaas, 1988) and equivalent mixed-model equations, which yields the same solutions for  $\mathbf{b}$  and  $\mathbf{u}$ , is derived as in Quaas (1988) as

$$\begin{bmatrix} \mathbf{X}'\mathbf{X} & \mathbf{X}'\mathbf{Z} & \mathbf{0} \\ \mathbf{Z}'\mathbf{X} & \mathbf{Z}'\mathbf{Z} + \mathbf{H}^{-1}\lambda & -\mathbf{H}^{-1}\mathbf{Q}\lambda \\ \mathbf{0} & -\mathbf{Q}'\mathbf{H}^{-1}\lambda & \mathbf{Q}'\mathbf{H}^{-1}\mathbf{Q}\lambda + \alpha^{-1}\lambda \end{bmatrix} \times \begin{bmatrix} \mathbf{b} \\ \mathbf{u} \\ \mu \end{bmatrix} = \begin{bmatrix} \mathbf{X}'\mathbf{y} \\ \mathbf{Z}'\mathbf{y} \\ 0 \end{bmatrix}.$$

The above expression is simplified by computing the product  $\mathbf{Q}'\mathbf{H}^{-1}$  as

$$\begin{aligned} -\mathbf{Q}'\mathbf{H}^{-1} &= -\begin{bmatrix} \mathbf{1}'\mathbf{A}_{22}^{-1}\mathbf{A}_{21} & \mathbf{1} \end{bmatrix} \begin{bmatrix} \mathbf{A}^{11} & \mathbf{A}^{12} \\ \mathbf{A}^{21} & \mathbf{A}^{22} + \mathbf{G}^{-1} - \mathbf{A}_{22}^{-1} \end{bmatrix} \\ &= \begin{bmatrix} -\mathbf{1}'\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{11} - \mathbf{1}\mathbf{A}^{21} & -\mathbf{1}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{12} - \mathbf{1}(\mathbf{A}^{22} + \mathbf{G}^{-1} - \mathbf{A}_{22}^{-1}) \end{bmatrix} \\ &= \begin{bmatrix} -\mathbf{1}'\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{11} - \mathbf{1}\mathbf{A}^{21} & -\mathbf{1}(\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{12} + \mathbf{A}^{22} - \mathbf{A}_{22}^{-1}) - \mathbf{1}\mathbf{G}^{-1} \end{bmatrix}. \end{aligned}$$

procedures even in the presence of selection. The corrected  $\mathbf{G}$  is an appropriate methodological solution that takes into account the effect of non-random genotyping (due to strong selection) on prediction. The results clearly showed that a single-step genomic

Note that  $\mathbf{A}^{21} = -\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{11}$  and  $\mathbf{A}^{22} = \mathbf{A}_{22}^{-1} - \mathbf{A}_{21}\mathbf{A}_{22}\mathbf{A}_{22}^{-1} = \mathbf{A}_{22}^{-1} - \mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}^{12}$ . We can see that  $-\mathbf{Q}'\mathbf{H}^{-1} = -\begin{bmatrix} \mathbf{0} & \mathbf{1}'\mathbf{G}^{-1} \end{bmatrix}$  and the product  $\mathbf{Q}'\mathbf{H}^{-1}\mathbf{Q} = \mathbf{1}'\mathbf{G}^{-1}\mathbf{1}$ , which is simply the sum of its elements.



Thus, the mixed-model equations can be expressed as

$$\begin{bmatrix} \mathbf{X}'\mathbf{X} & \mathbf{X}'\mathbf{Z} \\ \mathbf{Z}'\mathbf{X} & \mathbf{Z}'\mathbf{Z} + \mathbf{H}^{*-1}\lambda \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ \boldsymbol{\theta} \end{bmatrix} = \begin{bmatrix} \mathbf{X}'\mathbf{y} \\ \mathbf{Z}'\mathbf{y} \end{bmatrix}, \quad (6)$$

with  $\mathbf{Z}'\mathbf{Z}$  appropriately expanded with zeros,  $\boldsymbol{\theta} = \begin{bmatrix} \mathbf{u} \\ \boldsymbol{\mu} \end{bmatrix}$  and setting  $\mathbf{H}^{*-1}$  as

$$\mathbf{H}^{*-1} = \begin{bmatrix} \mathbf{H}^{11} & \mathbf{H}^{12} & \mathbf{0} \\ \mathbf{H}^{21} & \mathbf{H}^{22} & -\mathbf{G}^{-1}\mathbf{1} \\ \mathbf{0} & -\mathbf{1}'\mathbf{G}^{-1} & \mathbf{1}'\mathbf{G}^{-1}\mathbf{1} + \alpha^{-1} \end{bmatrix} \\ = \begin{bmatrix} \mathbf{A}^{11} & \mathbf{A}^{12} & \mathbf{0} \\ \mathbf{A}^{21} & \mathbf{A}^{22} + \mathbf{G}^{-1} - \mathbf{A}_{22}^{-1} & -\mathbf{G}^{-1}\mathbf{1} \\ \mathbf{0} & -\mathbf{1}'\mathbf{G}^{-1} & \mathbf{1}'\mathbf{G}^{-1}\mathbf{1} + \alpha^{-1} \end{bmatrix}.$$

Absorption of  $\boldsymbol{\mu}$  in (6) gives mixed-model equations with  $\mathbf{H}^{\dagger-1}$  (eqn (2)) using expression (2) in Henderson & Searle (1981). Computing  $\mathbf{H}^{*-1}$  is similar to the creation of  $\mathbf{A}^{-1}$  with genetic groups as reported by Quaas (1988).

The expression (6) is of interest because there is an explicit estimate of  $\boldsymbol{\mu}$ . In addition, mixed-model equations in (6) have a straightforward interpretation. The genetic value of a genotyped individual is conditional on its mean  $\boldsymbol{\mu}$  and relatives through genomic relationships. The genetic value of a non-genotyped individual is conditional on its relatives through pedigree relationships, including relatives with genotypes.

## References

- Aguilar, I., Misztal, I., Johnson, D. L., Legarra, A., Tsuruta, S. & Lawlor, T. J. (2010). A unified approach to utilize phenotypic, full pedigree, and genomic information for genetic evaluation of Holstein final score. *Journal of Dairy Science* **93**, 743–752.
- Chen, C. Y., Misztal, I., Aguilar, I., Legarra, A. & Muir, W. M. (2011). Effect of different genomic relationship matrices on accuracy and scale. *Journal of Animal Science*, in press.
- Christensen, O. F. & Lund, M. S. (2010). Genomic prediction when some animals are not genotyped. *Genetics Selection Evolution* **42**, 2.
- Cockerham, C. C. (1969). Variance of gene frequencies. *Evolution* **23**, 72–84.
- Cockerham, C. C. (1973). Analysis of gene frequencies. *Genetics* **74**, 679–700.
- Daetwyler, H. D., Pong-Wong, R., Villanueva, B. & Woolliams, J. A. (2010). The impact of genetic architecture on genome-wide evaluation methods. *Genetics* **185**, 1021–1031.
- Fernando, R. & Gianola, D. (1986). Optimal properties of the conditional mean as a selection criterion. *Theoretical and Applied Genetics* **72**, 822–825.
- Garrick, D. J., Taylor, J. F. & Fernando, R. L. (2009). Deregressing estimated breeding values and weighting information for genomic regression analyses. *Genetics Selection Evolution* **41**, 55.
- Goddard, M. E. & Hayes, B. J. (2009). Mapping genes for complex traits in domestic animals and their use in breeding programmes. *Nature Reviews Genetics* **10**, 381–391.
- Hayes, B. J., Vissler, P. M. & Goddard, M. E. (2009). Increased accuracy of artificial selection by using the realized relationship matrix. *Genetics Research* **91**, 47–60.
- Henderson, C. R. (1973). Sire evaluation and genetic trends. In: *Proceedings of the Animal Breeding and Genetics Symposium in Honor of Dr. J. L. Lush*. pp. 10–41. Champaign, IL: American Society of Animal Science and American Dairy Science Association.
- Henderson, C. R., Kempthorne, O., Searle, S. R. & von Krosigk, C. M. (1959). The estimation of environmental and genetic trends from records subject to culling. *Biometrics* **15**, 192–218.
- Henderson, C. R. & Searle, S. R. (1981). On deriving the inverse of a sum of matrices. *SIAM Review* **23**, 53–60.
- Hill, W. G. (2010). Understanding and using quantitative genetic variation. *Philosophical Transactions of the Royal Society B* **365**, 73–85.
- Im, S., Fernando, R. L. & Gianola, D. (1989). Likelihood inferences in animal breeding under selection: a missing-data theory viewpoint. *Genetics Selection Evolution* **21**, 399–414.
- Kennedy, B. W., Schaeffer, L. R. & Sorensen, D. A. (1988). Genetic properties of animal models. *Journal of Dairy Science* **71**, 17–26.
- Legarra, A., Aguilar, I. & Misztal, I. (2009). A relationship matrix including full pedigree and genomic information. *Journal of Dairy Science* **92**, 4656–4663.
- Luan, T., Woolliams, J. A., Lien, S., Kent, M., Svendsen, M. & Meuwissen, T. H. E. (2009). The accuracy of genomic selection in Norwegian red cattle assessed by cross-validation. *Genetics* **183**, 1119–1126.
- Mäntysaari, E., Liu, Z. & VanRaden, P. (2010). Interbull validation test for genomic evaluations. *Interbull Bulletin* **41**, 5 p.
- Meuwissen, T. H. E., Hayes, B. J. & Goddard, M. E. (2001). Prediction of total genetic value using genome-wide dense marker maps. *Genetics* **157**, 1819–1829.
- Oliehoek, P. A., Winding, J. J., van Arendonk, J. A. M. & Bijma, P. (2006). Estimating relatedness between individuals in general populations with a focus on their use in conservation programs. *Genetics* **173**, 483–496.
- Patry, C. & Ducrocq, V. (2011). Evidence of biases in genetic evaluations due to genomic pre-selection in dairy cattle. *Journal of Dairy Science* **94**, 1011–1020.
- Powell, J. E., Vissler, P. M. & Goddard, M. E. (2010). Reconciling the analysis of IBD and IBS in complex trait studies. *Nature Reviews Genetics* **11**, 800–805.
- Quaas, R. L. (1988). Additive genetic models with groups and relationships. *Journal of Dairy Science* **71**, 1338–1345.
- Roehe, R. & Kennedy, B. W. (1993). The influence of maternal effects on accuracy of evaluation of litter size in swine. *Journal of Animal Science* **71**, 2353–2364.
- Sargolzaei, M. & Schenkel, F. (2009). QMSim: a large-scale genome simulator for livestock. *Bioinformatics* **25**, 680–681.
- Solberg, T. R., Sonesson, A. K., Woolliams, J. A. & Meuwissen, T. H. E. (2008). Genomic selection using different marker types and densities. *Journal of Animal Science* **86**, 2447–2454.
- Sorensen, D. A. & Kennedy, B. W. (1984). Estimation of response to selection using least squares and mixed

- model methodology. *Journal of Animal Science* **58**, 1097–1103.
- VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *Journal of Dairy Science* **91**, 4414–4423.
- VanRaden, P. M., Van Tassell, C. P., Wiggans, G. R., Sonstegard, T. S., Schnabel, R. D., Taylor, J. F. & Schenkel, F. S. (2009a). Invited review: Reliability of genomic predictions for North American Holstein bulls. *Journal of Dairy Science* **92**, 16–24.
- VanRaden, P. M., Tooker, M. E. & Cole, J. B. (2009b). Can you believe those genomic evaluations for young bulls? *Journal of Dairy Science* **92**(E-Suppl. 1), 314 (Abstr.).