

ON WEYL'S THEOREM FOR TENSOR PRODUCTS

C. S. KUBRUSLY

Catholic University of Rio de Janeiro, Rio de Janeiro, 22453-900, RJ, Brazil
e-mail: carlos@ele.puc-rio.br

and B. P. DUGGAL

Department of Mathematics, University of Niš, P.O. Box 224, Niš, Serbia
e-mail: bpduggal@yahoo.co.uk

(Received 23 August 2011; accepted 9 May 2012; first published online 2 August 2012)

Abstract. Let A and B be operators acting on infinite-dimensional spaces. In this paper we prove that if A and B are isoloid, satisfy Weyl's theorem, and the Weyl spectrum identity holds, then $A \otimes B$ satisfies Weyl's theorem.

1991 *Mathematics Subject Classification.* Primary 47A80; Secondary 47A53.

1. Notation and terminology. By an *operator* we mean a *bounded* linear transformation of a Hilbert space into itself. We work in a Hilbert space setting, although the results in this paper hold in a Banach space setting with essentially the same proofs. Let T be an operator, and let $\mathcal{N}(T)$ and $\mathcal{R}(T)$ denote kernel and range of T , respectively. Consider the classical partition $\{\sigma_P(T), \sigma_R(T), \sigma_C(T)\}$ of the spectrum $\sigma(T)$, where $\sigma_P(T) = \{\lambda \in \mathbb{C} : \mathcal{N}(\lambda I - T) \neq \{0\}\}$ is the point spectrum (i.e. the set of all eigenvalues of T), $\sigma_R(T) = \sigma_P(T^*)^* \setminus \sigma_P(T)$ is the residual spectrum (where T^* denotes the adjoint of T and $\Lambda^* = \{\bar{\lambda} \in \mathbb{C} : \lambda \in \Lambda\}$ denotes the set of all complex conjugates from a subset Λ of \mathbb{C}), and $\sigma_C(T) = \sigma(T) \setminus (\sigma_P(T) \cup \sigma_R(T))$ is the continuous spectrum. Let $\sigma_w(T) = \{\lambda \in \mathbb{C} : \lambda I - T \text{ is not a Fredholm operator of index zero}\}$ be the Weyl spectrum of T , which is a subset of the whole spectrum $\sigma(T)$; that is,

$$\sigma_w(T) = \{\lambda \in \sigma(T) : \lambda I - T \text{ is not a Fredholm operator of index zero}\}.$$

The set

$$\sigma_0(T) = \{\lambda \in \sigma_P(T) : \mathcal{R}(\lambda I - T) \text{ is closed and} \\ \dim \mathcal{N}(\lambda I - T) = \dim \mathcal{N}(\bar{\lambda} I - T^*) < \infty\}$$

is precisely a complement of the Weyl spectrum $\sigma_w(T)$ in the whole spectrum $\sigma(T)$ (see e.g. [5] or [6, Section 5.3]). Hence,

$$\sigma_w(T) = \sigma(T) \setminus \sigma_0(T),$$

and so $\{\sigma_w(T), \sigma_0(T)\}$ forms another partition of spectrum $\sigma(T)$. Set $\sigma_{PF}(T) = \{\lambda \in \sigma_P(T) : \dim \mathcal{N}(\lambda I - T) < \infty\}$; the set of all eigenvalues of T of finite multiplicity so that $\sigma_0(T) \subseteq \sigma_{PF}(T)$ and $\sigma_R(T) \cup \sigma_C(T) \cup (\sigma_P(T) \setminus \sigma_{PF}(T)) \subseteq \sigma_w(T)$. Set

$$\pi_{00}(T) = \sigma_{\text{iso}}(T) \cap \sigma_{PF}(T),$$

where $\sigma_{\text{iso}}(T)$ denotes a set of all isolated points of spectrum $\sigma(T)$. One says that an operator T satisfies *Weyl's theorem* if

$$\sigma_0(T) = \pi_{00}(T),$$

and it is said to *satisfy Browder's theorem* if

$$\sigma_0(T) \subseteq \pi_{00}(T).$$

An operator T is *isoloid* if $\sigma_{\text{iso}}(T) \subseteq \sigma_p(T)$ (i.e. if every isolated point of the spectrum is an eigenvalue).

2. The question. Consider a tensor product $A \otimes B$ of a pair of operators A and B . It is known from [1] that the spectrum of a tensor product coincides with the product of the spectra of factors

$$\sigma(A \otimes B) = \sigma(A) \cdot \sigma(B).$$

For the Weyl spectrum it was proved in [3] that the inclusion

$$\sigma_w(A \otimes B) \subseteq \sigma_w(A) \cdot \sigma(B) \cup \sigma(A) \cdot \sigma_w(B)$$

holds. However, since then it remained as an open question whether the preceding inclusion is an identity or not. That is, it was not known if there existed a pair of operators A and B for which the above inclusion was proper. This question was solved quite recently by using a counterexample from [4, Section 3] (which exhibits a pair of operators that satisfy Weyl's theorem whose tensor product does not satisfy Browder's theorem) together with Corollary 6 from [7] (which says that Browder's theorem is transferred from a pair of operators to their tensor product if and only if the above inclusion is an identity). Thus, there exists a pair of operators for which the above inclusion is proper. If a pair of operators A, B is such that

$$\sigma_w(A \otimes B) = \sigma_w(A) \cdot \sigma(B) \cup \sigma(A) \cdot \sigma_w(B),$$

then we say that the *Weyl spectrum identity holds* for A and B . It was shown in [7, Proposition 5] that if the Weyl spectrum identity holds for A and B , then

$$\sigma_0(A \otimes B) \subseteq \sigma_0(A) \cdot \sigma_0(B),$$

and it was also shown that the above inclusion may be proper even if the Weyl spectrum identity holds for A and B , with A, B and $A \otimes B$ being isoloid operators that satisfy Weyl's theorem (see [7, Remark 2]).

The problem of transferring Weyl's theorem from isoloid operators A and B to their tensor product $A \otimes B$ was considered in this journal in [8], where it was stated that *if A and B are isoloid operators that satisfy Weyl's theorem, then $A \otimes B$ satisfies Weyl's theorem*. Their proof stands only if it is further assumed that the Weyl spectrum identity holds for A and B (which is not true in general), and also that $\sigma_0(A) \cdot \sigma_0(B) \subseteq \sigma_0(A \otimes B)$ (which may fail even if the Weyl spectrum identity holds and all operators are isoloid and satisfy Weyl's theorem). However, under these extra two assumptions, the problem of transferring Weyl's theorem from isoloid operators A and B to their tensor product $A \otimes B$ survives.

PROPOSITION 1. *Let A and B be operators acting on infinite-dimensional spaces. If*

- (a) *A and B are isoloid,*
- (b) *A and B satisfy Weyl's theorem, and*
- (c) *the Weyl spectrum identity holds for A and B ,*

then the tensor product $A \otimes B$ satisfies Weyl's theorem if, in addition,

- (d) $\sigma_0(A \otimes B) = \sigma_0(A) \cdot \sigma_0(B)$.

Proof. [7, Corollary 5]. □

Question: Is condition (d) necessary in Proposition 1? In other words, is the next conjecture true?

CONJECTURE 1. *If A and B are isoloid, satisfy Weyl's theorem, and the Weyl spectrum identity holds, then $A \otimes B$ satisfies Weyl's theorem.*

We shall show in the next section (Theorem 1) that the above conjecture is, in fact, a theorem: It holds true for operators acting on infinite-dimensional spaces. (If the spaces are finite-dimensional, where every subspace is closed and every operator is isoloid with spectrum consisting only of eigenvalues of finite multiplicity, then the conjecture holds trivially.) The proof is carried out by using results originally proved in a Hilbert space setting, which clearly still holds in a Banach space setting.

3. The answer. Before proving Theorem 1 we need the following auxiliary results. Let A , B and T be arbitrary operators.

LEMMA 1. *If A and B satisfy Browder's theorem, and if the Weyl spectrum identity holds, then the tensor product $A \otimes B$ satisfies Browder's theorem.*

Proof. [7, Proposition 7(a)]. □

LEMMA 2. *If T satisfies Browder's theorem but not Weyl's theorem, then*

$$\sigma_w(T) \cap \pi_{00}(T) \neq \emptyset.$$

Proof. T satisfies Browder's theorem but not Weyl's theorem if and only if the proper inclusion $\sigma_0(T) \subset \pi_{00}(T)$ holds true, which implies that there exists an isolated eigenvalue of finite multiplicity not in $\sigma_0(T)$ (i.e. in $\sigma_w(T) = \sigma(T) \setminus \sigma_0(T)$). □

LEMMA 3. *If A and B are isoloid operators on infinite-dimensional spaces, then*

$$\pi_{00}(A \otimes B) \subseteq \pi_{00}(A) \cdot \pi_{00}(B).$$

Proof. [7, Proposition 4]. □

LEMMA 4. *If A and B are operators on infinite-dimensional spaces, then*

$$0 \notin \sigma_{PF}(A \otimes B).$$

Proof. [7, Proposition 1]. □

LEMMA 5. Suppose $\sigma_{\text{iso}}(A) \neq \emptyset$ and $\sigma_{\text{iso}}(B) \neq \emptyset$. If $\lambda \in \sigma(A)$, $\mu \in \sigma(B)$, and $\lambda\mu \in \sigma_{\text{iso}}(A \otimes B)$, then

$$\lambda \in \sigma_{\text{iso}}(A) \quad \text{and} \quad \mu \in \sigma_{\text{iso}}(B).$$

Proof. [7, Proof of Proposition 3(a)]. □

LEMMA 6. If $\lambda \in \sigma_P(A)$, $\mu \in \sigma_P(B)$, and $\lambda\mu \in \sigma_{PF}(A \otimes B)$, then

$$\lambda \in \sigma_{PF}(A) \quad \text{and} \quad \mu \in \sigma_{PF}(B).$$

Proof. [7, Proof of Proposition 2(a)]. □

Suppose A and B are operators acting on infinite-dimensional spaces.

THEOREM 1. If A and B are isoloid, satisfy Weyl's theorem, and the Weyl spectrum identity holds, then $A \otimes B$ satisfies Weyl's theorem.

Proof. Suppose A and B satisfy Weyl's theorem. Thus, A and B satisfy Browder's theorem. Therefore, if the Weyl's spectrum identity holds, then $A \otimes B$ satisfies Browder's theorem according to Lemma 1. Suppose $A \otimes B$ does not satisfy Weyl's theorem, then it follows by Lemma 2 that

$$\sigma_w(A \otimes B) \cap \pi_{00}(A \otimes B) \neq \emptyset.$$

Since the Weyl's spectrum identity holds, that is, since

$$\sigma_w(A \otimes B) = \sigma_w(A) \cdot \sigma(B) \cup \sigma(A) \cdot \sigma_w(B),$$

it follows that every product $\lambda\mu$ with $\lambda \in \sigma_w(A)$ and $\mu \in \sigma(B)$ or with $\lambda \in \sigma(A)$ and $\mu \in \sigma_w(B)$ lies in $\sigma_w(A \otimes B)$. Thus, $v \in \sigma_w(A \otimes B)$ if and only if $v = \lambda\mu$ for arbitrary λ, μ such that $(\lambda, \mu) \in \sigma_w(A) \times \sigma(B)$ or $(\lambda, \mu) \in \sigma(A) \times \sigma_w(B)$. First suppose $v \in \sigma_w(A \otimes B)$ is such that

$$v = \lambda\mu \quad \text{for an arbitrary pair } (\lambda, \mu) \in \sigma_w(A) \times \sigma(B).$$

If $v \in \pi_{00}(A \otimes B)$, then

$$v \in \pi_{00}(A) \cdot \pi_{00}(B)$$

by Lemma 3 because A and B are isoloid, so that

$$v = \lambda'\mu' \quad \text{for some pair } (\lambda', \mu') \in \pi_{00}(A) \times \pi_{00}(B).$$

Thus, if

$$v \in \sigma_w(A \otimes B) \cap \pi_{00}(A \otimes B),$$

then it follows by Lemma 4 that $0 \neq v = \lambda\mu = \lambda'\mu'$, with

$$\lambda = \frac{v}{\mu} \in \sigma_w(A), \quad \lambda' = \frac{v}{\mu'} \in \pi_{00}(A) = \sigma_0(A),$$

$$\mu = \frac{v}{\lambda} \in \sigma(B), \quad \mu' = \frac{v}{\lambda'} \in \pi_{00}(B) = \sigma_0(B),$$

because A and B satisfy Weyl's theorem. Thus, $\lambda \neq \lambda'$, since they live in complementary sets, and so $\mu \neq \mu'$. Since

$$\sigma_{\text{iso}}(A) \neq \emptyset \quad \text{and} \quad \sigma_{\text{iso}}(B) \neq \emptyset$$

(because $\lambda' \in \pi_{00}(A) \subseteq \sigma_{\text{iso}}(A)$ and $\mu' \in \pi_{00}(B) \subseteq \sigma_{\text{iso}}(B)$), and since

$$v = \lambda\mu \in \pi_{00}(A \otimes B) \subseteq \sigma_{\text{iso}}(A \otimes B),$$

it follows by Lemma 5 that

$$\lambda \in \sigma_{\text{iso}}(A) \quad \text{and} \quad \mu \in \sigma_{\text{iso}}(B).$$

Recalling again that A and B are isoloid, we get

$$\lambda \in \sigma_P(A) \quad \text{and} \quad \mu \in \sigma_P(B).$$

Thus, since $\lambda\mu = v \in \pi_{00}(A \otimes B) \subseteq \sigma_{PF}(A \otimes B)$, it follows by Lemma 6 that

$$\lambda \in \sigma_{PF}(A) \quad \text{and} \quad \mu \in \sigma_{PF}(B).$$

Therefore, since A satisfies Weyl's theorem,

$$\lambda \in \sigma_w(A) \cap \sigma_{\text{iso}}(A) \cap \sigma_{PF}(A) = \sigma_w(A) \cap \pi_{00}(A) = \sigma_w(A) \cap \sigma_0(A) = \emptyset,$$

which is a contradiction. On the other hand, if $v \in \sigma_w(A \otimes B)$ is such that

$$v = \lambda\mu \quad \text{for an arbitrary pair } (\lambda, \mu) \in \sigma(A) \times \sigma_w(B),$$

then, similarly and symmetrically,

$$\mu \in \sigma_w(B) \cap \sigma_{\text{iso}}(B) \cap \sigma_{PF}(B) = \sigma_w(B) \cap \pi_{00}(B) = \sigma_w(B) \cap \sigma_0(B) = \emptyset,$$

which is again a contradiction because B satisfies Weyl's theorem as well. Therefore, if A and B are isoloid, both satisfy Weyl's theorem, and the Weyl spectrum identity holds, then $A \otimes B$ must satisfy Weyl's theorem. \square

REMARK. The a -version of Theorem 1 was considered in [2]: If A and B are a -isoloid, satisfy a -Weyl's theorem and the a -Weyl spectrum identity holds, then $A \otimes B$ satisfies a -Weyl's theorem (where the prefix ' a ' means that spectrum is replaced with approximate point spectrum in every definition).

REFERENCES

1. A. Brown and C. Pearcy, Spectra of tensor products of operators, *Proc. Amer. Math. Soc.* **17** (1966), 162–166.
2. B. P. Duggal, S. V. Djordjević and C. S. Kubrusly, On the a -Browder and a -Weyl spectra of tensor products, *Rend. Circ. Mat. Palermo* **59** (2010), 473–481.
3. T. Ichinose, Spectral properties of linear operators I, *Trans. Amer. Math. Soc.* **235** (1978), 75–113.
4. D. Kitson, R. Harte and C. Hernandez, Weyl's theorem and tensor products: A counterexample, *J. Math. Anal. Appl.* **378** (2011), 128–132.
5. C. S. Kubrusly, Fredholm theory in Hilbert space – a concise introductory exposition, *Bull. Belg. Math. Soc. Simon Stevin* **15** (2008), 153–177.

6. C. S. Kubrusly, *Spectral theory of operators on Hilbert spaces* (Birkhäuser/Springer, New York, 2012).
7. C. S. Kubrusly and B. P. Duggal, On Weyl and Browder spectra of tensor product, *Glasgow Math. J.* **50** (2008), 289–302.
8. Y.-M. Song and A.-H. Kim, Weyl's theorem for tensor products, *Glasgow. Math. J.* **46** (2004), 301–304.