

concept scale (KK Scale). The total costs of treatment in the past year were estimated including all outpatient services used within the last year.

**Results:** The costs of treatment amounted on average 31000 DM/year in the year before index admission and 43000 DM/year in the following year (including index admission). The costs were highly correlated with social disability (t2) ( $p < 0.0001$ ) and moderately correlated with negative symptoms (t1, t2) ( $p < .05$ ), but not with positive or general symptoms. Negative symptoms were highly correlated with social disability (t1, t2) ( $p < .01$ ), positive and general symptoms only moderately (t2) ( $p < .05$ ). The t1-t2-correlation of the KKS was pretty good with mean  $r = .4$  in the 7 subscales. A negative correlation was found between drug confidence and the costs of treatment in the following year ( $p < .001$ ).

**Conclusions:** The use of medical and social support in chronic schizophrenic patients is highly determined by illness-related factors as negative symptoms and social disability. This supports the assumption that the inpatient and outpatient care system meets the needs of patients.

### Wed-P16

A LONGITUDINAL EVALUATION OF TWO PSYCHIATRIC INPATIENT UNITS ATTACHED TO A COMMUNITY MENTAL HEALTH CENTRE. OUTCOMES AND COSTS

A.P. Boardman<sup>2\*</sup>, R. Hodgson<sup>1</sup>, M. Lewis<sup>1</sup>, K. Allen<sup>1</sup>, A. Haycox<sup>3</sup>. <sup>1</sup>Guy's UMDS, London; <sup>2</sup>Keele University; <sup>3</sup>Liverpool University, UK

**Background:** The inpatient care of those with psychiatric disorder is little evaluated despite the central role of inpatient units in catchment area services. Two innovative 8-bedded psychiatric inpatient units attached to two Community Mental Health Centres (CMHCs) were evaluated. These units have potential advantages for the assessment and after care of those recovering from severe mental illnesses.

**Method:** All patients admitted to the acute psychiatric wards serving the two CMHCs were assessed for suitability for inclusion. Those transferred to the community beds or admitted directly to these beds constituted the experimental group. A control group consisted of those patients admitted to the acute wards serving two demographically similar catchment areas but whose community services did not have associated community beds. Clinical and social measures were made during the first week of admission and repeated 6 and 12 months later. Public and private resource consumption was also measured.

**Results:** 177 individuals met the inclusion criteria. The experimental and control groups were matched on demographic and clinical preliminary variables. The experimental group made significantly greater improvement on most outcome measures including symptom and social measures, user satisfaction, length of acute inpatient stay, number of readmissions, improved after care contact. The overall costs of the community units were greater than that of traditional inpatient units but there was significant variation of costs between the two community units.

**Conclusions:** The attachment of community inpatient units to Community Mental Health Centres has theoretical and practical advantages. The use of such community beds may have significant benefits for clinical and social outcomes for the patients with severe mental illnesses. Higher costs of the units were not inevitable and may depend on the way in which such units are integrated into existing care provision. The model of provision of inpatient services within a Community Mental Health Centre may be worth adopting more widely.

### Wed-P17

REPROVISION OF 'REMNANT' SCHIZOPHRENIC PATIENTS: CAN ALL PATIENTS LEAVE HOSPITAL?

M. Franz<sup>1\*</sup>, F. Ehlers<sup>1</sup>, M. Knoll<sup>2</sup>, T. Meyer<sup>1</sup>, B. Gallhofer<sup>1</sup>. <sup>1</sup>Centre for Psychiatry, Justus-Liebig-University Giessen; <sup>2</sup>Department of hospital planning and development, LWV Hessen, Germany

Several studies on deinstitutionalisation of long-stay patients have brought attention to a group of 'remnant' patients, apparently difficult to move into the community. This has led to pessimistic conclusions about the chances of fully closing long-stay wards of psychiatric hospitals. A remarkable reduction of hospital beds in the 70s and 80s in the German state of Hesse was followed by a political decision in 1992 to close long-stay wards of mental hospitals. The object of this study is (1) to display the characteristics of a remaining population of schizophrenic patients ( $n = 266$ ) that had not been deinstitutionalised despite an ongoing re-provision-program until 1995. (2) Moreover, selection criteria for community re-provision within this 'remnant' group of patients should be identified.

(1) Patients were found to be extremely disabled. Mean duration of stay was 30 years. (2) Only 50 of the 'remnant' patients have been moved into the community since 1995. Significant differences between discharged and remaining 'remnant' patients were found regarding gender, age, and duration of hospitalisation at baseline (1995), while the age of onset of illness was equal in both groups. The patients left in hospital had been less engaged in spare-time activities and less likely to want to leave hospital at baseline. No differences could be found with regard to psychopathology and quantitatively assessed quality of life. However, results of content analysis of open-ended quality of life interviews indicate differences regarding the internal orientation to various life domains between the newly discharged patients and the still remaining patients.

Results indicate that even within this 'remnant' group of patients a selection bias for community re-provision operates. It is discussed whether the 'hard core' of remnant patients could be resistant to re-provision or possibly needs a special intensive care in the community.

### Wed-P18

SCHIZOPHRENIC PATIENTS' NORMATIVE NEEDS FOR CARE IN THE YEAR AFTER DISCHARGE FROM THE HOSPITAL

Th.W. Kallert<sup>1\*</sup>, M. Leißle<sup>1</sup>. <sup>1</sup>Department of Psychiatry, Technical University, 01307 Dresden, Germany

**Introduction:** As part of a public health research project that evaluates restructured psychiatric community care for chronic patients after the German reunification in Saxony a group of ICD-10 schizophrenic patients limited to the Dresden area was examined.

**Methods:** Data were collected 1, 6 and 12 months after discharge from the hospital using the MRC Needs for Care Assessment.

**Characteristics of the Study's Clientele:** ( $n = 112$ ): sex-ratio 1:1; mean age 42 y; 60% unmarried; 75% recipients of a pension; mean duration of disorder 14.4 y; mean GAF-Score 41.8 ( $\pm 11.4$ ), mean BPRS-Score 43.7 ( $\pm 12.5$ ).

**Results:** 1 month after discharge normative psychiatric needs for care (mainly based on informations of psychiatrists, social workers and relatives) with current clinical significance were rated in 2.1 ( $\pm 1.3$ ) clinical and 1.6 ( $\pm 2.3$ ) social NCA-areas (esp. concerning negative symptoms, psychosocial distress, social interaction skills,