



11 mcg (3MIU), 44 mcg (12MIU) lyophilized powder for injection
22 mcg (6MIU)/0.5ml, 44 mcg (12MIU)/0.5ml, liquid formulation for injection

THERAPEUTIC CLASSIFICATION

Immunomodulator

ACTIONS AND CLINICAL PHARMACOLOGY

Description: Rebif® (Interferon beta-1a) is a purified, sterile glycoprotein product produced by recombinant DNA techniques and formulated for use by injection. The active ingredient of Rebif® is produced by genetically engineered Chinese Hamster Ovary (CHO) cells. Interferon beta-1a is a highly purified glycoprotein that has 166 amino acids and an approximate molecular weight of 22,500 daltons. It contains a single N-linked carbohydrate moiety attached to Asn-80 similar to that of natural human Interferon beta. The specific activity of Rebif® is approximately 0.27 million international units (MIU)/mcg Interferon beta-1a. The unit measurement is derived by comparing the antiviral activity of the product to an in-house natural hFN-beta NIH standard that is obtained from human fibroblasts (BILS 11), which has been calibrated against the NIH natural hFN-beta standard (GB 23-902-531). General: Interferons are a family of naturally occurring proteins, which have molecular weights ranging from 15,000 to 21,000 daltons. Three major classes of interferons have been identified: alpha, beta, gamma Interferon beta, Interferon alpha and Interferon gamma have overlapping yet distinct biologic activities.

Interferon beta-1a acts through various mechanisms:

- Immunomodulation through the induction of cell membrane components of the major histocompatibility complex i.e., MHC Class I antigens, an increase in natural killer (NK) cell activity, and an inhibition of IFN-gamma induced MHC Class II antigen expression, as well as a sustained reduction in TNF level.
Antiviral effect through the induction of proteins like 2'-5' oligoadenylate synthetase and p78.
Antiproliferative effect through direct cytostatic activity and indirect through antitumoral immune response enhancement.

The mechanism of action of Rebif® in relapsing-remitting multiple sclerosis is still under investigation.

Relapsing-Remitting Multiple Sclerosis

Two pivotal studies, including a total of 628 patients, evaluated the long-term safety and efficacy of Rebif® when administered subcutaneously three times weekly to relapsing-remitting multiple sclerosis patients. The results indicate that Rebif® alters the natural course of relapsing-remitting multiple sclerosis. Efficacy was demonstrated with respect to the 3 major aspects of this disease: disability (patients EDSS 0-5), exacerbations, and burden of disease and activity as measured by MRI scans.

PRISMS STUDY

In the larger trial, a total of 560 patients diagnosed with clinically definite or laboratory-supported relapsing-remitting multiple sclerosis EDSS 0-5 with at least a 1-year history before study entry, were enrolled and randomized to the 3 treatments (placebo, 22 mcg (6MIU) Rebif®, or 44 mcg (12MIU) Rebif®) in a ratio of 1:1:1. About 90% of patients completed the 2 years of treatment, and very few patients withdrew from the study due to adverse events.

The main criteria for inclusion were:

- history of 2 or more acute exacerbations in the 2 years prior to study entry
no previous systemic treatment with interferons
no treatment with corticosteroids or ACTH in the 2 months preceding study entry
no exacerbation in the 8 weeks prior to study entry

Patients were evaluated at 3-month periods, during exacerbations and coinciding with MRI scanning. Each patient underwent cranial proton density/T2-weighted (PD/T2) MRI scans at baseline and every 6 months during the study. A subset of patients underwent PD/T2 and T1-weighted (T1) Gd-MRI scans one month before the start of treatment, at baseline and then monthly until the end of the first 9 months of treatment. Of those, another subset of 39 continued with the monthly scans throughout the 24 month treatment period.

This study demonstrated that Rebif® at a total dose of 66 or 132 mcg weekly, significantly improved all 3 major outcomes, including exacerbation rate, disease activity and burden of disease as measured by MRI scanning and progression of disability. In addition, the study showed that Rebif® is effective in delaying the progression in disability in patients with an EDSS of 4.0 or higher who are known to progress more rapidly. Also, the drug reduced the requirements for steroids to treat multiple sclerosis and, at 132 mcg weekly Rebif® reduced the number of hospitalizations for multiple sclerosis.

Effect on exacerbation

Table with 5 columns: Efficacy parameters, Placebo, Rebif 22 mcg/wk, Rebif 44 mcg/wk, Rebif 66 mcg/wk vs placebo, Rebif 132 mcg/wk vs placebo. Rows include Mean # exacerbations over 2 year study, % of exacerbation-free patients, Median time to first exacerbation, Median time to second exacerbation, and Mean # of moderate and severe exacerbations during 2 year period.

* Median time to second exacerbation not reached in 132 mcg/wk dose group

The results after one year of treatment were also significant.

Effect on time to first progression in disability

Table with 5 columns: Efficacy parameters, Placebo, Rebif 22 mcg/wk, Rebif 44 mcg/wk, Rebif 66 mcg/wk vs placebo, Rebif 132 mcg/wk vs placebo. Rows include Time to confirmed progressive disability (first quartile) and Median change in EDSS score at 2 years.

Effect on multiple sclerosis pathology as detected by MRI scans

Table with 5 columns: Efficacy parameters, Treatment Groups (Placebo, Rebif 22 mcg/wk, Rebif 44 mcg/wk, Rebif 66 mcg/wk vs placebo, Rebif 132 mcg/wk vs placebo), p-value. Rows include Burden of disease (BD) Median % change, MRI activity (All patients, Patients with monthly MRIs), and Number active lesions (per month) for patients with monthly MRIs throughout the study (2 years).

Requirement for steroids: The proportion of patients requiring steroids for MS (excluding non-MS indications) was higher in the placebo group (more than 50%) than in either of the 2 Rebif® groups (around 40% in each group).

Hospitalization for multiple sclerosis: The observed mean numbers of hospitalizations for MS in the Rebif® 66 and 132 mcg weekly groups represented reductions of 21% and 48%, respectively, from that in the placebo group.

Cohort of patients with high baseline EDSS (baseline EDSS >3.5):

Additional analyses were conducted in order to study the efficacy of Rebif® in populations of patients with adverse prognostic outcome factors, who were likely to be at higher risk for progression in disability. The primary prognostic factor examined was baseline EDSS >3.5. Patients in this cohort have a more severe degree of disability and are at higher risk for progression than those with lower EDSS: natural history studies have shown that patients at EDSS levels of 4.0 to 5.0 spend less time at these EDSS levels than at lower levels of disability. Treatment with Rebif® at both doses significantly reduced the mean exacerbation count per patient compared to placebo treatment. Progression in this group of patients is of particular concern, as it involves development of difficulty in ambulation. The 132 mcg weekly dose significantly prolonged time to confirmed progression whereas the 66 mcg weekly dose did not. Both doses of Rebif® significantly affected percent change from baseline in MRI burden of disease in the high-EDSS cohort, and the 132 mcg weekly dose significantly reduced the number of T2 active lesions in this population. The efficacy results in this cohort of patients with established disability confirms that the 132 mcg weekly dose has a marked effect on progression in disability and the underlying pathology of the disease.

Effect on exacerbation (High-EDSS cohort)

Table with 4 columns: Efficacy parameters, Placebo, Rebif 66 mcg/week, Rebif 132 mcg/week. Rows include Mean # exacerbations, % of exacerbation-free patients, and p-value (Rebif vs placebo).

* Log-linear model.

Progression in disability by one point on the EDSS (High-EDSS cohort)

Table with 4 columns: Treatment Group, % of progressors, # patients, Median (days), Q1 (days). Rows include Placebo, Rebif 66 mcg weekly, and Rebif 132 mcg weekly.

* Excludes patients lost to follow-up without progression.

Progression in disability: statistical comparisons

Table with 3 columns: Test, Group Comparison, p-value. Rows include Log-rank test for 66 mcg weekly vs placebo and 132 mcg weekly vs placebo.

MRI Burden of Disease: % Change (High-EDSS cohort)

Table with 4 columns: Burden of disease - Median % change, Placebo, Rebif 66 mcg/week, Rebif 132 mcg/week. Rows include Burden of disease - Median % change, Burden of disease - Mean % change, and p-value (Rebif vs placebo).

* ANOVA on the ranks.

Number of T2 Active Lesions (High-EDSS cohort)

Table with 4 columns: Treatment Group, Median, Mean, p-value**. Rows include Placebo, Rebif 66 mcg weekly, and Rebif 132 mcg weekly.

* ANOVA on the ranks.

CROSS-OVER STUDY

The other study was an open cross-over design, with MRI evaluations conducted in a blinded fashion. Enrolled in this study were 68 patients between the ages of 15 and 45 years, with clinically definite and/or laboratory supported relapsing-remitting MS for up to 10 years in duration. The main inclusion criteria included:
at least 2 relapses in the previous 2 years
EDSS score between 1-5

- no corticosteroid or plasmapheresis treatments or administration of gamma globulins within the 3 months prior to study
no immunomodulating or immunosuppressive therapy for the 6 months prior to the study
absence of HBSAg and HIV antibodies.

Once enrolled, patients remained under clinical observation for 6 months with assessments of their neurological status and other parameters, and extensive monitoring of exacerbations. Patients were then randomized to treatment with either 11 mcg (3MIU) (n=35) or 33 mcg (9MIU) (n=33) of Rebif®, self-administered subcutaneously three times per week. The total dose was therefore 33 or 99 mcg weekly.

Six-months observation vs six-months treatment:

Treatment with Rebif® at both doses used in this study, achieved a statistically significant reduction in both the MRI evidence of MS activity in the brain and the clinical relapse rate versus the corresponding observation periods. This pattern of improvement was also reflected in additional MRI measures. In the biannual T2-weighted scans, a reduction in the mean number of new lesions and in the mean number of enlarging lesions was demonstrated.

Table with 5 columns: Outcome, Observation period, Treatment period, Reduction %, p-value. Rows include Exacerbation rate, # exacerbation-free patients, # of monthly lesions, Volume of lesions, Total mean # of T2 lesions, and Total mean # of T2 enlarged lesions.

Two-year results: At the end of this study, 62 patients continued treatment for a further 18 months. Each of these patients continued to receive the dose to which they were randomized. Validation of the results of the 2 year treatment period is ongoing, however, the results from the continuation of treatment at both doses demonstrate that Rebif® maintained its dose-dependent effect in reducing the relapse rate and the brain lesion volume detected by T2 weighted MRI scans compared to the observation period, which corroborates the findings of the longer, placebo-controlled study.

Condyloma acuminatum: The results from four double-blind, placebo-controlled studies, including 349 patients (aged 17-62), each reveal that Rebif®, when injected intralesionally at a dose of 3.67 mg (1MIU)/lesion 3 times per week for 3 weeks, is efficacious in the treatment of condyloma acuminatum in men and women. This efficacy is evidenced by both the induction of complete disappearance of lesions as well as the reduction in the area of lesions. The majority of treated patients in these studies had recurrent warts that had failed previous treatments. The number of lesions treated per patient was between 3 and 8, as stated in the summary table below.

Table with 4 columns: Study, # patients (%), # lesions treated, Results. Rows include study 1, 2, 3, and 4 with details on patient numbers, lesion counts, and efficacy results.

INDICATIONS AND CLINICAL USE

Multiple Sclerosis: Rebif® (Interferon beta-1a) is indicated for the treatment of relapsing-remitting multiple sclerosis in patients with an EDSS between 0 and 5.0, to reduce the number and severity of clinical exacerbations, slow the progression of physical disability, reduce the requirement for steroids, and reduce the number of hospitalizations for treatment of multiple sclerosis. The efficacy has been confirmed by T1-Gd enhanced and T2 (burden of disease) MRI evaluations. Evidence of efficacy beyond 2 years is not known since the primary evidence of efficacy derives from 2-year trials. Condyloma acuminatum: Rebif® is best suited for the patient who has less than nine lesions, and who has failed several prior treatments. In the case of patients with nine or more lesions, if the first Rebif® treatment is successful, the remaining lesions could be treated with a second course of Rebif® therapy. Rebif® should also be considered for the treatment of condyloma acuminatum in patients for whom the side-effects from other treatments, e.g., scarring, are of concern. While not all patients who were treated with Rebif® attained a complete response, patients whose lesions decreased in size and had at least a partial response may have also benefited from treatment because lesion shrinkage may facilitate subsequent management with other therapies, as has been reported with IFN-alpha.

CONTRAINDICATIONS:

Rebif® (Interferon beta-1a) is contraindicated in patients with a known hypersensitivity to natural or recombinant interferon beta, albumin (human), or any other component of the formulation.

WARNINGS:

Rebif® (Interferon beta-1a) should be used under the supervision of a physician.

Relapsing-Remitting Multiple Sclerosis: Depression and suicidal ideation are known to occur at an increased frequency in the multiple sclerosis population. The use of Rebif® has not been associated with an increase in the incidence and/or severity of depression, or with an increased incidence of suicide attempts or suicide. In the relapsing-remitting multiple sclerosis study, a similar incidence of depression was seen in the placebo-treated group and in the two Rebif® patient groups. Nevertheless, patients with depression should be closely monitored for signs of significant worsening of depression or suicidal ideation. The first injection should be performed under the supervision of an appropriately qualified health care professional. Condyloma: All injections should be administered by a qualified health care professional.

PRECAUTIONS

General: Patients should be informed of the most common adverse events associated with interferon beta administration, including symptoms of the flu-like syndrome (see Adverse Reactions). These symptoms tend to be most prominent at the initiation of therapy and decrease in frequency and severity with continued treatment.

Based on the results of clinical trials of Rebif® in MS, in which more than 500 patients were randomized to drug treatment, there is no indication of an increased risk of seizure disorder with Rebif® therapy. However, since seizures have been reported with other interferon therapies, caution should be exercised when administering interferon-beta-1a to patients with pre-existing seizures disorder. For patients without a pre-existing seizure disorder who develop seizures during therapy, an etiologic basis should be established and appropriate anti-convulsant therapy instituted prior to considering resuming treatment with Rebif®. The effect of Rebif® administration on the medical management of patients with seizure disorder is unknown. Serum neutralising antibodies against Rebif® (interferon beta-1a) may develop. The precise incidence and clinical significance of antibodies is as yet uncertain (see ADVERSE REACTIONS).

Hypersensitivity reactions, both local and systemic, have developed during therapy with Rebif®. Intralesional injections can be painful to some patients treated for condyloma acuminatum. In such cases an anesthetic cream such as lidocaine-prilocaine can be used.

Pregnancy and Lactation: Rebif® should not be administered in case of pregnancy and lactation. There are no studies of interferon beta-1a in pregnant women. At high

Table 3: Treatment-Emergent Adverse Events in Oral Placebo-Controlled Clinical Trials Reported by at Least 1% of Patients with Migraine

	Placebo	IMITREX® 25 mg	IMITREX® 50 mg	IMITREX® 100 mg**
Number of Patients	690	351	723	2021
Number of Migraine Attacks Treated	1187	945	1889	14750
Symptoms of Potentially Cardiac Origin				
• Chest Sensations*	0.6%	2.3%	2.6%	3.2%
• Neck/Throat/Jaw Sensations*	1.4%	2.3%	3.5%	5.2%
• Upper Limb Sensations*	1.2%	1.4%	2.5%	3.6%
• Palpitations	0.6%	0.3%	1.0%	1.1%
Neurological				
• Head/Face Sensations*	1.3%	2.3%	2.5%	4.7%
• Dizziness	2.5%	3.1%	3.3%	6.2%
• Headache	3.3%	4.0%	2.2%	3.3%
• Vertigo	0.6%	1.1%	1.1%	1.0%
• Drowsiness	1.6%	1.1%	1.2%	2.1%
• Tremor	0.4%	0.9%	0.4%	1.1%
Gastrointestinal				
• Nausea	5.8%	2.8%	4.4%	11.0%
• Hyposalivation	1.2%	1.4%	1.1%	1.2%
• Vomiting	2.9%	4.3%	1.1%	4.4%
• Gastrointestinal Discomfort & Pain	1.4%	1.1%	0.8%	2.0%
• Abdominal Discomfort & Pain	0.3%	NR	0.4%	1.2%
• Diarrhea	0.9%	0.3%	0.6%	1.1%
Musculoskeletal				
• Musculoskeletal Pain	0.7%	2.3%	0.4%	1.4%
• Muscle Pain	0.3%	0.9%	0.1%	1.0%
• Muscle Atrophy Weakness & Tiredness	NR	0.6%	0.4%	1.4%
Ear, Nose & Throat				
• Infections	0.6%	0.6%	1.1%	1.4%
• Nasal Signs & Symptoms	0.7%	1.4%	0.8%	1.0%
• Throat & Tonsil Symptoms	0.6%	NR	0.4%	2.3%
Respiratory				
• Viral Infection	0.3%	1.1%	0.1%	1.0%
Non-Site Specific				
• Limb Sensations*	0.4%	1.1%	0.4%	1.5%
• Sensations* (body region unspecified)	4.5%	5.7%	8.0%	9.0%
• Malaise/Fatigue	5.1%	3.7%	2.6%	9.5%
• Sweating	0.4%	0.6%	0.6%	1.6%

*The term "sensations" encompasses adverse events described as pain & discomfort, pressure, heaviness, constriction, tightness, heat/burning sensation, paresthesia, numbness, tingling, and strange sensations.
**Includes patients receiving up to 3 doses of 100 mg
NR = Not Reported

Table 5: Treatment-Emergent Adverse Events in Intranasal Placebo-Controlled Clinical Trials Reported by at Least 1% of Patients with Migraine

	Placebo	IMITREX® 5 mg	IMITREX® 10 mg	IMITREX® 20 mg**
Number of Patients	741	496	1007	1638
Number of Migraine Attacks Treated	1047	933	1434	2070
Symptoms of Potentially Cardiac Origin				
• Chest Sensations*	0.3%	1.0%	0.7%	0.6%
• Neck/Throat/Jaw Sensations*	1.2%	0.6%	1.6%	2.3%
Neurological				
• Head/Face Sensations*	0.8%	1.4%	2.4%	2.4%
• Dizziness	1.2%	1.6%	1.5%	1.2%
• Headache	0.7%	1.4%	0.9%	0.8%
• Migraine	2.6%	3.2%	2.4%	1.8%
Gastrointestinal				
• Nausea	10.4%	14.3%	9.6%	8.3%
• Vomiting	7.6%	11.1%	9.6%	6.8%
Ear, Nose & Throat				
• Sensitivity to Noise	3.1%	4.4%	2.5%	1.5%
• Nasal Signs & Symptoms	1.3%	3.0%	1.6%	1.8%
• Infections	0.9%	1.8%	1.3%	0.5%
• Upper Respiratory Inflammation	0.5%	1.0%	0.6%	0.7%
• Throat & Tonsil Symptoms	0.8%	0.2%	1.0%	0.7%
Non-Site Specific				
• Sensations* (body region unspecified)	1.8%	2.4%	2.7%	2.4%
• Malaise/Fatigue	1.3%	1.8%	1.3%	0.8%
• Descriptions of odor or taste	1.8%	15.3%	20.2%	20.8%

*The term "sensations" encompasses adverse events described as pain & discomfort, pressure, heaviness, constriction, tightness, heat/burning sensation, paresthesia, numbness, tingling, and strange sensations.
**Includes patients receiving up to 3 doses of 20 mg

IMITREX® is generally well tolerated. Most of the events were transient in nature and resolved within 45 minutes of subcutaneous administration and within 2 hours of oral or intranasal administration. Of the 3630 patients treated with IMITREX® Nasal Spray in clinical trials, there was one report of a coronary vasospasm related to IMITREX® administration. Minor disturbances of liver function tests have occasionally been observed with sumatriptan treatment. There is no evidence that clinically significant abnormalities occurred more frequently with sumatriptan than with placebo. Patients treated with IMITREX® rarely exhibit visual disorders like flickering and diplopia. Additionally, cases of nystagmus, scotoma and reduced vision have been observed. Very rarely a transient loss of vision has been reported. However, visual disorders may also occur during a migraine attack itself.

DOSE AND ADMINISTRATION

General:
IMITREX® (sumatriptan succinate/sumatriptan) is indicated for the acute treatment of migraine headache with or without aura. Sumatriptan should not be used prophylactically. Sumatriptan may be given orally, subcutaneously or as a nasal spray. The safety of treating an average of more than four headaches in a 30 day period has not been established.

In selecting the appropriate formulation for individual patients, consideration should be given to the patient's preference for formulation and the patient's requirement for rapid onset of relief. Significant relief begins about 10-15 minutes following subcutaneous injection, 15 minutes following intranasal administration and 30 minutes following oral administration.

In addition to relieving the pain of migraine, sumatriptan (all formulations) has also been shown to be effective in relieving associated symptoms of migraine (nausea, vomiting, photophobia, photophobia). Sumatriptan is equally effective when administered at any stage of a migraine attack. Long term (12-24 months) clinical studies with maximum recommended doses of sumatriptan indicate that there is no evidence of the development of tachyphylaxis, or medication-induced (rebound) headache.

Tablets:
The minimal effective single adult dose of IMITREX® Tablets is 25 mg. The maximum recommended single dose is 100 mg.

The optimal dose is a single 50 mg tablet. However, depending on individual patient's needs and response to treatment, some patients may require 100 mg. Clinical trials have shown that approximately 50 - 75% of patients have headache relief within two hours after oral dosing with 100 mg, and that a further 15 - 25% have headache relief by 4 hours. Comparator studies have shown similar efficacy rates with the 50 mg and 100 mg tablets. There is evidence that doses of 50 and 100 mg may provide greater effect than 25 mg. If the migraine headache returns, or if a patient has a partial response to the initial dose, the dose may be repeated after 2 hours. Not more than 200 mg should be taken in any 24 hour period.

If a patient does not respond to the first dose of IMITREX® Tablets, a second dose should not be taken for the same attack, as it is unlikely to be of clinical benefit. IMITREX® may be taken to treat subsequent migraine attacks.

The tablet should be swallowed whole with water, not crushed, chewed or split.
Hepatic Impairment: In patients with mild or moderate hepatic impairment, plasma sumatriptan concentrations up to two times those seen in healthy subjects have been observed. Therefore, a 25 mg dose (single tablet) may be considered in these patients (see PRECAUTIONS). Sumatriptan should not be administered to patients with severe hepatic impairment (see CONTRAINDICATIONS).

Injection:
IMITREX® injection should be injected subcutaneously (on the outside of the thigh or in the upper arm) using an autoinjector.

The recommended adult dose of sumatriptan is a single 6 mg subcutaneous injection. Clinical trials have shown that approximately 70-72% of patients have headache relief within one hour after a single subcutaneous injection. This number increases to 82% by 2 hours.

If the migraine headache returns, or if a patient has a partial response to the

initial dose, the dose may be repeated after 1 hour. Not more than 12 mg (two 6mg injections) should be taken in any 24 hour period.
If a patient does not respond to the first dose of IMITREX® injection, a second dose should not be taken for the same attack, as it is unlikely to be of clinical benefit. IMITREX® may be taken for subsequent attacks.
Administration during migraine aura prior to other symptoms occurring may not prevent the development of a headache.
Patients should be advised to read the patient instruction leaflet regarding the safe disposal of syringes and needles.

Nasal Spray:
The minimal effective single adult dose of sumatriptan nasal spray is 5 mg. The maximum recommended single dose is 20 mg.

If the migraine headache returns, or if a patient has a partial response to the initial dose, the dose may be repeated after 2 hours. Not more than 40 mg should be taken in any 24 hour period.

If a patient does not respond to the first dose of IMITREX® Nasal Spray, a second dose should not be taken for the same attack, as it is unlikely to be of clinical benefit. IMITREX® may be taken for subsequent attacks.

Placebo-controlled clinical trials revealed the following incidence of headache relief, defined as a decrease in migraine severity from severe or moderate to mild or no pain, within 2 hours after treatment with intranasal sumatriptan at doses of 5, 10 or 20 mg (see Table 6 below).

TABLE 6. Percentage of patients with headache relief at 2 hours

Study	Placebo (n)	5 mg (n)	10 mg (n)	20 mg (n)
Study 1*	35% (40)	67% ^v (42)	67% ^v (39)	78% ^v (40)
Study 2*	42% (31)	45% (33)	66% ^v (35)	74% ^v (39)
Study 3	25% (63)	49% ^v (122)	46% ^v (115)	64% ^v † (119)
Study 4	25% (151)	-	44% ^v (288)	55% ^v † (292)
Study 5	32% (198)	44% ^v (297)	54% ^v (293)	60% ^v † (288)
Study 6*	35% (100)	-	54% ^v (106)	63% ^v (202)
Study 7*	29% (112)	-	43% (109)	62% ^v (215)

Headache relief was defined as a decrease in headache severity from severe or moderate to mild or none.

n = total number of patients who received treatment
* comparisons between sumatriptan doses not conducted
^v p < 0.05 versus placebo † p < 0.05 versus lower sumatriptan doses
* p < 0.05 vs 5 mg - not evaluated

As shown in the table above, optimal rates of headache relief were seen with the 20 mg dose. Single doses above 20 mg should not be used due to limited safety data and lack of increased efficacy relative to the 20 mg single dose. Within the range of 5-20 mg, an increase in dose was not associated with any significant increase in the incidence or severity of adverse events other than taste disturbance (see ADVERSE REACTIONS).

The nasal spray should be administered into one nostril only. The device is a ready to use single dose unit and must not be primed before administration. Patients should be advised to read the patient instruction leaflet regarding the use of the nasal spray device before administration.

AVAILABILITY OF DOSAGE FORMS

IMITREX® Tablets are available as pink 100 mg, white 50 mg, or white 25 mg film-coated tablets in blister packs containing 6 tablets. Four blister packs are placed in a carton.

IMITREX® Injection (6 mg total volume = 0.5 mL) is available in pre-filled syringes placed in a larger-evident carrying case/disposal case. Two pre-filled syringes plus the IMITREX® STATdose Pen™ autoinjector are packed in an IMITREX® STATdose System™ autoinjector kit. A refill pack is available containing 2 pre-filled syringes in a carton.

IMITREX® Injection is also available to physicians or hospitals in a single dose vial (6 mg total volume = 0.5mL). There are 5 vials per carton.

IMITREX® Nasal Spray 5 mg and 20 mg are each supplied in boxes of 6 nasal spray devices (3 x 2 devices). Each unit dose spray supplies 5 and 20 mg respectively.

Product Monograph available to physicians and pharmacists upon request.

Please contact GlaxoSmithKline Inc., 7333 Mississauga Road N., Mississauga, Ontario L5N 6L4

IMITREX® is a registered trademark, used under license by GlaxoSmithKline Inc. *The appearance, namely the colour, shape, and size of the IMITREX® Nasal Spray device and IMITREX® STATdose System are trademarks, used under license by GlaxoSmithKline Inc.

Date of preparation: January 17, 1992
Date of revision: February 14, 2003

References: 1. Product Monograph of "IMITREX® (sumatriptan succinate/sumatriptan); GlaxoSmithKline Inc. February 2003. 2. Cady R, McNeal S, O'Quinn S, Putman G. Effect of early intervention with sumatriptan on migraine pain: Retrospective analyses of data from three clinical trials. *Clinical Therapeutics* 2000;22(9):1035-1048.



GlaxoSmithKline
7333 Mississauga Road North
Mississauga, Ontario L5N 6L4



*The term "sensations" encompasses adverse events described as pain & discomfort, pressure, heaviness, constriction, tightness, heat/burning sensation, paresthesia, numbness, tingling, and strange sensations.
NR = Not Reported

Table 3
Dose-Related Adverse Events From Placebo-Controlled, Add-On Trials in ADULTS

Adverse Event	TOPAMAX Dosage (mg/day)			
	Placebo (n=216)	200 (n=45)	400 (n=68)	600–1,000 (n=414)
Fatigue	13.4	11.1	11.8	29.7
Nervousness	7.4	13.3	17.6	19.3
Difficulty with Concentration/Attention	1.4	6.7	8.8	14.5
Confusion	4.2	8.9	10.3	13.8
Depression	5.6	8.9	7.4	13.0
Anorexia	3.7	4.4	5.9	12.3
Language problems	0.5	2.2	8.8	10.1
Anxiety	6.0	2.2	2.9	10.4
Mood problems	1.9	0.0	5.9	9.2

In six double-blind clinical trials, 10.6% of subjects (n=113) assigned to a topiramate dosage of 200 to 400 mg/day in addition to their standard AED therapy discontinued due to adverse events, compared to 5.8% of subjects (n=69) receiving placebo. The percentage of subjects discontinuing due to adverse events appeared to increase at dosages above 400 mg/day. Overall, approximately 17% of all subjects (n=527) who received topiramate in the double-blind trials discontinued due to adverse events, compared to 4% of the subjects (n=216) receiving placebo.

Pediatrics

Adverse events associated with the use of topiramate at dosages of 5 to 9 mg/kg/day in worldwide pediatric clinical trials that were seen at greater frequency in topiramate-treated patients were: fatigue, somnolence, anorexia, nervousness, difficulty with concentration/attention, difficulty with memory, aggressive reaction, and weight decrease.

Table 4 lists treatment-emergent adverse events that occurred in at least 2% of children treated with 5 to 9 mg/kg/day topiramate in controlled trials that were numerically more common than in patients treated with placebo.

Table 4

Incidence (%) of Treatment-Emergent Adverse Events in Worldwide Pediatric Clinical Trials Experience (2-16 years of Age)^a
(Events that Occurred in ≥2% of Topiramate-Treated Patients and Occurred More Frequently in Topiramate-Treated Than Placebo-Treated Patients)

Body System/ Adverse Event	Placebo (N=101)	Topiramate (N=98)
Body as a Whole - General Disorders		
Fatigue	5	16.3
Injury	12.9	14.3
Allergic Reaction	1	2
Central & Peripheral Nervous System Disorders		
Gait Abnormal	5	8.2
Ataxia	2	6.1
Hyperkinesia	4	5.1
Dizziness	2	4.1
Speech Disorders/Related Speech Problems	2	4.1
Convulsions Aggravated	3	3.1
Hyporeflexia	0	2
Gastrointestinal System Disorders		
Nausea	5	6.1
Saliva Increased	4	6.1
Constipation	4	5.1
Gastroenteritis	2	3.1
Metabolic and Nutritional Disorders		
Weight Decrease	1	9.2
Thirst	1	2
Platelet, Bleeding, & Clotting Disorders		
Purpura	4	8.2
Epistaxis	1	4.1
Nervous Disorders		
Somnolence	15.8	25.5
Anorexia	14.9	24.5
Nervousness	6.9	14.3
Personality Disorder (Behavior Problems)	8.9	11.2
Difficulty with Concentration/Attention	2	10.2
Aggressive Reaction	4	9.2
Insomnia	6.9	8.2
Mood Problems	6.9	7.1
Difficulty with Memory NOS	0	5.1
Emotional Lability	5	5.1
Confusion	3	4.1
Psychomotor Slowing	2	3.1
Reproductive Disorders, Female		
Leukorrhea	0.0	2.3
Resistance Mechanism Disorders		
Infection Viral	3.0	7.1
Infection	3.0	3.1
Respiratory System Disorders		
Upper Respiratory Tract Infection	36.6	36.7
Pneumonia	1.0	5.1
Skin and Appendages Disorders		
Skin Disorder	2.0	3.1
Alopecia	1.0	2.0
Dermatitis	0.0	2.0
Hypertichiasis	1.0	2.0
Rash Erythematous	0.0	2.0
Urinary System Disorders		
Urinary Incontinence	2.0	4.1
Vision Disorders		
Eye Abnormality	1.0	2.0
Vision Abnormal	1.0	2.0
White Cell and RES Disorders		
Leukopenia	0.0	2.0

^a Patients in these add-on trials were receiving 1 to 2 concomitant antiepileptic drugs in addition to TOPAMAX topiramate or placebo.

^b Values represent the percentage of patients reporting a given adverse event. Patients may have reported more than one adverse event during the study and can be included in more than one adverse event category.

^c Not Otherwise Specified.

None of the pediatric patients who received topiramate adjunctive therapy at 5 to 9 mg/kg/day in controlled clinical trials discontinued due to adverse events. In open extensions of the controlled clinical trials, approximately 9% of the 303 pediatric patients who received topiramate at dosages up to 30 mg/kg/day discontinued due to adverse events. Adverse events associated with discontinuing therapy included aggravated convulsions (2.3%), language problems (1.3%), and difficulty with concentration/attention (1.3%).

In adult and pediatric patients, nephrolithiasis was reported rarely. Isolated cases of thromboembolic events have also been reported; a causal association with the drug has not been established.

When the safety experience of patients receiving TOPAMAX topiramate as adjunctive therapy in both double-blind and open-label trials (1,446 adults and 303 children) was analyzed, a similar pattern of adverse events emerged.

Post-Marketing Adverse Reactions

The most frequently reported adverse events in spontaneous post-marketing reports on topiramate include:

Psychiatric: somnolence or sedation, hallucination(s), depression, anorexia, aggressive reaction, psychosis, thinking abnormal, paranoid reaction, insomnia, emotional lability, suicide attempt, delusion

Central and Peripheral Nervous System: confusion, convulsions aggravated, paresthesia, agitation, speech disorder, ataxia, dizziness, convulsions, amnesia, headache, hyperkinesia

Metabolic and Nutritional: weight decrease

Autonomic Nervous System: vomiting

Vision: vision abnormal

Gastrointestinal: nausea, diarrhea, abdominal pain, constipation

Body as a Whole - General Disorders: fatigue

Urinary System: renal calculus

Skin and Appendages: rash

SYMPTOMS AND TREATMENT OF OVERDOSAGE

In acute TOPAMAX topiramate overdose, if the ingestion is recent, the stomach should be emptied immediately by lavage or by induction of emesis. Activated charcoal has not been shown to adsorb topiramate *in vitro*. Therefore, its use in overdose is not recommended. Treatment should be appropriately supportive.

Hemodialysis is an effective means of removing topiramate from the body. However, in the few cases of acute overdose reported, including doses of over 20 g in one individual, hemodialysis has not been necessary.

DOSAGE AND ADMINISTRATION

General: TOPAMAX Tablets or Sprinkle Capsules can be taken without regard to meals. Tablets should not be broken. TOPAMAX Sprinkle Capsules may be swallowed whole or may be administered by carefully opening the capsule and sprinkling the entire contents on a small amount (teaspoon) of soft food. This drug/food mixture should be swallowed immediately and not chewed. It should not be stored for future use. The sprinkle formulation is provided for those patients who cannot swallow tablets, e.g. pediatric and the elderly.

Adults (Age 17 years and older): It is recommended that TOPAMAX topiramate as adjunctive therapy be initiated at 50 mg/day, followed by titration as needed and tolerated to an effective dose. At weekly intervals, the dose may be increased by 50 mg/day and taken in two divided doses. Some patients may benefit from lower initial doses, e.g. 25 mg and/or a slower titration schedule. Some patients may achieve efficacy with once-daily dosing.

The recommended total daily maintenance dose is 200 mg-400 mg/day in two divided doses. Doses above 400 mg/day have not been shown to improve responses and have been associated with a greater incidence of adverse events. The maximum recommended dose is 800 mg/day. Daily doses above 1,600 mg have not been studied.

Children (Ages 2-16 years): It is recommended that TOPAMAX topiramate as adjunctive therapy be initiated at 25 mg (or less, based on a range of 1 to 3 mg/kg/day) nightly for the first week followed by titration as needed and tolerated to an effective dose. The dosage should then be increased at 1- or 2-week intervals by increments of 1 to 3 mg/kg/day (administered in two divided doses). Some patients may benefit from lower initial doses and/or a slower titration schedule.

The recommended total daily maintenance dose is approximately 5 to 9 mg/kg/day in two divided doses. Daily doses up to 30 mg/kg/day have been studied and were generally well tolerated.

Geriatrics

See **PRECAUTIONS** section.

Patients with Renal Impairment

In renally impaired subjects (creatinine clearance less than 70 mL/min/1.73m²), one-half of the usual adult dose is recommended. Such patients will require a longer time to reach steady-state at each dose.

Patients Undergoing Hemodialysis

Topiramate is cleared by hemodialysis at a rate that is 4 to 6 times greater than a normal individual. Accordingly, a prolonged period of dialysis may cause topiramate concentration to fall below that required to maintain an antiseizure effect. To avoid rapid drops in topiramate plasma concentration during hemodialysis a supplemental dose of topiramate may be required. The actual adjustment should take into account 1) the duration of dialysis, 2) the clearance rate of the dialysis system being used, and 3) the effective renal clearance of topiramate in the patient being dialyzed.

Patients with Hepatic Disease

In hepatically impaired patients, topiramate plasma concentrations are increased approximately 30%. This moderate increase is not considered to warrant adjustment of the topiramate dosing regimen. Initiate topiramate therapy with the same dose and regimen as for patients with normal hepatic function. The dose titration in these patients should be guided by clinical outcome, i.e. seizure control, and avoidance of adverse effects. Such patients will require a longer time to reach steady-state at each dose.

AVAILABILITY OF DOSAGE FORMS

TOPAMAX topiramate is available as embossed tablets in the following strengths as described below:

- 25 mg: white, round, coated tablets containing 25 mg topiramate.
- 100 mg: yellow, round, coated tablets containing 100 mg topiramate.
- 200 mg: salmon-colored, round, coated tablets containing 200 mg topiramate.

TOPAMAX topiramate Sprinkle Capsules contain small white to off-white spheres. The gelatin capsules are white and clear. They are marked as follows:

- 15 mg: "TOP" and "15 mg" on the side.
- 25 mg: "TOP" and "25 mg" on the side.

Supplied: Bottles of 60 tablets with desiccant.
Bottles of 60 capsules without desiccant.

TOPAMAX is a Schedule F Drug.

Product Monograph available to physicians and pharmacists upon request.



JANSSEN-ORTHO Inc.

Janssen-Ortho Inc., Toronto, Ontario M3C 1L9

Date of Issuance: April 2000
TXPD01013A

© 2000 JANSSEN-ORTHO Inc.
* All trademark rights used under license.



Nervous System

Dizziness	28%	35%
Hypertonia	24%	26%
Depression	24%	25%
Anxiety	13%	15%
Nervousness	5%	8%
Somnolence	3%	6%
Confusion	2%	4%
Speech disorder	1%	3%
Convulsion	0%	2%
Hyperkinesia	0%	2%
Amnesia	0%	2%

Respiratory System

Sinusitis	26%	36%
Dyspnea*	2%	8%
Laryngitis	2%	6%

Skin and Appendages

Sweating*	11%	23%
Alopecia	2%	4%

Special Senses

Conjunctivitis	10%	12%
Abnormal vision	4%	7%

Urogenital System

Dysmenorrhea	11%	18%
Menstrual disorder*	8%	17%
Metrorrhagia	8%	15%
Cystitis	4%	8%
Breast pain	3%	7%
Menorrhagia	3%	6%
Urinary urgency	2%	4%
Fibrocystic breast	1%	3%
Breast neoplasm	0%	2%

* significantly associated with BETASERON treatment (p<0.05)

It should be noted that the figures cited in Table 1 cannot be used to predict the incidence of side effects in the course of usual medical practice where patient characteristics and other factors differ from those that prevailed in the clinical trials. The cited figures do provide the prescribing physician with some basis for estimating the relative contribution of drug and nondrug factors to the side effect incidence rate in the population studied.

2. Secondary-progressive MS: The incidence of adverse events that occurred in at least 2% of patients treated with 8 MIU BETASERON or placebo for up to three years, or where an adverse event was reported at a frequency at least 2% higher with BETASERON than that observed for placebo-treated patients in the secondary-progressive study, is presented in Table 2. Adverse events significantly associated with BETASERON compared to placebo (p<0.05) are also indicated in Table 2.

Table 2: Incidence of Adverse Events ≥ 2% or > 2% Difference (BETASERON vs. Placebo) in the Secondary Progressive MS Study

Adverse Event	Placebo n=358	0.25 mg (8 MIU) n=360
Body as a Whole		
Asthenia	58%	63%
Flu syndrome*	40%	61%
Pain	25%	31%
Fever*	13%	40%
Back pain	24%	26%
Accidental injury	17%	14%
Chills*	7%	23%
Pain In Extremity	12%	14%
Infection	11%	13%
Abdominal pain*	6%	11%
Malaise	5%	8%
Neck pain	6%	5%
Abscess*	2%	4%
Laboratory test abnormal	1%	3%
Allergic reaction	3%	2%
Chills and fever*	0%	3%
Thorax pain	2%	1%

Cardiovascular System

Vasodilatation	4%	6%
Peripheral vascular disorder	5%	5%
Chest pain	4%	5%
Migraine	3%	4%
Hypotension	4%	2%
Hypertension*	2%	4%
Palpitation	3%	2%
Syncope	3%	2%
Hemorrhage	2%	2%
Tachycardia	1%	2%

Digestive System

Nausea	13%	13%
Constipation	12%	12%
Diarrhea	10%	7%
Gastroenteritis	5%	6%
Vomiting	6%	4%
Dysphagia	5%	4%
Gastrointestinal disorder	5%	4%
Tooth disorder	4%	4%
Dyspepsia	4%	4%
Anorexia	2%	4%
Fecal incontinence	3%	2%
Liver function test abnormal	1%	3%
Gastritis	2%	2%
Flatulence	1%	3%
Sore throat	1%	2%
Colitis	2%	0%
Gastrointestinal pain	0%	2%
Gingivitis	0%	2%

Hemic and Lymphatic System

Leukopenia*	5%	10%
Anemia	5%	2%
Eosinophilia	2%	1%
Lymphadenopathy	1%	3%

Injection Site

Injection site reaction*	10%	46%
Injection site inflammation*	4%	48%
Injection site pain	5%	9%
Injection site necrosis*	0%	5%
Injection site hemorrhage	2%	2%

Metabolic and Nutritional Disorders

Peripheral edema	7%	7%
Weight loss	3%	2%
SGPT increased	2%	2%
Hypercholesterolemia	2%	1%

Musculoskeletal System

Myasthenia	40%	39%
Arthralgia	20%	20%
Myalgia*	9%	23%
Bone fracture (not spontaneous)	5%	3%
Muscle cramps	3%	3%
Spontaneous bone fracture	3%	3%
Arthritis	1%	2%
Joint disorder	1%	2%

Nervous System

Headache	41%	47%
Neuropathy	41%	38%
Paresthesia	39%	35%
Hypertonia*	31%	41%
Abnormal gait	34%	34%
Depression	31%	27%
Ataxia	23%	19%
Dizziness	14%	14%
Incoordination	13%	11%
Insomnia	8%	12%
Vertigo	12%	8%
Emotional lability	11%	8%
Paralysis	10%	8%
Somnolence	8%	8%
Tremor	9%	6%
Sweating increased	6%	6%
Neuralgia	7%	5%
Movement disorder	6%	5%
Sleep disorder	5%	6%
Anxiety	5%	6%
Hypesthesia	4%	6%
Nervousness	3%	4%

Speech disorder	5%	2%
Dysarthria	4%	2%
Spastic paralysis	1%	3%
Convulsion	2%	2%
Hyperesthesia	2%	2%
Amnesia	3%	1%
Dry mouth	2%	1%
Hemiplegia	2%	1%
Thinking abnormal	2%	1%
Myoclonus	2%	0%

Respiratory System

Rhinitis	32%	38%
Pharyngitis	20%	16%
Bronchitis	12%	9%
Cough increased	10%	5%
Sinusitis	6%	6%
Pneumonia	5%	5%
Dyspnea	2%	3%
Upper respiratory tract infection	2%	3%
Asthma	2%	1%
Voice alteration	2%	1%

Skin and Appendages

Rash*	12%	20%
Pruritus	6%	6%
Skin disorder	4%	4%
Eczema	4%	2%
Herpes simplex	2%	3%
Alopecia	2%	2%
Acne	2%	2%
Dry skin	3%	1%
Subcutaneous hematoma	3%	1%
Breast pain	2%	1%
Herpes zoster	2%	1%
Seborrhea	2%	1%

Special Senses

Abnormal vision	35%	11%
Amblyopia	10%	7%
Diplopia	9%	7%
Eye pain	5%	4%
Otitis media	3%	2%
Conjunctivitis	3%	2%
Eye disorder	2%	2%
Deafness	2%	1%
Optic neuritis	2%	2%
Ear disorder	2%	1%
Tinnitus	2%	1%

Urogenital System

Urinary tract infection	25%	22%
Urinary incontinence	15%	8%
Urinary tract disorder	10%	7%
Cystitis	9%	7%
Urinary urgency	7%	8%
Menstrual disorder	13%	9%
Increased urinary frequency	5%	6%
Metrorrhagia	8%	12%
Urinary retention	6%	4%
Vaginitis	4%	3%
Armenorrhoea	4%	3%
Dysuria	2%	2%
Impotence	4%	7%
Menopause	4%	2%
Menorrhagia	4%	2%
Nocturia	1%	2%
Vaginal moniliasis	2%	2%
Kidney pain	2%	0%
Pyelonephritis	0%	2%
Prostatic disorder	1%	2%

*significantly associated with BETASERON treatment (p<0.05)

Seventy-four (74) patients discontinued treatment due to adverse events (23 on placebo and 51 on BETASERON) injection site reactions were significantly associated with early termination of treatment in the BETASERON group compared to placebo (p<0.05). The highest frequency of adverse events leading to discontinuation involved the nervous system, of which depression (7 on placebo and 11 on BETASERON) was the most common. Significantly more patients on active therapy (14.4% vs. 4.7% on placebo) had elevated ALT (SGPT) values (>5 times

baseline value). Elevations were also observed in AST (SGOT) and gamma-GT values in the BETASERON group throughout the study. In the BETASERON group, most ALT (SGPT) abnormalities resolved spontaneously with continued treatment whereas some resolved upon dose reduction or temporary discontinuation of treatment.

Lymphopenia (<1500/mm³) was observed in 90.9% of BETASERON patients compared to 74.3% of placebo patients and neutropenia (<1400/mm³) was noted in 18.0% BETASERON and 5.1% placebo patients.

DOSE AND ADMINISTRATION
FOR SUBCUTANEOUS USE ONLY

BETASERON (interferon beta-1b) should only be prescribed by (or following consultation with) clinicians who are experienced in the diagnosis and management of multiple sclerosis.

The recommended dose of BETASERON for both relapsing-remitting and secondary-progressive MS patients is 0.25 mg (8 MIU) injected subcutaneously every other day. Limited data regarding the activity of a lower dose in relapsing-remitting MS patients are presented above (see ACTION AND CLINICAL PHARMACOLOGY, Clinical Trials).

PHARMACOLOGY, Clinical Trials:

In the secondary-progressive MS study patients initiated treatment with half the dose (4 MIU s.c. every other day) for a period of 2 weeks prior to escalating to the recommended dose of 8 MIU (s.o. every other day).

Efficacy of treatment for longer than 2 years has not been substantially demonstrated in relapsing-remitting multiple sclerosis. For secondary progressive multiple sclerosis, safety and efficacy data beyond 3 years are not available.

To reconstitute lyophilized BETASERON for injection, use a sterile syringe and needle to inject 1.2 mL of the diluent supplied. Sodium Chloride, 0.54% Solution, into the BETASERON vial. Gently swirl the vial of BETASERON to dissolve the drug completely; do not shake. Inspect the reconstituted product visually and discard the product before use if it contains particulate matter or is discolored. After reconstitution with accompanying diluent, each mL of solution contains 0.25 mg (8 MIU) interferon beta-1b, 13 mg Albumin Human USP, and 13 mg Mannitol USP.

Withdraw 1 mL of reconstituted solution from the vial into a sterile syringe fitted with a 27-gauge 1/2-inch needle and inject the solution subcutaneously. Sites for self-injection include abdomen, buttocks and thighs. A vial is suitable for single use only; unused portions should be discarded (See BETASERON® [interferon beta-1b] INFORMATION FOR THE PATIENT section for SELF-INJECTION PROCEDURE.)

AVAILABILITY OF DOSAGE FORMS

BETASERON (interferon beta-1b) is presented in single-use vials of lyophilized powder containing 0.3 mg (9.6 MIU) interferon beta-1b, 15 mg Albumin Human USP, and 15 mg Mannitol USP. BETASERON is supplied in cartons containing 15 vials of medication and 15 vials of diluent (2 mL of Sodium Chloride 0.54% solution, per vial).

Product Monograph available upon request
B10204E5

REFERENCES:

1. Data on file, Berlex Canada Inc., 1999.
2. Product Monograph of BETASERON® (interferon beta-1b), Berlex Canada, June 1999.
3. The FNB Multiple Sclerosis Study Group and the University of British Columbia MS/MRI Analysis Group. Interferon beta-1b in the treatment of multiple sclerosis. Final outcome of the randomized, controlled trial. *Neurology* 1995; 45:1227-1235.

2260 32nd Avenue, Lachine, Québec H8T 3H4



NEW
Keppra®
levetiracetam
CONNECTING EXCELLENT PROFILES IN
EFFICACY AND TOLERABILITY

PRESCRIBING INFORMATION

Tablets of 250 mg, 500 mg, and 750 mg
Therapeutic classification: Antiepileptic

ACTIONS AND CLINICAL PHARMACOLOGY

Mechanism of Action

Levetiracetam is a drug of the pyrrolidine class chemically unrelated to existing antiepileptic drugs (AEDs). Levetiracetam exhibits anti-seizure and antiepileptogenic activity in several models of chronic epilepsy in both mice and rats, while being devoid of anticonvulsant activity in the classical screening models of acute seizures. The mechanism of action of levetiracetam has not yet been fully established, however, it appears to be unlike that of the commonly used AEDs. *In vitro* studies show that levetiracetam, at concentrations of up to 10 μM did not result in significant ligand displacement at known receptor sites such as benzodiazepine, GABA (gamma-aminobutyric acid), glycine, NMDA (N-methyl-D-aspartate), reuptake sites or second messenger systems. Furthermore, levetiracetam does not modulate neuronal voltage-gated sodium and T-type calcium currents and does not induce conventional facilitation of the GABAergic system.

Pharmacokinetics

Summary: Single- and multiple-dose pharmacokinetics of levetiracetam have included healthy volunteers, adult and pediatric patients with epilepsy, elderly subjects, and subjects with renal and hepatic impairment. Results of these studies indicate that levetiracetam is rapidly and almost completely absorbed after oral administration. The pharmacokinetic profile is linear with low intra- and inter-subject variability. There is no modification of the clearance after repeated administration. Food does not affect the extent of absorption of levetiracetam, although the rate is decreased. Levetiracetam is not protein-bound (< 10% bound) and its volume of distribution is close to the volume of intracellular and extracellular water. Sixty-six percent (66%) of the dose is renally excreted unchanged. The major metabolic pathway of levetiracetam (24% of the dose) is an enzymatic hydrolysis of the acetamide group. It is not liver cytochrome P450 dependent. The metabolites have no known pharmacodynamic activity and are renally excreted. Plasma half-life of levetiracetam across studies is 6-8 hours. Plasma half-life is increased in subjects with renal impairment, and in the elderly primarily due to impaired renal clearance.

Based on its pharmacokinetic characteristics, levetiracetam is unlikely to produce or to be subject to metabolic interactions. The pharmacokinetic profile is comparable in healthy volunteers and in patients with epilepsy.

Due to its complete and linear absorption, plasma levels can be predicted from the oral dose of levetiracetam expressed as mg/kg bodyweight. Therefore, there is no need for plasma level monitoring of levetiracetam.

Human Pharmacology

Pharmacokinetics: The pharmacokinetics of levetiracetam have been characterized in single and multiple dose PK studies, with doses up to 5000 mg; these studies included healthy volunteers (n = 98), patients with epilepsy (n = 58 adult patients and n = 24 pediatric patients), elderly subjects (n = 16) and subjects with renal and hepatic impairment (n = 36 and 16, respectively).

Absorption and Distribution: Levetiracetam is rapidly and almost completely absorbed after oral administration. The oral bioavailability of levetiracetam tablets is 100%. Plasma peak concentrations (C_{max}) are achieved at 1.3 hours after dosing. The extent of absorption is independent of both dose and the presence of food, but the latter delays T_{max} by 1.5 hours and decreases C_{max} by 20%. The pharmacokinetics of levetiracetam are linear over the dose range of 500 – 5000 mg. Steady-state is achieved after two days of a twice daily administration schedule. Mean peak concentrations (C_{max}) are 31 and 43 μg/mL, respectively, following a single 1000 mg dose, and a repeated 1000 mg twice daily dose.

Neither levetiracetam nor its primary metabolite is significantly bound to plasma proteins (< 10%). The volume of distribution of levetiracetam is approximately 0.5 to 0.7 L/kg, a value that is close to the total body water volume. No tissue distribution data for humans are available.

Metabolism: Levetiracetam is not extensively metabolized in humans. The major metabolic pathway is the enzymatic hydrolysis of the acetamide group, which produces the pharmacologically inactive carboxylic acid metabolite, ucb L057 (24% of dose). The production of this metabolite is not dependent on any liver cytochrome P450 isoenzymes and is mediated by serine esterase(s) in various tissues, including blood cells. Two minor metabolites were identified as the product of hydroxylation of the 2-oxo-pyrrolidine ring (2% of dose) and opening of the 2-oxo-pyrrolidine ring in position 5 (1% of dose). There is no evidence for enantiomeric interconversion of levetiracetam or its major metabolite.

Elimination: Levetiracetam plasma half-life in adults is 7 ± 1 hours and was unaffected by dose, route of administration or repeated administration. Levetiracetam is eliminated from the systemic circulation by renal excretion as unchanged drug, which represents 66% of administered dose. The total body clearance is 0.96 mL/min/kg and the renal clearance is 0.6 mL/min/kg. Approximately 93% of the dose was excreted within 48 hours. The mechanism of excretion is glomerular filtration with subsequent partial tubular reabsorption. The primary metabolite, ucb L057, is excreted by glomerular filtration and active tubular secretion with a renal clearance of 4 mL/min/kg. Levetiracetam elimination is correlated to creatinine clearance and clearance is thus reduced in patients with impaired renal function (See PRECAUTIONS and DOSAGE AND ADMINISTRATION).

Special Populations: Elderly: Pharmacokinetics of levetiracetam were evaluated in 16 elderly patients, ranging in age from 61-88 years, with 11 of the 16 patients aged 75 years of age or over with creatinine clearance ranging from 30 to 74 mL/min. Following oral administration of 500 mg bid for 10 days, total body clearance decreased by 38% and the half-life was increased about 40% (10 to 11 hours) when compared to healthy adults. This is most likely due to the decrease in renal function in these subjects. **Pediatrics (6 to 12 years):** Pharmacokinetics of levetiracetam were evaluated in 24 pediatric patients (age 6-12 years) after a single dose. The apparent clearance of levetiracetam adjusted to body weight was approximately 40% higher than in epileptic adults. **Gender:** Levetiracetam C_{max} and AUC were 20% higher in women (n = 11) compared to men (n = 12). However, clearances adjusted for body weight were comparable. **Race:** Formal pharmacokinetic studies of the effects of race have not been conducted. Because levetiracetam is primarily renally excreted and there are no known important racial differences in creatinine clearance, significant pharmacokinetic differences due to race are not expected.

Renal Impairment: Single dose pharmacokinetics were performed in 20 subjects with renal impairment (n = 7 mild/CL_{CR} of 50-79 mL/min; n = 8 moderate/CL_{CR} of 30-49 mL/min; n = 5 severe/CL_{CR} < 30 mL/min), and n = 11 matching healthy volunteers. Clearance of levetiracetam is correlated with creatinine clearance and levetiracetam pharmacokinetics following repeat administration were well predicted from single dose data. The apparent body clearance of the parent drug levetiracetam is reduced in patients with impaired renal function by approximately 40% in the mild group, 50% in the moderate group, and 60% in the severe renal impairment group. For the primary metabolite ucb L057, the decrease in clearance values from baseline was greater than that seen for the parent drug in all subject groups.

In anuric (end stage renal disease) patients, the apparent body clearance was approximately 30% compared to that of normal subjects. Approximately 50% of the pool of levetiracetam in the body is removed during a standard 4-hour hemodialysis procedure. Dosage should be reduced in patients with impaired renal function receiving levetiracetam, and supplemental doses should be given to patients after dialysis (see PRECAUTIONS and DOSAGE AND ADMINISTRATION).

Hepatic Impairment: A single-dose pharmacokinetic study was performed in 16 subjects with hepatic impairment (n = 5 mild/Child-Pugh Grade A; n = 6 moderate/Grade B; n = 5 severe/Grade C vs 5 healthy controls). For the mild and moderate subgroups neither mean nor individual pharmacokinetic values were clinically different from those of controls. In patients with severe hepatic impairment, mean apparent body clearance was 50% that of normal subjects, with decreased renal clearance accounting for most of the decrease. Patients with severe hepatic impairment thus require a reduced dosage of Keppra® (See PRECAUTIONS and DOSAGE AND ADMINISTRATION).

INDICATIONS AND CLINICAL USE

Keppra® (levetiracetam) is indicated as adjunctive therapy in the management of patients with epilepsy who are not satisfactorily controlled by conventional therapy.

CONTRAINDICATIONS

This product should not be administered to patients who have previously exhibited hypersensitivity to levetiracetam or any of the inactive ingredients in Keppra® (levetiracetam) tablets.

WARNINGS

Central Nervous System Adverse Events

Keppra® (levetiracetam) use is associated with the occurrence of central nervous system (CNS) adverse events; the most significant of these can be classified into the following categories: 1) somnolence and fatigue, 2) behavioral/psychiatric symptoms and 3) coordination difficulties.

There was no clear dose response relationship for any of the three categories of CNS adverse events, within the recommended dose range of up to 3000 mg/day. Somnolence/asthenia and coordination difficulties occurred most frequently within the first four weeks of treatment and usually resolved while patients remained on treatment. In the case of behavioral/psychiatric symptoms (including such adverse events as aggression, agitation, anger, anxiety, emotional lability, hostility, irritability), approximately half of the patients reported these events within the first four weeks, with the remaining events occurring throughout the duration of the trials. See also PRECAUTIONS, Central Nervous System Adverse Events.

Withdrawal of Anti-Epileptic Drugs

As with all antiepileptic drugs, Keppra® should be withdrawn gradually to minimize the potential of increased seizure frequency.

PRECAUTIONS

General

Hematological Abnormalities: Minor but statistically significant decreases compared to placebo were seen in total mean RBC count, mean hemoglobin, and mean hematocrit in Keppra®-treated patients in controlled trials. For hemoglobin values, the percentage of Keppra® or placebo treated patients with possibly clinically significant abnormalities were less than 0.5% each. For hematocrit values, a total of 5.1% of Keppra® treated versus 3.2% of placebo patients had at least one possibly significant decrease in hematocrit (≤ 37% in males and 32% in females).

For white blood cells (WBC), 2.9% of treated versus 2.3% of placebo patients had at least one possibly clinically significant decrease in WBC count (≤ 2.8 × 10⁹/L), while 2.6% of treated vs. 1.7% of placebo patients had at least one possibly significant decrease in neutrophil count (≤ 1.0 × 10⁹/L). Of the Keppra®-treated patients with a low neutrophil count, all but one rose towards or reached baseline with continued treatment. No patient was discontinued secondary to low neutrophil counts.

Central Nervous System Adverse Events (See WARNINGS): Keppra® (levetiracetam) use is associated with the occurrence of central nervous system (CNS) adverse events; the most significant of these can be classified into the following categories: 1) somnolence and fatigue, 2) behavioral/psychiatric symptoms and 3) coordination difficulties.

The following CNS adverse events were observed in controlled clinical trials.

Table 1:
Total Combined Incidence Rate for Each of the Three Categories of CNS Adverse Events in Placebo-controlled Add-on Clinical Trials.

Category of CNS adverse event	Keppra® + AED therapy (n = 672)	Placebo + AED therapy (n = 351)
Somnolence and fatigue		
Somnolence	15%	10%
Asthenia	14%	10%
Behavioral/psychiatric symptoms		
Nonpsychotic ¹	14%	6%
Psychotic ²	1%	0%
Coordination difficulties ³	3%	2%

¹Reflects Keppra® doses of 1000 mg, 2000 mg, 3000 mg, and 4000 mg per day.

²Non-psychotic behavioral/psychiatric symptoms¹ encompasses the following terms: agitation, antisocial reaction, anxiety, apathy, depersonalization, depression, emotional lability, euphoria, hostility, nervousness, neurosis, personality disorder and suicide attempt.

³Psychotic behavioral/psychiatric symptoms² encompasses the following terms: hallucinations, paranoid reaction, psychosis and psychotic depression.

⁴Coordination difficulties³ encompasses the following terms: ataxia, abnormal gait, incoordination.

See ADVERSE EVENTS, Table 2, for incidence rate of individual AEs contained within the categories.

Behavioral/psychiatric symptoms (including agitation, emotional lability, hostility, anxiety, etc.) have been reported approximately equally in patients with and without a psychiatric history.

There was no clear dose response relationship for any of the three categories of CNS adverse events, within the recommended dose range of up to 3000 mg/day. In a controlled study including a dose of 4000 mg, administered without titration, the incidence rate of somnolence during the first four weeks of treatment for patients receiving the high dose was 42%, compared to 21% for patients receiving 2000 mg/day.

Special Populations

Patients with Renal Impairment: Renal excretion of unchanged drug accounts for approximately 66% of administered levetiracetam dose. Consistent with this, pharmacokinetic studies in renally-impaired patients indicate that apparent clearance is significantly reduced in subjects with renal impairment (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations).

In patients with renal impairment Keppra® dosage should be appropriately reduced. Patients with end stage renal disease, i.e. those undergoing dialysis, should be given supplemental doses after dialysis (See DOSAGE AND ADMINISTRATION).

Pregnancy and Nursing: There are no adequate and well-controlled studies on the use of Keppra® in pregnant women. Levetiracetam and/or its metabolites cross the placental barrier in animal species. In reproductive toxicity studies in rats and rabbits, levetiracetam induced developmental toxicity at exposure levels similar to or greater than the human exposure. There was evidence of increased skeletal variations/minor anomalies, retarded growth, embryonic death, and increased pup mortality. In the rat, fetal abnormalities occurred in the absence of overt maternal toxicity. The systemic exposure at the observed no effect level in the rabbit was about 4 to 5 times the human exposure. The potential risk for humans is unknown. Keppra® should not be used during pregnancy unless potential benefits to mother and fetus are considered to outweigh potential risks to both. Discontinuation of antiepileptic treatments may result in disease worsening, which can be harmful to the mother and the fetus.

Pregnancy Exposure Registry: To facilitate monitoring of fetal outcomes of pregnant women exposed to Keppra®, physicians should encourage patients to register, before fetal outcome is known (e.g., ultrasound, results of amniocentesis, etc.), in the Antiepileptic Drug Pregnancy Registry by calling (888) 233-2334 (toll free).

Nursing Mothers: Levetiracetam is excreted in breast milk. Therefore, there is a potential for serious adverse reactions from Keppra® in nursing infants. Recommendations regarding nursing and epilepsy medication should take into account the importance of the drug to the mother, and the as yet uncharacterized risks to the infant. Typically, recommendations are made in the context of the necessary prior risk-benefit judgement, regarding pregnancy and epilepsy medication.

Use in Pediatric Patients: Safety and efficacy in patients below the age of 18 have not been established.

Use in the Elderly: Renal function can be decreased in the elderly and levetiracetam is known to be substantially excreted by the kidney, the risk of adverse reactions to the drug may be greater in patients with impaired renal function. A pharmacokinetic study in 16 elderly subjects (age 61-88 years) showed a decrease in clearance by about 40% with oral administration of both single dose and 10 days of multiple twice-daily dosing. This decrease is most likely due to the expected decrease in renal function in these elderly subjects. Care should therefore be taken in dose selection for elderly patients, and it may be useful to monitor renal function.

There were insufficient numbers of elderly patients in controlled trials of epilepsy to adequately assess the efficacy or safety of Keppra® in these patients. Nine of 672 patients treated with Keppra® were 65 or over.

Drug Interactions

In Vitro Studies on Metabolic Interaction Potential *In vitro*, levetiracetam and its primary metabolite have been shown not to inhibit the major human liver cytochrome P450 isoforms (CYP3A4, 2A6, 2C8/9/10, 2C19, 2D6, 2E1 and 1A2), glucuronyl transferase (paracetamol UGT, i.e. UGT1A6, ethinyl estradiol UGT, i.e.UGT1A1, and p-nitrophenol UGT, i.e.UGT [p]16.2) and epoxide hydrolase activities. In addition, levetiracetam does not affect the *in vitro* glucuronidation of valproic acid. In human hepatocytes in culture, levetiracetam did not cause enzyme induction.

Levetiracetam circulates largely unbound (< 10% bound) to plasma proteins; therefore clinically significant interactions with other drugs through competition for protein binding sites are unlikely.

Thus *in-vitro* data, in combination with the pharmacokinetic characteristics of the drug, indicate that Keppra® is unlikely to produce, or be subject to, pharmacokinetic interactions.

Clinical Pharmacokinetic Data

Other Antiepileptic Drugs (AEDs): Potential drug interactions between Keppra® and other AEDs (phenytoin, carbamazepine, valproic acid, phenobarbital, lamotrigine, gabapentin and primidone) were assessed by evaluating the serum concentrations of levetiracetam and these AEDs during placebo-controlled clinical studies. These data suggest that levetiracetam may not significantly influence the plasma concentrations of these other AEDs, and that the other AEDs may not significantly influence the plasma concentrations of levetiracetam.

For two of these AEDs — phenytoin and valproate — formal pharmacokinetic interaction studies with Keppra® were performed. Keppra® was co-administered with either phenytoin or valproate at doses of 3000 mg/day and 1000 mg/day respectively. No clinically significant interactions were observed.

Other Drug Interactions

Oral Contraceptives: A pharmacokinetic clinical interaction study has been performed in healthy subjects between the oral contraceptive containing 0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel, and the lowest therapeutic dose of Keppra® (500 mg bid). No clinically significant pharmacokinetic interactions were observed.

However, pharmacokinetic interaction studies using Keppra® as adjunctive therapy and covering the recommended dosage range, have not been conducted. Therefore, physicians should advise their female patients to be alert to any irregular vaginal bleeding or spotting, and to immediately report to them any occurrences.

Digoxin: Keppra® (1000 mg bid) did not influence the pharmacokinetics and pharmacodynamics (ECG) of digoxin given as a 0.25 mg dose every day. Coadministration of digoxin did not influence the pharmacokinetics of levetiracetam.

Warfarin: Keppra® (1000 mg bid) did not influence the pharmacokinetics of R and S warfarin (2.5 mg, 5 mg, or 7.5 mg daily). Prothrombin time was not affected by levetiracetam. Coadministration of warfarin did not affect the pharmacokinetics of levetiracetam.

Probenecid: Probenecid, a renal tubular secretion blocking agent, administered at a dose of 500 mg four times a day, did not change the pharmacokinetics of levetiracetam 1000 mg bid. C_{0-24h} of the metabolite, ucb L057, was approximately doubled in the presence of probenecid and the renal clearance of the metabolite ucb L057 was decreased by 60%; this alteration is likely related to competitive inhibition of tubular secretion of ucb L057. The effect of Keppra® on probenecid was not studied.

ADVERSE EVENTS

Commonly Observed

In well-controlled clinical studies, the most frequently reported adverse events associated with the use of Keppra® in combination with other AEDs, not seen at an equivalent frequency among placebo-treated patients, were somnolence, asthenia, dizziness and infection. Of the most frequently reported adverse events, asthenia, somnolence and dizziness appeared to occur predominantly during the first four weeks of treatment with Keppra®.

Incidence of AEs in Controlled Clinical Trials

Table 2:
Incidence (%) of Treatment-emergent Adverse Events in Placebo-controlled, Add-on Studies by Body System. (Adverse Events Occurred in at least 1% of Keppra®-treated Patients and Occurred More Frequently than Placebo-treated Patients.)
(Studies N051, N052, N132 and N138)

Body system/ adverse event	Keppra® + AED therapy (n = 672) (%)	Placebo + AED therapy (n = 351) (%)
Body as a whole		
Asthenia	14	10
Infection*	13	7
Digestive system		
Tooth disorders	2	1
Hemic and lymphatic system		
Ecchymosis	2	1
Nervous system		
Amnesia	2	0
Anxiety	2	1
Ataxia	3	1
Depression	4	2
Dizziness	9	4
Emotional lability	2	0
Hostility	2	1
Nervousness	4	2
Personality disorders	1	0
Somnolence	15	10
Thinking abnormal	2	1
Vertigo	3	1
Respiratory system		
Pharyngitis	6	4
Rhinitis	4	3
Sinusitis	2	1

* In levetiracetam-treated patients, the majority of "infection" events (93%) were coded to reported terms of "common cold" or "infection upper respiratory".

Additional Events Observed in Placebo Controlled Trials

Lack of Dose-related Incidence within Therapeutic Range: Based on the data from the controlled clinical trials, there was no evidence of dose relationship within the recommended dose range of 1000 to 3000 mg/day.

Discontinuation or Dose Reduction in Well-controlled Clinical Studies: In well-controlled clinical studies, 14.3% of patients receiving Keppra® and 11.7% receiving placebo either discontinued or had a dose reduction as a result of an adverse event. The adverse events most commonly associated (>1%) with discontinuation or dose reduction in either treatment group are presented in Table 3.

Table 3:
Adverse Events Most Commonly Associated with Discontinuation or Dose Reduction in Placebo-controlled Studies in Patients with Epilepsy

	Keppra® (n = 672)	Placebo (n = 351)
Asthenia	9 (1.3%)	3 (0.9%)
Headache	8 (1.2%)	2 (0.6%)
Convulsion	16 (2.4%)	10 (2.8%)
Dizziness	11 (1.6%)	0
Somnolence	31 (4.6%)	6 (1.7%)
Rash	0	5 (1.4%)

The overall adverse experience profile of Keppra® was similar between females and males. There are insufficient data to support a statement regarding the distribution of adverse experience reports by age and race.

Post-marketing Experience

In post-marketing experience, nervous system and psychiatric disorders have most frequently been reported. In addition to adverse reactions during clinical studies, and listed above, the following adverse reactions have been reported in post-marketing experience. Data are insufficient to support an estimate of their incidence in the population to be treated.

Blood and lymphatic disorders: leukopenia, neutropenia, pancytopenia, thrombocytopenia.

SYMPTOMS AND TREATMENT OF OVERDOSE

Symptoms

The highest reported Keppra® overdose is approximately 10 times the therapeutic dose. In the majority of overdose cases, multiple drugs were involved. Somnolence, agitation, aggression, depressed level of consciousness, respiratory depression, and coma were observed with Keppra® overdoses. The minimal lethal oral dose in rodents is at least 233 times the maximum clinically studied dose.

Treatment

There is no antidote for overdose with Keppra®; treatment is symptomatic and may include hemodialysis. If indicated, elimination of unabsorbed drug should be attempted by emesis or gastric lavage; usual precautions should be observed to maintain airway. General supportive care of the patient is indicated including monitoring of vital signs and observation of the clinical status of the patient.

Standard hemodialysis procedures result in significant removal of levetiracetam (approximately 50% in 4 hours) and should be considered in cases of overdose. Although hemodialysis has not been performed in the few known cases of overdose, it may be indicated by the patient's clinical state or in patients with significant renal impairment.

DOSAGE AND ADMINISTRATION

General

Renal excretion of unchanged drug accounts for approximately 66% of administered levetiracetam dose. Consistent with this, reduced doses are recommended for patients with renal impairment. Keppra® is given orally with or without food.

Adults

Treatment should be initiated at a dose of 1000 mg/day, given as twice daily dosing (500 mg bid). Depending on clinical response and tolerability, the daily dose may be increased every two weeks by increments of 1000 mg, to a maximum recommended daily dose of 3000 mg.

In clinical trials, daily doses of 1000 mg, 2000 mg, and 3000 mg, given as twice a day dosing, were shown to be effective. Although there was a tendency toward greater response rate with higher dose, a consistent statistically significant increase in response with increased dose has not been shown. There are limited safety data from controlled clinical trials at doses higher than 3000 mg/day (approximately 40 patients), therefore these doses are not recommended.

Patients with Impaired Renal Function

Keppra® dosage should be reduced in patients with impaired renal function (see Table 4 below). Patients with end stage renal disease should receive supplemental doses following dialysis. To use this dosing table, an estimate of the patient's CL_{Cr} in mL/min is needed. CL_{Cr} in mL/min may be estimated from serum creatinine (mg/dL) determination using the following formula:

$$CL_{Cr} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dL)}} \times 0.85 \text{ (for female patients)}$$

Table 4:
Dosing Adjustment for Patients with Impaired Renal Function

Group	Creatinine clearance (mL/min)	Dosage and frequency
Normal	≥ 80	500 to 1500 mg twice daily
Mild	50-79	500 to 1000 mg twice daily
Moderate	30-49	250 to 750 mg twice daily
Severe*	< 30	250 to 500 mg twice daily
End-stage renal disease patients undergoing dialysis†	—	500 to 1000 mg once daily

* Following dialysis, a 250 to 500 mg supplemental dose is recommended, or according to best clinical judgement.

Patients with Impaired Hepatic Function

No dose adjustment is needed in patients with mild-to-moderate hepatic impairment. In patients with severe hepatic impairment, the creatinine clearance may underestimate the renal insufficiency. Therefore a 50% reduction of the daily maintenance dose is recommended when the creatinine clearance is < 70 mL/min.

Elderly Patients

Dose selection and titration should proceed cautiously in elderly patients, as renal function decreases with age.


PHARMACEUTICAL INFORMATION

Drug Substance

U.S.A.N.: levetiracetam

Chemical Name: (-)-(S)-α-ethyl-2-oxo-1-pyrrolidine acetamide

Structural Formula:



Molecular Formula: $C_8H_{14}N_2O_2$

Molecular Weight: 170.21

Physical Form: A white to off-white crystalline powder with a faint odor and a bitter taste.

Solubility: It is very soluble in water (104.0 g/100 mL). It is freely soluble in chloroform (65.3 g/100 mL) and in methanol (53.6 g/100 mL), soluble in ethanol (16.5 g/100 mL), sparingly soluble in acetonitrile (5.7 g/100 mL) and practically insoluble in n-hexane.

pKa and pH values: The pKa of levetiracetam is < -2 and cannot be determined with accuracy due to the chemical instability of the protonated form.

The protonation of ucb L059 starts at pH_0 values between -1 and -2. Partition Co-efficient: $\Delta \log P$ ($\log P_{\text{octanol}} - \log P_{\text{cyclohexane}}$) was calculated at pH 7.4 using phosphate buffered saline and at pH 1.0 using KCl/HCl. The $\Delta \log P$ at pH 7.4 is 3.65 and at pH 1.0 is 3.10.

Melting Range: 115-119°C

Composition: Keppra® tablets contain the labeled amount of levetiracetam. Inactive ingredients include colloidal silicon dioxide, corn starch, hydroxypropyl methylcellulose, magnesium stearate, polyethylene glycol 4000, povidone, talc, titanium dioxide and coloring agents.

The individual tablets contain the following coloring agents:

250 mg tablets: FD&C Blue No. 2

500 mg tablets: FD&C Blue No. 2 and yellow iron oxide.

750 mg tablets: FD&C Blue No. 2, FD&C Yellow No. 6

and red iron oxide.

Stability and Storage Recommendations

Store between 15-30°C (59-86°F).

AVAILABILITY OF DOSAGE FORMS

Keppra® (levetiracetam) tablets, 250 mg are blue, oblong-shaped, film-coated tablets debossed with "ucb" and "250" on one side. They are supplied in bottles of 120 tablets.

Keppra® (levetiracetam) tablets, 500 mg are yellow, oblong-shaped, film-coated tablets debossed with "ucb" and "500" on one side. They are supplied in bottles of 120 tablets.

Keppra® (levetiracetam) tablets, 750 mg are orange, oblong-shaped, film-coated tablets debossed with "ucb" and "750" on one side. They are supplied in bottles of 120 tablets.

For more information, please refer to the complete Keppra® Product Monograph.

References: 1. Cereghino JJ, Bitton V, Abou-Khalil B, et al. Levetiracetam for partial seizures: results of a double-blind, randomized clinical trial. *Neurology* 2000;55:236-4. 2. Keppra Product Monograph. UCB Pharma, Inc.



Keppra is a registered trademark of UCB S.A. Keppra® is distributed by Lundbeck Canada Inc., 413 St-Jacques St. West, Suite FB-230, Montreal, Quebec H2Y 1N9

COPAXONE®

(glatiramer acetate injection)

20 mg, single use vials and 20 mg/1.0 mL, pre-filled syringes for Subcutaneous Injection
THERAPEUTIC CLASSIFICATION Immunomodulator

ACTION AND CLINICAL PHARMACOLOGY

COPAXONE® (glatiramer acetate for injection [formerly known as copolymer-1]) is a sterile, lyophilized mixture of synthetic polypeptides containing four naturally occurring amino acids: L-glutamic acid, L-alanine, L-tyrosine and L-lysine with an average molar fraction of 0.141, 0.427, 0.095 and 0.338, respectively.

The mechanism(s) by which glatiramer acetate exerts its effect on Multiple Sclerosis (MS) is (are) not fully elucidated. However, it is thought to act by modifying immune processes that are currently believed to be responsible for the pathogenesis of MS. This hypothesis is supported by findings of studies that have been carried out to explore the pathogenesis of experimental allergic encephalomyelitis (EAE), a condition induced in animals that is generally accepted as an experimental model of MS.

Studies in animals and *in vitro* systems suggest that upon its administration glatiramer acetate specific suppressor T cells are induced and activated in the periphery. Because the immunological profile of glatiramer acetate remains to be fully elucidated, concerns exist about its potential to alter naturally occurring immune responses (see PRECAUTIONS).

Pharmacokinetics: Results obtained in pharmacokinetic studies performed in humans (healthy volunteers) and animals support the assumption that a substantial fraction of the therapeutic dose delivered to patients subcutaneously is hydrolyzed locally. Nevertheless, larger fragments of glatiramer acetate can be recognized by glatiramer acetate reactive antibodies. Some fraction of the injected material, either intact or partially hydrolyzed, is presumed to enter the lymphatic circulation, enabling it to reach regional lymph nodes, and, some, may enter the systemic circulation intact.

Clinical Studies: The efficacy of COPAXONE® (glatiramer acetate for injection) was evaluated in two placebo-controlled trials in patients with Relapsing-Remitting MS (RR-MS). In a third placebo-controlled study the effects of glatiramer acetate on MRI parameters were assessed. In these studies, a dose of 20 mg/day was used. No other dose or dosing regimen has been studied in placebo-controlled trials of RR-MS.

The first trial was a pilot study Trial I (Final BR-1) which was conducted at a single-center and was a double-blind, randomized, matched-pair, parallel group placebo-controlled trial. Fifty patients with RR-MS were randomized to receive 20 mg/day glatiramer acetate (n=25) or placebo (n=25) subcutaneously. The protocol-specified primary outcome measure was the proportion of patients who were relapse free during the 2-year duration of the trial, but two additional relevant outcomes were also specified as endpoints: frequency of attacks during the trial, and the change in the number of attacks compared to the rate of attacks in the 2 years prior to study entry. Results from this study (see Table 1) show that there was a statistically significant effect of glatiramer acetate on number of relapses.

TABLE 1 – Trial BR-1: Efficacy Results

Outcome	Trial I*		
	Glatiramer acetate n=25	Placebo n=25	p-Value
% Relapse Free Patients	14/25 (56%)	7/25 (28%)	0.085
Mean Relapse Frequency	0.6/2 years	2.4/2 years	0.005
Reduction in Relapse Rate compared to pre-study	3.2	1.6	0.025
Median Time to First Relapse (days)	>700	150	0.03
% of Progression-Free† Patients	20/25 (80%)	13/25 (52%)	0.07

* The primary efficacy measure for Trial I was the proportion of patients who were relapse free during the 2 year duration of the trial (% Relapse Free). Analyses were based on the intent-to-treat population.

† Progression defined as an increase of at least 1 point on the EDSS that persists for at least 3 consecutive months.

Trial II (01-9001) was a multicenter double-blind, randomized, placebo-controlled trial. Two hundred and fifty-one patients with RR-MS were randomized to receive 20 mg/day glatiramer acetate (n=125) or placebo (n=126) subcutaneously. Patients were diagnosed with RR-MS by standard criteria, and had at least 2 exacerbations during the 2 years immediately preceding enrollment. Patients had a score of no more than 5 on the Kurtzke Expanded Disability Scale Score (EDSS), a standard scale ranging from 0 (normal) to 10 (death due to MS). A score of 5 is defined as one at which a patient is still ambulatory but for whom full daily activities are impaired due to disability, a score of 6 is defined as one at which the patient is still ambulatory but requires assistance and a score of 7 on this scale means that the patient requires a wheelchair.

Patients were seen every 3 months for 2 years, as well as within several days of a presumed exacerbation. In order for an exacerbation to be confirmed, a blinded neurologist had to document objective neurologic signs, as well as document the existence of other criteria (e.g., the persistence of the lesion for at least 48 hours).

The protocol-specified primary outcome measure was the mean number of relapses during treatment. Table 2 shows results of the analysis of primary as well as several secondary outcome measures at two years based on the intent-to-treat population.

TABLE 2 – Core (24-month) Double-Blind Study: Effect on Relapse Rate

Outcome	Trial II†		
	Glatiramer acetate n=125	Placebo n=126	p-Value
Mean No. of Relapses/2 years‡	1.19	1.68	0.055
% Relapse Free Patients	42/125 (34%)	34/126 (27%)	0.25
Median Time to First Relapse (days)	287	198	0.23
% of Patients Progression Free†	98/125 (78%)	95/126 (75%)	0.48
Mean Change in EDSS	-0.05	+0.21	0.023

* The primary efficacy measure for Trial II was the number of relapses during treatment. Analyses were based on the intent-to-treat population.

† Baseline adjusted mean.

‡ Progression defined as an increase of at least 1 point on the EDSS that persists for at least 3 consecutive months.

The effects of glatiramer acetate on relapse severity were not evaluated in either trial. Both studies showed a beneficial effect of glatiramer acetate on relapse rate, and on this basis glatiramer acetate is considered effective.

The third study (9003) was a multi-national, multi-center, MRI-monitored study. A total of 239 patients with RR-MS (119 on glatiramer acetate and 120 on placebo) were randomized. Inclusion criteria were similar to those in Trial II (Study 01-9001) with the additional criteria that patients had to have at least one Gd-enhancing lesion on the screening MRI. The patients were treated initially in a double-blind manner for nine months, during which they underwent monthly MRI scanning. The primary endpoint for the double-blind phase was the total cumulative number of T1 Gd-enhancing lesions over nine months. Other MRI parameters were assessed as secondary endpoints. Table 3 summarizes the results for the parameters monitored during the nine-month double-blind phase for the intent-to-treat cohort. Because the link between MRI findings and the clinical status of patients is contentious, the prognostic value of the following statistically significant findings is unknown.

TABLE 3 – Nine-Month Double-Blind Phase: MRI Endpoints – Results

No.	Outcome	Glatiramer acetate n=113	Placebo n=115	p-Value
Primary Endpoint				
1.	Medians of the Cumulative Number of T1 Gd-Enhancing Lesions	12	17	0.0037
Secondary Endpoints				
2.	Medians of the Cumulative Number of New T1 Gd-Enhancing Lesions	9	14	0.0347
3.	Medians of the Cumulative Number of New T2 Lesions	5	8	0.01
4.	Medians of the Cumulative Change from Baseline in volumes (mL) of T1 Gd-Enhancing Lesions	-0.309	0	0.0248
5.	Medians of the Cumulative Change from Baseline in volumes (mL) of T2 Lesions	8.852	13.566	0.0229
6.	Medians of the Cumulative Change from Baseline in volumes (mL) of T1 Hypointense Lesions	1.642	1.829	0.7311
7.	Proportion of T1 Gd-Enhancing Lesion-Free Patients	46.4%	32.2%	0.0633

The mean number of relapses in this 9-month study was 0.50 for the COPAXONE® group and 0.77 for the placebo group (p=0.0077).

INDICATIONS AND CLINICAL USE

For use in ambulatory patients with Relapsing-Remitting Multiple Sclerosis to reduce the frequency of relapses. The safety and efficacy of COPAXONE® in chronic progressive MS have not been established.

CONTRAINDICATIONS

COPAXONE® (glatiramer acetate for injection) is contraindicated in patients with known hypersensitivity to glatiramer acetate or mannitol.

WARNINGS

The only recommended route of administration of COPAXONE® (glatiramer acetate for injection) injection is the subcutaneous route. COPAXONE® should not be administered by the intravenous route.

Symptoms of Potentially Cardiac Origin: Approximately 26% of COPAXONE® patients in the pre-marketing multicenter controlled trial (compared to 10% of placebo patients) experienced at least one episode of what was described as transient chest pain (see ADVERSE REACTIONS: Chest Pain). While some of these episodes occurred in the context of the Immediate Post-Injection Reaction (see ADVERSE REACTIONS: Immediate Post-Injection Reaction), many did not. The pathogenesis of this symptom is unknown. Patients in controlled clinical trials were free of significant cardiovascular problems (New York Heart Association Class I and II) and thus the risks associated with COPAXONE® treatment for Multiple Sclerosis patients with comorbid cardiovascular disease are unknown.

COPAXONE® has been associated with an Immediate Post-Injection Reaction consisting of a constellation of symptoms appearing immediately after injection that could include flushing, chest pain, palpitations, anxiety, dyspnea, constriction of the throat and urticaria (see ADVERSE REACTIONS: Immediate Post-Injection Reaction).

COPAXONE® has not been studied in patients with a history of severe anaphylactoid reactions, obstructive pulmonary disease or asthma, nor in patients under treatment for either of these two latter conditions. Particular caution is therefore advised regarding the use of COPAXONE® in such patients.

Anaphylactoid reactions associated with the use of COPAXONE® have been reported in rare instances (<1/1000) during the post-marketing period. Some cases required treatment with epinephrine and other appropriate medical treatment.

PRECAUTIONS

General: Patients should be instructed in aseptic reconstitution and self-injection techniques to assure the safe administration of COPAXONE® (glatiramer acetate for injection) (see INFORMATION FOR THE PATIENT). The first injection should be performed under the supervision of an appropriately qualified health care professional. Patient understanding and use of aseptic self-injection techniques and procedures should be periodically re-evaluated. Patients should be cautioned against the re-use of needles or syringes and instructed in safe disposal procedures. A puncture-resistant container for disposal of used needles and syringes should be used by the patient. Patients should be instructed on the safe disposal of full containers.

Considerations Involving the Use of a Product Capable of Modifying Immune Responses: COPAXONE® is an antigenic substance and thus it is possible that detrimental host responses can occur with its use. Whether COPAXONE® can alter normal human immune responses, such as the recognition of foreign antigens is unknown. It is therefore possible that treatment with COPAXONE® may undermine the body's defenses against infections and tumor surveillance. Systematic assessments of these risks have not been done. Continued alteration of cellular immunity due to chronic treatment with glatiramer acetate might result in untoward effects.

Glatiramer acetate-reactive antibodies are formed in practically all patients exposed to daily treatment with the recommended dose. Studies in both the rat and monkey have suggested that immune complexes are deposited in the renal glomeruli. Furthermore, in a controlled clinical trial of 125 RR-MS patients given glatiramer acetate 20 mg for 2 years, serum IgG levels reached at least 3 times baseline values in 80% of patients by 3 months of initiation of treatment. By 12 months of treatment, however, 30% of patients still had IgG levels at least 3 times baseline values, and 90% had levels above baseline by 12 months. The antibodies are exclusively of the IgG subtype – and predominantly of the IgG-1 subtype. No IgE type antibodies could be detected in any of the 94 sera tested. Nevertheless, anaphylaxis can be associated with the administration of almost any foreign substance and, therefore, this risk cannot be excluded.

Preclinical studies to assess the carcinogenic potential of glatiramer acetate in mice and rats do not suggest any evidence of carcinogenic potential related to glatiramer acetate administered subcutaneously at dose levels of up to 30 mg/kg/day in rats and 60 mg/kg/day in mice (see TOXICOLOGY: Carcinogenicity). The relevance of these findings for humans is unknown (see PRECAUTIONS: Considerations Involving the Use of a Product Capable of Modifying Immune Responses).

Drug Interactions: Interactions between COPAXONE® and other drugs have not been fully evaluated. Results from existing clinical trials do not suggest any significant interactions of COPAXONE® with therapies commonly used in MS patients. This includes the concurrent use of corticosteroids for up to 28 days. COPAXONE® has not been formally evaluated in combination with interferon beta. However, 246 patients who failed on or who did not tolerate therapy with interferon beta and were later treated with COPAXONE® within the framework of an open clinical trial did not report any serious or unexpected adverse events thought to be related to treatment.

Use in Pregnancy: There are no adequate and well-controlled studies in pregnant women. No evidence of reproductive toxicity was observed in preclinical studies (see TOXICOLOGY: Reproduction and Teratology). Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed. During pre-marketing clinical trials with COPAXONE®, seven women conceived while being treated with the active drug. One case was lost to follow-up. Three of the patients electively discontinued pregnancy. Three patients stopped treatment 1, 1.5 and 2 months after learning they were pregnant; all delivered healthy babies.

Nursing Mothers: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, treating a nursing woman with COPAXONE® should only be considered after careful risk/benefit assessment and be used with caution.

Use in Children: The safety and effectiveness of COPAXONE® have not been established in individuals below 18 years of age. Use in the Elderly: COPAXONE® has not been studied in the elderly (>65 years old).

Use in Patients with Impaired Renal Function: The pharmacokinetics of COPAXONE® in patients with impaired renal function have not been determined.

ADVERSE REACTIONS

In the pre-marketing clinical trials, approximately 900 individuals have received at least one dose of COPAXONE® (glatiramer acetate for injection) in controlled and uncontrolled clinical trials. Total patient exposure to COPAXONE® in clinical trials ranged from 6 months (693 patients) to 2 years (306 patients), and to over 7 years (69 patients) at a daily dose of 20 mg.

In controlled clinical trials, the most commonly observed adverse events associated with the use of COPAXONE® which occurred at a higher frequency than in placebo treated patients were injection site reactions, vasodilation, chest pain, asthenia, infection, pain, nausea, arthralgia, anxiety and hypertension.

Of a total of 844 patients who could be evaluated for safety, approximately 8% discontinued treatment due to an adverse event. The adverse events most commonly associated with discontinuation were: injection site reaction (6.5%), vasodilation, unintended pregnancy, depression, dyspnea, urticaria, tachycardia, dizziness and tremor. Treatment discontinuation due to a serious adverse event considered by investigators to be related to COPAXONE® treatment included a case of life-threatening serum sickness.

Immediate Post-Injection Reaction: Approximately 10% of Multiple Sclerosis patients exposed to COPAXONE® in pre-marketing studies reported a post-injection reaction immediately following subcutaneous injection of COPAXONE®. Symptoms experienced could include flushing, chest pain, palpitations, anxiety, dyspnea, constriction of the throat and urticaria. These symptoms were invariably transient, self-limited, did not require specific treatment and in general, arose several months after initiation of treatment, although they may occur earlier in the course of treatment. A given patient may experience one or several episodes of these symptoms during treatment with COPAXONE®. Whether these episodes are mediated by an immunologic or non-immunologic mechanism, and whether several similar episodes seen in a given patient have identical mechanisms is unknown. In fact, whether or not this constellation of symptoms actually represents a specific syndrome is unknown. During the post-marketing period, there have been reports of patients with similar symptoms who received emergency medical care (see WARNINGS).

Chest Pain: Approximately 26% of glatiramer acetate patients in the multicenter pre-marketing controlled trial (compared to 10% of placebo patients) experienced at least one episode of what was described as transient chest pain. While some of these episodes occurred in the context of the Immediate Post-Injection Reaction described above, many did not. The temporal relationship of the chest pain to an injection of glatiramer acetate was not always known, although the pain was transient (usually lasting only a few minutes), often unassociated with other symptoms, and appeared to have no important clinical sequelae. Some patients experienced more than one such episode, and episodes usually began at least 1 month after the initiation of treatment. The pathogenesis of this symptom is unknown. There has been only one episode of chest pain during which a full ECG was performed; the ECG showed no evidence of ischemia. Patients in clinical trials were free of significant cardiovascular disease (New York Heart Association Class I or II); therefore, the risks associated with glatiramer acetate treatment for Multiple Sclerosis patients with comorbid cardiovascular disease are unknown (see WARNINGS: Symptoms of Potentially Cardiac Origin).

Table 4 lists the adverse experiences after up to 35 months of treatment (>27-33 months: COPAXONE®, n=84; Placebo, n=75; >33 months: COPAXONE®, n=12; Placebo, n=24) in the pre-marketing multicenter placebo-controlled study (Trial II) in relapsing-remitting Multiple Sclerosis patients that occurred at an incidence of at least 2% among patients who received COPAXONE® and at an incidence that was at least 2% more than that observed in the same trial for placebo patients regardless of their causal relationship to treatment. No laboratory adverse experiences that met these criteria were reported.

It should be noted that the figures cited in Table 4 cannot be used to predict the incidence of side effects during the course of usual medical practice, where patient characteristics and other factors differ from those that prevailed in the clinical trials. However, the cited figures do provide the prescribing physician with some basis for estimating the relative contribution of drug and non-drug factors to the adverse event incidence rate in the population studied.



PRESCRIBING INFORMATION

THERAPEUTIC CLASSIFICATION
Immunomodulator

INDICATIONS AND CLINICAL USE

AVONEX® (Interferon beta-1a) is indicated for the treatment of relapsing forms of multiple sclerosis to slow the progression of disability, decrease the frequency of clinical exacerbations, and reduce the number and volume of active brain lesions identified on Magnetic Resonance Imaging (MRI) scans. Safety and efficacy have not been evaluated in patients with chronic progressive multiple sclerosis.

CONTRAINDICATIONS

AVONEX® (Interferon beta-1a) is contraindicated in patients with a history of hypersensitivity to natural or recombinant interferon beta, human albumin, or any other component of the formulation.

WARNINGS

AVONEX® (Interferon beta-1a) should be used with caution in patients with depression. Depression and suicide have been reported to occur in patients receiving other interferon compounds. Depression and suicidal ideation are known to occur at an increased frequency in the MS population. A relationship between the occurrence of depression and/or suicidal ideation and the use of AVONEX® has not been established. An equal incidence of depression was seen in the placebo-treated and AVONEX®-treated patients in the placebo-controlled relapsing MS study. Patients treated with AVONEX® should be advised to report immediately any symptoms of depression and/or suicidal ideation to their prescribing physicians. If a patient develops depression, antidepressant therapy or cessation of AVONEX® therapy should be considered.

PRECAUTIONS

General

Caution should be exercised when administering AVONEX® (Interferon beta-1a) to patients with pre-existing seizure disorder. In the placebo-controlled study, 4 patients receiving AVONEX® experienced seizures, while no seizures occurred in the placebo group. Of these 4 patients, 3 had no prior history of seizure. It is not known whether these events were related to the effects of MS alone, to AVONEX®, or to a combination of both. For patients with no prior history of seizure who developed seizures during therapy with AVONEX®, an etiologic basis should be established and appropriate anti-convulsant therapy instituted prior to considering resumption of AVONEX® treatment. The effect of AVONEX® administration on the medical management of patients with seizure disorder is unknown.

Patients with cardiac disease, such as angina, congestive heart failure, or arrhythmia, should be closely monitored for worsening of their clinical condition during initiation of therapy with AVONEX®. AVONEX® does not have any known direct-acting cardiac toxicity; however, symptoms of flu syndrome seen with AVONEX® therapy may prove stressful to patients with severe cardiac conditions.

Laboratory Tests

In addition to those laboratory tests normally required for monitoring patients with MS, complete blood cell counts and white blood cell differential, platelet counts, and blood chemistries, including liver and thyroid function tests, are recommended during AVONEX® therapy. During the placebo-controlled study, complete blood cell counts and white blood cell differential, platelet counts, and blood chemistries were performed at least every 6 months. There were no significant differences between the placebo and AVONEX® groups in the incidence of thyroid abnormalities, liver enzyme elevation, leukopenia, or thrombocytopenia (these are known to be dose-related laboratory abnormalities associated with the use of interferons). Patients with myelosuppression may require more intensive monitoring of complete blood cell counts, with differential and platelet counts.

Drug Interactions

No formal drug interaction studies have been conducted with AVONEX®. In the placebo-controlled study, corticosteroids or ACTH were administered for treatment of exacerbations in some patients concurrently receiving AVONEX®. In addition, some patients receiving AVONEX® were also treated with anti-depressant therapy and/or oral contraceptive therapy. No unexpected adverse events were associated with these concomitant therapies.

Other interferons have been noted to reduce cytochrome P-450 oxidase-mediated drug metabolism. Formal hepatic drug metabolism studies with AVONEX® in humans have not been conducted. Hepatic microsomes isolated from AVONEX®-treated rhesus monkeys showed no influence of AVONEX® on hepatic P-450 enzyme metabolism activity.

As with all interferon products, proper monitoring of patients is required if AVONEX® is given in combination with myelosuppressive agents.

Use in Pregnancy

If a woman becomes pregnant or plans to become pregnant while taking AVONEX®, she should be informed of the potential hazards to the fetus, and it should be recommended that the woman discontinue therapy. The reproductive toxicity of AVONEX® has not been studied in animals or humans. In pregnant monkeys given interferon beta at 100 times the recommended weekly human dose (based upon a body surface area comparison), no teratogenic or other adverse effects on fetal development were observed. Abortifacient activity was evident following 3 to 5 doses at this level. No abortifacient effects were observed in monkeys treated at 2 times the recommended weekly human dose (based upon a body surface area comparison).

Although no teratogenic effects were seen in these studies, it is not known if teratogenic effects would be observed in humans. There are no adequate and well-controlled studies with interferons in pregnant women.

Nursing Mothers

It is not known whether AVONEX® is excreted in human milk. Because of the potential of serious adverse reactions in nursing infants, a decision should be made to either discontinue nursing or to discontinue AVONEX®.

Pediatric Use

Safety and effectiveness have not been established in pediatric patients below the age of 18 years.

Information to Patients

Patients should be informed of the most common adverse events associated with AVONEX® administration, including symptoms associated with flu syndrome (see **Adverse Events** and **Information for the Patient**). Symptoms of flu syndrome are most prominent at the initiation of therapy and decrease in frequency with continued treatment. In the placebo-controlled study, patients were instructed to take 650 mg acetaminophen immediately prior to injection and for an additional 24 hours after each injection to modulate acute symptoms associated with AVONEX® administration.

Patients should be cautioned to report depression or suicidal ideation (see **Warnings**).

When a physician determines that AVONEX® can be used outside of the physician's office, persons who will be administering AVONEX® should receive instruction in reconstitution and injection, including the review of the injection procedures (see **Information for the Patient**). If a patient is to self-administer, the physical ability of that patient to self-inject intramuscularly should be assessed. If home use is chosen, the first injection should be performed under the supervision of a qualified health care professional. A puncture-resistant container for disposal of needles and syringes should be used. Patients should be instructed in the technique and importance of proper syringe and needle disposal and be cautioned against reuse of these items.

ADVERSE EVENTS

The safety data describing the use of AVONEX® (Interferon beta-1a) in MS patients are based on the placebo-controlled trial in which 158 patients randomized to AVONEX® were treated for up to 2 years.

The 5 most common adverse events associated (all p<0.075) with AVONEX® treatment were flu-like symptoms (otherwise unspecified), muscle ache, fever, chills, and asthenia. The incidence of all 5 adverse events diminished with continued treatment.

One patient in the placebo group attempted suicide; no AVONEX®-treated patients attempted suicide. The incidence of depression was equal in the 2 treatment groups. However, since depression and suicide have been reported with other interferon products, AVONEX® should be used with caution in patients with depression (see **Warnings**).

In the placebo-controlled study, 4 patients receiving AVONEX® experienced seizures, while no seizures occurred in the placebo group. Of these 4 patients, 3 had no prior history of seizure. It is not known whether these events were related to the effects of MS alone, to AVONEX®, or to a combination of both (see **Precautions**).

Table 1 enumerates adverse events and selected laboratory abnormalities that occurred at an incidence of 2% or more among the 158 patients with relapsing MS treated with 30 mcg of AVONEX® once weekly by IM injection. Reported adverse events have been classified using standard COSTART terms. Terms so general as to be uninformative or more common in the placebo-treated patients have been excluded.

AVONEX® has also been evaluated in 290 patients with illnesses other than MS. The majority of these patients were enrolled in studies to evaluate AVONEX® treatment of chronic viral hepatitis B and C, in which the doses studied ranged from 15 mcg to 75 mcg, given subcutaneously (SC), 3 times a week, for up to 6 months. The incidence of common adverse events in these studies was generally seen at a frequency similar to that seen in the placebo-controlled MS study. In these non-MS studies, inflammation at the site of the SC injection was seen in 52% of treated patients. In contrast, injection site inflammation was seen in 3% of MS patients receiving AVONEX®, 30 mcg by IM injection. SC injections were also associated with the following local reactions: injection site necrosis, injection site alopecia, injection site edema, and injection site hemorrhage. None of the above was observed in the MS patients participating in the placebo-controlled study.

Table 1
Adverse Events and Selected Laboratory Abnormalities
in the Placebo-Controlled Study

Adverse Event	Placebo (N = 143)	AVONEX® (N = 158)
Body as a Whole		
Headache	57%	67%
Flu-like symptoms (otherwise unspecified)*	40%	61%
Pain	20%	24%
Fever*	13%	23%
Asthenia	13%	21%
Chills*	7%	21%
Infection	6%	11%
Abdominal pain	6%	9%
Chest pain	4%	6%
Injection site reaction	1%	4%
Malaise	3%	4%
Injection site inflammation	0%	3%
Hypersensitivity reaction	0%	3%
Ovarian cyst	0%	3%
Ecchymosis injection site	1%	2%
Cardiovascular System		
Syncope	2%	4%
Vasodilation	1%	4%

Table 1
Adverse Events and Selected Laboratory Abnormalities
in the Placebo-Controlled Study

Adverse Event	Placebo (N = 143)	AVONEX® (N = 158)
Digestive System		
Nausea	23%	33%
Diarrhea	10%	16%
Dyspepsia	7%	11%
Anorexia	6%	7%
Hemic and Lymphatic System		
Anemia*	3%	8%
Eosinophils ≥ 10% HCT (%) ≤ 32 (females) or ≤ 37 (males)	4%	5%
	1%	3%
Metabolic and Nutritional Disorders		
SGOT ≥ 3 x ULN	1%	3%
Musculoskeletal System		
Muscle ache*	15%	34%
Arthralgia	5%	9%
Nervous System		
Sleep difficult	16%	19%
Dizziness	13%	15%
Muscle spasm	6%	7%
Suicidal tendency	1%	4%
Seizure	0%	3%
Speech disorder	0%	3%
Ataxia	0%	2%
Respiratory System		
Upper respiratory tract infection	28%	31%
Sinusitis	17%	18%
Dyspnea	3%	6%
Skin and Appendages		
Urticaria	2%	5%
Alopecia	1%	4%
Nevus	0%	3%
Herpes zoster	2%	3%
Herpes simplex	1%	2%
Special Senses		
Otitis media	5%	6%
Hearing decreased	0%	3%
Urogenital		
Vaginitis	2%	4%

* Significantly associated with AVONEX® treatment (p ≤ 0.05).

DOSAGE AND ADMINISTRATION

The recommended dosage of AVONEX® (Interferon beta-1a) for the treatment of relapsing forms of multiple sclerosis is 30 mcg injected intramuscularly once a week.

AVONEX® is intended for use under the guidance and supervision of a physician. Patients may self-inject only if their physician determines that it is appropriate and with medical follow-up, as necessary, after proper training in IM injection technique.

AVAILABILITY OF DOSAGE FORMS

AVONEX® (Interferon beta-1a) is available as:

Package (Administration Pack) containing 4 Administration Dose Packs (each containing one vial of AVONEX®, one 10 mL (10 cc) diluent vial, two alcohol wipes, one gauze pad, one 3 cc syringe, one Micro Pin®, one needle, and one adhesive bandage).

Product Monograph Available upon request.

REFERENCES

- Galetta SL, Markowitz C, Lee AG. Immunomodulatory agents for the treatment of relapsing multiple sclerosis. *Arch Intern Med* 2002; 162:2161-2169.
- Bertolotto A, Malucchi S, Sala A, et al. Differential effects of three interferon betas on neutralizing antibodies in patients with multiple sclerosis: a follow-up study in an independent laboratory. *J Neuro Neurol Psychiatry* 2002; 73:148-153.
- Giovannoni G, Munschauer FE, Disensteinhammer F. Neutralizing antibodies to interferon beta during the treatment of multiple sclerosis. *J Neuro Neurol Psychiatry* 2002; 73:465-469.
- Rudick RA, Simonian NA, Alam JA. Incidence and significance of neutralizing antibodies to interferon beta-1a in multiple sclerosis. *Neurology* 1998; 50:1266-1272.
- AVONEX product monograph, 2000.
- Rudick RA, Fisher E, Lee JC, et al. Use of the brain parenchymal fraction to measure whole brain atrophy in relapsing-remitting MS. *Neurology* 1999; 53:1698-1704.
- Data on file, Biogen, Inc.



3 Robert Speck Parkway, Suite 300, Mississauga, Ontario L4Z 2G5
© 2003 Biogen Canada, Inc. All rights reserved.

Biogen Canada is a trademark of Biogen, Inc.
AVONEX is a registered trademark of Biogen, Inc.
Rebif is a registered trademark of Serono Canada Inc.
Betaseron is a registered trademark of Berlex Canada Inc.
Micro Pin® is a registered trademark of B. Braun Medical Inc.



Ropinirole (as ropinirole hydrochloride)

TABLETS: 0.25 mg, 1.0 mg, 2.0 mg, 5.0 mg

THERAPEUTIC CLASSIFICATION: AntiParkinsonian Agent / Dopamine Agonist
INDICATIONS AND CLINICAL USE: REQUIP® (ropinirole hydrochloride) is indicated in the treatment of the signs and symptoms of idiopathic Parkinson's disease. REQUIP® can be used both as early therapy, without concomitant levodopa and as an adjunct to levodopa. Three year and five year active-comparator controlled clinical trials have been conducted.

CONTRAINDICATIONS: REQUIP® (ropinirole hydrochloride) is contraindicated in patients with a known hypersensitivity to ropinirole hydrochloride or the excipients of the drug product.

WARNINGS: Sudden Onset of Sleep – Patients receiving treatment with REQUIP® (ropinirole hydrochloride), and other dopaminergic agents have reported suddenly falling asleep while engaged in activities of daily living, including operating a motor vehicle, which has sometimes resulted in accidents. Although some of the patients reported somnolence while on REQUIP®, others perceived that they had no warning signs, such as excessive drowsiness, and believed that they were alert immediately prior to the event. Physicians should alert patients of the reported cases of sudden onset of sleep, bearing in mind that these events are NOT limited to initiation of therapy. Patients should also be advised that sudden onset of sleep has occurred without warning signs. If drowsiness or sudden onset of sleep should occur, patients should immediately contact their physician. Until further information is available on the management of this unpredictable and serious adverse event, patients should be warned not to drive or engage in other activities where impaired alertness could put themselves and others at risk of serious injury or death (e.g., operating machines). Episodes of falling asleep while engaged in activities of daily living have also been reported in patients taking other dopaminergic agents, therefore, symptoms may not be alleviated by substituting these products. Presently, the precise cause of this event is unknown. It is known that many Parkinson's disease patients experience alterations in sleep architecture, which results in excessive daytime sleepiness or spontaneous dozing, and that dopaminergic agents can also induce sleepiness. There is insufficient information to determine whether this event is associated with REQUIP®, all dopaminergic agents or Parkinson's disease itself.

Orthostatic Symptoms – Dopamine agonists appear to impair the systemic regulation of blood pressure with resulting orthostatic symptoms of dizziness or lightheadedness, with or without documented hypotension. These symptoms appear to occur especially during dose escalation. Therefore, patients treated with dopamine agonists should be carefully monitored for signs and symptoms of orthostatic hypotension, especially during dose escalation (see DOSAGE and ADMINISTRATION) and should be informed of this risk. **Hallucinations – Early Therapy:** In placebo-controlled trials, REQUIP® (ropinirole hydrochloride) caused hallucination in 5.1% of patients during early therapy (1.4% in the placebo group). Hallucination was of sufficient severity that it led to discontinuation in 1.3% of patients. The incidence of hallucination was dose-dependent. In a 5-year study comparing REQUIP® with levodopa in early Parkinson's patients, the overall incidence of hallucinations was 17.3% (31/179) for patients treated with REQUIP® and 5.6% (5/89) for levodopa patients. Hallucinations led to discontinuation of the study treatment in 5.0% of REQUIP® and 2.2% of levodopa patients. In a 3-year study comparing REQUIP® with another dopamine agonist, the overall incidence of hallucinations was 9.5% (16/168) for patients treated with REQUIP® and 9.0% (15/167) for patients receiving active comparator. Hallucinations led to discontinuation of the study treatment in 2.4% of REQUIP® patients and 3.0% of comparator patients. Concomitant Selegiline: In a 5-year study, REQUIP® patients receiving concomitant selegiline reported a higher incidence of hallucinations (23.5%) than did those without (12.2%), this subpopulation effect was not seen in the L-dopa arm (hallucinations with concomitant selegiline = 2.0% vs hallucinations without selegiline = 8.0%).

Adjunct Therapy: Hallucinations were experienced by 10.1% of patients receiving REQUIP® and levodopa, compared to 4.2% receiving placebo and levodopa. Hallucinations were of sufficient severity that it led to discontinuation in 1.9% of patients. The incidence of hallucinations was dose dependent.

PRECAUTIONS: Cardiovascular – Since REQUIP® (ropinirole hydrochloride) has not been studied in patients with a history or evidence of significant cardiovascular disease including myocardial infarction, unstable angina, cardiac decompensation, cardiac arrhythmias, vaso-occlusive disease (including cerebral) or cardiomyopathy, it should be used with caution in such patients. There is limited experience with REQUIP® in patients treated with antihypertensive and antiarrhythmic agents. Consequently, in such patients, the dose of REQUIP® should be titrated with caution. **Orthostatic Symptoms** - Orthostatic symptoms of dizziness or lightheadedness as well as somnolence may occur during REQUIP® therapy. **Neuroleptic Malignant Syndrome** – A symptom complex resembling the neuroleptic malignant syndrome (characterized by elevated temperature, muscular rigidity, altered consciousness, and autonomic instability), with no other obvious etiology, has been reported in association with rapid dose reduction, withdrawal of, or changes in anti-Parkinsonian therapy. A single spontaneous report of a symptom complex resembling the neuroleptic malignant syndrome has been observed in a 66 year old diabetic male patient with Parkinson's disease, who developed fever, muscle stiffness, and drowsiness 8 days after beginning REQUIP® treatment. The patient also experienced acute bronchitis, which did not respond to antibiotic treatment. REQUIP® was discontinued three days

before the patient died. The reporting physician considered these events to be possibly related to REQUIP® treatment. (see DOSAGE AND ADMINISTRATION). A single spontaneous report of severe muscle pain has been reported in a 66 year old male patient around his thigh. The reporting physician considered the event to be probably related to REQUIP® treatment.

Retinal Pathology in Rats – In a two year carcinogenicity study in albino Sprague-Dawley rats, retinal atrophy was observed at incidences of 0%, 1.4%, 1.4% and 10% of male rats and 0%, 4.4%, 2.9% and 12.9% of female rats dosed at 0, 1.5, 15 and 50 mg/kg/day respectively. The incidence was significantly higher in both male and female animals dosed at 50 mg/kg/day. The 50 mg/kg/day dose represents a 2.8 fold greater exposure (AUC) and a 13.1 fold greater exposure (C_{max}) to ropinirole in rats than the exposure would be in humans at the maximum recommended dose of 24 mg/day. The relevance of this finding to humans is not known. **Pregnancy** – The use of REQUIP® during pregnancy is not recommended. REQUIP® given to pregnant rats during organogenesis (gestation days 8 through 15) resulted in decreased fetal body weight at 60 mg/kg/day (approximately 3–4 times the AUC at the maximal human dose of 8 mg t.i.d.), increased fetal death at 90 mg/kg/day (approximately 5 times the AUC at the maximal human dose of 8 mg t.i.d.) and digital malformations at 150 mg/kg/day (approximately 8–9 times the AUC at the maximal human dose of 8 mg t.i.d.). These effects occurred at maternally toxic doses. There was no indication of an effect on development of the conceptus at a maternally toxic dose of 20 mg/kg/day in the rabbit. In a perinatal-postnatal study in rats, 10 mg/kg/day of REQUIP® (approximately 0.5 – 0.6 times the AUC at the maximal human dose of 8 mg t.i.d.) impaired growth and development of nursing offspring and altered neurological development of female offspring.

Nursing Mothers – Since REQUIP® suppresses lactation, it should not be administered to mothers who wish to breast-feed infants. Studies in rats have shown that REQUIP® and/or its metabolites cross the placenta and are excreted in breast milk. Consequently, the human fetus and/or neonate may be exposed to dopamine agonist activity. **Use in Women Receiving Estrogen Replacement Therapy** – In female patients on long-term treatment with conjugated estrogens, oral clearance was reduced and elimination half-life prolonged compared to patients not receiving estrogens. In patients, already receiving estrogen replacement therapy, REQUIP® may be titrated in the recommended manner according to clinical response. However, if estrogen replacement therapy is stopped or introduced during treatment with REQUIP®, adjustment of the REQUIP® dosage may be required. **Pediatric Use** – Safety and effectiveness in the pediatric population have not been established. **Renal and Hepatic Impairment** – No dosage adjustment is needed in patients with mild to moderate renal impairment (creatinine clearance of 30 to 50 mL/min). Because the use of REQUIP® in patients with severe renal impairment or hepatic impairment has not been studied, administration of REQUIP® to such patients is not recommended. **Drug Interactions – Psychotropic Drugs:** Neuroleptics and other centrally active dopamine antagonists may diminish the effectiveness of REQUIP®. Therefore, concomitant use of these products is not recommended. Based on population pharmacokinetic assessment, no interaction was seen between REQUIP® and tricyclic antidepressants or benzodiazepines. **Anti-Parkinson Drugs:** Based on population pharmacokinetic assessment, there were no interactions between REQUIP® and drugs commonly used to treat Parkinson's disease, i.e., selegiline, amantadine, and anticholinergics. **Levodopa:** The potential pharmacokinetic interaction of levodopa/carbidopa (100 mg/10 mg b.i.d.) and REQUIP® (2 mg t.i.d.) was assessed in levodopa naive (de novo) male and female patients with Parkinson's disease (n=30, mean age 64 years). The rate and extent of availability of REQUIP® at steady state were essentially the same with or without levodopa. Similarly, the rate and extent of availability of levodopa, as well as its elimination half-life, were essentially the same in the presence and absence of REQUIP®. **Inhibitors of CYP1A2: Ciprofloxacin:** The effect of ciprofloxacin (500 mg b.i.d.) on the pharmacokinetics of REQUIP® (2 mg t.i.d.) was studied in male and female patients with Parkinson's disease (n=12, mean age 55 years). The extent of systemic availability of REQUIP® was significantly increased when coadministered with ciprofloxacin (AUC increased by 1.84 fold). Thus, in patients already receiving CYP1A2 inhibitors such as ciprofloxacin, REQUIP® therapy may be instituted in the recommended manner and the dose titrated according to clinical response. However, if therapy with a drug known to be an inhibitor of CYP1A2 is stopped or introduced during treatment with REQUIP®, adjustment of the REQUIP® dosage will be required. **Substrates of CYP1A2: Theophylline:** The effect of oral theophylline (300 mg b.i.d.) on the pharmacokinetics of REQUIP® (2 mg t.i.d.) was studied in male and female patients with Parkinson's disease (n=12, mean age 59 years). There was no marked change in the rate or extent of availability of REQUIP® when coadministered with theophylline. Similarly, coadministration of REQUIP® with intravenous theophylline (5 mg/kg) did not result in any marked change in the pharmacokinetics of theophylline. It is therefore unlikely that substrates of CYP1A2 would significantly alter the pharmacokinetics of REQUIP®, and vice-versa. **Digoxin:** The effect of REQUIP® (2 mg t.i.d.) on the pharmacokinetics of digoxin (0.125–0.25 mg o.d.) was studied in male and female patients with Parkinson's disease (n=10, mean age 72 years). Coadministration at steady state with REQUIP® resulted in a 10% decrease in digoxin AUC although mean trough digoxin plasma concentrations were unaltered. However, the effect of higher recommended doses of REQUIP® on the pharmacokinetics of digoxin is not known. **Alcohol:** No information is available on the potential for interaction between REQUIP® and alcohol. As with other centrally active medications, patients should be cautioned against taking REQUIP® with alcohol. **Psycho-Motor Performance** – (see WARNINGS-Sudden Onset of Sleep).

ADVERSE REACTIONS: Adverse Reactions Associated with Discontinuation of Treatment – Of 1599 patients who received REQUIP® (ropinirole hydrochloride) during the premarketing clinical trials, 17.1% in

early-therapy studies and 17.3% in adjunct-therapy studies discontinued treatment due to adverse reactions. The events resulting in discontinuation of REQUIP® in 1% or more of patients were as follows: **Early therapy:** nausea (6.4%), dizziness (3.8%), aggravated Parkinson's disease (1.3%), hallucination (1.3%), headache (1.3%), somnolence (1.3%) and vomiting (1.3%). **Adjunct therapy:** dizziness (2.9%), dyskinesia (2.4%), confusion (2.4%), vomiting (2.4%), hallucination (1.9%), nausea (1.9%), anxiety (1.9%), and increased sweating (1.4%). Patients over 75 years of age (n=130) showed slightly higher incidences of withdrawal due to hallucination, confusion and dizziness than patients less than 75 years of age. **Most Frequent Adverse Events** – Adverse events occurring with an incidence of greater than, or equal to, 10% were as follows: **Early therapy:** nausea, dizziness, somnolence, headache, peripheral edema, vomiting, syncope, fatigue and viral infection. **Adjunct therapy:** dyskinesia, nausea, dizziness, somnolence and headache. Dopamine agonists, with an ergoline chemical structure have been associated with adverse experiences such as retroperitoneal fibrosis, erythromelalgia and pulmonary reactions. REQUIP® has a novel, non-ergoline chemical structure and no reports of such events have been observed in clinical trials. **Incidence of Adverse Events in Placebo Controlled Trials** – The incidence of postural hypotension, an event commonly associated with initiation of dopamine agonist therapy, was not notably different from placebo in clinical trials. However, decreases in systolic blood pressure to < 90 mmHg have been observed in 13% (<65 years), 16% (65 – 75 years) and 7.6% (>75 years) of patients treated with REQUIP®. **Table 2** lists adverse events that occurred at an incidence of 1% or more among REQUIP®-treated patients who participated in placebo-controlled trials for up to one year. Patients were dosed in a range of 0.75 mg to 24 mg/day. Reported adverse events were classified using a standard World Health Organization (WHO)-based dictionary terminology. The prescriber should be aware that these figures can not be used to predict the incidence of adverse events in the course of usual medical practice where patient characteristics and other factors differ from those which prevailed in the clinical trials. Similarly, the cited frequencies can not be compared with figures obtained from other clinical investigations involving different treatments, uses and investigators. The cited figures, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and non-drug factors to the adverse events incidence rate in the population studied.

TABLE 2
Adverse events with incidence ≥1% from all placebo-controlled early and adjunct therapy studies

	Early Therapy		Adjunct Therapy	
	REQUIP® N = 157 % occurrence	Placebo N = 147 % occurrence	REQUIP® N = 208 % occurrence	Placebo N = 128 % occurrence
Autonomic Nervous System				
Sweating increased	6.4	4.1	7.2	1.7
Mouth Dry	5.1	3.4	5.3	0.8
Flushing	3.2	0.7	1.4	0.8
Body as a Whole General				
Peripheral Edema	13.4	4.1	3.9	2.5
Fatigue	10.8	4.1	–	–
Injury	–	–	10.6	9.2
Pain	7.6	4.1	5.3	3.3
Asthenia	6.4	1.4	–	–
Drug Level Increased	4.5	2.7	6.7	3.3
Chest Pain	3.8	2.0	–	–
Melaise	3.2	0.7	1.4	0.8
Therapeutic Response				
Decreased	1.9	0.7	–	–
Cellulitis	1.3	0.0	–	–
Influenza-like Symptoms	–	–	1.0	0.0
Fever	–	–	1.4	0.0
Cardiovascular General				
Syncope	11.5	1.4	2.9	1.7
Hypotension Postural	6.4	4.8	–	–
Hypertension	4.5	3.4	3.4	3.3
Hypotension	1.9	0.0	2.4	0.8
Cardiac Failure	–	–	1.0	0.0
Central and Peripheral Nervous System				
Dizziness	40.1	21.8	26.0	15.8
Dyskinesia	–	–	33.7	12.5
Headache	17.2	17.0	16.8	11.7
Ataxia (Falls)	–	–	9.6	6.7
Tremor	–	–	6.3	2.5
Paresthesia	–	–	5.3	2.9
Hypersensitivity	3.8	2.0	–	–
Dystonia	–	–	4.3	4.2
Hypokinesia	–	–	5.3	4.2
Paresis	–	–	2.9	0.0
Speech Disorder	–	–	1.0	0.0
Vertigo	1.9	0.0	–	–
Carpal Tunnel Syndrome	1.3	0.7	–	–
Gastrointestinal System				
Nausea	59.9	21.8	29.8	18.3
Vomiting	12.1	6.8	7.2	4.2
Dyspepsia	9.6	4.8	–	–
Constipation	9.3	7.5	5.8	3.3
Abdominal Pain	6.4	2.7	8.7	7.5
Diarrhea	–	–	4.8	2.5
Anorexia	3.8	1.4	–	–
Fatulence	2.5	1.4	1.9	0.8
Tooth Disorder	1.9	0.7	1.0	0.8
Saliva Increased	–	–	2.4	0.8
Colitis	1.3	0.0	–	–
Dysphagia	1.3	0.0	2.4	0.8
Periodontitis	1.3	0.0	1.4	0.8
Erectation	–	–	1.4	0.0
Facial Incontinence	–	–	1.0	0.0
Hemorrhoids	–	–	1.0	0.0
Gastroesophageal Reflux	–	–	1.0	0.0
Gastrointestinal Disorder (NOS)	–	–	1.0	0.0
Tooth Ache	–	–	1.0	0.0
Hearing and Vestibular				
Tinnitus	1.3	0.0	–	–
Heart Rate and Rhythm				
Palpitation	3.2	2.0	2.9	2.5

	Early Therapy		Adjunct Therapy	
	REQUIP [®] N = 157 % occurrence	Placebo N = 147 % occurrence	REQUIP [®] N = 208 % occurrence	Placebo N = 120 % occurrence
Heart Rate and Rhythm				
Extrasystoles	1.9	0.7	—	—
Tachycardia	1.9	0.0	1.0	0.0
Fibrillation Atrial	1.9	0.0	—	—
Tachycardia Supraventricular	1.3	0.0	—	—
Bradycardia	—	—	1.0	0.0
Liver and Biliary System				
Gamma - GT Increased	1.3	0.7	1.0	0.0
Hepatic Enzymes Increased	1.3	0.0	—	—
Metabolic and Nutritional				
Alkaline Phosphate Increased	2.5	1.4	1.0	0.0
Weight Decrease	—	—	2.4	0.8
Hypoglycemia	1.3	0.0	—	—
Musculoskeletal System				
Arthralgia	—	—	6.7	5.0
Arthritis	—	—	2.9	0.8
Arthritis Aggravated	1.3	0.0	1.4	0.0
Myocardial, Endocardial, Pericardial Valve				
Myocardial Ischemia	1.3	0.7	—	—
Psychiatric				
Somnolence	40.1	6.1	20.2	8.3
Anxiety	—	—	6.3	3.3
Confusion	5.1	1.4	8.7	1.7
Hallucination	5.1	1.4	10.1	4.2
Nervousness	—	—	4.8	2.5
Yawning	3.2	0.0	—	—
Amnesia	2.5	1.4	4.8	0.8
Dreaming Abnormal	—	—	2.9	1.7
Depersonalization	—	—	1.4	0.0
Paranoid Reaction	—	—	1.4	0.0
Asthenia	1.3	0.7	1.0	0.0
Concentration Impaired	1.9	0.0	1.0	0.0
Illusion	1.3	0.0	—	—
Thinking Abnormal	—	—	1.4	0.8
Apathy	—	—	1.0	0.0
Increased Libido	—	—	1.0	0.0
Personality Disorder	—	—	1.0	0.0
Red Blood Cell				
Anemia	—	—	2.4	0.0
Reproductive Male				
Impotence	2.5	1.4	—	—
Prostatic Disorder	—	—	1.0	0.0
Penis Disorder	—	—	1.3	0.0
Resistance Mechanism				
Upper Respiratory Tract Infection	—	—	8.7	8.3
Infection Viral	10.8	3.4	7.2	6.7
Respiratory System				
Pharyngitis	6.4	4.1	—	—
Rhinitis	3.8	2.7	—	—
Sinusitis	3.8	2.7	—	—
Dyspnea	3.2	0.0	2.9	1.7
Bronchitis	2.5	1.4	—	—
Respiratory Disorder	1.9	1.4	1.9	0.0
Pneumonia	1.3	0.7	1.0	0.8
Coughing	—	—	1.4	0.8
Skin/Appendages				
Pruritis	—	—	1.0	0.0
Urinary System				
Urinary Tract Infection	5.1	4.1	6.3	2.5
Cystitis	1.3	0.7	—	—
Micturition Frequency	—	—	1.4	0.0
Pyuria	—	—	1.9	0.8
Urinary Incontinence	—	—	1.9	0.8
Urinary Retention	1.3	0.7	—	—
Dysuria	—	—	1.0	0.0
Vascular Extracardiac				
Peripheral Ischemia	2.5	0.0	—	—
Vision				
Vision Abnormal	5.7	3.4	—	—
Eye Abnormality	3.2	1.4	—	—
Diplopia	—	—	1.9	0.8
Xerophthalmia	1.9	0.0	1.4	0.8
Cataract	—	—	1.4	0.8
Lacrimation Abnormal	—	—	1.4	0.0
White Cell and Reticuloendothelial System				
Eosinophilia	—	—	1.4	0.0

a: Incidence of adverse event <1%.

Post-Marketing Experience - Patients treated with REQUIP[®] have rarely reported suddenly falling asleep while engaged in activities of daily living, including operation of motor vehicles which has sometimes resulted in accidents (see WARNINGS).

DOSAGE AND ADMINISTRATION: REQUIP[®] (ropinirole hydrochloride) should be taken three times daily. While administration of REQUIP[®] with meals may improve gastrointestinal tolerance, REQUIP[®] may be taken with or without food. The recommended starting dosage is 0.25 mg three times daily. Based on individual patient response, dosage should then be titrated by weekly increments of 0.25 mg per dose as described in the table below. After week 4, daily dosage may be increased by 0.5 to 1.0 mg per dose on a weekly basis until an optimal therapeutic response is established. Smaller dose increments are recommended for patients who may be at risk for orthostatic symptoms.

	Week			
	1	2	3	4
Unit Dose (mg)	0.25	0.5	0.75	1.0
Total Daily Dose (mg)	0.75	1.5	2.25	3.0


In clinical trials, initial benefits were observed with 3 mg/day and higher doses. Doses greater than 24 mg/day have not been included in clinical trials. In a 5-year, double-blind study of early therapy in Parkinson's disease patients, the average daily dose of REQUIP[®] (based on the observed data set) was 10.1 mg at 6 months (median dose = 9.0 mg), 14.4 mg at 3 years (median dose = 15.0 mg), and 16.6 mg at 5 years (median dose = 18.0 mg), regardless of levodopa supplementation. When REQUIP[®] is administered as adjunct therapy to levodopa, the dose of levodopa may be decreased gradually as tolerated once a therapeutic effect with REQUIP[®] has been observed. REQUIP[®] should be



discontinued gradually over a 7-day period. The frequency of administration should be reduced from three times daily to twice daily for 4 days. For the remaining 3 days, the frequency should be reduced to once daily prior to complete withdrawal of REQUIP[®]. **Renal and Hepatic Impairment:** In patients with mild to moderate renal impairment, REQUIP[®] may be titrated in the recommended manner according to clinical response. Patients with severe renal impairment or on hemodialysis have not been studied and administration of REQUIP[®] to such patients is not recommended. Patients with hepatic impairment have not been studied and administration of REQUIP[®] to such patients is not recommended. **Estrogen Replacement Therapy:** In patients already receiving estrogen replacement therapy, REQUIP[®] may be titrated in the recommended manner according to clinical response. However, if estrogen replacement therapy is stopped or started during treatment with REQUIP[®], adjustment of the REQUIP[®] dosage may be required. **AVAILABILITY OF DOSAGE FORM:** REQUIP[®] is supplied as a pentagonal film-coated 'Tiltab' tablet with beveled edges containing ropinirole (as ropinirole hydrochloride) as follows: 0.25 mg - white imprinted with SB and 4890; 1.0 mg - green imprinted with SB and 4892; 2.0 mg - pale pink imprinted with SB and 4893; 5.0 mg - blue tablets imprinted with SB and 4894. REQUIP[®] is available in bottles in the pack size of 100 tablets. Full Product Monograph available to practitioners upon request.

GlaxoSmithKline Inc.
7333 Mississauga Road North
Mississauga, Ontario
L5N 6L4

REQUIP[®] is a registered trademark, used under license by GlaxoSmithKline Inc.
Date of preparation: June 18, 2001
Date of revisions: July 31, 2002

1. Rascol O *et al.* A five-year study of the incidence of dyskinesia in patients with early Parkinson's disease who were treated with ropinirole or levodopa. *N Eng J Med* 2000;342(20):1484-1491. 2. Product Monograph of ReQUIP[®] (ropinirole hydrochloride), GlaxoSmithKline, July 31, 2002.

 GlaxoSmithKline



KING MEDICAL

THE CANADIAN ELECTRODE PLACE

- AMBU Blue Sensor • Neuroline
- CHALGREN Bar/Ring/Clip Electrodes
- CLASS A Monopolar Needles
- KENDALL Adhesive • NuTab
- KING MEDICAL Cables & Adapters
- MAVIDON Lemon & Green Skin Prep
- MEDTRONIC Mono/Conc. Needles
- PARKER LAB. Electrode Paste
- 3M CANADA Micropore • Transpore
- D.O. WEAVER Ten20 • NuPrep

Bo-ject™ • Chalgren • Inoject™
Large stock of Hypodermic Needles

Tel 905-833-3545

Fax 905-833-3543

E-mail: soren@kingmedical.com

Web Site: www.kingmedical.com

King Medical Ltd.
145 Kingsworth Road
King City • Ontario L7B 1K1

ADVERTISERS INDEX

- Aventis
Altace - A-10, A-11, A-40
- Bayer
Gamunex - A-9
- Berlex
Betaseron - A-6, A-7, A-14, A-15,
A-33, A-34
- Biogen
Avonex - A-20, A-21, A-41
- GlaxoSmithKline
Imitrex - IBC, A-29, A-30
Requip - A-17, A-43, A-44
- Janssen-Ortho
Topamax - A-4, A-5, A-24, A-25,
A-31, A-32
- Lundbeck
Corporate - A-12, A-13
Keppra - A-18, A-19, A-35, A-36
- Novartis
Exelon - OBC, A-39
- Pfizer
Aricept - A-23, A-42
- Serono
Rebif - IFC, A-26, A-27, A-28
- Teva Neuroscience
Copaxone - A-3, A-37, A-38
- Classified Ads - A-44, A-46, A-47



Fellowship in Stereotactic & Functional Neurosurgery



Dalhousie University

The Division of Neurosurgery at Dalhousie University is offering a one year Clinical Fellowship in Stereotactic & Functional Neurosurgery. All functional neurosurgical procedures for Atlantic Canada (population 2,500,000) are performed at the QEII Health Sciences Center/Dalhousie University. Fellows will participate in the evaluation and treatment of patients with a broad range of functional neurosurgical disorders including:

- movement disorders
- angina
- complex pain
- epilepsy
- spasticity

Fellows will have training in different techniques including:

- Deep brain stimulation, with and without microelectrode recording
- Neurotransplantation
- Spinal cord stimulation
- Intrathecal therapy
- Ablative procedures
- Selective mesial temporal resections
- Extratemporal resections for epilepsy
- Vagus nerve stimulation

Candidates must have completed their neurosurgical training and be eligible for licensure in Nova Scotia, commencing July 1, 2004. Candidates must have successfully completed the MCCEE or USMLE qualifying exams to be eligible for licensure. Interested candidates should send a current Curriculum Vitae and three letters of reference along with their cover letter outlining why they wish to study stereotactic and functional neurosurgery, by November 15, 2003, to:

Dr. Robert Brownstone, MD, PhD, FRCS
 c/o Mrs. Patricia Turner, Division of Neurosurgery,
 QEII Health Sciences Center, 3844 - 1796 Summer Street,
 Halifax, NS B3H 3A7.
 Phone: (902) 473-8919; Fax: (902) 473-6609;
 e-mail: patricia.turner@dal.ca



Parkinson Society Canada Soci t  Parkinson Canada

**Do you see patients
with Parkinson's?**

**If so, Parkinson Society
Canada is here to help.**

We are the national voice of Canadians living with Parkinson's. Our purpose is to ease the burden and find a cure through research, education, advocacy and support. We can offer your patients:

- an Information & Referral Centre, at 1-800-565-3000, to answer questions and link them to support services across Canada
- a current, informative website, www.parkinson.ca
- a wide variety of free brochures, in English and French
- *Parkinson Post/L'Actualit  Parkinson*, a quarterly magazine for Canadians living with Parkinson's

To request a sample Patient Information Package, please call us at 1-800-565-3000, ext. 225.

Ease the Burden; Find a Cure
www.parkinson.ca ■ 1-800-565-3000

Subscribe Now!

YES! Please send me a one-year subscription to the Journal, now including the ONLINE version at www.cjns.org

Complete the enclosed subscription form and send by mail, fax or e-mail to:

Subscriptions, Canadian Journal of Neurological Sciences
 PO Box 5456, Stn. A
 Calgary AB Canada T2H 1X8
 Tel: (403) 229-9575 Fax: (403) 229-1661
 E-mail: journal@cjns.org



SUBSCRIPTION RATES (Print and Online) 2003

	Canada*	US	Other Countries
Non-Member Individual	\$C 80.00	\$US 80.00	\$US 85.00
Non-Member Student	\$C 40.00	\$US 40.00	\$US 42.50
Institutions	\$C 90.00	\$US 90.00	\$US 95.00

* add 7% GST or 15% HST

Agent discount 5%

Subscription Rate	\$
* Tax: GST 7% / HST 15%	\$
TOTAL	\$

PAYMENT METHODS:

- Cheque (*\$C drawn on a Canadian bank and \$US drawn on a US bank*)
 Visa Mastercard

Credit Card No.: _____

Expiry date: _____

Signature: _____

Telephone: _____

Last Name: _____

First Name: _____

Address: _____

City: _____

Province/State: _____ Country: _____

Postal Code: _____



Queen's
UNIVERSITY

Chair, Division of Neurology: The Department of Medicine,

Queen's University, seeks a Chair of the Division of Neurology. Candidates will have the skills and the professional experience to lead and direct the Division in fulfilling the academic and clinical missions of Neurology within the Southeastern Ontario Health Sciences Centre. Successful candidates must be eligible for licensure in Ontario.

The division provides a full spectrum of neurological consultative services for patients in Southeastern Ontario. Specialized outpatient clinics in several areas of subspecialty neurology as well as general clinics have been established along with several outreach clinics. Inpatient neurology and neurosurgery services are integrated at Kingston General Hospital (KGH) and they include a state-of-the-art neurosciences critical care unit. The division provides clinical neurophysiology services for the teaching hospitals and several community hospitals. KGH is the site of a Regional Stroke Program, funded by the Ministry of Health.

The division offers a 5-year program in Neurology in accordance with Royal College training guidelines. Members also participate in teaching programs at both the undergraduate and postgraduate level, as well as regional CME Programs.

The incumbent will help develop and direct collaborative clinical and basic science research activities in conjunction with Queen's Centre for Neuroscience Studies (CNS). Queen's is internationally recognized for its integrated approach to basic and clinical neurological research. The newly established CNS draws together over 48 faculty members from across campus with expertise in molecular, cellular, systems, behavioural, cognitive, clinical and perceptual aspects of neuroscience.

An attractive, competitive compensation package is available through an innovative 'alternative funding plan' of Queen's University. Queen's University is top ranked among the "doctorate-level" universities in Canada; successfully merging the intimacy of a medium-sized university with an academic rigor that has enabled members of the Health Sciences Faculty to achieve world-class distinction.

Kingston is situated in the heart of historic Southeastern Ontario on the shores of beautiful Lake Ontario, midway between Toronto and Montréal. The city combines the charms of smaller city living with the cultural advantages conferred by the presence of two universities, a community college, a number of public services institutions, along with reasonable costs of living.

All qualified candidates are encouraged to apply. However, Canadian citizens and permanent residents will be given priority. Queen's University is committed to employment equity and welcomes applications from all qualified candidates, including women, aboriginal people, people with disabilities and visible minorities.

Interested applicants should send a curriculum vitae and the names and addresses of three referees to:

Dr. J. L. McCans
Professor and Head
Department of Medicine
Etherington Hall, Rm. 3041
Queen's University
Kingston, Ontario
Canada K7L 3N6
Fax: (613) 533-6695
Email: koenn@post.queensu.ca

Review of applications will commence on October 1, 2003 and will continue until the position is filled.

Movement Disorder Neurologist

Toronto, Ontario

We are looking for a second Neurologist with clinical expertise in Movement Disorders to join our unique multi-disciplinary group at the Centre for Movement Disorders in the Toronto area. The position primarily includes the provision of clinical services focused on Parkinson's and Huntington's disease together with nurse specialists, neuropsychologist and social work services. Research opportunities will include collaborative studies with the Centre's existing projects in clinical trials and observational studies, epidemiology, positron emission tomography, health service delivery and post-mortem neurochemistry projects as well as the opportunity to initiate new projects and programs. This program is supported by the Ministry of Health and Long-Term Care.

The successful candidate must be eligible to be licensed in the Province of Ontario, to obtain certification by the Royal College of Physicians and Surgeons of Canada as well as being a citizen of Canada or eligible for landed immigrant status.

Please fax curriculum vitae to Mark Guttman MD, FRCPC at 905-472-6270.

NEUROPATHOLOGIST

The Department of Pathology and Laboratory Medicine of the Capital District Health Authority (CDHA) in Halifax, Nova Scotia, seeks a neuropathologist to fill an academic position at this centre.

The CDHA is a tertiary health care institution affiliated with Dalhousie University. It has active neurological and neurosurgical clinical units with strong links to a productive basic neurosciences investigative group. The successful candidate will share the adult neuropathology clinical service for the district with the existing neuropathologist. This role entails handling of brain, spinal cord and peripheral nerve biopsies with performance of intraoperative consultations when relevant. Expertise in diagnostic peripheral nerve and muscle biopsies is required to support an active clinical neuromuscular group. Duties include neuropathological examination of hospital autopsy brains, with performance of specialized neuropathological autopsies in pertinent cases. Participation in a regional neuropathological consultation service would be expected. The successful candidate will contribute to training of anatomic pathology, neurosurgery and neurology residents, and to undergraduate and graduate programs. Significant academic productivity would be anticipated in relation to basic or translational research, including participation in a supervisory capacity in graduate student training, if appropriate.

Canadian certification in neuropathology is a requirement. The successful candidate will hold a faculty position in the Dalhousie University Department of Pathology.

Persons interested in this position are invited to apply to:

Dr Michael Moss,
Professor and Head, Department of Pathology, Room 11B2,
Sir Charles Tupper Medical Building,
5859 University Avenue,
Halifax, NS, B3H 4H7

Canadian Headache Society- GlaxoSmithKline Headache Fellowship

This fellowship has been created to support research and clinical training in the field of headache in Canada. The fellowship is valued at \$50,000 and will be awarded for a one year period. The award will be tenable as of July 1st, 2004.

Candidates must have an MD or PhD degree. Preference will be given to those who have completed a specialty program approved by the Royal College of Physicians and Surgeons of Canada, but others are welcome to apply and will be considered. Applications must contain a research proposal relevant to headache. The proposed research must be done in Canada.

Applications must be received by December 31, 2003.

Further details and instructions for applicants may be obtained from:

Canadian Headache Society
Dr. Michel Aubé, President,
CUSM – Montreal Neurological Hospital
3801 rue Université
Montreal, QC H3A 2B4

Société canadienne des céphalées Bourse de recherche clinique en céphalée

Cette bourse a été créée afin de soutenir la recherche clinique dans le domaine de la céphalée au Canada. D'une valeur de 50 000 \$, la bourse sera attribuée pour une période d'un an et prendra effet le 1er juillet 2004.

Les candidats doivent être titulaire d'un diplôme de médecine ou d'un doctorat de 3^{ième} cycle. Une préférence sera donnée à ceux qui sont inscrits à un programme de spécialité approuvé par le Collège royal des médecins et chirurgiens du Canada. Tous les autres candidats seront les bienvenus et leurs demandes seront considérées. Les demandes doivent contenir un projet de recherche dans le domaine de la céphalée. La recherche proposée doit être entreprise au Canada.

La date limite de réception des demandes de bourse : le 31er décembre 2003.

Pour obtenir plus de précisions, écrire à l'adresse suivante:


Canadian Headache Society
Dr. Michel Aubé, President,
CUSM – Montreal Neurological Hospital
3801 rue Université
Montreal, QC H3A 2B4

OUR GOAL: ZERO PAIN™[†] IN MIGRAINE



- IMITREX® is effective at any stage of migraine pain^{1‡}
- IMITREX® at the first sign of pain reduced most attacks to ZERO^{2‡}
- Nearly 1 out of 2 attacks reduced to ZERO PAIN™ at 1 hour
- 2 out of 3 attacks reduced to ZERO PAIN™ at 2 hours
- 9 out of 10 attacks reduced to ZERO PAIN™ at 4 hours

% of Attacks Reduced to Zero Pain¹⁽¹⁾
(Based on a retrospective Analysis)^{2*}




Adapted from Cady et al. (p<0.05 at 1, 2 & 4 hours)

**2 and 4 hour post dose time points were the primary endpoints.

SUCCESS WITH JUST ONE DOSE²

- When IMITREX® was taken at first sign of pain[‡], 80% of migraine attacks did not require a second dose²
- Reduced need for a second dose = Potential cost savings (acquisition cost only)

% of attacks that did not require a second dose when treated early with IMITREX®



IMITREX® (sumatriptan succinate/sumatriptan) is a selective 5-HT₁ receptor agonist indicated for the acute treatment of migraine attacks with or without aura.¹ IMITREX® is not indicated for prophylactic therapy of migraine, or for the management of hemiplegic, basilar or ophthalmoplegic migraine. Safety and efficacy have not been established for cluster headache.¹

IMITREX® is contraindicated in patients with history, symptoms, or signs of ischemic cardiac, cerebrovascular, or peripheral vascular syndromes, valvular heart disease or cardiac arrhythmias. In addition, patients with other significant underlying cardiovascular diseases should not receive IMITREX®. IMITREX® is also contraindicated in patients with uncontrolled or severe hypertension.

The most common adverse events with IMITREX® 100 mg tablets included: nausea (11.0% vs. 5.8% placebo), malaise/fatigue (9.5% vs. 5.1% placebo), sensations (body regions unspecified) (9.0% vs. 4.5% placebo).

[†] Refers to 0 (zero) on a 4 point pain scale where 0=no pain, 1=mild pain, 2=moderate pain and 3=severe pain.²

¹ IMITREX® should not be used prophylactically. Administration during migraine aura prior to other symptoms occurring may not prevent the development of a headache.²

[‡] Early intervention = treatment initiated at first sign of pain - when pain was mild, before progression to moderate-severe pain.

[#] Based on a retrospective analysis, 92 patients treated 118 headaches at first sign of pain, where the original prospective study did not pre-define this end-point. Further investigation using prospective analysis is required to prove clinical significance.

IMITREX® is a registered trademark, used under license by GlaxoSmithKline Inc. TMused under license by GlaxoSmithKline Inc. Product Monograph available to health care professionals upon request.



*A faster way back.*TM

* Onset of action: 10-15 min. subcutaneous, 15 min. nasal spray, 30 min. tablet.

For brief prescribing information see pages A-29, A-30



Exelon can make a difference in patients with Alzheimer Disease



The only dual cholinesterase inhibitor[†]

EXELON can help enhance cholinergic activity in the brain by inhibiting acetylcholinesterase. In addition, EXELON also inhibits butyrylcholinesterase.

Proven efficacy[‡] in 3 key domains – the ABCs of Alzheimer Disease

Activities of daily living were maintained or improved with a mean difference of more than 3 points vs. placebo on the PDS ($p < 0.05$).[§]

Behaviour and other parameters of global functioning assessed on the CIBIC-Plus were significantly improved vs. placebo ($p < 0.05$).[¶]

Cognitive function was maintained or enhanced by a mean difference of almost 5 points vs. placebo on the ADAS-Cog ($p < 0.001$).^{††}

EXELON can help many of your patients with Alzheimer Disease look forward to staying at home a while longer.

EXELON (rivastigmine as the hydrogen tartrate salt) is indicated for the symptomatic treatment of mild to moderate dementia of the Alzheimer type. EXELON has not been studied in controlled clinical trials for longer than 6 months. There is no evidence that rivastigmine alters the course of the underlying dementing process. EXELON should only be prescribed by (or following a consultation with) clinicians who are experienced in the diagnosis and management of AD.

The most common adverse events (>5%) during the maintenance phase in three pivotal trials were nausea (15%), vomiting (14%), dizziness (10%) and diarrhea (9%).

[†] Comparative clinical significance has not yet been established.

[‡] Based on EXELON dosages of 6-12 mg/day.

[§] Double-blind, randomized, placebo-controlled, international multicentre clinical trial; $n=725$. PDS=Progressive Deterioration Scale.

[¶] Pooled results from three prospective, randomized, double-blind, placebo-controlled, international multicentre clinical trials; $n=2,126$. CIBIC-Plus=Clinician's Interview-Based Impression of Change Scale.

^{††} Prospective, randomized, double-blind, placebo-controlled, clinical trial; $n=699$. ADAS-Cog=Alzheimer Disease Assessment Scale, Cognitive Subscale.

1. Rösler M, Anand R, Cicin-Sain A et al. *BMJ* 1999;318:633-40.

2. Schneider LS, Anand R, Farlow MR. *Int J Ger Psychopharm* 1998;Suppl(1):S1-S34.

3. Corey-Bloom J, Anand R, Veach J. *Int J Ger Psychopharm* 1998;1:55-65.

Product Monograph available upon request.

*Exelon is a registered trademark.
EXE-03-05-7151E



Novartis Pharmaceuticals Canada Inc.
Dorval, Quebec H9S 1A9

www.novartis.ca



DUAL INHIBITING
EXELON[®]
(rivastigmine)

For brief prescribing information see page A-39