

RADIOCARBON IN THE AIR OF CENTRAL EUROPE: LONG-TERM INVESTIGATIONS

I Svetlík¹ • P P Povinec² • M Molnár³ • M Vána⁴ • A Šivo² • T Bujtás⁵

ABSTRACT. Regional levels of radiocarbon have been monitored in order to investigate the impact of fossil fuel combustion on the activity of atmospheric ¹⁴CO₂ in central Europe. Data from atmospheric ¹⁴CO₂ monitoring stations in the Czech Republic, Slovakia, and Hungary for the period 2000–2008 are presented and discussed. The Prague and Bratislava monitoring stations showed a distinct local Suess effect when compared to the Jungfraujoch clean-air monitoring station. However, during the summer period, statistically insignificant differences were observed between the low-altitude stations and the high-mountain Jungfraujoch station. ¹⁴C data from the Hungarian monitoring locality at Dunaföldvár and the Czech monitoring station at Košetice, which are not strongly affected by local fossil CO₂ sources, indicate similar grouping and amplitudes, typical for a regional Suess effect.

INTRODUCTION

Radiocarbon is a long-lived radionuclide (half-life 5730 yr) of global occurrence produced either naturally or artificially. It is produced naturally by interactions of cosmic rays mainly with atmospheric nitrogen in the nuclear reaction ¹⁴N(n,p)¹⁴C. The mean natural ¹⁴C production rate is about 2.5 atoms cm⁻² s⁻¹ at the Earth's surface (UNSCEAR 2000). ¹⁴C is subsequently oxidized to CO₂, which is then transferred to the troposphere, where ¹⁴C becomes a part of the carbon environmental cycle. Inorganic and organic chemical forms of ¹⁴C are deposited into oceanic and terrestrial carbon reservoirs. Although the natural ¹⁴C production and deposition rates have been relatively stable, the ¹⁴C levels in the atmosphere and biosphere during the pre-industrial period showed multiple changes (“wiggles” in the ¹⁴C calibration curve).

In the last century, nuclear weapons tests were important sources of anthropogenic ¹⁴C. Consequently, ¹⁴C concentration in the atmosphere of the Northern Hemisphere was double the natural level in 1963 (Nydal and Lövseth 1965; Meijer et al. 1995). Since the nuclear moratorium on atmospheric nuclear bomb tests was signed in 1963, the ¹⁴C concentration in the atmosphere has been decreasing due to its intensive transfer to oceanic and terrestrial carbon reservoirs (Levin et al. 1980, 1995; Segl et al. 1983; Burchuladze et al. 1989; Hesshaimer et al. 1994; Levin and Kromer 1997, 2004). Currently, ¹⁴C activity is gradually approaching the level that was seen before the nuclear age. The observed ¹⁴C levels give information about carbon ¹⁴C sinks and exchange processes (stratosphere-troposphere, troposphere-ocean, troposphere-biosphere mixing). The decreasing bomb ¹⁴C curve has also been used for dating samples that originated during the nuclear period (Reimer et al. 2004).

In the last few decades, discharges from nuclear facilities (nuclear fuel reprocessing and nuclear power plants) have also been a significant artificial source of ¹⁴C. Their contributions have been estimated to be ~10% of natural production (UNSCEAR 2000). These releases have been increasing ¹⁴C concentration in the atmosphere, and several studies have been devoted to investigating local anthropogenic effects (Chudy and Povinec 1982; Kunz 1985; Cimbák et al. 1986; McCartney et al.

¹Department of Radiation Dosimetry, Nuclear Physics Institute AS CR, Na Truhlárce 39/64, CZ-180 86 Prague, Czech Republic. Corresponding author. Email: svetlik@ujf.cas.cz.

²Faculty of Mathematics, Physics and Informatics, Comenius University, SK-842 48 Bratislava, Slovakia.

³Institute of Nuclear Research of the Hungarian Academy of Sciences (ATOMKI), Bem tér 18/c, 4026 Debrecen, Hungary.

⁴Czech Hydrometeorological Institute, CZ-394 22 Košetice Observatory, Czech Republic.

⁵Paks nuclear power plant, Paks, Hungary.

1986; Obelić et al. 1986; Povinec et al. 1986a,b, 2008; Levin et al. 1988; Hertelendi et al. 1989b; Loosli and Oeschger 1989; Otlet et al. 1990; Uchrin et al. 1992, 1998; Milton et al. 1995; Roussel-Debet et al. 2006).

There is another significant anthropogenic influence on ^{14}C levels in the atmosphere and biosphere, the Suess effect (Suess 1955; Burchuladze et al. 1980; Segl et al. 1983; Levin et al. 1995, 2003; Kuc and Zimnoch 1998; Levin and Hesshaimer 2000). This effect causes a relative decrease of the ^{14}C activity on global, regional, and local scales as a result of the dilution of the carbon isotopic mixture by fossil carbon. ^{14}C monitoring continental stations might shed more light on the (spatial and temporal) distribution of fossil fuel emissions, mainly by providing information about summer versus winter ^{14}C levels in comparison with “nearby” background (clean-air) sampling stations (Levin et al. 1988; Meijer et al. 1995).

SAMPLES AND METHODS

Samples of atmospheric $^{14}\text{CO}_2$ have been collected in 1-month periods at 4 localities in central Europe (see Figure 1). The intentions of monitoring were the following (Svetlik et al. 2006; Molnár et al. 2007):

1. To determine ^{14}C reference levels in the environment and their seasonal and annual changes in locations with lesser (Košetice, Dunaföldvár) and greater (Prague, Bratislava) local anthropogenic influences.
2. To find robust ^{14}C reference parameters, which are minimally influenced by the Suess effect, e.g. spring/summer $^{14}\text{CO}_2$ activities, or activity concentrations of atmospheric $^{14}\text{CO}_2$ in air in the relation to the Suess effect (Svetlik et al. 2010).
3. To provide corrections for ^{14}C dating of various types of samples originating after 1960 (e.g. utilizing CaliBomb, <http://intcal.qub.ac.uk/CALIBomb/>). It can be expected that many of the samples will come from areas affected by a local and regional Suess effect.

Czech Republic (CZ)

Monitoring of atmospheric $^{14}\text{CO}_2$ at the Prague-Bulovka (50°07'N, 14°27'E) site began in 2001. This site is near the border of the urban agglomeration of Prague, in the vicinity of a heavily used motorway; therefore, a local load from fossil fuel combustion can be expected there. The second monitoring station was launched in 2004 at the Košetice Meteorological Observatory (49°35'N, 15°05'E; a part of the Czech Hydrometeorological Institute), which is situated in the Czech-Moravian Highlands. The surroundings of this site have an agricultural-forestry character, without significant local sources of fossil CO_2 .

Monthly CO_2 samples were collected by bubbling the air through a 0.7M NaOH solution in 1.9-L flasks (the final amount of carbonates did not exceed 30% of the saturation capacity of the solution) located at the Prague-Bulovka and Košetice sampling stations. The samples were processed by acid carbonate decomposition, CO_2 purification, benzene preparation, and measurement by a low-background liquid scintillation spectrometer, Quantulus 1220TM (Svetlik et al. 2006; Molnár et al. 2007). For calibration purposes, oxalic acid NIST (National Institute of Standards and Technology, Gaithersburg, USA) SRM 4990C, has been used. A few mL of exposed NaOH solution, collected from samples, were analyzed using a stable isotope mass spectrometer for determination of the $^{13}\text{C}/^{12}\text{C}$ isotopic ratio. Measured data were statistically evaluated and the resulting $^{14}\text{CO}_2$ activities were $\delta^{13}\text{C}$ -corrected and reported as $\Delta^{14}\text{C}$ following Stuiver and Polach (1977).



Figure 1 ¹⁴CO₂ sampling stations in the Czech Republic (1 and 2), Slovakia (3), and Hungary (4)

Slovakia (SK)

Samples in Bratislava were collected on the roof of the Physics building, which has been situated in the center of the town since 1967 (Povinec et al. 1968; Usačev et al. 1973). Since 1975, samples of CO₂ in Bratislava air were collected on the roof (~15 m above the ground) of the Physics building at the new University campus at Mlynská dolina (48°09'N, 17°07'E), about 3 km from downtown. The campus is situated on a small hill about 164 m asl. Bratislava, with almost 500,000 inhabitants, represents an industrialized region that is expected to be influenced by fossil fuel CO₂ emissions, mainly from cars and fossil fuel energy sources. Since 1987, samples of atmospheric CO₂ were collected by bubbling the air through two 2.5-L flasks filled with 0.5M NaOH solution in 1-month periods (the final amount of carbonates did not exceed 54% of the saturation capacity of the solution). The sample from the exposed solution was precipitated as BaCO₃, and CO₂ was liberated in a vacuum line by adding H₃PO₄. The amount of CO₂ gas was measured volumetrically in a calibrated volume. Further, CH₄ was prepared from the CO₂ sample (Povinec 1972) for filling the low-level proportional counter, which was used for counting ¹⁴C decays (Povinec 1978). A few mL of CO₂ collected from samples were analyzed using a stable isotopic mass spectrometer for determination of the ¹³C/¹²C isotopic ratio. ¹⁴C results are presented as Δ¹⁴C values (standard deviations ±5‰) relative to the NIST oxalic acid standard SRM 4990C, and corrected for isotopic fractionation according to convention (Stuiver and Polach 1977).

Hungary (H)

The excess of ¹⁴C in ¹⁴CO₂ and ¹⁴CH₄ chemical forms has been measured in Hungary in the vicinity of the Paks nuclear power plant (NPP) by sampling environmental air since 1991 (Veres et al. 1995).

However, for this work, we used data from another Hungarian sampling site, situated near Dunaföldvár city (46°47'N, 18°57'E), which has been used as a reference environmental monitoring station since 1996. The station is located about 20 km north of the NPP in an agricultural area (Molnár et al. 2007).

^{14}C was collected in the form of CO_2 by bubbling the air through 0.6-L flasks filled with 0.4 L of 3M NaOH solution (the final amount of carbonates did not exceed 15% of the saturation capacity of the solution). To extract CO_2 from the samples, concentrated (75%) sulfuric acid was added to the NaOH solution. The liberated CO_2 was purified over charcoal, then frozen into a CO_2 trap with liquid nitrogen at -196°C , and the remaining non-condensable components were removed by a vacuum pump. The activity of the samples was measured using a gas proportional counter (Csongor et al. 1982; Csongor and Hertelendi 1986; Hertelendi et al. 1989a; Molnár et al. 2010). The standard deviation of a single $\Delta^{14}\text{C}$ measurement applying this method was $\pm 5\%$ (Hertelendi 1990). $\delta^{13}\text{C}$ corrections were measured by a stable isotope mass spectrometer. $\delta^{13}\text{C}$ -corrected $\Delta^{14}\text{C}$ data are given relative to the NIST SRM 4990 (oxalic acid) standard (Stuiver and Polach 1977).

RESULTS

Figure 2 shows the $\Delta^{14}\text{C}$ results from monitoring stations in the Czech Republic, Slovakia, and Hungary. For comparison, data from Jungfraujoch (JFJ) in Switzerland, a high-mountain background (clean-air) monitoring station with only a global Suess effect, are also included (Levin and Kromer 2004; Levin et al. 2008). For estimation of the anthropogenic impact from a local and regional Suess effect, a difference between the Jungfraujoch monitoring station and a local monitoring station was calculated according to the formula:

$$S_{l+r} = \frac{a_{JF} - a_{obs}}{1 + 0.001 \cdot a_{JF}} [\text{‰}] \quad (1)$$

where S_{l+r} is the local and regional Suess effect (reported in per mil of fossil contribution in the carbon isotopic mixture), a_{JF} is the fit with harmonics for the seasonal cycle of the reference ^{14}C activity measured in the Jungfraujoch monitoring station, and a_{obs} is the value observed in a given monitoring station (Nakazawa et al. 1997; Svetlik et al. 2006; Levin et al. 2008). All activities utilized in this formula are given in per mil of $\Delta^{14}\text{C}$ (Stuiver and Polach 1977). Diagrams of the calculated local and regional Suess effect together with mean monthly temperatures are shown in Figure 3.

A local Suess effect may be important in urban areas, in the vicinity of motorways, coal electric power, or heat plants, etc. A regional Suess effect may be prevailing only in areas without the presence of greater local fossil carbon sources, where the local Suess effect is minimal. If relevant reference data from a monitoring station charged predominantly by regional Suess effect are available, a local Suess effect for bigger fossil carbon sources, such as urban areas, can be estimated using a similar formula:

$$S_l = \frac{a_{reg} - a_{obs}}{1 + 0.001 \cdot a_{JF}} [\text{‰}] \quad (2)$$

where S_l is the local Suess effect (reported in per mil of fossil contribution in the carbon isotopic mixture), and a_{reg} is the regional reference activity observed in the corresponding location where local sources of fossil carbon can be neglected (Svetlik et al. 2006). To minimize distortions given by different atmospheric conditions, the time behavior of mean monthly temperature data should be similar in the regional reference monitoring station and in the compared locality. Similarly as for




Figure 2 Atmospheric ¹⁴CO₂ levels in the Czech Republic (a), Slovakia (b), and Hungary (c), compared with the Jungfraujoch Alpine (clean-air) station.




Figure 3 Local and regional Suess effect S_{l+r} in the Prague-Bulovka (a), Bratislava (b), and Dunafoldvár and Košetice stations with similar time behavior given by prevailing regional Suess effect (c), calculated using the Jungfrauoch (JFJ) harmonic fit as reference values (Levin and Kromer 2004; Levin et al. 2008). Data reported in % of fossil C amount from local and regional sources in carbon isotopic mixture.

S_{l+r} , the local Suess effect is normalized to the a_{JF} value (supposing that S_l is a part of S_{l+r} observed in a given site). Using the data presented here, the local Suess effect for the Prague-Bulovka site can be calculated using regional reference values from the Košetice monitoring station (Figure 4). The seasonal cycles of the regional reference data have significantly larger amplitudes than those observed at the JFJ station; hence, most of the time, the high Alpine monitoring station at Jungfraujoch is sampling free tropospheric air (Levin and Kromer 2004; Meijer et al. 2006).




Figure 4 Local Suess effect S_l in the monitoring area Prague-Bulovka, calculated using the Košetice data as reference regional values. Reported in % of fossil C amount from local sources in carbon isotopic mixture.

DISCUSSION

The time behavior of observed ¹⁴C activities from the monitoring localities demonstrates seasonal changes and a gradual interannual decrease (Figure 2). The most distinct seasonal changes are observed in both urban monitoring localities at Prague-Bulovka and Bratislava-Mlynská dolina, where a strong local impact from fossil fuel combustion is expected. The resulting ¹⁴C concentrations at Dunaföldvár are similar to those observed at Košetice, where the local Suess effect is expected to be negligible. The amplitudes of seasonal changes in Dunaföldvár and Košetice are similar; however, they are considerably greater than variations observed at the Jungfraujoch monitoring station.

The time behavior of calculated values S_{l+r} for each locality and S_l for the Prague-Bulovka station highlights seasonal changes, with maximum values during winter (Figures 3 and 4). Emissions of fossil carbon were amplified by abundant occurrence of atmospheric inversions that occurred during the winter months. A lack of atmospheric mixing during an inversion has a much larger influence on the ¹⁴C signal (as well as on the CO₂ concentration) than an increase of fossil CO₂ emissions during winter as a result of enhanced heating. This effect is clearly visible in the diurnal cycle, but also in the seasonal cycle. This implies that the seasonal effect is much less sensitive to CO₂ fossil sources and sinks (photosynthesis) in summer than in winter. This effect is called the “seasonal rectifier effect” (Denning et al. 1999; Palstra et al. 2008).

A comparison with the temperature curve shows that a small fossil carbon asymmetry between the spring and autumn parts seems to be visible, although the temperatures are similar. This effect may be connected with modestly different conditions for fossil CO₂ dissipation in the atmosphere during spring and autumn. The S_{f+r} values observed in the urban Prague-Bulovka site have the greatest amplitudes, which are connected with minimum temperatures. By contrast, the time behavior of greater values of the Suess effect in the urban Bratislava-Mlynská dolina station has a longer duration during each year. Such a dissimilarity could be caused by different dissipation conditions in the surroundings of the Bratislava-Mlynská dolina monitoring station and also partly by higher seasonal stability of fossil CO₂ sources (Povinec et al. 2009). In both urban localities, with larger loads from fossil fuel combustion, the observed Suess effect values (S_{f+r}) seem to be close to or below the significance level during the summer period (i.e. June–July), although frequently motorways, which are relatively stable sources of fossil carbon throughout the year, are situated in the vicinity of these sampling stations (Povinec et al. 2008).

If the data of the CO₂ mixing ratio are absent when calculating fossil CO₂ concentration (in ppm), there is the possibility of calculating the local and regional Suess effect S_{f+r} , corresponding to a percentage (per mil) of fossil carbon amount in the carbon isotopic mixture (from local and regional sources). Using S_{f+r} values, corrections for ¹⁴C activity influenced by fossil carbon sources can be estimated (see Figure 5). Mean monthly values of S_{f+r} from both urban (Prague and Bratislava) and both regional sites (Košetice and Dunaföldvár) were statistically compared (Table 1, paired *t* test, $\alpha = 0.05$, $|t_{obs}| < t_{crit} = > \mu_1 = \mu_2$) and mean values for “cities” and “regions” were subsequently calculated (see Table 2).

Table 1 Statistical comparison of mean monthly S_{f+r} values for monitoring stations Prague-Bulovka, Bratislava-Mlynská dolina, Košetice, and Dunaföldvár (calculated according to Goulden [1956]).

	Prague-Bulovka	Bratislava-Mlynská dolina
Mean (μ)	25.51	25.56
Variance (δ^2)	131.15	47.57
Minimum	8.0	17.2
Maximum	41.6	37.3
Number of observations	12	12
2-sample paired <i>t</i> test for equal means, $\alpha = 0.05$		
$ t_{obs} $		0.0300
Critical value for <i>t</i> (bilateral)		2.2010
	Košetice	Dunaföldvár
Mean (μ)	10.12	10.62
Variance (δ^2)	22.72	31.03
Minimum	4.2	4.0
Maximum	18.9	19.3
Number of observations	12	12
2-sample paired <i>t</i> test for equal means, $\alpha = 0.05$		
$ t_{obs} $		0.4885
Critical value for <i>t</i> (bilateral)		2.2010

If we want to apply the S_{f+r} corrections for dating of samples originating in the last decades, the investigated samples should originate from areas with similar loads from fossil fuel combustion. Such corrections will also depend on a sample ingrowth period. For samples with ingrowth during summer months, such a correction can be insignificant, as evident from S_{f+r} seasonal courses (see Figures 3 and 5). In the case of a continual sample accumulation throughout the year (e.g. into car-

Table 2 Calculated mean values of S_{I+T} for sites with prevailing regional Suess effect (Dunaföldvár and Košetice) and in urban areas (Prague-Bulovka and Bratislava-Mlyská dolina).

Month	Urban areas			Regional sites		
	Average value	Standard deviation	Nr of observations	Average value	Standard deviation	Nr of observations
I	34.0	11.0	15	16.8	8.6	12
II	32.6	10.6	15	11.9	7.3	12
III	27.3	9.8	15	11.2	4.9	12
IV	23.4	10.4	15	6.2	8.1	12
V	23.5	11.8	15	6.1	8.7	12
VI	13.9	8.5	14	5.7	8.3	12
VII	15.1	8.5	15	5.8	10.5	13
VIII	15.1	9.7	16	5.4	8.1	13
IX	18.9	12.1	16	8.6	7.1	13
X	27.9	10.6	16	10.3	8.1	13
XI	36.4	12.5	16	17.6	7.4	13
XII	39.5	6.8	16	19.2	5.4	13




Figure 5 Calculated mean monthly values of S_{I+T} for sites with prevailing regional Suess effect (Dunaföldvár and Košetice) and in urban areas (Prague-Bulovka and Bratislava-Mlyská dolina). The average ratio of S_{I+T} in cities to regional reference sites is 2.64 ± 0.61 , ranging from 2.0 (January) to 3.8 (April and May).

bonate layers), it can be beneficial to estimate the connected uncertainty and to calculate a Suess effect correction (see Table 2). In the case of a uniform sample ingrowth, an annual mean value can be applied. In the opposite case, the weighted mean value can be estimated if changes in a sample accumulation rate during the season are known.

CONCLUSIONS

The results of atmospheric ^{14}C monitoring carried out in the Czech Republic, Slovakia, and Hungary showed that at the Prague-Bulovka and Bratislava-Mlynská dolina monitoring stations, a distinct local Suess effect was observed when compared to the values observed at the Jungfraujoch background monitoring station. This effect has a distinct seasonal behavior. Nevertheless, during the summer period (June to August), small differences were observed between the low-altitude stations and the high Alpine Jungfraujoch station, which is supposed to be minimally affected by local fossil fuel combustion in Europe. The data from the Hungarian monitoring station at Dunaföldvár and the Czech station at Košetice indicate similar grouping and amplitudes, which seems typical for the regional Suess effect.

In general, on the basis of our data it appears that values of $\Delta^{14}\text{C}$ in the summer period are minimally charged by a local and regional Suess effect, even in the sites with relatively intensive local fossil fuel combustion due to better atmospheric mixing. ^{14}C activities during summer are close to Jungfraujoch values with interannual changes corresponding to a linear decrease in the last few years.

From the point of view of ^{14}C dating utilizing a bomb-peak curve, in the areas with greater sources of fossil CO_2 (e.g. from combustion of natural gas, gasoline, oil, and coal) an influence of the local and regional Suess effect can be more significant in the case of relatively uniform yearly carbon accumulation (carbonates) than in the case of samples accumulating carbon mainly during the summer period (biota).

ACKNOWLEDGMENTS

This study was partially supported by the International Atomic Energy Agency, the Scientific Grant Agency of the Slovak Republic VEGA (project No. 1/108/08), institutional funding of Nuclear Physics Institute AS CR (AV0Z 10480505), and by funding of the State Office for Nuclear Safety of the Czech Republic (SUJB 3/2006).

REFERENCES

- Burchuladze AA, Pagava SV, Povinec P, Togonidze GI, Usačev S. 1980. Radiocarbon variations with the 11-year solar cycle during the last century. *Nature* 287(5780):320–2.
- Burchuladze AA, Chudy M, Eristavy IV, Pagava SV, Povinec P, Šivo A, Togonidze GI. 1989. Anthropogenic ^{14}C variations in atmospheric CO_2 and wines. *Radiocarbon* 31(3):771–6.
- Chudy M, Povinec P. 1982. Radiocarbon production in CO_2 coolant of a nuclear reactor. *Acta Universitatis Comenianae, Physica* 22:127–31.
- Cimbák Š, Čechová A, Grgula M, Povinec P, Šivo A. 1986. Anthropogenic radionuclides ^3H , ^{14}C , ^{85}Kr , and ^{133}Xe in the atmosphere around nuclear power reactors. *Nuclear Instruments and Methods in Physics Research B* 17(5–6):560–3.
- Csongor É, Hertelendi E. 1986. Low-level counting facility for ^{14}C dating. *Nuclear Instruments and Methods in Physics Research B* 17(5–6):493–5.
- Csongor É, Szabó I, Hertelendi E. 1982. Preparation of counting gas of proportional counters for radiocarbon dating. *Radiochemical and Radioanalytical Letters* 55:303–7.
- Denning AS, Takahashi T, Friedlingstein P. 1999. Can a strong atmospheric CO_2 rectifier effect be reconciled with a “reasonable” carbon budget? Keynote Perspective. *Tellus B* 51(2):249–53.
- Goulden CH. 1956. *Methods of Statistical Analysis*. 2nd edition. New York: Wiley. p 50–55.
- Hertelendi E. 1990. Sources of random error in the Debrecen radiocarbon laboratory. *Radiocarbon* 32(3):283–7.
- Hertelendi E, Csongor É, Záborszky L, Molnár J, Gál J, Györfi M, Nagy S. 1989a. A counter system for high-precision ^{14}C dating. *Radiocarbon* 31(3):399–406.
- Hertelendi E, Uchrin G, Ormai P. 1989b. ^{14}C releases in various chemical forms with gaseous effluents from the Paks nuclear power plant. *Radiocarbon* 31(3):754–61.
- Hesshaimer V, Heimann V, Levin I. 1994. Radiocarbon evidence for a smaller oceanic carbon dioxide sink than previously believed. *Nature* 370(6486):201–3.
- Kuc T, Zimnoch M. 1998. Changes of the CO_2 sources and sinks in a polluted urban area (southern Poland) over the last decade, derived from the carbon isotope composition. *Radiocarbon* 40(1):417–23.

- Kunz C. 1985. Carbon-14 discharge at three light-water reactors. *Health Physics* 49(1):25–35.
- Levin I, Hesshaimer V. 2000. Radiocarbon—a unique tracer of global carbon cycle dynamics. *Radiocarbon* 46(1):69–80.
- Levin I, Kromer B. 1997. Twenty years of high-precision atmospheric ¹⁴CO₂ observations at Schauinsland station, Germany. *Radiocarbon* 39(2):205–18.
- Levin I, Kromer B. 2004. The tropospheric ¹⁴CO₂ level in mid-latitudes of the Northern Hemisphere (1959–2003). *Radiocarbon* 46(3):1261–72.
- Levin I, Münnich KO, Weiss W. 1980. The effect of anthropogenic CO₂ and ¹⁴C sources on the dilution of ¹⁴C in atmosphere. *Radiocarbon* 22(2):379–81.
- Levin I, Kromer B, Barabas M, Münnich KO. 1988. Environmental distribution and long-term dispersion of reactor ¹⁴CO₂ around two German nuclear power plants. *Health Physics* 54:149–56.
- Levin I, Graul R, Trivett NBA. 1995. Long-term observations of atmospheric CO₂ and carbon isotopes at continental sites in Germany. *Tellus B* 47:23–34.
- Levin I, Kromer B, Schmidt M, Sartorius H. 2003. A novel approach for independent budgeting of fossil fuel CO₂ over Europe by ¹⁴CO₂ observations. *Geophysical Research Letters* 30(23):2194, doi:10.1029/2002GL018477.
- Levin I, Hammer S, Kromer B, Meinhardt F. 2008. Radiocarbon observations in atmospheric CO₂: determining fossil fuel CO₂ over Europe using Jungfraujoch observations as background. *Science of the Total Environment* 391(2–3):211–6.
- Loosli HH, Oeschger H. 1989. ¹⁴C in the environment of Swiss nuclear installations. *Radiocarbon* 31(3):747–53.
- McCartney P, Baxter MS, McKay K, Scott EM. 1986. Global and local effects of ¹⁴C discharges from the nuclear fuel cycle. *Radiocarbon* 28(2A):634–43.
- Meijer HAJ, van der Plicht J, Gislefoss JS, Nydal R. 1995. Long-term atmospheric records near Groningen, Fruholmen, and Izaña. *Radiocarbon* 37(1):39–50.
- Meijer HAJ, Pertuisot MH, van der Plicht J. 2006. High-accuracy ¹⁴C measurements for atmospheric CO₂ samples by AMS. *Radiocarbon* 48(3):355–72.
- Milton GM, Kramer SJ, Brown RM, Repta CJW, King KJ, Rao RR. 1995. Radiocarbon dispersion around Canadian nuclear facilities. *Radiocarbon* 37(2):485–96.
- Molnár M, Bujtás T, Svingor É, Futó I, Svetlik I. 2007. Monitoring of atmospheric excess ¹⁴C around Paks nuclear power plant, Hungary. *Radiocarbon* 49(2):1031–43.
- Molnár M, Haszpra L, Svingor É, Major I, Svetlik I. 2010. Atmospheric fossil fuel CO₂ measurement using a field unit in a central European city during the winter of 2008/09. *Radiocarbon* 52(2–3):835–45.
- Nakazawa T, Ishizawa M, Higuchi K, Trivett NBA. 1997. Two curve fitting methods applied to CO₂ flask data. *EnvironMetrics* 8(3):197–218.
- Nydal R, Lövsæth K. 1965. Distribution of radiocarbon from nuclear tests. *Nature* 206(4988):1029–31.
- Obelić B, Krajcar-Bronić I, Srdoč D, Horvatinčić N. 1986. Environmental ¹⁴C levels around the 632MWe nuclear power plant Krško in Yugoslavia. *Radiocarbon* 28(2A):644–8.
- Otlet RL, Walker AJ, Fulker MJ. 1990. Survey of the dispersion of ¹⁴C in the vicinity of the UK reprocessing site at Sellafield. *Radiocarbon* 32(1):23–30.
- Palstra SWL, Karstens U, Streurman HJ, Meijer HAJ. 2008. Wine ethanol ¹⁴C as a tracer for fossil fuel CO₂ emissions in Europe: measurements and model comparison. *Journal of Geophysical Research* 113(D21):D21305, doi:10.1029/2008JD010282.
- Povinec P. 1972. Very low background proportional counter for tritium dating. *Nuclear Instruments and Methods* 101(3):613–4.
- Povinec P. 1978. Multiwire proportional counters for low-level ¹⁴C and ³H measurements. *Nuclear Instruments and Methods in Physics Research* 156(3):441–5.
- Povinec P, Saro S, Chudy M, Seliga M. 1968. Rapid method of carbon-14 counting in atmospheric carbon dioxide. *International Journal of Applied Radiation and Isotopes* 19(12):877–81.
- Povinec P, Šivo A, Chudy M. 1986a. Seasonal variations of anthropogenic radiocarbon in the atmosphere. *Nuclear Instruments and Methods in Physics Research B* 17(5–6):556–9.
- Povinec P, Chudy M, Šivo A. 1986b. Anthropogenic radiocarbon: past, present and future. *Radiocarbon* 28(2A):668–72.
- Povinec PP, Šivo A, Šimon J, Holy K, Chudy M, Richtáriková M, Morávek J. 2008. Impact of the Bohunice Nuclear Power Plant on atmospheric radiocarbon. *Applied Radiation and Isotopes* 66(11):1686–90.
- Povinec PP, Chudy M, Šivo A, Šimon J, Holy K, Richtáriková M. 2009. Forty years of atmospheric radiocarbon monitoring around Bohunice nuclear power plant, Slovakia. *Journal of Environmental Radioactivity* 100(2):125–30.
- Reimer PJ, Brown TA, Reimer RW. 2004. Discussion: reporting and calibration of post-bomb ¹⁴C data. *Radiocarbon* 46(3):1299–304.
- Roussel-Debet S, Gontier G, Siclet F, Fournier M. 2006. Distribution of carbon 14 in the terrestrial environment close to French nuclear power plants. *Journal of Environmental Radioactivity* 87(3):246–59.
- Segl M, Levin I, Schoch-Fischer H, Münnich M, Kromer B, Tschiersch J, Münnich KO. 1983. Anthropogenic ¹⁴C variations. *Radiocarbon* 25(2):583–92.
- Stuiver M, Polach HA. 1977. Discussion: reporting of ¹⁴C data. *Radiocarbon* 19(3):355–63.
- Suess HE. 1955. Radiocarbon concentration in modern wood. *Science* 122(3166):415–7.

- Svetlik I, Molnár M, Svingor E, Futó I, Pintér T, Rulík P, Michálek V. 2006. Monitoring of atmospheric $^{14}\text{CO}_2$ in central European countries. *Czechoslovak Journal of Physics* 56D(4):291–7.
- Svetlik I, Povinec PP, Molnár M, Meinhardt F, Michálek V, Simon J, Svingor E. 2010. Estimation of long-term trends in the tropospheric $^{14}\text{CO}_2$ activity concentration. *Radiocarbon* 52(2–3):815–22.
- Uchrin G, Csaba E, Hertelendi E, Ormai P, Barnabás I. 1992. ^{14}C release from a Soviet-designed pressurized water reactor nuclear power plant. *Health Physics* 63: 651–5.
- Uchrin G, Hertelendi E, Volent G, Slávik O, Morávek J, Kobal I, Vokal B. 1998. ^{14}C measurements at PWR-type nuclear power plants in three middle European countries. *Radiocarbon* 40(1):439–46.
- UNSCEAR (United Nations Scientific Committee on the Effects of Atomic Radiation). 2000. Exposures from natural and man-made sources of radiation. Report to the General Assembly 1. New York: United Nations.
- Usačev S, Povinec P, Chudy M, Šeliga M. 1973. Bratislava radiocarbon measurements I. *Radiocarbon* 15(3):443–50.
- Veres M, Uchrin G, Hertelendi E, Csaba E, Barnabás I, Ormai P. 1995. Concentration of radiocarbon and its chemical forms in gaseous effluents, environmental air, nuclear waste and primary water of a pressurized water reactor power plant in Hungary. *Radiocarbon* 37(2):497–504.