

P01-236 - ADHD SYMPTOMATOLOGY IN BIPOLAR DISORDER AND DESCENDANT

M.D. Ortega García¹, C. Martínez Martínez², R. Gómez Martínez², C. Dominguez Martín³, C. Salazar Li¹, G. Rodríguez Carretero³, P. Blanco del Valle⁴, A. Agúndez Modino⁵, M.A. Alonso de la Torre², A. Rodríguez Campos⁶

¹Psychiatry, Complejo Hospitalario La Mancha Centro, Alcazar de San Juan, ²Psychiatry, Complejo Hospitalario de León, Leon, ³Psychiatry, Hospital Clínico Universitario de Valladolid, Valladolid, ⁴Psychiatry, Centro de Salud Mental Nou Barris, Barcelona, ⁵Psychiatry, Hospital Universitario Río Hortega, Valladolid, ⁶Departamento Psiquiatría, Complejo Hospitalario León, Leon, Spain

Objective: Describe Attention Deficit Hyperactive Disorder's (ADHD) prevalence in Bipolar Disorders (BD) and relatives.

Methods: 78 admissions for Bipolar Disorder (DSM-IV) in Inpatient Psychiatric Unit, in *Hospital Clínico Universitario* of Valladolid (Spain). Only 36/78 patients participate in study. Demographic, social and clinical information were registered. ADHD symptomatology was evaluated from patient and descendant (Conners short version).

Results: ADHD symptomatology suggestive in childhood/adolescence were detected in 13,9% (5/36). Conners score were negative (below 15) in all case.

ADHD symptomatology suggestive in their children were detected in 6,25% (n=3). Conner score were positive in 2,1%. Family psychiatry history in 72,2% (n=26), affective disorder in 60,52% (n=23). No family history with ADHD diagnosis. Only one case (2,8%) with symptomatology suggestive of ADHD in relatives.

Conclusions: The ADHD prevalence in our sample of BD and relatives weren't higher than general population.

Bibliography:

- *Frontiers Between Attention Deficit Hyperactivity Disorder and Bipolar Disorder*. Cathryn A. Galanter, MDa, Ellen Leibenluft, MD. *Child Adolesc Psychiatric Clin N Am* 17 (2008) 325-346.

- *Co-occurrence of bipolar and attention-deficit hyperactivity disorders in children*.

Singh MK, DelBello MP, Kowatch RA, Strakowski SM. Review Article Bipolar Disorders 2006: 8: 710-720