

The non-synchronous response of Rabots Glaciär and Storglaciären, northern Sweden, to recent climate change: a comparative study

Keith A. BRUGGER

*Geology Discipline, University of Minnesota, Morris, 600 E. 4th Street, Morris, MN 56267, USA
E-mail: bruggeka@morris.umn.edu*

ABSTRACT. Rabots Glaciär and Storglaciären, two small valley glaciers in the Swedish Arctic, have not behaved synchronously in response to recent climate change. Both glaciers advanced late in the 19th century and then began to retreat in response to a $\sim 1^\circ\text{C}$ warming that occurred around 1910. By the mid-1980s the terminus and volume of Storglaciären had essentially stabilized, so it may have completed its response to the earlier warming. In contrast, ongoing thinning and retreat of Rabots Glaciär are substantial and suggest its response time is considerably longer. A time-dependent numerical model was used to investigate each glacier's response to perturbations in mass balance. This modeling suggests that, for small perturbations, volume timescales for Storglaciären and Rabots Glaciär are ~ 125 and ~ 215 years, respectively. Another measure of response time (i.e. length response time) yields somewhat lower values for each glacier; however, what is significant is that by either measure and accounting for uncertainties, the response time for Rabots Glaciär is consistently about 1.5 times longer than that for Storglaciären. This implies that their non-synchronous behavior is likely due to differences in response times. The latter ultimately result from markedly different longitudinal geometries (particularly near the termini), velocity profiles and specific net balance gradients.

INTRODUCTION

Projections have suggested that global warming will disproportionately affect polar regions (e.g. Houghton and others, 2001; Holland and Bitz, 2003); consequently the behavior of Arctic (and high-latitude) glaciers may be among the first clear indicators of the onset and severity of climate change. Some recent observations and analyses of glacier behavior (e.g. Arendt and others, 2002; Meier and others, 2003; Oerlemans, 2005) support this conclusion. Other studies, however, point to a lack of coherent regional trends or otherwise suggest that the behavior of some Arctic glaciers with respect to enhanced greenhouse warming is equivocal (Dowdeswell and others, 1997; Dyrgerov and Meier, 2000; Serreze and others, 2000; Abdalati and others, 2004). To reduce this apparent ambiguity, therefore, improvements in the observational database are needed, specifically increases in both the number and geographic distribution of those glaciers being monitored (Dyrgerov and Meier, 2000), and in understanding how glacier behavior may lag climatic forcings (e.g. Oerlemans and others, 1998; Oerlemans, 2005). Because of the latter, however, glaciers within a particular geographic area do not always show a consistent pattern of advance and retreat. Such non-synchronous response to climate change is exemplified by the behaviors of Rabots Glaciär and Storglaciären, two small valley glaciers in the Swedish Arctic (Fig. 1). The purpose of this paper is to present the results of numerical modeling that indicate this non-synchronous behavior is due, in large part, to differences in individual glacier dynamics, and ultimately in their response times.

RECENT BEHAVIOR OF THE GLACIERS

Changes in glacier length and ice volume for Rabots Glaciär and Storglaciären were compared by Brugger and others

(2005) and are summarized in Figure 2. Both glaciers advanced late in the 19th century and then retreated in response to a warming of $\sim 1^\circ\text{C}$ at the beginning of the 20th century (Holmlund, 1987, 1993; Holmlund and Jansson, 1999). The terminus position and geometry of Storglaciären more-or-less stabilized in the mid-1980s, by which time ice volume had decreased by about $131 \times 10^6 \text{ m}^3$. Subsequently the glacier experienced a slight increase in volume without any significant change in terminus position. These post-1980 volume increases may be due to contemporaneous increases in (winter) precipitation and the development of a more maritime climatic regime in the region (Dowdeswell and others, 1997; Pohjola and Rogers, 1997). The reaction of Rabots Glaciär in response to the same




Fig. 1. Location of Storglaciären and Rabots Glaciär. Note the similarity in size of the glaciers. Dotted lines on the glaciers show the approximate positions of the velocity profiles shown in Figure 3.




Fig. 2. Measured and/or estimated changes in ice volume and terminus position of Storglaciären and Rabots Glaciär. Data for volume changes on Storglaciären are from Holmlund (1987, 1988) and updated using recent mass-balance measurements. Volume changes are estimated prior to 1946. Volume changes for Rabots Glaciär (shown with uncertainties) are estimated largely from measured changes in ice surface elevations or those depicted in maps (see Brugger and others, 2005). All other data are from unpublished reports of the Tarfala Research Station and Brugger (1992). (After Brugger and others, 2005.)

warming lagged behind that of Storglaciären, and its initial rates of retreat were low ($\sim 2 \text{ m a}^{-1}$). Retreat generally increased to $>11 \text{ m a}^{-1}$ by 1959 and has remained relatively constant thereafter. By the mid-1980s, concomitant reductions in the volume of Rabots Glaciär ($\sim 135 \times 10^6 \text{ m}^3$) were comparable to those of Storglaciären, but more noteworthy is the fact that since then the glacier has lost an additional $20 \times 10^6 \text{ m}^3$ of ice. Thus as noted by Holmlund (1988, 1993) and Brugger (1997), it appears that Storglaciären is, or was, in quasi-equilibrium, at least with respect to mass/volume changes, while Rabots Glaciär continues to adjust to the earlier climatic warming.

GLACIER CHARACTERISTICS AND MASS BALANCES

Longitudinal profiles based on their 1980 geometries and surface velocities of Storglaciären and Rabots Glaciär are contrasted in Figure 3a and b. Mean ice thickness (~ 100 and 85 m respectively), maximum thickness (250 and 175 m), and thickness variations are all greater for Storglaciären, resulting in distinctly different variations of horizontal velocities. This is perhaps the first hint that these two glaciers should behave rather differently. Hypsometries of the glaciers (Fig. 3c) are similar.

The consistent increases of summer over annual velocities on Rabots Glaciär seen in Figure 3b are attributed to meltwater production that facilitates basal sliding (Brugger, 1992). More pertinent for the present study, however, is the close similarity of the winter and annual velocity profiles. The assumption made here, as it has been elsewhere (e.g. MacGregor and others, 2005), is that basal sliding is minimal during the winter, so the contribution of sliding to annual velocities is negligible. Measured seasonal variations of surface velocities on Storglaciären (Hooke and others,

1989; R.LeB. Hooke, unpublished data) suggest that sliding might account for 5–10% of annual velocities. Moreover, Hanson (1995) and Albrecht and others (2000) found that little or no sliding was required to fit modeled surface velocities to those observed on Storglaciären.

Very few meteorological or climatological data are available for the valley of Rabots Glaciär, making it difficult to directly compare the respective microclimates under which each glacier exists. Alternatively, specific net mass-balance curves (Fig. 3c) might be compared, as these essentially capture the meteorological parameters (temperature, precipitation, radiation balances and so forth) relevant to glacier regime. Using the method of Kuhn (1984), Brugger (1992) showed that the specific net-balance curves for each glacier are generally similar in form from year to year (see also Rasmussen, 2004). It is obvious (Fig. 3c) that mass-balance gradients (db/dz , where b is specific net mass balance and z is altitude) differ between the glaciers. The steeper gradient on Storglaciären ($\sim 0.008 \text{ m w.e. m}^{-1}$) indicates a greater mass turnover, and therefore perhaps also hints at its quicker response time and greater climatic sensitivity (Oerlemans, 2005). In contrast, db/dz on Rabots Glaciär ($\sim 0.005 \text{ m w.e. m}^{-1}$) suggests a slower response time.

Interannual variations in the net mass balance of the glaciers are well correlated ($r^2 = 0.78$, $n = 23$, $p < 0.0001$). Thus one might expect the climatic ‘events’ that forced the retreat of the glacier from their ~ 1910 maximums were synchronous; in other words, the glaciers reacted to a regional climate signal. This is corroborated by historical evidence (photographs, unpublished reports, etc.) that indicates both glaciers began retreating at the same time. Because of local filtering of that regional signal (by interactions between/among topography and local energy and mass balances), it could be argued that each glacier experienced climate change of different magnitude. Regressions between net mass balance and temperature (e.g. Brugger, 1992; Holmlund and others, 1996) suggest, however, that the climatic forcing induced by the warming was comparable for both glaciers, being on the order of -0.3 to $-0.4 \text{ m w.e. a}^{-1}$. The underlying assumption is that no strong feedback occurs between changes in glacier geometry (particularly surface elevation) and mass balance (cf. Harrison and others, 2001).

Glacier geometry and mass balance allow first-order estimates of response time τ to be calculated in several ways. Jóhannesson and others (1989a) presented a simple geometric argument that suggests

$$\tau = \frac{\bar{h}_0}{|b_0(l_0)|}, \quad (1)$$

where \bar{h}_0 is a thickness scale (variously taken as mean ice thickness, maximum ice thickness or some other measure) and $b_0(l_0)$ is the mass balance at the terminus that characterizes the glacier’s datum, or equilibrium state (subscript ‘0’). However, in light of potential difficulties in assigning precise datum-state values of \bar{h}_0 and $b_0(l_0)$ for both Storglaciären and Rabots Glaciär, a simplified formulation of the form

$$\tau_L = 13.6 \beta_0^{-1} s_0^{-1} (1 + 20s_0)^{-1} L_0^{-\frac{1}{2}} \quad (2)$$

is used (Oerlemans, 2001, 2005), where β is the mass-balance gradient, db/dz , s is the surface slope and L is the glacier length. Here τ_L is a ‘length response time’ (Oerlemans, 2001). As noted previously, it is reasonable to assume

the mass-balance gradients are relatively constant, and furthermore that the 1910 geometries of each glacier reflected some quasi-equilibrium. Thus for Storglaciären and Rabots Glaciär, respectively, representative values for β_0 are ~ 0.008 and $0.005 \text{ m w.e. m}^{-1}$, $s_0 \sim 0.130$ and 0.115 , and $L_0 \sim 4065$ and 4610 m yield estimates of τ_L of ~ 60 and 105 years. Despite the generality of Equation (2), these results imply a significantly longer (length) response time for Rabots Glaciär.

THE MODEL

A one-dimensional, time-dependent flow model was used to estimate the response times of the glacier to small perturbations in mass balance. The model, discussed in detail by Bindschadler (1982), is a finite-difference approximation to the continuity equation

$$\frac{\partial S}{\partial t} + \frac{\partial Q}{\partial x} = bW, \tag{3}$$

where Q is the volume flux through a glacier cross-section of area S , t is time, x is the center-line distance, b is the specific net balance and W is glacier width. Net mass balance is subsequently assumed to consist of a steady- or datum-state value b_0 upon which is superimposed a perturbation b_1 . Glacier width and cross-sectional area are parameterized as a function of ice thickness H along x using the empirical relations

$$W = DH^{\frac{1}{2}} + EH \tag{4}$$

and

$$S = \frac{2}{3}DH^{\frac{3}{2}} + \frac{1}{2}EH^2 + F. \tag{5}$$

The constants D , E and F as well as values for H and W were determined from existing topographic maps and radio-echo soundings. Volume flux is related to the center-line surface velocity U_s through use of a flux shape factor f^* that yields a velocity averaged over S , hence

$$Q = f^* U_s S. \tag{6}$$

Surface velocity is

$$U_s = \frac{2A}{n+1} (f\rho g \sin \alpha) H^{n+1}, \tag{7}$$

where A and n ($=3$) are flow-law parameters, f is a shape factor used to account for drag imparted by valley walls (Nye, 1965), ρ is the density of ice ($=900 \text{ kg m}^{-3}$), g is gravitational acceleration ($=9.8 \text{ ms}^{-2}$) and α is the slope of the ice surface. In the model, surface slopes were longitudinally averaged over distances substantially greater than ice thicknesses. Equation (7) ignores basal sliding, but, as discussed above, sliding is not thought to be important in the long-term mean velocities of either glacier. However, this is a point that is returned to in a subsequent section of this paper.

The finite-difference implementation together with specified boundary conditions uses an iterative predictor-corrector technique to approximate the true solution to Equations (3–7) within some defined level of accuracy (expressed in terms of maximum allowable changes between iterations). Only the lower reach of each glacier could be accurately modeled because of both the complications arising from the confluence of tributaries, and, in the case of Rabots Glaciär, sparse velocity and mass-balance data for the tributaries. Thus a constant-head boundary condition (Bindschadler, 1980, 1982; Bindschadler and Rasmussen, 1983; Brugger, 1997) was imposed such that




Fig. 3. (a, b) Longitudinal thickness and velocity profiles for (a) Storglaciären (see Hooke and others, 1989) and (b) Rabots Glaciär (Brugger, 1992). Thickness profiles are based on 1980 glacier geometries, and velocities were measured in the mid-1980s. Approximate positions of the velocity profiles are shown in Figure 1. All vertical scales and horizontal scales are the same. (c) Individual glacier hypsometry and average mass balance for the interval 1981–2003. (Data from unpublished reports of the Tarfala Research Station.) Net-balance gradients are based on linear regressions.

the volume flux is

$$Q_{\text{head}} = A_{u_0}(b_0 + b_1), \tag{8}$$

where A_{u_0} is the surface area in the datum state up-glacier from the modeled reach. While this boundary condition

Table 1. Model-derived volume timescales calculated for three mass-balance perturbations

	Storglaciären			Rabots Glaciär			
A_0 , model area (m^2)	3.46×10^6			3.91×10^6			
b_1 , mass-balance perturbation (m w.e. a^{-1})	-0.01	-0.03	-0.05	-0.01	-0.03	-0.05	
Terminus retreat (m)	108	296	368	138	296	437	
$V_1(\infty)$, final volume change (10^6 m^3) in model section only	-4.06	-10.72	-16.75	-6.25	-17.52	-29.97	
τ_v , volume timescale (years) from Equation (9)	94	95	89	127	125	136	
Mean and std dev.	93 \pm 3			129 \pm 6			
Ratio (Mean _{Rabots} /Mean _{Storglaciären})							1.4
Estimated additional volume change (10^6 m^3) up-glacier of model boundary	-0.67	-2.01	-3.35	-1.86	-7.12	-14.06	
$V_1(\infty)$, total final volume change (10^6 m^3)	-4.73	-12.73	-20.10	-8.11	-24.67	-44.03	
τ_v , volume timescale (years) from Equation (9)	137	127	116	207	216	225	
Mean and std dev.	127 \pm 11			214 \pm 10			
Ratio (Mean _{Rabots} /Mean _{Storglaciären})							1.7
$V_1(\infty)$, total volume change (10^6 m^3) in model with 10% sliding	-4.38	-11.99	-19.52				
τ_v , volume timescale (years) from Equation (9)	127	116	113				
Mean and std dev.	119 \pm 7						
$V_1(\infty)$, total volume change (10^6 m^3) in model with 50% sliding	-3.22	-8.74	-14.54				
τ_v , volume timescale (years) from Equation (9)	93	84	84				
Mean and std dev.	87 \pm 5						

preserves mass continuity, it does not capture the time-dependent behavior of the unmodeled sections of the glaciers. Instead it implies that these regions of the glaciers respond instantaneously to mass-balance perturbations, whereas in reality changes in geometry upstream of the boundary would result in changes in volume flux that would propagate more slowly through each glacier. Consequently, to minimize the influence of this boundary condition on glacier behavior, small mass-balance perturbations were used so that glacier geometry (i.e. H and α) remains relatively constant in this region and thus the volume flux imposed is comparable to that which would be given by Equation (6). Nevertheless, the potential effects of this boundary condition are evaluated below.

Datum states for the glaciers were defined by their mid-1980s geometries, as these correspond to a time when detailed measurements of the surface velocity field and mass balance of Rabots Glaciär were carried out (Brugger, 1992). Values for f were tuned in accordance with measured surface velocities (Fig. 3a and b). Because of this tuning, modeling results are not sensitive to the choice of the flow-law parameter A . Once tuned, a set of $b_0(x)$ for each glacier was found that would maintain these geometries, and the model was then run (with $b_1(x) = 0$) to insure a steady-state configuration. For that reason, minor adjustments in initial glacier geometry occur, but differences between actual and modeled datum-state ice thicknesses did not exceed 0.5 m except at the termini where they were on the order of 5 m. It bears mentioning that the values of $b_0(x)$ required to maintain datum-state geometry of Storglaciären were in most cases comparable to the values shown in Figure 3c, again suggesting that the glacier is close to being in equilibrium with existing climate. $b_0(x)$ values required to preserve the mid-1980s geometry of Rabots Glaciär were generally 0.5–0.75 m w.e. a⁻¹ more positive than those shown in Figure 3c, except close to the terminus where they are ~ 1.25 m w.e. a⁻¹ more positive. It must be emphasized that the particular datum-state glacier geometry, mass-balance parameters and tuning criteria were selected

in order to determine the response times of the glaciers. No attempt is made here to either simulate historic variation in glacier length (and volume) or predict future behavior in response to ongoing climate warming.

RESPONSE TIMES DEDUCED FROM NUMERICAL MODELING

Steady-state mass balances were uniformly ($b_1 \neq f(z)$) altered using step-like perturbations of -0.01, -0.03 and -0.05 m w.e. a⁻¹, and the model was allowed to run until new steady-state profiles were established. In practice, steady state was achieved when changes in ice thickness between successive iterations were <1%. Results of the modeling experiments are summarized in Table 1. Figure 4 shows examples of the temporal evolution of glacier geometry in response to such perturbations (in this case, $b_1 = 0.05$ m w.e. a⁻¹) and the corresponding changes in terminus position. These examples are representative of all modeling runs, but also show that even for the largest b_1 used the change in ice thicknesses at the head boundary is small for Storglaciären ($\sim 2\%$) and slightly larger for Rabots Glaciär ($\sim 10\%$). From Figure 4 it can be seen that the new equilibrium profile of Storglaciären is established more quickly than that for Rabots Glaciär. However, the time periods required to achieve new steady state (varying between ~ 200 and 500 years) are, to some extent, an artifact of the modeling and reflect the termination of the approach to equilibrium specified by desired level of accuracy. These numbers are therefore less significant for the present discussion.

What is more significant is that a characteristic response time for each glacier can be estimated from modeled changes in volume. Expressed in terms of a volume timescale τ_v (Jóhannesson and others, 1989b), the response time is

$$\tau_v = \frac{V_1(\infty)}{A_0 b_1}, \quad (9)$$

where $V_1(\infty)$ is the change in ice volume in the glacier's final steady state ($t = \infty$) and A_0 is the total area of the glacier in its datum state. Although thickness changes at the model boundaries are relatively small (Fig. 4), they do imply volume changes up-glacier, which might be especially significant for Rabots Glaciär. Volume timescales calculated directly from Equation (9) using only volume changes over the model reaches are therefore *minimum* estimates. Mean values for τ_v for Storglaciären and Rabots Glaciär (Table 1) are 106 ± 10 and 154 ± 6 years respectively.

Obviously, more realistic estimates of each glacier's volume timescale need to account for volume loss in the areas up-glacier of the head boundaries that could not be modeled. These quantities were thus estimated by first extrapolating final thickness changes in the model reaches into the unmodeled regions. Excluding the large changes that occur near the termini, these vary linearly with distance ($r^2 > 0.98$). When multiplied by the (unmodeled) area affected by these changes and added to modeled volume losses, better approximations of $V_1(\infty)$ are obtained (Table 1). This procedure tends, however, to overestimate volume losses in the unmodeled regions for two principal reasons. Firstly it ignores the small reductions in surface area that would be expected there. Secondly, in the regions of tributary confluence, the effective cross-sectional area (defined as being perpendicular to flow) is wider than it is down-valley. Decreased ice flux could therefore be accommodated by smaller decreases in ice thickness. If instead decreased flux were wholly accommodated by decreases in velocity, *no* decreases in thickness would occur. Using these estimates of additional volume loss thus yields approximate upper limits to the calculated response times.

Not surprisingly, volume losses in the unmodeled area calculated in this manner are significantly larger for Rabots Glaciär and increase total volume loss by 30–47%. For Storglaciären, increases in total volume loss are more modest, ranging from about 17% to 20%. Concomitant increases in each glacier's response time are given in Table 1. The modeling thus suggests that the volume timescale of Storglaciären to step-like perturbations in mass balance is on the order of 120–140 years. Rabots Glaciär, on the other hand, is characterized by a longer response time of ~210–230 years. The difference in the glaciers' volume timescales using these estimates is substantially greater than that determined when changes in volume up-glacier of the head boundary are ignored (these being about 90 and 50 years respectively). This suggests that the effect of including the latter *increases* the calculated difference between τ_v for Storglaciären and Rabots Glaciär. Because of the aforementioned uncertainties involving the boundary condition, it is the relative magnitudes of τ_v that are deemed important here rather than the actual values. As a measure of response, then, the volume timescale for Rabots Glaciär is at least ~1.5 times that of Storglaciären, and perhaps as much as ~1.7 times greater.

DISCUSSION AND CONCLUSIONS

It should be stressed that the volume timescales determined in this study are derived from a relatively simple model simulating a hypothetical and highly idealized climate-change scenario. Nevertheless, it is worthwhile to briefly consider the significance of τ_v estimated here with the observed recent behavior of Storglaciären in particular,




Fig. 4. An example of the temporal evolution of the new steady-state profiles for (a) Storglaciären and (b) Rabots Glaciär in response to a step-like perturbation in mass balance b_1 (here $b_1 = -0.05$ m w.e. a^{-1}). Intermediate profiles are shown at 50 year intervals. All vertical scales and horizontal scales are the same. Insets show details of the terminus retreat.

since measurements of it are more abundant and detailed. As noted previously, it has been suggested that Storglaciären is in quasi-equilibrium with existing climate (Holmlund, 1988, 1993; Brugger, 1992, 1997). If it is also taken to be true that the glacier was in or near steady state at the 1910 maximum then its response time, defined by Nye (1960) as the time required for glacier geometry to be within $1 - (1/e)$ of its final steady-state value, is estimated to be ~45 years ($= 0.63 \times (1980 - 1910)$; Holmlund, 1988). Fitting the measured (or estimated, as opposed to modeled) time variations of volume $V_1(t)$ (Holmlund, 1988) to

$$V_1(t) = V_1(\infty) \left(1 - e^{-\frac{t}{\tau_v}} \right) \quad (10)$$

(Jóhannesson and others 1989b) might also yield estimates of the glacier's volume timescale. Again under the assumption that Storglaciären is now in steady state and $V_1(\infty) \sim 1.31 \times 10^8$ m³, τ_v is ~33 years (Fig. 5). If it is assumed based on the foregoing discussion that τ_v is ~45 years, then $V_1(\infty)$ is $\sim 1.57 \times 10^8$ m³. Allowing both τ_v and $V_1(\infty)$ to be free parameters, a best fit to Equation (10) is obtained with $V_1(\infty) \sim 1.89 \times 10^8$ m³ and $\tau_v \sim 63$ years. For comparison,




Fig. 5. Observed or estimated variations in ice volume for Storglaciären fitted to an equation of the form $V_1(t) = V_1(\infty) \times (1 - e^{-t/\tau_v})$. The different curves correspond to different assumptions made regarding the value of $V_1(\infty)$ or τ_v given in the explanation; curves are labeled according to estimated parameters (see text for discussion).

requiring τ_v to be ~ 127 years as suggested by the modeling results, $V_1(\infty)$ would be $\sim 3.02 \times 10^8 \text{ m}^3$. Although all the fitted curves shown in Figure 5 are statistically significant ($r^2 > 0.92$), none appear to fit the observations very well.

There are a number of factors that are likely to be contributing to the discrepancy between these estimates of τ_v and that based on the modeling in this study (and others made for Storglaciären (e.g. Raper and others, 1996; Hooke, 2005, p. 381)), the simplicity of the model notwithstanding. These include:

1. Climatic change early in the 20th century was only approximately step-like, and subsequent variations have occurred. Thus the glacier's behavior undoubtedly reflects a more complex forcing (cf. Raper and others, 1996).
2. That Storglaciären was in equilibrium around 1910 is questionable (Stroeven, 1996), therefore introducing the potential for yet more complexity in its response. In particular, Brugger and others (2005) speculated that had Storglaciären not fully completed its response to the climate change driving its Little Ice Age advance, (a) the onset of retreat would have been delayed, and (b) the actual magnitude of post-1910 retreat and reductions in volume would be less than what might be expected. In effect, this would reduce both $V_1(\infty)$ and the apparent time to complete its response to the $\sim 1^\circ\text{C}$ warming during the early 20th century.
3. The mass-balance perturbation associated with this warming was probably not small (Hooke, 2005). As stated previously, this perturbation was probably about $-0.4 \text{ m.w.e. a}^{-1}$, nearly an order of magnitude larger than the largest used in the modeling experiments. This is significant because Equation (9) strictly holds for small perturbations only (Jóhannesson and others, 1989b), so

volume timescales derived from observed volume changes, driven by such a large forcing, might not be compatible with those determined from the modeling in the present study.

4. The characteristic response time (or volume timescale) of Storglaciären in its 1910 configuration may have been shorter than in its present state. Bahr and others (1998) showed that τ_v might decrease with increasing glacier size (or volume; Raper and others, 1996) precisely because larger glaciers extend to lower elevations where their termini experience greater rates of ablation. For the specific case of Storglaciären, the steep bed slope combined with the mass-balance gradient could result in a substantially larger value for $b_0(l_0)$ at its 1910 terminus and therefore a shorter τ_v via Equation (1). In particular, assuming again that db/dz is constant, a terminus elevation of 1050 m and an equilibrium-line altitude of ~ 1400 m (very roughly corresponding to a steady-state mass balance for the glacier's 1910 geometry), $b_0(l_0)$ would have been on the order of $2.8 \text{ m.w.e. a}^{-1}$. Thus an estimate of τ_v for the glacier in 1910 is $\sim 0.35 \bar{h}_{0,1910}$. Regardless of the thickness scale used, as a first approximation it is reasonable to expect that \bar{h}_0 varies proportionally to mean ice thickness, so that $\bar{h}_{0,modern}$ is $\sim 0.8 \bar{h}_{0,1910}$. Since $b_0(l_0)$ is ~ 1.0 for the interval 1981–2003, the glacier's current response time would also be $\sim 0.8 \bar{h}_{0,1910}$, more than double that for 1910. In the modeling, where a specific net-balance curve is used that will maintain Storglaciären's mid-1980s geometry, $b_0(l_0)$ is $\sim 0.75 \text{ m.w.e. a}^{-1}$ so that τ_v is $\sim 1.1 \bar{h}_{0,1910}$. The foregoing analysis, however crude, might reconcile the longer response deduced from the modeling with that suggested by the observed behavior. It furthermore underscores the importance of the bed slope near the terminus in the glacier's response to climate change (presumably through local dynamics of flow), as has been noted elsewhere (e.g. Rasmussen and Conway, 2001).
5. It was mentioned earlier that during recent decades this region of the Arctic seems to be experiencing more winter precipitation, which may mean Storglaciären's approach to its current 'steady state' was hastened, or that the observed quasi-equilibrium is in fact transient.
6. Finally the model's failure to incorporate sliding could also explain, to some extent, the resultant longer volume timescale compared with those based on its observed changes in geometry. Brugger (1992) considered the effects of sliding in an admittedly ad hoc manner by modifying f^* in Equation (4), and these results are included in Table 1. In the model, sliding is invariant along x , so the values of τ_v shown are probably not quantitatively correct. However, these results do show qualitatively that (1) by virtue of increased mass turnover, sliding shortened the volume timescales as would be expected; (2) the values of τ_v determined for both Storglaciären and Rabots Glaciär in this study are robust even if sliding contributes $\sim 10\%$ to mean annual surface velocity; and (3) the magnitude of sliding required to reconcile τ_v inferred from the model with that based on observed volume changes on Storglaciären would need to be substantial ($\sim 50\%$ or more), and hence arguably not reasonable.

In conclusion, numerical modeling indicates that in their mid-1980s configurations and for small mass-balance perturbations the volume timescale of Rabots Glaciär is ~215 years while that of nearby Storglaciären is ~125 years. In view of the vagaries of 'real-world' climate signals and actual as opposed to perceived steady states, it is difficult to assess whether these response times can, at present, be verified by observations of the glaciers' respective behaviors. Moreover, the glaciers' response time may not have been constant through time. By virtue of a rather steep bed slope, the advanced and thin terminus of Storglaciären in 1910 extended to a much lower elevation and hence was subjected to substantially greater ablation. Storglaciären would have therefore been particularly sensitive to the reduced ice flux brought about by climate warming, and this is reflected in the shorter response time estimated for its 1910 geometry. The same cannot be said for Rabots Glaciär, because of its relatively flat bed slope. Thus the difference in response times of the two glaciers may have been even greater in the past. However, what is emphasized here is not a particular individual response time, but their relative magnitudes. A longer response time explains why Rabots Glaciär lagged behind Storglaciären in its initial reaction to the warming that occurred early in the 20th century, and why significant ice retreat and mass loss continues today. When considering their response to CO₂-induced global warming, Storglaciären might be expected to react before Rabots Glaciär, and any response by Rabots Glaciär could be superimposed on its final adjustments to the earlier warming.

ACKNOWLEDGEMENTS

Fieldwork on Rabots Glaciär in the 1980s was carried out under the supervision of R.LeB. Hooke, who first suggested this problem, and was supported by US National Science Foundation grant DPP83-05935. R. Bindschadler provided the code for the numerical model and patiently answered questions concerning its implementation. Comments by R.LeB. Hooke, A. Rasmussen and two anonymous reviewers improved the manuscript. The continued support, cooperation and hospitality of personnel of the Tarfala Field Station and Stockholms Universitet, especially P. Jansson and P. Holmlund, are gratefully acknowledged.

REFERENCES

- Abdalati, W. and 9 others. 2004. Elevation changes of ice caps in the Canadian Arctic Archipelago. *J. Geophys. Res.*, **109**(F4), F04007. (10.1029/2003JF000045.)
- Albrecht, O., P. Jansson and H. Blatter. 2000. Modelling glacier response to measured mass-balance forcing. *Ann. Glaciol.*, **31**, 91–96.
- Arendt, A.A., K.A. Echelmeyer, W.D. Harrison, C.S. Lingle and V.B. Valentine. 2002. Rapid wastage of Alaska glaciers and their contribution to rising sea level. *Science*, **297**(5580), 382–386.
- Bahr, D.B., W.T. Pfeffer, C. Sassolas and M.F. Meier. 1998. Response time of glaciers as a function of size and mass balance. 1. Theory. *J. Geophys. Res.*, **103**(B5), 9777–9782.
- Bindschadler, R. 1980. The predicted behavior of Griesgletscher, Wallis, Switzerland, and its possible threat to a nearby dam. *Z. Gletscherkd. Glazialgeol.*, **16**(1), 45–59.
- Bindschadler, R. 1982. A numerical model of temperate glacier flow applied to the quiescent phase of a surge-type glacier. *J. Glaciol.*, **28**(99), 239–265.
- Bindschadler, R.A. and L.A. Rasmussen. 1983. Finite-difference model predictions of the drastic retreat of Columbia Glacier, Alaska. *USGS Prof. Pap.* 1258-D.
- Brugger, K.A. 1992. A comparative study of the response of Rabots Glaciär and Storglaciären to recent climatic change. (PhD thesis, University of Minnesota.)
- Brugger, K.A. 1997. Predicted response of Storglaciären, Sweden, to climatic warming. *Ann. Glaciol.*, **24**, 217–222.
- Brugger, K.A., K.A. Refsnider and M.F. Whitehill. 2005. Variation in glacier length and ice volume of Rabots Glaciär, Sweden, in response to climate change, 1910–2003. *Ann. Glaciol.*, **42**, 180–188.
- Dowdeswell, J.A. and 10 others. 1997. The mass balance of circum-Arctic glaciers and recent climate change. *Quat. Res.*, **48**(1), 1–14.
- Dyrgerov, M.B. and M.F. Meier. 2000. Twentieth century climate change: evidence from small glaciers. *Proc. Natl. Acad. Sci. USA (PNAS)*, **97**(4), 1406–1411.
- Hanson, B. 1995. A fully three-dimensional finite-element model applied to velocities on Storglaciären, Sweden. *J. Glaciol.*, **41**(137), 91–102.
- Harrison, W.D., D.H. Elsberg, K.A. Echelmeyer and R.M. Krimmel. 2001. On the characterization of glacier response by a single timescale. *J. Glaciol.*, **47**(159), 659–664.
- Holland, M.M. and C.M. Bitz. 2003. Polar amplification of climate change in coupled models. *Climate Dyn.*, **21**(3–4), 221–232.
- Holmlund, P. 1987. Mass balance of Storglaciären during the 20th century. *Geogr. Ann.*, **69A**(3–4), 439–44.
- Holmlund, P. 1988. Is the longitudinal profile of Storglaciären, northern Sweden, in balance with the present climate? *J. Glaciol.*, **34**(118), 269–273.
- Holmlund, P. 1993. Surveys of post-Little Ice Age glacier fluctuations in northern Sweden. *Z. Gletscherkd. Glazialgeol.*, **29**(1), 1–13.
- Holmlund, P. and P. Jansson. 1999. The Tarfala mass balance programme. *Geogr. Ann.*, **81**(4), 621–631.
- Holmlund, P., W. Karlén and H. Grudd. 1996. Fifty years of mass balance and glacier front observations at the Tarfala research station. *Geogr. Ann.*, **78A**(2–3), 105–114.
- Hooke, R.LeB. 2005. *Principles of glacier mechanics. Second edition.* Upper Saddle River, NJ, Prentice Hall.
- Hooke, R.LeB., P. Calla, P. Holmlund, M. Nilsson and A. Stroeven. 1989. A 3 year record of seasonal variations in surface velocity, Storglaciären, Sweden. *J. Glaciol.*, **35**(120), 235–247.
- Houghton, J.T. and 7 others, eds. 2001. *Climate change 2001: the scientific basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge, etc., Cambridge University Press.
- Jóhannesson, T., C. Raymond and E. Waddington. 1989a. A simple method for determining the response time of glaciers. In Oerlemans, J., ed. *Glacier fluctuations and climatic change.* Dordrecht, etc., Kluwer Academic Publishers, 343–352.
- Jóhannesson, T., C. Raymond and E. Waddington. 1989b. Timescale for adjustment of glaciers to changes in mass balance. *J. Glaciol.*, **35**(121), 355–369.
- Kuhn, M. 1984. Mass budget imbalances as criterion for a climatic classification of glaciers. *Geogr. Ann.*, **66A**(3), 229–238.
- MacGregor, K.R., C.A. Riihimaki and R.S. Anderson. 2005. Spatial and temporal evolution of rapid basal sliding on Bench Glacier, Alaska, USA. *J. Glaciol.*, **51**(172), 49–63.
- Meier, M.F., M.B. Dyrgerov and G.J. McCabe. 2003. The health of glaciers: recent changes in glacier regime. *Climatic Change*, **59**(1–2), 123–135.
- Nye, J.F. 1960. The response of glaciers and ice-sheets to seasonal and climatic changes. *Proc. R. Soc. London, Ser. A*, **256**(1287), 559–584.
- Nye, J.F. 1965. The flow of a glacier in a channel of rectangular, elliptic or parabolic cross-section. *J. Glaciol.*, **5**(41), 661–690.

- Oerlemans, J. and 10 others. 1998. Modelling the response of glaciers to climate warming. *Climate Dyn.*, **14**(4), 267–274.
- Oerlemans, J. 2001. *Glaciers and climate change*. Lisse, etc., A.A. Balkema.
- Oerlemans, J. 2005. Extracting a climate signal from 169 glacier records. *Science*, **308**(5722), 675–677.
- Pohjola, V.A. and J.C. Rogers. 1997. Coupling between the atmospheric circulation and extremes of the mass balance of Storglaciären, northern Scandinavia. *Ann. Glaciol.*, **24**, 229–233.
- Raper, S.C.B., K.R. Briffa and T.M.L. Wigley. 1996. Glacier change in northern Sweden from AD 500: a simple geometric model of Storglaciären. *J. Glaciol.*, **42**(141), 341–351.
- Rasmussen, L.A. 2004. Altitude variation of glacier mass balance in Scandinavia. *Geophys. Res. Lett.*, **31**(13), L13401. (10.1029/2004GL020273.)
- Rasmussen, L.A. and H. Conway. 2001. Estimating South Cascade Glacier (Washington, U.S.A.) mass balance from a distant radiosonde and comparison with Blue Glacier. *J. Glaciol.*, **47**(159), 579–588.
- Serreze, M.C. and 9 others. 2000. Observational evidence of recent change in the northern high-latitude environment. *Climatic Change*, **46**(2), 159–207.
- Stroeven, A.P. 1996. The robustness of one-dimensional, time-dependent, ice-flow models: a case study from Storglaciären, northern Sweden. *Geogr. Ann.*, **78A**(2–3), 133–146.