

lowed for 26 months after cancer diagnosis. They and their partners were assessed using semistructured interviews conducted by trained interviewers. Period prevalence rates were calculated for the 26 months following cancer diagnosis for DSM-III-R major depressive disorder (MDD) and total affective disorders (i.e. DSM-III-R MDD, DSM-III generalised anxiety disorder and minor affective disorders). Potential risk factors were assessed.

The 26 month period prevalence of total affective disorder (TAD) in female partners (12/50, 24.0%) did not differ significantly to that in male cancer patients (13/50, 26.0%), McNemar, $p = 1.000$. The prevalence of TAD in male partners (8/118, 6.8%) was significantly less than in female cancer patients (27/118, 22.9%), McNemar, $p < 0.001$. In both male and female partners the majority of affective disorders were accounted for by MDD. The prevalence of MDD in female partners was particularly high (10/50, 20.0%).

Most disorders in partners commenced 14–26 months after cancer diagnosis, significantly later than for cancer patients (McNemar, $p < 0.001$). The median duration of disorder in partners was 12 weeks, though this was a conservative estimate as nearly half of disorders were ongoing at the end of the study. Most of the 20 partners with affective disorder regarded their spouses' cancer as the cause of their disorder but only 12 consulted a doctor. Affective disorder in a partner was significantly associated with affective disorder in the corresponding cancer patient. Multivariate analysis showed three independent contributions to partners' affective disorder i.e. lower social class, female sex and past psychiatric history.

This is the only existing study to have used a standardised interview and operational criteria to compare rates of psychiatric disorder in cancer patients and their partners. Partners, particularly females, are at high risk of affective disorders. This raises the question of whether relatives/carers of patients with other chronic illnesses also suffer from high rates of psychiatric morbidity. Nearly half of partners with affective disorder did not receive treatment emphasising the importance of education for the public and health care professionals. The identification of risk factors may facilitate primary and secondary prevention of these disorders.

A DOUBLE-BLIND PLACEBO CONTROLLED TREATMENT TRIAL OF FLUOXETINE AND GRADED EXERCISE FOR CHRONIC FATIGUE SYNDROME (CFS)

R. Morriss, A. Wearden, R. Mullis, P. Strickland, L. Appleby, I. Campbell, D. Pearson. University of Manchester, Department of Psychiatry, Medicine, Manchester M20 8LR United Kingdom; Anaesthetics, Withington Hospital, Nell Lane, West Didsbury, Manchester M20 8LR United Kingdom

Objectives: To test the efficacy and acceptability to patients of two widely available treatments for chronic fatigue syndrome (CFS), with or without DSM-III-R depressive disorders, fluoxetine (20 mg/day) and graded exercise.

Methods: Six month prospective placebo and time with therapist controlled randomised trial with allocation to one of four treatment cells: exercise and fluoxetine; exercise and placebo drug; appointments and fluoxetine; appointments and placebo drug. The drug treatment was double-blind; patients were blind to the exercise programme. Graded exercise was delivered by a physiotherapist. 136 patients meeting Oxford research criteria for CFS were recruited from a University department of medicine outpatient clinic.

Results: 90 (66%) patients completed the trial. Patients were more likely to drop out of exercise than non-exercise treatment ($p = 0.05$). There was a non-specific treatment effect on most measures. When patients complied fully with exercise, there was a significant effect of exercise on fatigue ($p = 0.04$), functional work capacity ($p < 0.001$) and health perception ($p = 0.03$). In patients who complied, fluoxetine had a significant effect on fatigue ($p = 0.02$) and depression (p

$= 0.03$). On intention to treat, 12 (18%) patients had fully recovered from their fatigue symptoms in the exercise groups at six months compared to four (6%) who did not receive exercise ($p = 0.004$). Significant improvement with exercise only occurred in patients with DSM-III-R depressive disorders who were prescribed fluoxetine.

Conclusions: CFS patients benefit from graded exercise and fluoxetine if they can tolerate these treatments. Depressed CFS patients do not benefit from graded exercise unless they also receive an antidepressant. Graded exercise and fluoxetine have clinically distinct effects in CFS patients.

PREVALENCE OF DEPRESSION DURING HOSPITALIZATION FOR BONE MARROW TRANSPLANTATION: EFFECTS OF DIAGNOSTIC CRITERIA

J.M. Prieto, J. Blanch, J. Atala, C. Gastó, E. Cirera. Department of Consultation-Liaison Psychiatry, University of Barcelona, Villarroel 170, 08036, Barcelona, Spain

Objective: This study used different approaches to case identification to examine prevalence rates for major (MD) and minor depression (md) in a group of patients with hematological malignancies.

Method: A consecutive series of 103 patients hospitalized for bone marrow transplantation at the Hospital Clinic in Barcelona were evaluated on admission and weekly until discharge. **Results:** Diagnosis according to DSM-IV (exclude somatic symptoms related to a physical condition), DSM-III (include all somatic symptoms), and Endicott's revised criteria (replace somatic symptoms with non-somatic ones) are shown in the following table:

Diagnosis	DSM-IV		DSM-III		Endicott Criteria	
	N	%	N	%	N	%
MD	3	3	26	25	10	10
md	31	30	16	15	25	24

Taking in account the existing literature, a low rate of major depression was obtained when using DSM-IV criteria. The higher rate obtained with DSM-III can be related to the prevalence of somatic symptoms. After receiving intensive treatment and during the following 4 weeks, 95–81% of patients reported an energy loss $\geq 30\%$, with 98–78% reporting a moderate decrease in appetite.

Conclusions: When somatic symptoms are prevalent, an approach that relies more in psychological features can increase the diagnostic accuracy for mood disorders.

PSYCHOSOCIAL ASSESSMENT OF TRANSPLANT PATIENTS. THE GREEK EXPERIENCE

M. Syngelakis, A. Vidalis, G. Kaprinis. Psychiatric Department, Hippocratio General Peripheral Hospital, Konstantinoupoleos 49, TK 546 42 Thessaloniki, Greece

Research on psychiatric aspects of transplantation has been undertaken as transplantation is linked with increased psychiatric morbidity and also due to evidence that psychosocial factors influence the clinical outcome and the quality of life. The biopsychosocial model seems to be a necessary approach to these patients.

Since October 1994 we have evaluated and supported transplant patients referred from the two major Transplant Units in Northern Greece.

Our sample consist of all heart- ($n = 25$), lung- ($n = 5$), liver- ($n = 22$) candidates and recipients as well as a number of renal transplant patients ($n = 60$).

Results are presented using an 10-item rating scale for psychosocial screening of transplant patients (Psychosocial Assessment of Candidates for Transplantation, Olbrisch 1989). Mental health

was evaluated with clinical interviews as well as with psychometric testing instruments [Symptom Checklist-90-R, Mini Mental State Examination..., projective tests, family support assessment...] Further information were obtained from relatives and from medical records.

The socio-economic level of the patients was found to be low. Family support system was adequate. Regression, paranoid ideation and splitting were very common 'defences' used in our sample. Almost 50% of the patients warranted a psychiatric diagnosis (DSM-IV). Healthy Lifestyle and Compliance reported higher than acceptable level. During of presurgical period candidates were placing magical expectations in the surgical treatment, and also the majority of heart-, lung-, and liver- patients view transplant as cure and had no long range picture. In our country the relevant easy approach to identity of the donor, seems to promote identifications and make more difficult the rehabilitation of the recipient. Significant differences between the types of transplantation were found in the items of psychopathology and understanding of transplantation, in our sample.

The present study approaches for the first time the peculiarities presented in our country regarding the psychological, social and cultural aspects of transplantation. Additionally we describe some preliminary 'normative data' for Greek population in a scaled format that allows comparisons with other transplant programmes.

NR8. Substance and alcohol abuse

Chairmen: H Ghodse, J Marshall

HEALTH PROMOTION IN THE DRUG SUBCULTURE

Werner J. Fuchs, Peter J. Grob. Psychiatric Center Wetzikon/Zurich, Spitalstrasse 60, CH-8620 Wetzikon, Switzerland

A harm reduction approach in an open drug scene is described. The effectiveness of a needle exchange scheme can be increased if it is part of primary medical care. Besides the basic documentation on services provided through various studies, data on the population of an open drug scene were obtained. Viro-epidemiological studies showed a 100–300 times higher prevalence of hepatitis A, B, and C compared with the Swiss population in general. A downward tendency of HIV prevalence could be demonstrated. An overall reduction of needle sharing between 1989 and 1992 was noted. However risk behaviour still exists in some subgroups of intravenous drug users, especially in beginners, where a high prevalence of hepatitis B was found. Methodological considerations of research in open drug scenes are discussed.

VALIDATION OF THE ALCOHOL USE DISORDERS IDENTIFICATION TEST (AUDIT) — ROMANIAN VERSION FOR ICD-10 DIAGNOSES

Aurora Jianu. Psychiatric Clinic Timisoara, Romania

The AUDIT has been developed from WHO collaborative project as a screening instrument discriminating harmful and hazardous alcohol consumption and alcohol dependence. This instrument is very useful in preventive medicine, primary health care and psychiatric epidemiology. Previously screening techniques for alcoholism were based only on the presence of abnormal physical findings. On the other side, a lot of studies have shown that AUDIT was suitable in different cultures.

The present study aimed to transfer this epidemiologic technique in Romania as a first step developing a real communitar policy against alcohol misuse. The author tried to find if the AUDIT could appropriately "read" the ICD-10 diagnoses and diagnostic criteria in a Romanian clinical and nonclinical sample.

Two groups of volunteers were recruited for this study: the clinical sample encompassed 70 alcoholics from alcoholism treatment units (mean age 43.22 ± 9.91 ; M/F: 57/13; years of education 11.96 ± 3.91) and healthy control group contained 74 subjects from general population (mean age 38.33 ± 11.58 ; M/F: 43/31; years of education: 13.77 ± 2.96).

All subjects were assessed by AUDIT Core (the first 10 questions) and AUDADIS (Alcohol Use Disorders and Associated Disabilities Interview Schedule) developed by Grant & Hasin (1991) from National Institute of Alcoholism and Alcohol Abuse-Rockville USA.

The discriminant validity of the AUDIT was performed determining the ability of this instrument to differentiate alcoholics vs non-alcoholics and various alcoholism diagnostic criteria vs non-criteria. The cutoff point of AUDIT score is the true threshold between the categories above mentioned.

The AUDIT score of 10 is a good cutoff point discriminating between alcoholics vs non-alcoholics diagnosed according to AUDADIS algorithm ($\kappa = 0.81$, sensitivity/specificity: 90.6/90.6) and between non-criteria and harmful ($\kappa = 0.65$, sensit./specif.: 83.5/81.0), tolerance ($\kappa = 0.40$, sensit./specif. 67.0/72.9), neglecting interests ($\kappa = 0.51$, sensit./specif. 63.5/91.5), and withdrawal/relief ($\kappa = 0.71$, sensit./specif. 88.2/83.0) criteria.

This study highlights that AUDIT is a suitable instrument that is able to distinguish between alcoholics and non-alcoholics and furnishes confident thresholds outlining diagnostic criteria. Contrary, AUDIT is not able to differentiate between harmful and dependency, the diagnostic categories from ICD-10.

EPIDEMIOLOGY OF ALCOHOLISM IN THE EUROPEAN RUSSIAN NORTH

Alexei Kalinin, Pavel Sidorov. Department of Psychiatry, Arkhangelsk State Medical Academy, 51, Troitsky Ave., Arkhangelsk, 163061, Russia

The main aim of the present study was to investigate the level of alcoholism among some professional groups of Arkhangelsk city (500 000 inhabitants) and the rural population of Arkhangelsk region as well.

The object of the study were: seamen of the Northern Shipping Company, pilots of Arkhangelsk Air Company and workers of different industrial plants, rural population of the remote districts.

The main methods of investigations were testing, interviewing, clinical and psychological analysis, including symptomatological and syndromological evaluation of alcoholic disease.

According to the investigation data the frequency of alcoholism among these groups varies from 10% to 30%. It is specially dangerous for transport specialities where even slight postintoxication disorders can essentially affect the quality of professional ability.

A COMPARATIVE STUDY ON THE MOTIVATION OF DRUG ADDICTS ADMITTED TO METHADONE MAINTENANCE TREATMENT (UK,LONDON SW — BULGARIA, SOFIA)

Igor Kutsenok, (Bulgaria, National Center for Addictions- Sofia). MSc student, St. George's Hospital Medical School, Department of Addictive Behaviour, Cranmer Terrace, London SW 17 OPZ, UK

Measures of motivation for drug use change and admission to methadone maintenance were examined among two groups drug