


Pum3 is dispensable for mouse oocyte maturation and embryo development *in vitro*

TingTing Zhao¹, Wei Huang² and Kaibo Lin³ 

Research Article

Cite this article: Zhao TT *et al.* (2023) *Pum3* is dispensable for mouse oocyte maturation and embryo development *in vitro*. *Zygote*, **31**: 359–365. doi: [10.1017/S0967199423000205](https://doi.org/10.1017/S0967199423000205)

Received: 3 November 2022
Revised: 30 March 2023
Accepted: 3 April 2023
First published online: 17 May 2023

Keywords:

Embryo development; Oocyte maturation; PUF family; Pum3; RNA-binding protein

Corresponding author:

Kaibo Lin, Department of Assisted Reproduction, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200021, China. Email: linkb@126.com. Wei Huang, Department of Reproductive Medicine, The First Affiliated Hospital of Hainan Medical University, Hainan Medical University, Hainan, 570102, China. Email: huang1213@hainmc.edu.cn

¹State Key Laboratory of Reproductive Medicine, Nanjing Medical University, Nanjing, Jiangsu, 211166, China; ²Department of Reproductive Medicine, The First Affiliated Hospital of Hainan Medical University, Hainan Medical University, Hainan, 570102, China and ³Department of Assisted Reproduction, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200021, China

Summary

Pumilio3 (*Pum3*), an evolutionarily distant homologue of the classical RNA-binding protein PUF (PUMILIO and FBF) family member, is also involved in the process of RNA metabolism through post-transcriptional regulation. However, the functions of *Pum3* in mouse oocyte maturation and preimplantation embryonic development have not been elucidated. By comparing RNA levels in different tissues, we found that *Pum3* was widely expressed in multiple tissues, but moderately predominant in the ovary. Histochemical staining suggested that the PUM3 protein exhibits positive signals in oocytes, granulosa cells and theca cells of different follicle stages. Oocyte immunofluorescence results showed a slightly higher level of PUM3 protein in metaphase II compared with the germinal vesicle (GV) stage. After knockdown of *Pum3* in GV oocytes using siRNA injection (siPUM3), no obvious defect was observed in the processes of GV breakdown and polar body extrusion during *in vitro* maturation (IVM) for the siPUM3 oocytes. Compared with the control group, the siPUM3 group displayed no significant abnormality in the cleavage and blastocyst formation rate of these fertilized oocytes. Therefore, we can conclude that depletion of *Pum3* does not affect mouse oocyte maturation and early embryonic development *in vitro*.

Introduction

The PUF proteins are a family of RNA-binding proteins with a highly conserved PUF domain, including the classical PUF proteins (PUM1/2 and FBF1/2) and two evolutionary distant homologues (PUM3 and NOP9; Wang *et al.*, 2018; Najdová *et al.*, 2020). The PUF family members are widely distributed in eukaryotes and specifically recognize and bind the motif sequence through PUF repeats. This post-transcriptional regulation is mediated using RNA-binding proteins, which can control RNA splicing, polyadenylation, capping, modification, localization, stability, and spatiotemporally specific translation, and is known as RNA metabolism (Blackinton and Keene, 2014). Generally, PUM proteins regulate mRNA stability and translation by directly binding to the target mRNAs through the PUM homology domain (PUM-HD; Edwards *et al.*, 2001; Wang *et al.*, 2001). Pum3 (Pumilio RNA binding family member 3, also known as KIAA0020 or PUF-A in humans) is conserved in a wide range of eukaryotes, including fungi, plants, protozoans and metazoans (Liang *et al.*, 2018; Joshna *et al.*, 2020; Najdová *et al.*, 2020; Son *et al.*, 2021). The PUM-HD in classical PUMs (PUM1 and PUM2) consists of eight tandem PUF repeats, 36 amino acids contain the motif, forming a crescent-shaped structure (Wang *et al.*, 2001); however, PUM3 consists of 11 PUF repeats in PUM domain, which is arranged in an L-shape and exhibits the ability to bind to DNA and single-stranded and double-stranded RNA without sequence specificity (Najdová *et al.*, 2020). Therefore, the non-classical PUM homologue PUM3 may also play a significant role at the transcriptional and post-transcriptional levels.

Extensive studies in mammals have revealed that classic PUMs (PUM1 and PUM2) proteins play multiple roles in various biological processes (Goldstrohm *et al.*, 2018). Previous studies in invertebrates have suggested that PUF family proteins are essential for germline stem cell maintenance and embryonic development in *Drosophila* and *C. elegans* (Lin and Spradling, 1997; Zhang *et al.*, 1997; Kraemer *et al.*, 1999; Parisi and Lin, 1999). Recent studies in mammals have also shown that PUMs have key roles in germ-cell development, neurodegeneration, early embryonic development and postnatal growth. In particular, PUMs have a strong correlation with a variety of human cancers, and also have a strong promoting role in the occurrence and development of cancer using post-transcriptional regulation (Miles *et al.*, 2012; Naudin *et al.*, 2017; Munschauer *et al.*, 2018; Elguindy and Mendell, 2021; Silva *et al.*, 2022; Yoon *et al.*, 2022). In conclusion, classical PUM proteins control various cellular functions by post-transcriptionally regulating the fate of target RNAs in both invertebrates and mammals.

© The Author(s), 2023. Published by Cambridge University Press.



However, the physiological function of PUM3 in mammals remains unclear. A previous study has shown that PUM3 (Puf-A) is essential for eye development and primordial germ-cell (PGC) migration in zebrafish (Kuo *et al.*, 2009). In mouse embryonic stem cells, the loss of PUM3 can increase the Zscan4-positive 2-cell embryo-like state cells in a p53-dependent manner (Takahiro *et al.*, 2021). Recently, studies have shown that PUM3 is located in the nucleolus and regulates ribosome biogenesis during the promotion of cancer progression (Fan *et al.*, 2013; Lin *et al.*, 2021; Cho *et al.*, 2022). To explore the role of *Pum3* in mouse development, we characterized the expression levels of *Pum3* among different tissues. The result showed that *Pum3* mRNA was widely expressed in multiple organs, but predominantly in the ovary. Furthermore, using immunohistochemical staining, we found that the PUM3 protein was abundantly expressed in follicles at different stages. MII oocytes exhibited a higher level of PUM3 in the cytoplasm. To reveal the function of *Pum3* during oocyte *in vitro* maturation (IVM), we found that knockdown of *Pum3* in GV oocytes did not affect GVBD and polar body extrusion (PBE). In addition, we detected preimplantation embryo development for PUM3-deficient oocytes and the control group. However, the zygotes in the knockdown group developed normally, as in the control group. Taken together, our data suggested that PUM3 is dispensable for mouse oocyte IVM and early embryo development after fertilization.

Materials and methods

Animals

Animals were housed under controlled environmental conditions in the animal facility at Nanjing Medical University, Nanjing, China, with free access to water and food. Illumination was provided between 8 a.m. and 8 p.m. daily.

Oocyte collection, siRNA microinjection and oocyte IVM

Female ICR mice were primed intraperitoneally with 5 IU PMSG to stimulate the development of multiple follicles. After 48 h, mice were sacrificed for GV oocyte collection or injected with 5 IU of hCG to induce ovulation. The ovaries were washed three times with 0.9% (w/v) NaCl. Cumulus–oocyte complexes (COCs) were aspirated from follicles (2–8 mm) in mHTF medium (Irvine Scientific, USA) containing 1.0 mM IBMX (Sigma, USA) using an 18-gauge needle attached to a 10-ml syringe.

Three *Pum3* (NM_177474.5) siRNAs and nonsense siRNA were designed and chemically synthesized by GenePharma (Shanghai, China) and mixed for injection. The siRNA target sequences are shown in Table 1. The COCs were digested in 0.2% hyaluronidase for 3 min at 37°C. When all cumulus cells were separated from oocytes, 10 µl of the siRNA mixture (300 nmol/l) and nonsense RNA were microinjected into the germinal vesicle (GV)-stage oocytes using the Eppendorf Cell Tram Vario system (Eppendorf, Hamburg, Germany). After microinjection, oocytes were cultured in KSOM medium (Millipore, Germany) containing 1.0 mM IBMX (Sigma, USA) under mineral oil at 37°C and 5% CO₂ in the incubator for 16 h.

After siRNA injection, GV oocytes were washed and cultured under normal KSOM medium without IBMX addition for 30, 45, 60, 75, or 90 min. Then, oocytes were scored as either germinal vesicle intact or having undergone GVBD. The percentage of either GV or GVBD was calculated for each time point. The polar body was counted and calculated.

Table 1. Sequence of primers used for RT-qPCR and siRNA

Primer	Primer sequence (5'→3')
siPum3-1	GCAAAGGACCGAAGGACAA
siPum3-2	CCACAAGTTGCGGAGATAA
siPum3-3	CCAGCATAGTTAATGACAA
mPum3-1-F	GCAAGAAAGGTGTGAAGCAG
mPum3-1-R	TTGGGTTTCTGGCTCCTG
mActin-F	CTAAGGCCAACCGTGAAG
mActin-R	ACCAGAGGCATACAGGGACA

Embryo culture *in vitro*

Matured oocytes were used for *in vitro* fertilization as previously described (Yan *et al.*, 2015). Approximately 6 h later, these two-pronuclear zygotes were then washed and cultured in mHTF medium drops. The siRNA mixture or nonsense siRNA was microinjected into zygotes immediately. Then, all embryos were transferred into KSOM medium and cultured under normal conditions. Embryos (from the 2-cell stage to blastocysts) were observed and recorded at 1.5, 2.5, 3.5, and 4.5 days post fertilization.

RNA isolation and real-time polymerase chain reaction (PCR)

RNA was extracted from ~50 oocytes or embryos with the RNeasy Pure Micro Kit (Qiagen, China) according to the manufacturer's instructions. Reverse transcription was performed using a PrimeScript RT Master Mix (Vazyme, China). Real-time PCR was performed using *Taq* polymerase (Vazyme, China) according to the product manual and gene-specific primers. The number of PCR cycles ranged from 22 to 35, depending on the linearity of the reaction. All gene expression analyses were performed with samples from at least three independent experiments. All primers for real-time RT-qPCR are listed in Table 1.

Histology, immunohistochemistry and immunofluorescence

Tissues were fixed overnight in Hartman's fixative (Sigma). Fixed ovaries were embedded in paraffin and cut into 5-µm-thick serial sections. Sections were processed for haematoxylin and eosin (H&E) staining and immunohistochemistry according to standard protocols. Immunostaining for PUM3 (abcam, 1:50) was performed, following citrate buffer antigen retrieval, by incubation with primary antibodies and detected using biotin–streptavidin horseradish peroxidase (HRP) detection systems (ZSGB-BIO). A minimum of three randomly chosen discontinuous sections were used to determine positive cells in tubules.

Quantification and statistical analysis

All data in bar and line graphs are expressed as means ± standard deviation (SD) of the mean. All experiments were repeated at least three times unless noted. Statistical significance between two groups of data was evaluated using Student's *t*-test (two-tailed) comparison using GraphPad Prism software 5 or Microsoft Excel *t*-test function. Statistical significance is indicated by ns for no significant difference, **P*-value <0.05, ***P*-value <0.01, ****P*-value <0.001.




Figure 1. Characterization of *Pum3* gene expression in the ovary. (A) RT-qPCR detection of *Pum3* mRNA levels in various tissues. (B) Expression of PUM3 protein was detected using immunohistochemistry staining in the adult wild-type ovary. Scale bar: 200 μ m. Red arrows: oocyte; yellow arrows: granulosa cells; blue arrows: theca cells; asterisk: no specific signal in stroma cells. (C) PUM3 expression in different stages of follicles: primordial follicle (PmF), primary follicle (PF), secondary follicle (SF) and antral follicle (AF). Scale bars: 20 μ m. (D) Immunofluorescence images of PUM3 in GV and MII oocytes. Dotted circle: antibody negative control group incubated with 5% BSA. Red arrowheads: MII oocytes; yellow arrowheads: GV oocytes; blue arrowheads: granulosa cells. Scale bars 5 μ m. (E) Fluorescence intensity of PUM3 in MII and GV oocytes; *** $P < 0.001$.

Results

High expression of PUM3 in oocytes and granulosa cells

To identify the expression patterns of *Pum3* RNA in different mouse tissues, we extracted total RNA from various tissues of adult mice. *Pum3* RNA levels were detected using real-time quantitative reverse transcription PCR (RT-qPCR). Our results showed that *Pum3* RNA was widely expressed in multiple tissues, including the ovary, epididymis, testis, kidney, liver, thymus, lung, spleen, brain, uterus and intestine (Figure 1A), especially with the highest expression in the ovary tissue. Therefore, we speculated that *Pum3* may play an important role in the female reproductive system. Then we detected the expression of PUM3 protein in mouse ovary using immunohistochemistry (IHC) staining. Compared with the negative control (5% BSA), adult ovary sections stained with PUM3 antibody displayed special and strong positive signals in oocytes, granulosa cells and theca cells. However, a false-positive signal also appeared in the stroma cells of the negative control section, suggesting a non-specific signal (Figure 1B). At the same time, we also paid attention to the oocyte PUM3 expression in follicles at different stages. There were obvious signals in the oocyte cytoplasm from primordial follicles to mature antral follicles (Figure 1C). In addition, we also obtained GV-stage and MII-stage

oocytes by ovulation induction. Immunofluorescence staining showed that the MII oocyte, GV oocyte and its cumulus granulosa cells produced strong positive signals (Figure 1D). We also noticed that the PUM3 protein formed abundant granules in both GV and MII oocyte cortex. The nucleus had a weak signal and no fluorescent signal was detected in the nucleolus of GV oocytes. The nuclei of granulosa cells had a positive signal (Figure 1C, D). In MII oocytes, PUM3 does not appear to overlap the spindle. The fluorescence intensity of oocyte PUM3 at the MII stage was significantly higher (29%) than in the GV stage (Figure 1E). In summary, *Pum3* is highly expressed in the mouse ovary, especially in oocytes and granulosa cells, which is similar to the PUM1 expression pattern (Li *et al.*, 2022).

Pum3 knockdown oocytes display normal GVBD and PBE during in vitro maturation (IVM)

As previously reported, PUM1, a member of the classic PUM family, is a maternal factor regulating oocyte development in humans and mice (Mak *et al.*, 2016, 2018; Luo *et al.*, 2018). As PUM3 is highly expressed in oocytes at different stages, we suspected that PUM3 may play an important role during mouse oocyte development. To identify this we knocked down *Pum3* in GV oocytes using




Figure 2. IVM results of *Pum3* knockdown oocytes. (A) RT-qPCR of *Pum3* knockdown in oocyte cells at 48 h after siRNA infection. (B, C) Percentages of GVBD (B) and PBE (C) in the *Pum3* knockdown and control groups. *** $P < 0.001$; ns, not statistically significant, $P > 0.05$.




Figure 3. Effect of *Pum3* knockdown on preimplantation embryo development *in vitro*. (A) RT-qPCR detection of *Pum3* knockdown in the IVF blastocysts, *** $P < 0.001$. (B) Percentage of IVF embryos developed to the 2-cell, 4-cell, compaction and blastocyst stages in the *siPum3* and control groups.

microinjection siRNA. The *Pum3* mRNA was successfully knocked down in GV oocytes using siRNA (*siPum3*), and the knockdown efficiency of *siPum3* was as high as 96% (Figure 2A). By removing the nuclear maturation inhibitor IBMX from the culture medium, GV oocytes were released and then entered germinal vesicle breakdown (GVBD). Surprisingly, oocytes from the *siPum3* group displayed a similar speed and ratio of GVBD compared with the control group (Figure 2B). The proportion of oocytes entering GVBD was slightly higher in the knockdown group compared with the control group, but this was not significant at each test point and, finally, two groups reached a comparable level (96.37% in the knockdown group and 93.37% in the control group), therefore *Pum3* deletion did not affect the development of GVBD in mouse oocytes. In addition, we also assessed the rate of PBE and found that knockdown of *Pum3* did not affect the PBE process (Figure 2C). The PBE rates of the control and *siPum3* groups reached 68.46% and 77.74%, respectively. Taken together, these results suggested that PUM3 is not essential for *in vitro* maturation of mouse oocytes.

Pum3 knockdown does not affect early embryo development *in vitro*

Although *Pum3* knockdown did not affect the IVM process of mouse oocytes, its high expression in the oocyte cytoplasm implied that PUM3 might be an important regulator (Figure 1D), so we questioned that whether *Pum3* might affect early embryo development after fertilization. Therefore, we obtained a large number of zygotes by *in vitro* fertilization (IVF), and then knocked down *Pum3* in the fertilized oocytes by siRNA injection at day 0.5 (two-pronuclear stage, 2PN), and tracked the developmental progress of

embryos in each group. Our RT-qPCR results showed that the efficiency of *Pum3* knockdown in blastocysts reached 96% (Figure 3A). Next the zygote underwent cleavage to form the blastula. At 1.5 days, 96.9% of the control and 100.0% of *siPum3* embryos developed to the 2-cell stage; at 2.5 days, 89.7% of the control and 92.5% of *siPum3* embryos developed to the 4-cell stage; at 3.5 days, 91.8% of the control and 94.1% of *siPum3* embryos developed to the compaction stage; at 4.5 days, 91.8% of the control and 90.3% of *siPum3* embryos developed to the blastocyst stage. These data showed that *siPum3* may slightly promote zygote cleavage and development from the 2-cell stage to the compaction stage, but we also finally obtained a similar blastocyst rate to the control group at 4.5 days (Figure 3B). Overall, these data suggested that *Pum3* knockdown had little effect on early embryonic development *in vitro*.

Discussion

In this study, we knocked down *Pum3* by injecting siRNA into mouse GV-stage oocytes and obtained MII oocytes using IVM culture. We found that, although PUM3 was expressed at a high level in the oocyte at multiple stages of the follicle, deletion of PUM3 did not have significant negative effects on the GVBD and PBE of oocyte IVM. In addition, the knockdown of *Pum3* in zygotes at the 2PN stage did not seem to cause significant defects in early embryonic development *in vitro*, and these embryos developed normally to blastocysts. These results suggested that PUM3 is not essential for oocyte maturation and early embryonic development *in vitro*.

RNA-binding proteins are a large group of proteins that mediate post-transcriptional regulation by binding to RNA, regulating

the fate of RNA and affecting related gene expression and the corresponding cellular functions (Matoulkova *et al.*, 2012). The PUF family are evolutionarily conserved and widely distributed RNA-binding proteins in eukaryotes, such as yeast, green plants, invertebrates and vertebrates (Zhang *et al.*, 2017). PUF family proteins possess a crescent-shaped RNA-binding domain called the Pumilio homology domain (PUM-HD), which is formed from eight PUF repeats (Jenkins *et al.*, 2009; Wang *et al.*, 2001). This PUM-HD structure can specifically recognize the PUM-binding site in the 3'UTR of target RNAs, and this mode of binding is conserved and can be engineered for accurate identification of the designated RNA sequence (Lu *et al.*, 2009).

PUF family genes generally have multiple copies or variants in different species, which may be related to the number of target mRNAs that need to be regulated in different organisms (Najdrová *et al.*, 2020). The PUF family has two evolutionarily distant homologous proteins, PUM3 and NOP9. It is worth noting that PUM3 and NOP9 are also widely distributed in eukaryotes, but there is only one copy on average in different species, which is different from the gene copy number of the PUF family (Najdrová *et al.*, 2020). Both proteins contain a non-canonical PUM-HD domain consisting of 11 repeats, a U-shaped domain in NOP9 and an L-shaped domain in PUM3 (Wang *et al.*, 2018). In eukaryotes, NOP9 has the function of RNA binding and is mainly involved in the processing and folding of 18S rRNA in the nucleolus; PUM3 can bind to double-stranded DNA or RNA without sequence specificity and participate in the occurrence of large ribosomal subunits. In addition, PUM3 and PUF family proteins are more similar in sequence than NOP9 (Najdrová *et al.*, 2020). Therefore, PUM3 and NOP9 are evolutionarily orthologous proteins of the PUF family. Although they have similar PUM-HD domains and are capable of binding RNA, they have their own role in the biogenesis of ribosomal RNA. This may be completely different in function from the classical PUF family proteins.

The earliest discovered physiological functions of the PUF family are the regulation of early embryo and germline stem cell development in invertebrates (Nüsslein-Volhard, 1987; Lin and Spradling, 1997; Zhang *et al.*, 1997; Forbes and Lehmann, 1998). This raised the question of whether the PUF family, a conserved family of RNA-binding proteins, could also play an equally important role in mammals. The subsequent studies in mice found that PUM1 deletion can reduce mouse fertility (Chen *et al.*, 2012), whereas PUM2 deletion did not affect mouse fertility (Xu *et al.*, 2007). In the absence of both PUM1 and PUM2, mouse embryos died at 8.5 days due to defective gastrulation differentiation (Lin *et al.*, 2018), probably because there would be some complementation between two highly conserved paralogous proteins of the PUF family in mammals. This also showed that their functions were also conserved to a certain extent, except for the evolutionary conservation of the protein structure of the PUF family. Moreover, large numbers of studies have shown that the PUF family also has important functions in the nervous system of invertebrates and mammals (Chen *et al.*, 2008; Vessey *et al.*, 2010; Gennarino *et al.*, 2015). It has been previously reported that PUF-A plays an important role in the development of eyes and primordial germ cells in fish (Kuo *et al.*, 2009; Ko *et al.*, 2022), but it remains unclear whether PUM3 plays a role in mammalian germ cells. Recently, it has been reported that the PUF family members PUM1/2 may play an important role in the occurrence and development of multiple tumours, and their expression is significantly upregulated in cancer cells. PUM protein can bind to the mitosis-related cycle inhibitor

cdkn1b through post-transcriptional regulation, thereby promoting the proliferation of tumour cells (Li *et al.*, 2022; Silva *et al.*, 2022). At the same time, it has also been reported that PUM3 can promote the growth of tumour cells (Fan *et al.*, 2013; Cho *et al.*, 2022; Silva *et al.*, 2022), but the current research is relatively limited and this remains to be studied.

During female oocyte development, shortly after the resumption of meiosis at the GV stage *in vitro*, including those artificially maintained in an arrest, the nucleolus (or nucleoli) rapidly disappears as does detectable rRNA and tRNA transcription (Kresoja-Rakic and Santoro, 2019), as would be expected. Nucleolar dissolution is followed by GVBD and chromatin condensation into nascent bivalent chromosomes. Condensation is accompanied by a cessation of transcription that is not resumed at a significant level until approximately the 2-cell stage in mouse and is not fully recovered until full embryo genomic activation a few cleavage divisions later.

This early embryonic period is largely characterized by the degradation of inherited maternal transcripts and their replacement by those originating from the embryonic genome, when the PUF family might be expected to function. PUM1, as the highly conserved PUF family protein, is considered to be an important maternal factor. Maternal deletion of PUM1 can lead to poor quality oocytes. Loss of maternal and paternal PUM1 results in severe preimplantation embryonic lethality (Mak *et al.*, 2018). As PUM3 is a non-canonical member of the PUF family, it is expressed at high levels during the development of mouse oocytes. We wondered whether PUM3 was also involved in mammalian oocyte development and early embryonic development. However, our data implied that PUM3 was not essential for mouse oocyte maturation and preimplantation embryo development *in vitro*.

It would appear that *Pum3* does not play a regulatory role in reactivating transcription and would not be expected to have such a role from the GV stage until early cleavage, as translation is driven primarily by maternal RNAs and the post-translational modifications of existing proteins. Although PUM3 is thought to have the ability to bind DNA and RNA, fluorescence staining actually revealed that PUM3 is mainly localized in the cytoplasm of GV-phase eggs, with extremely low levels in the nucleus and no signal in the nucleolus (Figure 1D). This might explain why PUM3 may not be mainly involved in RNA transcriptional activities in the GV phase. Therefore, knockdown of PUM3 in the GV phase did not affect two significant biological events (GVBD and PBE) during oocyte maturation.

What is surprising is that PUM3 is seemingly not involved in RNA processing and expression during the preimplantation stages, implying that if this processing is occurring other RNA regulatory elements may also have this function, including other members of RNA-binding proteins and/or the PUF family. The redundant function of the PUM1 and PUM2 on embryogenesis, body size, and testis development has been reported previously (Lin *et al.*, 2018, 2019; Zhao *et al.*, 2022). Furthermore, earlier studies of preimplantation mouse embryogenesis have suggested that there may be long-lived maternal mRNAs that presumably occur at the GV stage and persist through to the blastocyst stage (Schultz *et al.*, 2018). If true, siRNA knockout at the GV stage may not have evident downstream developmental consequences, as these transcripts, suggested by some to be stored in a subcortical complex, already existed prior to siRNA treatment.

As oocytes originate from primordial germ cells in the fetal period, it takes a long time for female germ-cell growth during folliculogenesis and eventually maturation in the ovary. It would be

useful to know whether implantation occurs, and if fetal development is normal in blastocysts derived from oocytes from *Pum3* null mice or injected with siRNA at the GV stage prior to the cessation of transcription. Therefore, a *Pum3* knockout animal model is still required to more clearly and accurately display the physiological functions of *Pum3* in germ-cell and embryonic development in the future.

Acknowledgements. This research acknowledges the effort of all authors in planning the experimental design and interpreting the results.

Financial support. This work was supported by research grants from the Young Scientists Fund of the National Natural Science Foundation of China (Grant No. 32000599); the Hainan Provincial Natural Science Foundation of China (Grant No. 821RC686); and the Shanghai Ninth People's Hospital (Grant No. JC201806).

Conflict of interest. The authors declare none.

Ethical standards. All animal experiments were approved by the Animal Care and Use Committee (ACUC) of Nanjing Medical University and performed in accordance with institutional guidelines.

References

- Blackinton, J. G. and Keene, J. D. (2014). Post-transcriptional RNA regulons affecting cell cycle and proliferation. *Seminars in Cell and Developmental Biology*, **34**, 44–54. doi: [10.1016/j.semcd.2014.05.014](https://doi.org/10.1016/j.semcd.2014.05.014)
- Chen, G., Li, W., Zhang, Q. S., Regulski, M., Sinha, N., Barditch, J., Tully, T., Krainer, A. R., Zhang, M. Q. and Dubnau, J. (2008). Identification of synaptic targets of *Drosophila* pumilio. *PLoS Computational Biology*, **4**(2), e1000026. doi: [10.1371/journal.pcbi.1000026](https://doi.org/10.1371/journal.pcbi.1000026)
- Chen, D., Zheng, W., Lin, A., Uyhazi, K., Zhao, H. and Lin, H. (2012). Pumilio 1 suppresses multiple activators of p53 to safeguard spermatogenesis. *Current Biology: CB*, **22**(5), 420–425. doi: [10.1016/j.cub.2012.01.039](https://doi.org/10.1016/j.cub.2012.01.039)
- Cho, H. C., Huang, Y., Hung, J. T., Hung, T. H., Cheng, K. C., Liu, Y. H., Kuo, M. W., Wang, S. H., Yu, A. L. and Yu, J. (2022). Puf-A promotes cancer progression by interacting with nucleophosmin in nucleolus. *Oncogene*, **41**(8), 1155–1165. doi: [10.1038/s41388-021-02138-0](https://doi.org/10.1038/s41388-021-02138-0)
- Edwards, T. A., Pyle, S. E., Wharton, R. P. and Aggarwal, A. K. (2001). Structure of Pumilio reveals similarity between RNA and peptide binding motifs. *Cell*, **105**(2), 281–289. doi: [10.1016/s0092-8674\(01\)00318-x](https://doi.org/10.1016/s0092-8674(01)00318-x)
- Elguindy, M. M. and Mendell, J. T. (2021). NORAD-induced Pumilio phase separation is required for genome stability. *Nature*, **595**(7866), 303–308. doi: [10.1038/s41586-021-03633-w](https://doi.org/10.1038/s41586-021-03633-w)
- Fan, C. C., Lee, L. Y., Yu, M. Y., Tzen, C. Y., Chou, C. and Chang, M. S. (2013). Upregulated hPuf-A promotes breast cancer tumorigenesis. *Tumour Biology: The Journal of the International Society for Oncodevelopmental Biology and Medicine*, **34**(5), 2557–2564. doi: [10.1007/s13277-013-0801-4](https://doi.org/10.1007/s13277-013-0801-4)
- Forbes, A. and Lehmann, R. (1998). Nanos and Pumilio have critical roles in the development and function of *Drosophila* germline stem cells. *Development*, **125**(4), 679–690. doi: [10.1242/dev.125.4.679](https://doi.org/10.1242/dev.125.4.679)
- Gennarino, V. A., Singh, R. K., White, J. J., De Maio, A., Han, K., Kim, J. Y., Jafar-Nejad, P., di Ronza, A., Kang, H., Sayegh, L. S., Cooper, T. A., Orr, H. T., Sillitoe, R. V. and Zoghbi, H. Y. (2015). Pumilio1 haploinsufficiency leads to SCA1-like neurodegeneration by increasing wild-type ataxin1 levels. *Cell*, **160**(6), 1087–1098. doi: [10.1016/j.cell.2015.02.012](https://doi.org/10.1016/j.cell.2015.02.012)
- Goldstrohm, A. C., Hall, T. M. T. and McKenney, K. M. (2018). Post-transcriptional regulatory functions of mammalian pumilio proteins. *Trends in Genetics: TIG*, **34**(12), 972–990. doi: [10.1016/j.tig.2018.09.006](https://doi.org/10.1016/j.tig.2018.09.006)
- Jenkins, H. T., Baker-Wilding, R. and Edwards, T. A. (2009). Structure and RNA binding of the mouse Pumilio-2 PUF domain. *Journal of Structural Biology*, **167**(3), 271–276. doi: [10.1016/j.jsb.2009.06.007](https://doi.org/10.1016/j.jsb.2009.06.007)
- Joshna, C. R., Saha, P., Atugala, D., Chua, G. and Muench, D. G. (2020). Plant PUF RNA-binding proteins: A wealth of diversity for post-transcriptional gene regulation. *Plant Science: An International Journal of Experimental Plant Biology*, **297**, 110505. doi: [10.1016/j.plantsci.2020.110505](https://doi.org/10.1016/j.plantsci.2020.110505)
- Ko, C. F., Chang, Y. C., Cho, H. C. and Yu, J. (2022). The Puf-A protein is required for primordial germ cell development. *Cells*, **11**(9), 1476. doi: [10.3390/cells11091476](https://doi.org/10.3390/cells11091476)
- Kraemer, B., Crittenden, S., Gallegos, M., Moulder, G., Barstead, R., Kimble, J. and Wickens, M. (1999). NANOS-3 and FBF proteins physically interact to control the sperm-oocyte switch in *Caenorhabditis elegans*. *Current Biology: CB*, **9**(18), 1009–1018. doi: [10.1016/s0960-9822\(99\)80449-7](https://doi.org/10.1016/s0960-9822(99)80449-7)
- Kresoja-Rakic, J. and Santoro, R. (2019). Nucleolus and rRNA gene chromatin in early embryo development. *Trends in Genetics: TIG*, **35**(11), 868–879. doi: [10.1016/j.tig.2019.06.005](https://doi.org/10.1016/j.tig.2019.06.005)
- Kuo, M. W., Wang, S. H., Chang, J. C., Chang, C. H., Huang, L. J., Lin, H. H., Yu, A. L., Li, W. H. and Yu, J. (2009). A novel *puf-A* gene predicted from evolutionary analysis is involved in the development of eyes and primordial germ-cells. *PLoS ONE*, **4**(3), e4980. doi: [10.1371/journal.pone.0004980](https://doi.org/10.1371/journal.pone.0004980)
- Li, X., Zhu, M., Zang, M., Cao, D., Xie, Z., Liang, H., Bian, Z., Zhao, T., Hu, Z. and Xu, E. Y. (2022). PUMILIO-mediated translational control of somatic cell cycle program promotes folliculogenesis and contributes to ovarian cancer progression. *Cellular and Molecular Life Sciences: CMLS*, **79**(5), 279. doi: [10.1007/s00018-022-04254-w](https://doi.org/10.1007/s00018-022-04254-w)
- Liang, X., Hart, K. J., Dong, G., Siddiqui, F. A., Sebastian, A., Li, X., Albert, I., Miao, J., Lindner, S. E. and Cui, L. (2018). Puf3 participates in ribosomal biogenesis in malaria parasites. *Journal of Cell Science*, **131**(6). doi: [10.1242/jcs.212597](https://doi.org/10.1242/jcs.212597)
- Lin, H. and Spradling, A. C. (1997). A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the *Drosophila* ovary. *Development*, **124**(12), 2463–2476. doi: [10.1242/dev.124.12.2463](https://doi.org/10.1242/dev.124.12.2463)
- Lin, K., Zhang, S., Shi, Q., Zhu, M., Gao, L., Xia, W., Geng, B., Zheng, Z. and Xu, E. Y. (2018). Essential requirement of mammalian Pumilio family in embryonic development. *Molecular Biology of the Cell*, **29**(24), 2922–2932. doi: [10.1091/mbc.E18-06-0369](https://doi.org/10.1091/mbc.E18-06-0369)
- Lin, K., Qiang, W., Zhu, M., Ding, Y., Shi, Q., Chen, X., Zsiros, E., Wang, K., Yang, X., Kurita, T. and Xu, E. Y. (2019). Mammalian Pum1 and Pum2 control body size via translational regulation of the cell cycle inhibitor Cdkn1b. *Cell Reports*, **26**(9), 2434–2450.e6. doi: [10.1016/j.celrep.2019.01.111](https://doi.org/10.1016/j.celrep.2019.01.111)
- Lin, H. W., Lee, J. Y., Chou, N. L., Shih, T. W. and Chang, M. S. (2021). Phosphorylation of PUF-A/PUM3 on Y259 modulates PUF-A stability and cell proliferation. *PLoS ONE*, **16**(8), e0256282. doi: [10.1371/journal.pone.0256282](https://doi.org/10.1371/journal.pone.0256282)
- Lu, G., Dolgner, S. J. and Hall, T. M. (2009). Understanding and engineering RNA sequence specificity of PUF proteins. *Current Opinion in Structural Biology*, **19**(1), 110–115. doi: [10.1016/j.sbi.2008.12.009](https://doi.org/10.1016/j.sbi.2008.12.009)
- Luo, W., Ke, H., Liu, R., Qin, Y., Mak, W., Ma, J., Zhao, S. and Chen, Z. J. (2018). Variation analysis of *PUM1* gene in Chinese women with primary ovarian insufficiency. *Journal of Assisted Reproduction and Genetics*, **35**(4), 727–731. doi: [10.1007/s10815-017-1110-4](https://doi.org/10.1007/s10815-017-1110-4)
- Mak, W., Fang, C., Holden, T., Dratver, M. B. and Lin, H. (2016). An important role of pumilio 1 in regulating the development of the mammalian female germline. *Biology of Reproduction*, **94**(6), 134. doi: [10.1095/biolreprod.115.137497](https://doi.org/10.1095/biolreprod.115.137497)
- Mak, W., Xia, J., Cheng, E. C., Lowther, K. and Lin, H. (2018). A role of Pumilio 1 in mammalian oocyte maturation and maternal phase of embryogenesis. *Cell and Bioscience*, **8**, 54. doi: [10.1186/s13578-018-0251-1](https://doi.org/10.1186/s13578-018-0251-1)
- Matoulikova, E., Michalova, E., Vojtesek, B. and Hrstka, R. (2012). The role of the 3' untranslated region in post-transcriptional regulation of protein expression in mammalian cells. *RNA Biology*, **9**(5), 563–576. doi: [10.4161/rna.20231](https://doi.org/10.4161/rna.20231)
- Miles, W. O., Tschöp, K., Herr, A., Ji, J. Y. and Dyson, N. J. (2012). Pumilio facilitates miRNA regulation of the *E2F3* oncogene. *Genes and Development*, **26**(4), 356–368. doi: [10.1101/gad.182568.111](https://doi.org/10.1101/gad.182568.111)
- Munschauer, M., Nguyen, C. T., Sirokman, K., Hartigan, C. R., Hogstrom, L., Engreitz, J. M., Ulirsch, J. C., Fulco, C. P., Subramanian, V., Chen, J., Schenone, M., Guttman, M., Carr, S. A. and Lander, E. S. (2018). The NORAD lncRNA assembles a topoisomerase complex critical for genome stability. *Nature*, **561**(7721), 132–136. doi: [10.1038/s41586-018-0453-z](https://doi.org/10.1038/s41586-018-0453-z)
- Najdrová, V., Stairs, C. W., Vinopalová, M., Voleman, L. and Doležal, P. (2020). The evolution of the puf superfamily of proteins across the tree of eukaryotes. *BMC Biology*, **18**(1), 77. doi: [10.1186/s12915-020-00814-3](https://doi.org/10.1186/s12915-020-00814-3)

- Naudin, C., Hattabi, A., Michelet, F., Miri-Nezhad, A., Benyoucef, A., Pflumio, F., Guillonnet, F., Fichelson, S., Vigon, I., Dusanter-Fourt, I. and Lauret, E. (2017). PUMILIO/FOXP1 signaling drives expansion of hematopoietic stem/progenitor and leukemia cells. *Blood*, **129**(18), 2493–2506. doi: [10.1182/blood-2016-10-747436](https://doi.org/10.1182/blood-2016-10-747436)
- Nüsslein-Volhard, R. L. C. (1987). Involvement of the pumilio gene in the transport of an abdominal signal in the *Drosophila* embryo. *Nature*.
- Parisi, M. and Lin, H. (1999). The *Drosophila* pumilio gene encodes two functional protein isoforms that play multiple roles in germline development, gonadogenesis, oogenesis and embryogenesis. *Genetics*, **153**(1), 235–250. doi: [10.1093/genetics/153.1.235](https://doi.org/10.1093/genetics/153.1.235)
- Schultz, R. M., Stein, P. and Svoboda, P. (2018). The oocyte-to-embryo transition in mouse: Past, present, and future. *Biology of Reproduction*, **99**(1), 160–174. doi: [10.1093/biolre/iy013](https://doi.org/10.1093/biolre/iy013)
- Silva, I. L. Z., Kohata, A. A. and Shigunov, P. (2022). Modulation and function of Pumilio proteins in cancer. *Seminars in Cancer Biology*, **86**(3), 298–309. doi: [10.1016/j.semcancer.2022.03.010](https://doi.org/10.1016/j.semcancer.2022.03.010)
- Son, S. H., Jang, S. Y. and Park, H. S. (2021). Functions of PUF family RNA-binding proteins in *Aspergillus nidulans*. *Journal of Microbiology and Biotechnology*, **31**(5), 676–685. doi: [10.4014/jmb.2101.01011](https://doi.org/10.4014/jmb.2101.01011)
- Takahiro, M., Kimura, Y., Nakano, T. and Yamaguchi, S. (2021). Ribosomal stress induces 2-cell embryo-like state transition of the mouse ESCs through p53 activation. *Biochemical and Biophysical Research Communications*, **579**, 175–180. doi: [10.1016/j.bbrc.2021.09.068](https://doi.org/10.1016/j.bbrc.2021.09.068)
- Vessey, J. P., Schoderboeck, L., Gingl, E., Luzi, E., Riefler, J., Di Leva, F., Karra, D., Thomas, S., Kiebler, M. A. and Macchi, P. (2010). Mammalian Pumilio 2 regulates dendrite morphogenesis and synaptic function. *Proceedings of the National Academy of Sciences of the United States of America*, **107**(7), 3222–3227. doi: [10.1073/pnas.0907128107](https://doi.org/10.1073/pnas.0907128107)
- Wang, M., Ogé, L., Perez-Garcia, M. D., Hamama, L. and Sakr, S. (2018). The PUF protein family: Overview on PUF RNA targets, biological functions, and post transcriptional regulation. *International Journal of Molecular Sciences*, **19**(2). doi: [10.3390/ijms19020410](https://doi.org/10.3390/ijms19020410)
- Wang, X., Zamore, P. D. and Hall, T. M. (2001). Crystal structure of a Pumilio homology domain. *Molecular Cell*, **7**(4), 855–865. doi: [10.1016/s1097-2765\(01\)00229-5](https://doi.org/10.1016/s1097-2765(01)00229-5)
- Xu, E. Y., Chang, R., Salmon, N. A. and Reijo Pera, R. A. (2007). A gene trap mutation of a murine homolog of the *Drosophila* stem cell factor Pumilio results in smaller testes but does not affect litter size or fertility. *Molecular Reproduction and Development*, **74**(7), 912–921. doi: [10.1002/mrd.20687](https://doi.org/10.1002/mrd.20687)
- Yan, Z., Liang, H., Deng, L., Long, H., Chen, H., Chai, W., Suo, L., Xu, C., Kuang, Y., Wu, L., Lu, S. and Lyu, Q. (2015). Eight-shaped hatching increases the risk of inner cell mass splitting in extended mouse embryo culture. *PLOS ONE*, **10**(12), e0145172. doi: [10.1371/journal.pone.0145172](https://doi.org/10.1371/journal.pone.0145172)
- Yoon, D. S., Lee, K. M., Choi, Y., Ko, E. A., Lee, N. H., Cho, S., Park, K. H., Lee, J. H., Kim, H. W. and Lee, J. W. (2022). TLR4 downregulation by the RNA-binding protein PUM1 alleviates cellular aging and osteoarthritis. *Cell Death and Differentiation*, **29**(7), 1364–1378. doi: [10.1038/s41418-021-00925-6](https://doi.org/10.1038/s41418-021-00925-6)
- Zhang, B., Gallegos, M., Puoti, A., Durkin, E., Fields, S., Kimble, J. and Wickens, M. P. (1997). A conserved RNA-binding protein that regulates sexual fates in the *C. elegans* hermaphrodite germ line. *Nature*, **390**(6659), 477–484. doi: [10.1038/37297](https://doi.org/10.1038/37297)
- Zhang, M., Chen, D., Xia, J., Han, W., Cui, X., Neuenkirchen, N., Hermes, G., Sestan, N. and Lin, H. (2017). Post-transcriptional regulation of mouse neurogenesis by Pumilio proteins. *Genes and Development*, **31**(13), 1354–1369. doi: [10.1101/gad.298752.117](https://doi.org/10.1101/gad.298752.117)
- Zhao, T., Xiao, T., Cao, D., Xia, W., Gao, L., Cheng, L., Zang, M., Li, X. and Xu, E. Y. (2022). Sertoli cell PUMILIO proteins modulate mouse testis size through translational control of cell cycle regulators. *Biology of Reproduction*, **107**(1), 135–147. doi: [10.1093/biolre/iacl18](https://doi.org/10.1093/biolre/iacl18)