

## $\gamma^2$ Velorum revisited

Orsola De Marco and Werner Schmutz

*Institute of Astronomy, Federal Institute of Technology,  
Zürich, Switzerland*

Lars Koesterke and Wolf-Rainer Hamann

*Department of Physics, University of Potsdam, Germany*

**Abstract.** We have carried out a full spectroscopic analysis of the WC8 star in the  $\gamma^2$  Vel binary (WR 11). Through the binary radial-velocity curve the mass of the WR star can be determined, and hence its luminosity can be derived through the mass-luminosity relation for WR stars. This can be compared to the luminosity from the spectroscopic analysis. We find that the standard modeling methods underestimate the luminosity of the WR star.

### 1. Analysis

De Marco & Schmutz (1999) determined the parameters of the O star in the  $\gamma^2$  Vel binary, by fitting simultaneously the absorption lines of the O star and the WR/O light-ratio. In Table 1 the O star model parameters are summarized. The un-blended WR star spectrum is then recovered by subtracting the synthetic O star spectrum and scaling according to the WR/O light-ratio. The de-convolved spectra of the O and WR stars are shown by Schmutz & De Marco (these Proceedings).

The WR star is modeled using the Kiel-code which implements the ‘standard model’ in the co-moving frame (Koesterke & Hamann 1995). The final model parameters are listed in Table 1, while example fits to the de-convolved WR 11 spectrum are shown in Fig. 1. Helium lines are well fitted. The Balmer lines are under-fitted, which may point to the presence of hydrogen, possibly due to the contribution of the stellar wind from the O star. Not all lines of carbon are well fitted.




Figure 1. Example fits (solid line) to WR 11 (dotted line).

Table 1. O and WR stellar parameters. For the system is adopted:  $d = 258_{-31}^{+41}$  pc,  $M_V = -5.39$  mag,  $\Delta M = (1.47 \pm 0.13)$  mag and  $i = (63^\circ \pm 8^\circ)$ .

	O star	WR star
$M_V$ (mag)	$-5.14 \pm 0.16$	$-3.67 \pm 0.16$
$T_{\text{eff}}$ (K)	$35\,000 \pm 300$	$75\,900 \pm 500$
N(He) by number	0.087	0.875
N(C) by number	–	0.125
$R$ ( $R_\odot$ )	$12.4 \pm 1.7$	$1.84 \pm 0.2$
$\log(L/L_\odot)$	$5.32 \pm 0.2$	5.01
$\log(L/L_\odot)$ M-L	–	5.18
$\mathcal{M}$ ( $M_\odot$ )	$30 \pm 2^a$	$9 \pm 2$
$\log(\dot{M}/M_\odot \text{ yr}^{-1})$	$-6.75 \pm 0.09^a$	-4.33
$v_\infty$ ( $\text{km s}^{-1}$ )	$2500 \pm 250^a$	$1300 \pm 150$

<sup>a</sup>hydrodynamical wind solution

## 2. Conclusions

WR 11 appears to be particularly hot for its WC8 spectral class. The model luminosity is lower than that inferred from its mass and the mass-luminosity relation (Schaerer & Maeder 1992; Table 1). When introducing line-blanketing (a necessary step to determine the line-force), this will be lower still, indicating that the model has too low a luminosity. Increasing the luminosity of the models, while maintaining the same synthetic spectrum and  $V$  brightness is one of the achievements of the ‘photon loss’ mechanism (Schmutz 1997, see also ‘ $\gamma^2$  Vel, photon loss and the velocity field’ by De Marco *et al.*, these Proceedings).

## References

- De Marco, O., Schmutz, W. 1999, A&A 345, 163  
 Koesterke, L., Hamann, W.-R. 1995, A&A 299, 503  
 Schmutz, W. 1997, A&A 321, 268