

EPV1284

A cardinal approach to evaluating the effectiveness of psychosocial rehabilitation based on the analytical hierarchy process

V. Mitikhin*, T. Solokhina, G. Tiumenkova and M. Kuzminova

FSBSI “Mental Health Research Centre”, Department Of Mental Health Services, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1944

Introduction: The approach to evaluating the effectiveness of psychosocial rehabilitation (PSR) of mentally ill patients implies a multidimensional, hierarchical consideration of mental pathology with the inclusion of clinical and psychopathological, socio-environmental and personal indicators when setting goals and objectives of PSR

Objectives: Development of algorithms and models for evaluating the effectiveness of PSR taking into account: clinical, socio-demographic, psychological characteristics of the patient and the characteristics of the family environment, as well as factors of psychiatric care.

Methods: Clinical, statistical, algorithms of the analytical hierarchy process (AHP) [1].

Results: Numerical estimates of changes (before and after the PSR program) in the main areas of patient’s functioning disorders, such as motivation, cognition, compliance, coping strategies, family, skills, immediate environment, and others, are proposed as particular criteria for evaluating the effectiveness of PSR. Quantitative estimates for particular criteria are obtained on the basis of rank estimates, which are converted into numerical ones based on AHP algorithms [1]. Quantitative integral estimates of the effectiveness of PSR are obtained on the basis of partial estimates, taking into account the weight of the corresponding areas of impaired functioning of patients.

Conclusions: The developed approach opens up prospects for obtaining numerical, partial and integral estimates based on various rank scales, which are of interest from the point of view of forming criteria-indicators (markers) of the effectiveness of psychosocial, rehabilitation, psychoeducational and psychotherapeutic measures. References: 1. Mitikhin, V.G., Solokhina, T.A. S.S. *Korsakov Journal of Neurology and Psychiatry*. 2019; 119(2): 49-54

Disclosure: No significant relationships.

Keywords: rehabilitation; analytic hierarchy process; effectiveness; psychosocial

EPV1283

Profile of patients referred to the active rehabilitation unit in the health area of Soria (Spain)

R. Martínez Gallardo^{1,2*}, N. Palomar Ciria², B. Valiente Peña¹ and N. Hernandez Vicente¹

¹Complejo Hospitalario de Soria, Psiquiatra, Soria, Spain and

²Complejo Hospitalario, Psiquiatría, Soria, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1945

Introduction: Realizamos un estudio prospectivo en 15 pacientes, diagnosticados de psicosis, según el DSM V, y que son usuarios de una Unidad de Rehabilitación activa, estos pacientes utilizan las modalidades de hospitalización residencial o el programa de hospitalización parcial.

Objectives: El objetivo principal fue establecer el perfil del paciente (retrato de robot), que utiliza este tipo de dispositivo.

Methods: Se lleva a cabo un protocolo de evaluación que incluye la entrevista de evaluación psicopatológica (DSMV) y las escalas: impulsividad (Barrat), agresividad (Burke), calidad de vida (Woqol bref), actitud ante la medicación (DAI), Avd (RAI), riesgo de suicidio (Plutchick), riesgo de caídas (Downton), funcionalidad (EEAG) y hábitos de consumo (DAU).

Results: Con todos estos elementos y teniendo en cuenta la edad y sexo de los pacientes, intentamos establecer el tipo de perfil de paciente del área de Rehabilitación activa de nuestro servicio, luego luego de mediciones periódicas de todas estas variables estableceremos la influencia de Terapia de rehabilitación en la mejora o empeoramiento de nuestros pacientes.

Conclusions: CONCLUSIONES El perfil de tipo del paciente incluido en una Unidad de Rehabilitación Activa está compuesto por un varón de 42 años, con consumo esporádico de toxinas, con rasgos de personalidad donde predomina la agresividad, con un perfil bajo de efectos adversos y con conciencia parcial de la enfermedad. y mala adherencia a la medicación Cabe señalar que sus niveles en la escala de calidad de vida son altos, incluso comparables o en ocasiones superiores a los de la población general según la escala Woqol bref.

Disclosure: No significant relationships.

EPV1284

Hospital environment and patient recovery – a review

A. Lourenço*, M. Ribeiro and M. Lemos

Centro Hospitalar Lisboa Norte, Psychiatry, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1946

Introduction: Since the second half of the 20th century, studies have been carried out that prove the benefits of the hospital environment in the improvement and recovery of patients. In this way, it would be important to understand what has already been done within the reality of Psychiatry Department.

Objectives: To review the literature about the documentation of hospital environment influence in patients’ recovery.

Methods: We performed a MEDLINE search using the key words: hospital environment influence and patients’ recovery or patients healing. We only included studies with full text published in English.

Results: In the selected articles, we only found studies developed in the Surgery department; in one of them, the authors tested the presence of plants in the patients’ rooms and assessed lower blood pressure values, less pain and less anxiety than the control group; in another, they tested the presence of music and landscape, although there was no difference in terms of pain assessment, there was an improvement in the assessment of the postoperative experience. A review pointed some other aspects that patients linked with their recovery, such as: audio and visual environment; specifically, in anxiety, pain and stress. On the other hand, other studies address the influence of the hospital environment on the satisfaction of health care providers.

Conclusions: Although hospital environment has already demonstrated an impact on the patient recovery, none (in our review) was developed directly in a Psychiatric Department; further studies are needed to understand the impact on this kind of service.