

Effect of egg consumption on early childhood development: Evidence from *Un Oeuf* study

Helen Ernyey¹, Chhavi Tiwari², Heather Stark³, Emma Hunniford³, Aissata Wereme N'Diaye⁴, Yacouba Zare⁴, Anteneh Omer⁵, Sarah Lindley McKune^{2,6}

¹College of Medicine, University of Florida, Gainesville, FL, USA

²Department of Environmental and Global Health, College of Public Health and Health Professions, University of Florida, Gainesville, FL, USA

³Department of Epidemiology, College of Public Health and Health Professions, University of Florida, Gainesville, FL, USA

⁴Laboratoire de Recherche en Production et Santé Animales (LaRePSA), Institut de l'Environnement et de Recherches Agricoles (INERA), Centre National de Recherches Scientifique et Technologique –(CNRST) Ouagadougou, Burkina Faso

⁵Independent Nutrition Consultant, Hawassa, Ethiopia

⁶The Center for African Studies, University of Florida, Gainesville, FL, USA

Corresponding author: Sarah Lindley McKune, Mailing Address : 1225 Center Drive, PO BOX 100182, Gainesville, FL, USA 32611, Email: smckune@ufl.edu



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its

DOI 10.1017/S1368980024002490

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Short title : Egg consumption and early childhood development

Acknowledgements

We would like to acknowledge Drs. Patricia Snyder and Maureen Conroy from the University of Florida's Anita Zucker Center for Excellence in Early Child Development Studies for their contributions in modifying the ASQ for implementation in Burkina Faso. We would also like to acknowledge our study team and trained enumerators in Burkina Faso for their dedication and commitment to this study: Mahamoudoudo Ibriga, Zare Yacouoba, Iliasse Tiemtore, Christian Oueoraogo, and Aida Sanou.

Financial Support

This work was funded in whole or part by the United States Agency for International Development (USAID) Bureau for Resilience, Environment and Food Security under Agreement # AID-OAA-L-15-00003 as part of Feed the Future Innovation Lab for Livestock Systems. Additional funding was received from Bill & Melinda Gates Foundation OPP#060115. Any opinions, findings, conclusions, or recommendations expressed here are those of the authors alone.

Conflict of interest

We wish to confirm that there are no known conflicts of interest associated with this publication and that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

Authorship

HS, SM, AO and AW conceived of the study. HS, SM, AO and AW initiated the study design, and HS, SM, AO, AW and EM were involved in implementation. CT conducted statistical analyses and finalized tables. HE led the write up of this manuscript, with significant contribution by SM, HS, AO, CT, YZ and EH. All authors contributed to the refinement of the manuscript and approved the final manuscript.

Abstract

Objective: Recent studies have shown that inclusion of eggs in young children's diet can help meet nutritional requirements associated with cognitive development. This study aims to investigate the effect of egg consumption on Early Childhood Development (ECD) using Ages and Stages Questionnaire-3(ASQ-3) in Burkina Faso.

Design: The study presented here uses data collected during a follow-up of the *Un Oeuf*-a 3 arm clustered randomized controlled trial (RCT), conducted roughly four months after the end of the RCT.

Setting: This research was conducted in 18 rural villages within the Kaya Department of the Sanmatenga Province in Burkina Faso.

Participants: Participants of this study include a total of 244 children aged between 18-33 months, with 78 children in the full intervention group, 83 in the partial group, and 83 in the control group.

Results: Results show that children with consistent egg consumption (in all months) had a lower odd of falling below the cut-off scores in gross motor ($OR = 0.13, p = 0.02$) and personal social skills ($OR = 0.34, p = 0.05$). And a dose response was established; for each additional egg/week, a 1.9% increase in scores for problem-solving skills was observed.

Conclusions: Findings from this study contribute to a growing body of evidence that increasing egg consumption among children in LMICs can improve growth and development. The study highlights the need for additional research in LMICs to better understand the multifactorial relationship between diet and childhood development.

Keywords: early childhood development, ASQ-3, clustered randomized controlled trial, eggs, animal source food, infant and child nutrition, low-income countries

Introduction

Early childhood development (ECD) was included in the United Nations' Sustainable Development Goals (SDGs) in 2015, signifying a global commitment to ensure equitable access to early learning opportunities and childhood development for all children ⁽¹⁾. Early childhood offers a critical window of opportunity that can shape the lifetime trajectory of a child's holistic development, including academic achievement, occupational success, and social adjustment ^(2,3). Delay in ECD is more common in low- and middle-income countries (LMICs), with 250 million children-under-five (CU5) at risk of not achieving their developmental potential ⁽⁴⁾, and around 42% of children experiencing childhood delays living in West Africa ⁽⁵⁾.

ECD is impacted by myriad factors, including diet, environmental, and social risk exposures ^(6,7). One of the important underlying reasons for the high prevalence of ECD delays is underlying poor nutrition and food insecurity in LMICs, which emerges from lack of access to resources alongside inadequate and inequitable food production ⁽⁸⁾. Undernutrition during childhood leads to adverse health effects, including stunting, decreased cognitive functioning, and increased morbidity and mortality ^(7,9,10). In Burkina Faso, over 672,000 children suffer from chronic malnutrition ⁽¹¹⁾ with 22.6% children under 5 suffering from stunting ⁽¹²⁾. The high prevalence of stunting in Burkina Faso raises significant concerns for policymakers and necessitates additional research on nutrition and ECD in this context.

While meeting the nutritional requirement of early age is a challenge in most LMICs due to lack of resources, access to nutritious food ⁽¹³⁾, and suboptimal feeding practices ⁽¹⁴⁾, it has been shown that including animal source foods (ASF) such as eggs, meat, fish, or dairy in otherwise typical diets has the potential to improve a child's nutritional and health outcomes ⁽¹⁵⁾. However, the consumption of ASF in LMICs has remained low, likely due to poor availability, high relative cost ⁽¹⁶⁾, and restrictive food norms and taboos ⁽¹⁷⁾. Nutritional interventions and behavior change campaigns have been shown to reduce or remove the social and cultural barriers to consuming certain food items that may significantly impact early childhood growth and development ⁽¹⁸⁻²⁰⁾.

Like other ASF, eggs are rich in macro and micronutrients^(21,22). Consuming one egg per day for a child aged 6-24 months can provide adequate nutrients that are essential for child nutrition and brain development⁽²³⁾.

Based on evidence from Ecuador⁽²⁴⁾ that found significant improvements in child growth when infants were fed eggs, the *Un Oeuf* study—a clustered randomized controlled trial—was conducted in 18 rural villages of Burkina Faso in 2018-2019, with the primary objective of increasing egg consumption among infants and young children through a behavior change communication intervention⁽¹⁹⁾. In Kaya Department, the *Un Oeuf* study randomly selected 18 non-urban villages and then randomly assigned (1:1:1) the villages to one of the three intervention arms. In the full intervention arm, children were gifted three chickens from a community champion and a fourth chicken from their fathers, and caregivers received a behavior change communication (BCC) package, including monthly trainings on integrated nutrition and agriculture (INA). At a gifting ceremony, where the child received the chickens, village leaders presented the chickens as gifts to the child's flock, reinforcing an understanding that the chickens and any eggs produced belonged to the child. One egg per day from these chickens was advised to be fed to the child. While the project did not intervene on the sale or sharing of eggs, mothers were trained on the significance of diet during the first 1000 days and asked to see the egg as a treatment for or investment in the targeted child. They were also advised that only after the child's one egg/day "dose" had been achieved, might the mother consider feeding other children or selling the eggs. The partial intervention group received only the BCC package and did not receive any chickens. The control group did not receive any intervention; neither BCC nor chickens. The intervention, which included monthly BCC trainings for the full and partial intervention groups, continued for 9 months with baseline data collection in the month of July 2018 and the endline data collection in April 2019. The trial was registered at clinicaltrials.gov: NCT04135625. For a detailed protocol of the study design, methods, and baseline characteristics, see Stark et al., 2021. The *Un Oeuf* study population ate little to no eggs at baseline, however the intervention significantly increased egg consumption in the full and partial groups; the full intervention, in which children consumed around six eggs per week by end line, also significantly decreased wasting and underweight⁽¹⁹⁾. As the *Un Oeuf* study significantly improved egg consumption and nutritional status of children, additional funding was secured to

investigate the potential impact of the study on ECD. The study presented here uses data collected during a follow-up of the *Un Oeuf* study, conducted roughly four months after the end of the RCT and engaging the same children who participated in the initial study. This study aims to investigate the effect of egg consumption on ECD using the Ages and Stages Questionnaire (ASQ-3; Squires & Bricker, 2009), modified by the study team for the local environment.

Methods

Study Design

The follow-up study was conducted four months after the completion of the *Un Oeuf* study to assess the impact of egg consumption on ECD in Burkina Faso. During the 9-month intervention period of the *Un Oeuf* study, monthly surveys were used to collect data on egg consumption and anthropometric measurements of enrolled children. At the beginning of the study, children of age 6-12 months were enrolled. Upon conclusion, these children were assessed for development outcomes in the follow-up study, using a culturally modified ASQ-3 tool and anthropometric measurements of height, length, weight, and head circumference.

Variables

Child development measures

ASQ-3 is a widely used standardized screening instrument to measure development progress in children between the ages of one month to 66 months^(27–30). The research team selected and adapted the ASQ tool to assess ECD in Burkina Faso in collaboration with scientists at the Anita Zucker Center for Excellence in Early Child Development Studies. Guided by these experts, changes in the ASQ-3 tool were made to allow for appropriate assessment of an indicator, while utilizing materials that were more familiar to children in the Burkinabe context. These included substituting a stick that the children could use to write/draw in the dirt, since they did not have access to pencils and papers; using small pebbles instead of cheerios; and stepping onto logs instead of climbing steps. Researchers then trained the enumerators from the *Un Oeuf* study to use the modified ASQ-3 tool, after working with them to review and revise the proposed tool for cultural salience. The questionnaire was then pilot-tested with children of appropriate ages using set criteria established to assess the various questions to improve internal reliability and validity

prior to administering the ASQ-3 assessment. The ASQ-3 assessment was verbally administered to the child's mother. In accord with ASQ-3 instruction, the enumerator asked the mother to respond to the questions and, in some cases, asked the child to perform an activity. These data were recorded by the enumerator. ASQ items were observed by the enumerator or reported by a parent and consist of 30 items scored as *yes*, *sometimes*, or *not yet* across five domains: communication, gross motor, fine motor, problem-solving, and personal social skills. Within each domain, individuals can score from 0-60, with specific scores above or below the cut-off for expected childhood development. We examined cut-off scores in each domain of development. These domain-specific cut-off scores align with ASQ-3 guidelines, which use ASQ score and age of the child to determine the cut offs. A binary indicator was used to describe whether a child's score was below the cutoff value (1) or not (0). Following methodological approach of previous ECD research⁽²⁷⁾, we looked at domain-specific scores, as well as the overall total score (sum of the five domain-specific scores).

Egg consumption measures

After the baseline survey, the first two months' data quality was compromised due to heavy rains and inaccessibility to the study site. This study therefore used longitudinal information for egg consumption for seven months (months 3-9) with baseline variables measured at the beginning (month 0) and child development scores measured in the follow up (at month 12). In each month of the intervention period (month 3 through month 9), respondents were asked if they fed eggs to the child in the previous month. We created a categorical variable for consistency in egg consumption as: 1=*Never*, 2=*1-3 months*; 3=*4-6 months* and 4=*in all seven months*. Further, if/when respondents indicated that eggs were fed to the child, they were also asked how many eggs were fed in the previous week. Based on their responses, an average weekly egg consumption is calculated using seven months of data. The measure calculates on average, how many eggs the child ate per week in the last seven months. Therefore, we check for the effect of both consistency over time and quantity of egg consumption on child development outcomes.

As child development scores were only assessed after the trial ended, the analysis adjusted available confounding factors at baseline that are identified as determinants of child development in existing literature⁽³¹⁾. These confounders include child's gender, birth order, baseline weight

and height, mother's age at first birth, mother's education, household size and economic status (wealth index). The wealth index was calculated based on possession of assets and quality of housing using principal component analysis (PCA) following Demographic and Health Survey (DHS) guidelines and treated as tercile for modelling purposes. The items in the asset score included flooring material, cooking fuel, electricity, radio, television, cellphone, table, chair, mattress, solar panel, lamp, cycle, bike, and cart. Information for asset ownership was collected only in the seventh month of data collection. The study does not include child's own age as a confounder since Ages and Stages accounts for the age itself in score as well as the cut off values.

Statistical Analysis

All analysis was performed in STATA, v.17. Two separate regression analyses have been performed contingent on the nature of the dependent variable. First, logistic regressions were employed to calculate the odds of children falling below the domain-specific cutoff scores while accounting for potential confounders to egg consumption and child growth. Further, linear regression models were used for domain-specific scores and for the total scores. Logarithmic transformations of these scores were used to ensure normality in the distribution and to look at the percent change in scores due to egg consumption. In all models, standard errors were clustered at village level to account for village-level randomization and heterogeneity across villages. We also tested these models by including intervention arms, but since the intervention of the *Un Oeuf study* was specifically designed to increase egg consumption in children through behavior change, gifting chickens, and INA training, the intervention arms (full and partial) were found to be highly correlated with the egg consumption (correlation coefficients, $r = 0.92$ and $r = 0.70$, respectively), thereby, could not be included in the regression model. In addition, propensity score matching is used to ensure robustness of the results and deal with selection bias.

Results

Descriptive statistics

The final analysis included a total of 244 children aged between 18-33 months, with 78 children in the full intervention group, 83 in the partial group, and 83 in the control group. Descriptive

statistics for baseline characteristics, egg consumption, and child development outcomes across the three research arms are presented in Table 1.

In egg consumption variables, it was observed that children in the full intervention group consumed eggs more regularly than the partial intervention group and control group. In comparison, very few children consumed eggs in the control group. Average weekly egg consumption was also significantly higher in the full intervention group (6.4) as compared to the partial (1.76) and control group (0.22). Figure 1 shows a pattern of egg consumption across research arms in follow-up months.

Table 2 shows a comparison of average total ASQ score across research arms and the score was highest in the full intervention group (239.36), as compared to the partial intervention group (223.13) and the control group (223.37). Considering domain-specific outcomes, children in the full intervention group scored higher in almost all developmental domains (except personal social) than the children in the partial or control groups. Independent t-test showed that children in the full intervention arm had significantly higher mean scores in gross motor, fine motor, and problem-solving domains and in overall total scores compared to those in the control. However, in partial intervention arm, no significant difference in the development score was found when compared to the control arm.

Impact of egg consumption on early childhood development (ECD)

Table 3 reports adjusted Odds Ratios (OR) from logistic regressions (Col. 1-5) and coefficients (β) of linear regression models (Col. 6-11). After adjusting for baseline covariates (child's gender, birth order, child's anthropometry, mother's education, mother's age, household size, wealth index and baseline egg consumption), consistency in egg consumption was found to be statistically significant with lower odds of falling below the cut off in communication skills ($OR = 0.079, p = 0.06$), gross motor skills ($OR = 0.125, p < 0.05$) and for personal social skills ($OR = 0.34, p = 0.05$). Among the covariates, child's height at baseline is found to be associated with decreasing odds of falling below the cutoff scores in fine motor ($OR = 0.69, p < 0.001$), problem-solving ($OR = 0.82, p = 0.02$), and personal social ($OR = 0.82, = 0.06$) domains; and higher weight at baseline was associated with lower odds of falling below the cut off in communication skills ($OR = 0.37, p = 0.008$). The highest wealth index group was

associated with a significantly lower odds of falling below the cut off scores in gross motor ($OR = 0.19, p = 0.09$).

In multivariate regression models, there was no significant impact of consistency of egg consumption on domain specific ASQ scores or on total ASQ scores. Among the covariates, as in previous cases, child height at baseline remained a significant predictor of ASQ scores in children in all domains except problem-solving, with a 1 cm increase in height associated with 1 to 4% increase in development scores.

In the second stage of the analysis, we tested the quantity of egg consumption per week (Table 4). Weekly egg consumption was not found to be significantly associated with falling below the cut-off scores in any of the domain. Height was a significant predictor of early childhood development. Height was found to be significantly associated with lower odds of falling below the cut-off in fine motor, problem solving, and personal social skills.

For linear regression results (col 6-11), egg consumption was found to be significantly associated with problem solving skills where each increase in weekly egg consumption led to 1.9% increase in scores for problem solving skills.

In Figure 2, we present the predicted probability of falling below the cut off scores with respect to average weekly egg consumption based on logistic regressions. Results show that as weekly egg consumption increases, the probability of falling below the cut off scores for each domain specific ASQ scores declines. Noticeably, the impact is largest for gross motor, problem solving and personal social scores, with the steepest decline and with the largest negative change in the probability between 0 and 9 eggs per week corroborating our previous findings with consistency in egg consumption in Table 3.

Robustness check

Propensity score matching

Although we use data from a clustered randomized controlled trial where households in the two treatment arms and the control arm were similar at baseline with no significant differences in egg consumption among children, this study investigates the impact of higher egg consumption resulting from the intervention, not the intervention itself. Thus, there could be a potential of selection bias as the previous model does not account for the determinants of feeding eggs itself

and if there are systematic differences between households feeding eggs to the children and those who do not (e.g. more engagement with children, better educated, etc.). Therefore, to overcome this potential selection bias, the study uses Propensity Score Matching (PSM) to estimate the effect of treatment (egg consumption) on ECD. The treatment here is defined as binary and equal to “1” if the respondent reported feeding eggs to the child *in all the seven months*, and “0” otherwise. Propensity scores were generated using a multivariate logistic regression model with egg consumption in all the months as outcome variable and the baseline predictors as follows; produce enough eggs to feed child one egg per day, mother’s age at first birth, mother’s educational status, household size and wealth index. We also checked for some other determinants of feeding eggs related to mother’s knowledge, attitude, and practices (KAP) towards egg consumption. These include knowledge of nutritional importance of eggs, confidence in preparing eggs, difficulty in getting eggs, whether received nutritional knowledge about eggs, etc., but these factors did not appear to be statistically significant of feeding eggs, therefore were dropped from the model. PSM one to one matching was performed using nearest neighbor matching. The matching ensures that groups of households who fed their children and those who did not were systematically identical. For brevity (matching results are available on request)), here we only report the results for Average Treatment Effect on Treated (ATET) in Table 5. The coefficients in Table 5 indicate the difference in the scores among the treatment group (children who ate eggs in all the months) due to the treatment (egg consumption in all months). The significant increase in gross motor skills and fine motor skills aligns with our previously reported results in Table 3. However, we do not find significant increase in communication and personal social skills. Hence, the findings confirm the significant contribution of egg consumption on ECD outcomes after ensuring selection issues were controlled for.

Discussion

The empirical analysis in this paper provides supporting evidence for the promotion and inclusion of eggs in child diet to improve ECD. The challenges of food insecurity and malnutrition have been directly linked to the nutrient deficiencies and their contribution to learning and development deficits⁽³²⁾. Delays in ECD contribute to cognitive and motor development deficits, low educational attainment, and intergenerational transmission of poverty⁽³³⁾. Addressing malnutrition, especially in early childhood, is critical to the prevention of poor

cognitive and health outcomes. Findings highlighted the impact of full intervention and partial intervention as part of *Un Oeuf* trial on ECD scores. Children in the full intervention group who received chickens and BCC package reported highest ASQ scores in almost all domains (except personal social) than the children in partial intervention group (who only receive the BCC package) and the control group. Though the difference across partial and control group was not statistically significant indicating the importance of accessibility and availability of eggs over behavior change communication. Furthermore, the impact of eggs consumption in improving ECD is highlighted with this study with implications in LMICs.

The key findings of the paper hold statistical significance after baseline adjustment for three domains (communication, gross motor, and personal social) for consistency of egg consumption and for one domain (problem solving) for quantity of egg consumption. Result showed that each additional egg per week led to a 1.9% increase in problem solving scores in children establishing a dose response relationship between egg consumption and problem-solving skills scores.

Another important result was surrounding consistency of egg consumption. We find that children who consumed eggs consistently in all months (7) of intervention period were significantly less likely to fall below the cut-off score for communication, gross motor, and personal social skills, all of which are crucial elements of overall child development.

Our findings are consistent with Miller et al. (2020), who found egg consumption to be associated with lower odds of having ASQ score in the bottom quartile in Nepal. A study in Ethiopia also reported that children who had increased egg intake attained gross motor skills at a significantly earlier age compared to controls⁽³⁴⁾. Similar to ours, the study by Omer et al. (2022) focused on child owned poultry as compared to some other trials that distributed eggs directly to the children^(22,24). Similarly, a meta-analysis also found that micronutrients played a crucial role in cognitive performance among children aged 6-11 years⁽³⁵⁾, which suggests a possible mechanism underlying the findings reported here.

Our findings show that consistent consumption of eggs led to improved early childhood development in the domains of communication, gross motor, and personal social. Eggs contain numerous nutrients that play a vital role in neurocognitive development including iron, zinc, choline, folate, iodine and long-chain polyunsaturated fatty acids such as docosahexanoic acid (DHA)⁽³⁶⁾. The improved nutritional status as manifested by decreased wasting and underweight⁽¹⁹⁾ among the full intervention group might have also contributed to better ECD. Nutritional

status is found to have significant contribution on gross motor and fine motor skills though the findings have been mixed and not strong and consistent ^(37,38). Improved anthropometry is significantly associated with motor development and language skills ⁽³⁹⁾. Additionally previous studies ⁽⁷⁾ have noted significant association of motor development with executive functions, memory and later cognitive development in children. Therefore, the results of the study indicate the importance of better childhood nutrition and underscoring eggs as a complementary food option.

In Burkina Faso, undernutrition is one of the leading causes of morbidity in CU5 ⁽⁴⁰⁾. Behavioral and nutrition interventions, such as the *Un Oeuf* project, can be effective among vulnerable population for improved decision making, enhancing knowledge of nutrition, and improving access to livestock production resources. However, there are prevalent gaps in research as more information is needed to find appropriate and effective approaches to improve children's developmental and nutritional status at specific intervals of childhood development. Research conducted by Miller et al. (2020) states nutritional intervention not only affect the development outcomes immediately but also may be evident and permanent later in life. Thus, the extension of the study can be done by following the children in the later stages of childhood and adolescence and check how and if the effect of intervention continued over time.

This study also has a few limitations. First, ASQ-3 has its own reported rules ⁽⁴¹⁾ in capturing information and indicators of child development, including a stated purpose as a screening tool designed to recognize developmental delays but not affirm child development status. Like others who have used or modified the ASQ for research in LMICs, the research team believes that the ASQ provides meaningful information about different domains of development in children ^(28,30,42). Second, there could have been unique distractions or curiosities for children completing the evaluation that led to differences or biases in their recorded responses. For example, some children in these villages never had access to a mirror. Thus, for many, it was their first time seeing their reflection and could impact the ability to accurately assess childhood development. Third, as this study was a behavior change intervention, there is possibility of social desirability bias in reporting egg consumption especially in the full intervention group, although the questionnaire was administered to control for the same. Additionally, the weekly egg consumption measure is based on a 7 -day dietary recall with a potential to recall bias in the

dataset. Finally, other variables, such as environmental exposure, illness, other diet, maternal factors, water and sanitation etc., could have influenced child development during the study period, and were not directly measured in our study.

Conclusion

Early childhood development is closely related to the diets and nutritional status of children. Leveraging data collected during and just after the *Un Oeuf* intervention in Burkina Faso, this study found that both consistency and quantity of egg consumption led to observed improvements: greater consistency of egg consumption led to lower odds of falling below the ASQ-3 cut-off score for gross motor, personal social skills, and the total development score. The study also found that increase in the quantity of egg consumption led to an increase in scores in problem solving domain. These results underscore the importance of further research on dietary interventions in LMICs, with particular focus on complexity of nutrition as well as culturally appropriate tools that can measure child development in low resource settings in LMICs. Given the relative affordability and accessibility of eggs, integrating them into nutrition-focused public health programs could serve as a feasible and effective strategy to improve ECD outcomes in these countries

Ethical Standards Disclosure

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the University of Florida Institutional Review Board Committee of Ethics of the Government of Burkina. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

References

1. UNICEF (2023) Early Childhood Development Index 2030: A New Tool to Measure SDG Indicator 4.2.1. .
2. MAL-ED Network Investigators (2018) Early childhood cognitive development is affected by interactions among illness, diet, enteropathogens and the home environment: findings from the MAL-ED birth cohort study. *BMJ Glob Health* **3**, e000752.
3. Singh A, Yeh CJ & Boone Blanchard S (2017) Ages and Stages Questionnaire: a global screening scale. *Boletín Médico Del Hospital Infantil de México (English Edition)* **74**, 5–12.
4. Black MM, Walker SP, Fernald LCH, et al. (2017) Early childhood development coming of age: science through the life course. *The Lancet* **389**, 77–90.
5. Gil JD, Ewerling F, Ferreira LZ, et al. (2020) Early childhood suspected developmental delay in 63 low- and middle-income countries: Large within- and between-country inequalities documented using national health surveys. *Journal of Global Health* **10**, 010427.
6. Ibáñez-Alfonso JA, Company-Córdoba R, García De La Cadena C, et al. (2021) How Living in Vulnerable Conditions Undermines Cognitive Development: Evidence from the Pediatric Population of Guatemala. *Children* **8**, 90.
7. Larson LM, Martorell R & Bauer PJ (2018) A Path Analysis of Nutrition, Stimulation, and Child Development Among Young Children in Bihar, India. *Child Development* **89**, 1871–1886.
8. Company-Córdoba R, Sianes A, Simpson IC, et al. (2021) Cognitive interventions in children and adolescents from low socioeconomic status backgrounds: a systematic review protocol of randomized controlled trials. *Syst Rev* **10**, 187.
9. Nurliyana AR, Mohd Shariff Z, Mohd Taib MN, et al. (2016) Early nutrition, growth and cognitive development of infants from birth to 2 years in Malaysia: a study protocol. *BMC Pediatr* **16**, 160.

10. Larson LM & Yousafzai AK (2017) A meta-analysis of nutrition interventions on mental development of children under-two in low- and middle-income countries. *Maternal & Child Nutrition* **13**, e12229.
11. USAID (2018) *Burkina Faso: Nutrition Profile*. .
12. World Bank (2021) UNICEF, WHO, World Bank: Joint child Malnutrition Estimates (JME). .
13. Asare H, Rosi A, Faber M, et al. (2022) Animal-source foods as a suitable complementary food for improved physical growth in 6 to 24-month-old children in low- and middle-income countries: a systematic review and meta-analysis of randomised controlled trials. *Br J Nutr* **128**, 2453–2463.
14. Arabi M, Frongillo EA, Avula R, et al. (2012) Infant and Young Child Feeding in Developing Countries. *Child Development* **83**, 32–45.
15. Eaton JC, Rothpletz-Puglia P, Dreker MR, et al. (2019) Effectiveness of provision of animal-source foods for supporting optimal growth and development in children 6 to 59 months of age. *Cochrane Database of Systematic Reviews* **2019** [Cochrane Developmental, Psychosocial and Learning Problems Group, editor].
16. Headey D (2018) *Animal sourced foods and child nutrition in South Asia: Policy priorities*. LANSAs: Leveraging Institutions for Nutrition in South Asia.
17. Adesogan AT, Havelaar AH, McKune SL, et al. (2020) Animal source foods: Sustainability problem or malnutrition and sustainability solution? Perspective matters. *Global Food Security* **25**, 100325.
18. Gallegos-Riofrío CA, Waters WF, Salvador JM, et al. (2018) The Lulun Project’s social marketing strategy in a trial to introduce eggs during complementary feeding in Ecuador. *Maternal & Child Nutrition* **14**, e12700.
19. McKune SL, Stark H, Sapp AC, et al. (2020) Behavior Change, Egg Consumption, and Child Nutrition: A Cluster Randomized Controlled Trial. *Pediatrics* **146**, e2020007930.

20. Moore EV, Singh N, Serra R, et al. (2022) Household decision-making, women's empowerment, and increasing egg consumption in children under five in rural Burkina Faso: Observations from a cluster randomized controlled trial. *Front. Sustain. Food Syst.* **6**, 1034618.
21. Iannotti LL, Lutter CK, Bunn DA, et al. (2014) Eggs: the uncracked potential for improving maternal and young child nutrition among the world's poor. *Nutr Rev* **72**, 355–368.
22. Stewart ME (2018) The Effects of Consuming Eggs on the Physical and Cognitive Development of Food-Insecure Haitian Children. University of Arkansas.
23. Lutter CK, Iannotti LL & Stewart CP (2018) The potential of a simple egg to improve maternal and child nutrition. *Matern Child Nutr* **14**, e12678.
24. Iannotti LL, Lutter CK, Stewart CP, et al. (2017) Eggs in Early Complementary Feeding and Child Growth: A Randomized Controlled Trial. *Pediatrics* **140**, e20163459.
25. Stark H, Omer A, Wereme N'Diaye A, et al. (2021) The *Un Oeuf* study: Design, methods and baseline data from a cluster randomised controlled trial to increase child egg consumption in Burkina Faso. *Matern Child Nutr* **17**.
26. Squires J & Bricker D (2009) *Ages and Stages Questionnaire (ASQ): A Parent Completed Child Monitoring System*. 3rd ed. Baltimore, MD: Brooks Publishing Company.
27. Alvik A & Grøholt B (2011) Examination of the cut-off scores determined by the Ages and Stages Questionnaire in a population-based sample of 6 month-old Norwegian infants. *BMC Pediatr* **11**, 117.
28. Justino P, Leone M, Rolla P, et al. (2023) Improving Parenting Practices for Early Child Development: Experimental Evidence from Rwanda. *Journal of the European Economic Association* **21**, 1510–1550.
29. Miller AC, Rumaldo N, Soplapuco G, et al. (2021) Success at Scale: Outcomes of Community-Based Neurodevelopment Intervention (CASITA) for Children Ages 6–20 months With Risk of Delay in Lima, Peru. *Child Development* **92**, e1275–e1289.

30. Miller LC, Neupane S, Joshi N, et al. (2020) Diet quality over time is associated with better development in rural Nepali children. *Matern Child Nutr* **16**.
31. Mbabazi J, Pesu H, Mutumba R, et al. (2024) Correlates of early child development among children with stunting: A cross-sectional study in Uganda. *Maternal & Child Nutrition*, e13619.
32. Rosales FJ, Reznick JS & Zeisel SH (2009) Understanding the role of nutrition in the brain and behavioral development of toddlers and preschool children: identifying and addressing methodological barriers. *Nutritional Neuroscience* **12**, 190–202.
33. Drago F, Scharf RJ, Maphula A, et al. (2020) Psychosocial and environmental determinants of child cognitive development in rural south africa and tanzania: findings from the mal-ed cohort. *BMC Public Health* **20**, 505.
34. Omer A, Hailu D & Whiting SJ (2022) Egg consumption of children under two years of age through a child-owned poultry and nutrition intervention in rural Ethiopia: A community-based randomized controlled trial. *Journal of Agriculture and Food Research* **9**, 100354.
35. Meli AM, Ali A, Mhd Jalil AM, et al. (2021) Effects of Physical Activity and Micronutrients on Cognitive Performance in Children Aged 6 to 11 Years: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Medicina* **58**, 57.
36. Faber M, Malan L, Kruger HS, et al. (2022) Potential of Egg as Complementary Food to Improve Nutrient Intake and Dietary Diversity. *Nutrients* **14**, 3396.
37. Cavagnari BM, Guerrero-Vaca DJ, Carpio-Arias TV, et al. (2023) The double burden of malnutrition and gross motor development in infants: A cross-sectional study. *Clinical Nutrition* **42**, 1181–1188.
38. Zulkarnaen Z (2019) The Influence of Nutritional Status on Gross and Fine Motor Skills Development in Early Childhood. *ASS* **15**, 75.

39. Rico-González M, Ardigò LP, Ramírez-Arroyo AP, et al. (2024) Anthropometric Influence on Preschool Children's Physical Fitness and Motor Skills: A Systematic Review. *J Funct Morphol Kinesiol* **9**, 95.
40. Erismann S, Knoblauch AM, Diagbouga S, et al. (2017) Prevalence and risk factors of undernutrition among schoolchildren in the Plateau Central and Centre-Ouest regions of Burkina Faso. *Infect Dis Poverty* **6**, 17.
41. Rubio-Codina M, Araujo MC, Attanasio O, et al. (2016) Concurrent Validity and Feasibility of Short Tests Currently Used to Measure Early Childhood Development in Large Scale Studies. *PLoS ONE* **11**, e0160962 [Carpenter DO, editor].
42. Shrestha M, Ranjitkar S, Hysing M, et al. (2023) *The Relationship between the Ages and Stages Questionnaire, 3rd Edition Scores in Early Childhood and Future Cognitive Abilities in Young Nepalese Children*. SSRN.

Figure 1: Pattern of egg consumption in follow up months across research arms




Figure 2: Probability of falling below domain specific ASQ cut off by weekly egg consumption.




Table 1: Descriptive Statistics by intervention groups

	All (N=244)	Full (N=78)	Partial (N=83)	Control (N=83)
Baseline variables				
<i>Sex of the child, %(N)</i>				
Boy	52.05(127)	56.41(44)	44.58(37)	55.42(46)
Girl	47.95(117)	43.59(34)	55.42(46)	44.58(37)
Birth Order, Mean±SD (95% CI)	3.42±1.87 (3.2-3.7)	3.45±1.7 (3.1-3.8)	3.51±1.94 (3.1-3.9)	3.3±1.97 (2.9-3.7)
Weight (in kg), Mean±SD (95% CI)	7.9±1.09 (7.7-8)	7.92 ±1.34 (7.6-8.2)	7.7±0.98 (7.5-7.9)	8 ± 0.9 (7.8-8.2)
Height (in cm), Mean±SD (95% CI)	70.5±59.6 (69.9-70.9)	70.97±4.45 (69.9-71.9)	69.97 ±4.38 (69.01-70.9)	70.5±3.9 (69.6-71.3)
Mother's age at first				
birth, Mean±SD (95% CI)	18.2 ±2.1 (17.9-18.5)	18.2±2.2 (17.7-18.7)	18.5±2.1 (18.1-18.9)	17.9±1.8 (17.6-18.4)
<i>Education of mothers, %(N)</i>				
No formal education	79.8(194)	75.64(59)	76.83(63)	86.75(72)
Early primary education	5.35(13)	6.41(5)	6.1(5)	3.61(3)
Complimentary primary studies	4.94(12)	5.13(4)	6.1(5)	3.61(3)
Early secondary education	5.76(14)	6.41(5)	6.1(5)	4.82(4)
Koranic school	4.12(10)	6.41(5)	4.9(4)	1.2(1)
<i>HH size, %(N)</i>				
Less than 10	36.07(88)	26.92(21)	39.76(33)	40.96(34)
11-20 persons	31.56(77)	42.31(33)	28.92(24)	24.10(20)
21-30 persons	17.62(43)	16.67(13)	15.66(13)	20.48(17)
More than 30	14.75(36)	14.10(11)	15.66(13)	14.46(12)

Wealth Index^a, %(N)

Lowest	33.3(85)	38.46(30)	22.89(19)	38.55(32)
Middle	33.3(85)	32.05(25)	33.73(28)	34.94(29)
Highest	33.3(85)	29.49(23)	43.37(36)	26.51(22)

Egg consumption during the trial*Average Weekly Egg**Consumption,*

<i>Mean±SD</i>	<i>2.72±2.7</i>	<i>6.39±0.8</i>	<i>1.76±.09</i>	<i>0.22±0.48</i>
<i>(95% CI)</i>	<i>(2.4-3.1)</i>	<i>(6.2-6.6)</i>	<i>(1.5-1.9)</i>	<i>(0.12-2.33)</i>

Consistency in Egg Consumption, %(N)

Never	12.7(31)	0	0	37.35(31)
1-3 months	25.82(63)	0	19.28(16)	56.63(47)
4-6 months	26.64(65)	7.69(6)	66.27(55)	4.82(4)
in all months	34.84(85)	92.31(72)	14.46(12)	1.2(1)

^aCaptured only in month 7.

Table 2: Early Child Development (ECD) scores by intervention arm

	All (N=244)	Full (N=78)	Partial (N=83)	Control (N=83)	Mean difference (t-test) compared with control arm
Development					
score, Mean \pmSD (95% CI)					
Total ASQ					Full: 15.98**
Scores	228.4 \pm 48.92	239.36 \pm 42.6	223.13 \pm 52.3	223.37 \pm 49.4	Partial: 0.24
Range (0-300)	(222.5-234.3)	(230-248.7)	(212.3-233.9)	(213.1-233.7)	
Communication					Full: 2.48
Total	48.6 \pm 14.2	50.13 \pm 12.8	48.25 \pm 14.4	47.65 \pm 15.1	Partial: 0.60
Range (0-60)	(46.9-50.4)	(47.2-53.02)	(45.1-51.4)	(44.3-50.9)	
Gross Motor	49.9 \pm 9.79	51.73 \pm 8.9	49.27 \pm 9.0	48.67 \pm 11.1	Full: 3.06**
Range (0-60)	(48.6-51.1)	(49.7-53.7)	(47.3-51.2)	(46.3-51.1)	Partial: 0.60
Fine Motor	44.5 \pm 15.1	47.63 \pm 13.5	42.89 \pm 16.7	43.1 \pm 14.5	Full: 4.56**
Range (0-60)	(42.6-46.4)	(44.6-50.7)	(39.2-46.5)	(39.9-46.2)	Partial: 0.18
Problem Solving					Full: 6.36***
Solving	40.3 \pm 13.69	44.42 \pm 12.3	38.61 \pm 15.13	38.1 \pm 12.7	Partial: 0.54
Range (0-60)	(43.8-46.5)	(41.7-47.2)	(35.3-41.9)	(35.3-40.8)	
Person Social	45.1 \pm 10.9	45.5 \pm 10.8	44.09 \pm 10.9	45.9 \pm 10.7	Full: -0.45
Range (0-60)	(43.8-46.5)	(43-47.9)	(41.7-46.5)	(43.5-48.3)	Partial: -1.80

Table 3: Impact of consistency of egg consumption on ECD

		Logistic regression for falling below the cut-off values					Linear regression for log of total scores				
		Gross	Problem	Personal		Gross	Fine	Problem	Personal	Total	
		Motor	Fine Motor	Solving	Social	Communication	Motor	Motor	Solving	Social	Scores
		(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
		OR	OR	OR	OR	β	β	β	β	β	β
Adjusted											
Variables											
<i>Consistency of Egg Consumption</i>											
1-3 months	0.586 (0.485)	0.335 (0.334)	0.927 (0.633)	0.525 (0.361)	0.857 (0.679)	-0.064 (0.121)	0.015 (0.061)	-0.186 (0.114)	-0.087 (0.103)	-0.000 (0.090)	-0.068 (0.070)
4-6 months	0.584 (0.422)	0.327 (0.268)	0.864 (0.657)	0.743 (0.343)	0.609 (0.382)	0.020 (0.114)	-0.029 (0.061)	-0.207 (0.114)	-0.071 (0.081)	-0.030 (0.082)	-0.066 (0.060)
In all months	0.079* (0.106)	0.125** (0.112)	1.176 (0.795)	0.608 (0.334)	0.343* (0.190)	0.050 (0.087)	0.058 (0.052)	-0.044 (0.088)	0.041 (0.069)	0.004 (0.078)	0.007 (0.040)
Baseline variables											
Birth order	0.766 (0.138)	0.762** *	0.882 (0.111)	0.974 (0.100)	0.987 (0.158)	0.014 (0.010)	0.018 (0.011)	0.020 (0.019)	-0.009 (0.012)	-0.007 (0.012)	0.008 (0.009)
Baseline egg consumption (=1 if yes)	3.134 (3.794)	1.823 (1.949)	1.608 (1.609)	0.838 (0.698)	0.522 (0.543)	-0.011 (0.069)	-0.004 (0.052)	-0.098 (0.110)	-0.031 (0.080)	0.052 (0.055)	-0.013 (0.045)

Weight (kg)	0.371*** (0.138)	0.777 (0.365)	1.539 (0.428)	1.688* (0.473)	0.860 (0.310)	-0.014 (0.039)	0.013 (0.026)	-0.035 (0.037)	-0.06** (0.025)	0.012 (0.026)	-0.017 (0.017)
Height (cm)	0.849 (0.117)	0.924 (0.077)	0.694*** (0.047)	0.816** (0.073)	0.821* (0.085)	0.039*** (0.011)	0.01** (0.004)	0.010 (0.010)	0.020* (0.010)	0.015* (0.007)	0.02*** (0.005)
Girl	0.775 (0.443)	1.112 (0.546)	1.017 (0.482)	1.063 (0.412)	0.577 (0.268)	0.020 (0.050)	0.030 (0.047)	-0.096 (0.068)	-0.038 (0.046)	0.132*** (0.034)	-0.003 (0.030)
Mother's age at first birth	0.802* (0.093)	1.094 (0.132)	1.115 (0.123)	1.034 (0.091)	0.864 (0.138)	-0.004 (0.014)	-0.009 (0.009)	-0.003 (0.012)	0.001 (0.013)	-0.011 (0.009)	-0.004 (0.007)
<i>Education of mothers</i>											
Early primary education		2.329 (3.510)	1.062 (0.893)	1.291 (0.835)	1.582 (2.077)	-0.087 (0.093)	0.070 (0.053)	0.151 (0.180)	0.068 (0.074)	-0.059 (0.092)	0.014 (0.074)
Complimentary primary studies	1.718 (2.174)	1.174 (1.111)			2.608 (2.048)	0.038 (0.156)	-0.162 (0.197)	0.206* *	0.104* (0.055)	-0.169 (0.165)	0.030 (0.078)
Early secondary education			0.760 (0.557)			0.249*** (0.072)	0.055 (0.054)	0.313* *	0.038 (0.095)	0.070 (0.075)	0.143** (0.059)
<i>Kornaic school</i>					8.724* (10.13)	-0.121 (0.128)	0.011 (0.059)	0.106 (0.100)	0.079 (0.068)	-0.078 (0.082)	0.004 (0.025)
<i>HH size</i>											
11-20 persons	0.401 (0.317)	0.555 (0.518)	0.584 (0.354)	1.383 (0.685)	0.944 (0.506)	-0.056 (0.039)	0.019 (0.033)	-0.004 (0.094)	-0.025 (0.063)	0.001 (0.027)	-0.000 (0.033)

21-30 persons	1.158 (0.566)	2.370 (1.352)	0.941 (0.576)	0.856 (0.539)	1.314 (0.842)	-0.151** (0.056)	-0.063 (0.039)	-0.038 (0.082)	-0.119 (0.077)	-0.088 (0.052)	-0.067* (0.035)
More than 30	0.326 (0.360)	0.461 (0.451)	1.337 (0.885)	0.827 (0.481)		0.105 (0.081)	0.016 (0.034)	0.018 (0.111)	-0.041 (0.077)	0.037 (0.037)	0.033 (0.049)
<i>Wealth index</i> ^a											-0.000
Middle	0.981 (0.502)	0.949 (0.617)	1.322 (0.579)	1.136 (0.571)	1.466 (0.908)	0.109** (0.044)	0.021 (0.032)	0.059 (0.068)	-0.051 (0.072)	0.030 (0.054)	0.035 (0.028)
Highest	0.427 (0.441)	0.193* (0.188)	0.827 (0.534)	0.973 (0.490)	1.007 (0.946)	0.073 (0.063)	0.046 (0.040)	0.030 (0.070)	-0.023 (0.055)	0.046 (0.069)	0.030 (0.041)
Constant	3.033e+09** (2.612e+10)	229.76 (1,578.0 18)	1.126e+08 *** (4.573e+08)	4,144.5* (19,736.2 21)	6978048. 2** (4766598 1.083)	1.142 (0.759)	3.128* ** (0.229)	3.370* ** (0.653)	2.840*** (0.566)	2.763*** (0.465)	4.302** * (0.379)
Observations	199	212	214	201	189	234	234	234	230	234	234

^aCaptured only in month 7. *** p<0.01, ** p<0.05, * p<0.1. Clustered standard errors in parentheses.

Table 4: Impact of quantity of egg consumption on ECD

	Logistic regression for falling below the cut-off values					Linear regression for log of scores					
	Communication	Gross Motor	Fine Motor	Problem Solving	Personal Social	Communication	Gross Motor	Fine Motor	Problem Solving	Personal Social	Total Scores
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
	OR	OR	OR	OR	OR	β	β	β	β	β	β
Adjusted											
Variables											
<i>Average weekly</i>											
<i>egg consumption</i>	0.798 (0.121)	0.812 (0.107)	1.036 (0.08)	0.928 (0.074)	0.831 (0.094)	0.011 (0.008)	0.005 (0.006)	0.012 (0.01)	0.019** (0.008)	-0.001 (0.01)	0.008 (0.005)
Baseline variables											
Birth order	0.780 (0.128)	0.75*** (0.071)	0.885 (0.11)	0.960 (0.095)	0.982 (0.148)	0.014 (0.010)	0.018 (0.011)	0.017 (0.02)	-0.011 (0.011)	-0.007 (0.01)	0.007 (0.008)
Baseline egg consumption (=1 if yes)	2.358 (2.717)	1.624 (1.519)	1.602 (1.57)	0.764 (0.616)	0.537 (0.566)	-0.022 (0.067)	0.001 (0.053)	-0.104 (0.11)	-0.033 (0.08)	0.054 (0.05)	-0.015 (0.046)
Weight	0.383**	0.817	1.550	1.710*	0.817	-0.015	0.017	-0.022	-	0.014	-0.013

	(0.156)	(0.416)	(0.45)	(0.513)	(0.272)	(0.040)	(0.029)	(0.04)	0.056**	(0.021)	(0.03)	(0.017)
			0.69**									
Height	0.853	0.923	*	0.818**	0.826**	0.040***	0.009**	0.008	0.019*	0.015*	0.018***	
	(0.109)	(0.082)	(0.05)	(0.080)	(0.077)	(0.010)	(0.004)	(0.01)	(0.010)	(0.01)	(0.005)	
Girl	0.651	1.057	1.035	1.125	0.535	0.029	0.030	-0.083	-0.030	0.13***	0.003	
	(0.462)	(0.524)	(0.49)	(0.436)	(0.255)	(0.049)	(0.045)	(0.07)	(0.042)	(0.03)	(0.028)	
Mother's age at first birth	0.830	1.064	1.105	1.027	0.864	-0.005	-0.010	-0.008	-0.002	-0.012	-0.006	
	(0.126)	(0.137)	(0.11)	(0.084)	(0.142)	(0.014)	(0.009)	(0.01)	(0.013)	(0.01)	(0.007)	
<i>Education of mothers</i>												
Early primary education		2.416	1.118	1.376	1.528	-0.077	0.070	0.158	0.064	-0.059	0.017	
		(2.974)	(0.92)	(0.809)	(2.028)	(0.090)	(0.051)	(0.18)	(0.074)	(0.09)	(0.073)	
Complimentary primary studies	1.415	1.391			2.697	0.054	-0.156	0.23**	0.112*	-0.166	0.039	
	(1.787)	(1.375)			(2.233)	(0.153)	(0.199)	(0.09)	(0.061)	(0.17)	(0.084)	
Early secondary education			0.760			0.257***	0.056	0.31**	0.035	0.070	0.143**	
			(0.55)			(0.069)	(0.049)	(0.11)	(0.089)	(0.07)	(0.055)	
<i>Kornaic school</i>					8.387**	-0.109	0.022	0.135	0.095	-0.073	0.016	
					(8.375)	(0.128)	(0.051)	(0.12)	(0.064)	(0.08)	(0.032)	

<i>HH size</i>											
11-20 persons	0.399 (0.271)	0.568 (0.532)	0.584 (0.35)	1.499 (0.69)	0.955 (0.465)	-0.050 (0.040)	0.021 (0.034)	-0.002 (0.09)	-0.026 (0.061)	0.002 (0.02)	0.001 (0.034)
21-30 persons	1.170 (0.527)	2.720* (1.550)	0.963 (0.53)	0.939 (0.540)	1.392 (0.854)	-0.148** (0.057)	-0.062 (0.039)	-0.025 (0.08)	-0.115 (0.073)	-0.087 (0.05)	-0.063* (0.033)
More than 30	0.385 (0.471)	0.557 (0.626)	1.300 (0.87)	0.896 (0.542)		0.106 (0.074)	0.011 (0.035)	0.009 (0.11)	-0.049 (0.076)	0.036 (0.04)	0.029 (0.048)
<i>Wealth index^a</i>											
Middle	1.005 (0.569)	0.836 (0.508)	1.330 (0.57)	1.123 (0.55)	1.345 (0.848)	0.112** (0.047)	0.021 (0.031)	0.050 (0.07)	-0.051 (0.074)	0.029 (0.05)	0.034 (0.028)
Highest	0.432 (0.469)	0.170* (0.163)	0.824 (0.54)	0.979 (0.505)	0.956 (0.890)	0.077 (0.061)	0.046 (0.036)	0.019 (0.07)	-0.026 (0.054)	0.045 (0.07)	0.028 (0.039)
Constant	7.335e+08** (6.182e+09)	172.273 (1,068.426)	1.120e+08** (4.571 e+08)	2,694.569 (13,140.27)	560** (4477937.736)	1.119 (0.752)	3.17*** (0.21)	3.38** (0.72)	2.87*** (0.569)	2.8*** (0.46)	4.312*** (0.394)
Observations	199	212	214	201	189	234	234	234	230	234	234

^aCaptured only in month 7. *** p<0.01, ** p<0.05, * p<0.1. Clustered standard errors in parentheses.

Table 5: Effect of Egg consumption on child development outcomes - Average Treatment effect on the treated (ATET)

	Coefficient	Clustered Standard errors	95% confidence interval	
Total Scores	0.093**	0.037	0.0198	0.1670
Communication	0.094	0.069	-0.0412	0.2300
Gross Motor	0.106*	0.056	-0.0031	0.214
Fine Motor	0.134**	0.065	0.006	0.262
Problem Solving	0.059	0.057	-0.054	0.171
Personal Social	0.057	0.057	-0.056	0.169

*** p<0.01, ** p<0.05, * p<0.1.