

Original Research

Cite this article: Yoshida M, Miyatake H, Masunaga H, *et al.* Cross-sectional study on risk of flooding and landslides and their associated perceptions among home health-care patients living in Fukui prefecture, Japan. *Disaster Med Public Health Prep.* 17(e264), 1–8. doi: <https://doi.org/10.1017/dmp.2022.210>.






Keywords:

primary prevention; disaster medicine; disasters; floods; patient transfer

Corresponding author:

Hiroto Miyatake,
Email: hmiyatake@orangeclinic.jp.

Cross-Sectional Study on Risk of Flooding and Landslides and Their Associated Perceptions Among Home Health-Care Patients Living in Fukui Prefecture, Japan

Makoto Yoshida¹ , Hiroto Miyatake MD² , Hidehisa Masunaga MSc², Makoto Kosaka^{2,3}, Yudai Kaneda⁴ , Yasuhiro Kotera PhD⁵ , Yoshitaka Nishikawa MD, PhD^{2,6}, Chie Tsunetoshi², Izumi Yoshida MD⁷, Fubuki Ogawa², Akihiko Ozaki MD, PhD^{2,8}  and Hiroyuki Beniya MD²

¹Teikyo University School of Medicine, Tokyo, Japan; ²Orange Home-Care Clinic, Fukui, Japan; ³Faculty of Medicine, The University of Tokyo, Tokyo, Japan; ⁴Hokkaido University School of Medicine, Hokkaido, Japan; ⁵Faculty of Medicine and Health Sciences, University of Nottingham, Nottingham, United Kingdom; ⁶Department of Internal Medicine, Soma Central Hospital, Fukushima, Japan; ⁷Department of Dermatology and Allergology, University Hospital Augsburg, Augsburg, Germany and ⁸Department of Breast and Throid Surgery, Jyoban Hospital of Tokiwa Foundation, Fukushima, Japan

Abstract

Objective: This study aimed to evaluate a risk of flooding and landslides among home-care patients, to reveal an extent to which patients require support for evacuation, and to determine whether risk was accurately perceived among the patients.

Methods: This is a cross-sectional study targeting the patients who were actively treated at the home-care clinic in Fukui Prefecture, Japan. We collected data on the patients' socio-demographic and clinical characteristics. Additionally, we collected data on their risk of flooding and landslides through hazard maps and distributed a questionnaire to these patients regarding their risk awareness of flooding and landslides.

Results: Of the 199 eligible home-care patients, 84.9% (169 of 199) were at risk of flooding and/or landslides, and 58.6% (99 of 169) of them needs support during evacuation. Furthermore, of those who were at risk of flooding and/or landslides, 46.0% (45 of 99) had accurate risk assessments. Factors that resulted in inadequate risk awareness of flooding and landslides included: not placing importance on evacuation, not using medical equipment, and living on the first floor.

Conclusions: There was limited risk awareness of flooding and/or landslides among the home-care patients. The information of the risk factors regarding inadequate risk awareness of flooding and landslides should be used to sophisticate flooding and landslides evacuation strategy.

Home care is a service in which doctors and other health-care providers visit patients' houses, elderly housing, and nursing facilities to support their discharge from the hospital, support their daily medical care, deal with unexpected medical changes, and provide terminal care. The number of people choosing home care is increasing. Moreover, the home-care sector is expected to grow significantly with the aging of the global population and the increasing preference of patients for high quality care and is estimated to reach a market size of \$49.7 billion by the end of 2030, up from \$27.9 billion in 2022.¹

Persons with home-care services are considered a vulnerable population during flooding and landslides due to the need for continuous access to health care and declines in physical functions caused by aging and chronic diseases. Natural disasters are increasing all over the world² and have become a major health and socioeconomic problem worldwide, but home-care patients may not be fully prepared for disasters.^{3–6} In particular, flooding damage was concentrated among those with low activity of daily living.^{5,6}

In Japan, the number of home-care patients is increasing,⁷ and these patients face various difficulties in evacuation during flooding and landslides. In the case of Typhoon No. 21 on September 1, 2019, power outages caused several problems, including safety concerns for home-care patients using medical equipment such as ventilators, oxygen concentrators, and infusion pumps, as well as for older people who had difficulty in moving were exposed to the risk of heat stroke as a result of the lack of air conditioning.⁸ A wide range of flooding and landslides preparedness is necessary when considering flooding and landslides countermeasures for home-care patients.⁹

Our study purposes were as follows: (i) to determine the prevalence of home-care patients at risk of flooding and/or landslides, (ii) to reveal the extent to which patients require support for evacuation, and (iii) to clarify whether risk of flooding and landslides was accurately perceived among the participants.

Methods

Study Site and Participants

This study was conducted in Fukui Prefecture, located in the north-central part of Japan. The prefecture has a population of 766,863 and its population density is 183.0 people/km². The elderly population over 65 y old accounts for 30.6%, and approximately 17.8% of this elderly population is in need of nursing care. Fukui Prefecture experienced a flooding caused by the torrential rains in July 2004, which resulted in 4 deaths, 1 missing person, and 14,157 destroyed houses.¹⁰

The study site was Orange Home-Care Clinic in Fukui City, Fukui Prefecture, a detail of which was provided elsewhere.^{11–13} In short, it provides home-care services to patients living at home or in elderly facilities such as group homes. There are approximately 20 people running the clinic, including several doctors, nurses, and office staff. In this clinic, 1 doctor and 1 staff member (nurse or office staff) visit the patients as a team. Each team visits 8–10 patients/d on weekdays, and the visiting area covers a radius of 16 km from the clinic. Some of the patients are living in nearby cities that are more than 16 km away and are treated in cooperation with visiting nurses. Approximately 350 patients, from children to the elderly, are treated at this clinic every year. The clinic also provides medical care for patients in the terminal stages of cancer, chronic diseases such as neural intractable diseases and dementia, and severely handicapped children, including not only terminal care but also life support.

This is a cross-sectional study targeting the home-care patients using the services of the Orange Home-Care Clinic on December 1, 2020. We excluded facility users and hospitalized patients because we aimed to evaluate those who received home care.

Survey Sheet

The questionnaires were developed based on the experience of Typhoon Hagibis and other previous evidence as well as the local context of Fukui Prefecture. For example, in the course of treating patients affected by Typhoon Hagibis and in compiling data, it became clear that evacuation was difficult for those with low activity of daily living,^{5,6} and we assumed that this was also the case for home-care patients. There have been studies on specific needs among those with special needs care who stay at disaster reception centers,¹⁴ but to the best of our knowledge, there are no studies in Japan on risk of flooding and landslides of home-care patients. Therefore, we created a questionnaire on the topics we considered important.

The self-administered questionnaire consists of 4 sections: Section 1 asks about perceptions of evacuation in the event of flooding. It asked about the importance of evacuation, perception of disaster risk of flooding and landslides in their houses, attitudes toward evacuation, the range of needs for care and support, and the use of medical devices.

Section 2 focuses on the patient's home situation, such as the structure of the house and the bedroom floor.

Section 3 asks patients and their families about their actual feelings about evacuation, and their concerns and inadequacies in flooding measures through open-ended questions.

Section 4 asks the date the questionnaire was completed and who completed it.

The survey questions are described in greater detail in the Supplementary Materials Methods.

Data Collection

We collected information on sex, age, the range of care required, name of the primary disease, and address of the study subjects from the electronic medical records belonging to the Orange Home-Care Clinic. Age was calculated as of the time the questionnaire was distributed (December 1, 2020). The range of needs for care or support was classified as Not certified, Support level 1, Support level 2, Care level 1, Care level 2, Care level 3, Care level 4, and Care level 5 as described in the electronic medical record. As for the primary disease names, the diseases were classified based on the International Classification of Diseases, 11th Edition (ICD-11). For those with more than 1 disease name listed, the first disease name was given priority.

For the risk of flooding and landslides, we checked whether the home addresses of patients collected from their medical records were included in hazard areas on hazard maps issued by the Ministry of Land, Infrastructure, Transport and Tourism of Japan. We investigated the risk information of flooding and landslides of patients' houses based on the addresses listed on the electronic medical records. Risk Information of landslides was collected considering that landslides can be caused by flooding as well as earthquakes. For this survey, we used "Overlapping Hazard Map," which is a free website provided by the Ministry of Land, Infrastructure, Transport and Tourism of Japan that enables disaster risk information to be displayed on a map.¹⁵ The risk of flooding was categorized into 9 types based on the depth of flooding, and the risk of landslides was categorized into 7 types based on the area. We checked whether the patients' houses were in the warning areas or special warning areas. When we could not identify an address with "Overlapping Hazard Map", we searched for it on "Yahoo! Maps" and made the assessment as above.¹⁶ All checks were completed by at least 2 of us, and if there were differences in the classification of risk of flooding and landslides, we asked another person to check and make a decision.

Then, we conducted a questionnaire survey from December 1, 2020 to April 30, 2021 to determine whether their risk of flooding and landslides was accurately assessed. Self-administered questionnaires were sent to the participants from December 1, 2020, and the staff of the Orange Home-Care Clinic asked the patients and their families to complete the questionnaire. Its details will be described in the following section. To reduce response bias, participants were informed their participation was voluntary and their participation and response results would not affect their future services. The deadline for responses was set for April 30, 2021.

The completed self-administered questionnaires were collected by the Orange Home-Care Clinic, and the written contents of the questionnaires were tabulated and integrated with information on the presence or absence of responses, basic patient information obtained from electronic medical records, and information about risk of flooding and landslides.

Data Analysis

We calculated the absolute number of home-care patients whose houses were at risk of flooding and/or landslides and the percentage of these patients in the total number of home-care patients using the Orange Home-Care Clinic as of December 1, 2020.

Basic information on home-care patients such as sex, age, range of care needs, name of main disease, address (by city), depth of flooding, landslides information, and special warning areas were tabulated separately for participants and non-participants in the survey. Data from non-respondents are also included to provide a denominator for risk assessment and to calculate the number of persons requiring assistance who may need help during evacuation. In the respondent's group, information about the structure of their houses and the use of medical equipment were additionally tabulated.

Respondents were divided into 2 groups: those at risk of flooding and/or landslides and those not at risk. We tabulated their responses to questions 2 and 3 in section 1 of the questionnaire. Then, we investigated how many people are aware of risk of flooding and landslides of their houses accurately by comparing their risk assessment of flooding and landslides with that based on hazard maps. First, for those who have both flooding and landslide risks, only those who answered "located" for both in questions 2 and 3 in section 1 were considered to have the accurate perception. Second, for those with only flooding risk, only those who answered "located" in question 2 in section 1 were considered to have the accurate perception. Finally, for those with only landslide risk, only those who answered "located" in question 3 in section 1 were considered to have the accurate perception.

Among the questionnaire respondents who were at risk of flooding and/or landslides, we did univariable and multivariable logistic regression analysis to investigate the factors associated with those who are not fully aware of risk of flooding and landslides. In the univariable logistic regression, sex, age, primary disease, presence of flood risk, flooding depth, presence of landslide risk, presence of special alert, building count, home structure, presence of medical device use, importance of evacuation, presence of difficulties in evacuating alone, bedroom floor, presence of evacuation concern, respondents and number of disaster risks were used as the explanatory variable and inadequate comprehension of disaster risks as the independent variable. In the multivariable logistic regression analysis, we used explanatory variables that were significantly associated with the outcome in the univariable analysis. We considered *P* values of 0.05 or less to be significant.

For all analyses, the results of the answered questionnaires were recorded in Microsoft excel for Mac (ver. 16.53) for data extraction and tabulation. For logistic regression analysis, we used Stata/IC 15.

Results

Information for All Patients Receiving Home Care

There were 390 patients listed in the Orange Home-Care Clinic's electronic medical records as of December 1, 2020, including 7 (1.8%) patients who were hospitalized, 36 (9.2%) patients who had ended their services, 44 (11.3%) patients who had died, and 104 (26.7%) patients who had been institutionalized. These patients were excluded from the study because of the difficulty in having those patients respond. There were 199 home-care patients using the services of the Orange Home-Care Clinic, of which 86 (43.2%) were male, the median age was 78 (range,

2-105), and 112 (56.3%) were certified as requiring nursing care. 165 (82.9%) of the patients lived in Fukui City. The prevalent conditions were "diseases were of the nervous system" in 60 patients (30.2%) and "mental, behavioral, or neurodevelopmental disorders" in 57 patients (28.6%). Of the 199 home-care patients, 169 (84.9%) were at risk of flooding and/or landslides, and 99 (58.6%) of them were certified as requiring nursing care (Table 1).

Information of Questionnaire Respondents and Nonrespondents

Questionnaires were sent to 199 participants and 98 people filled out the questionnaire (response rate: 49.3%). Table 1 presents the sociodemographic and clinical characteristics of respondents and non-respondents. Of the 98 respondents, 25 (25.5%) responded by themselves, 58 (59.2%) by family members living with them, 14 (14.3%) by family members living separately, and 4 (4.1%) by others. Of the respondents, 86 (87.8%) lived in Fukui City, and 35 (35.7%) used medical equipment (Table 1).

Comparing the characteristics of Respondents and Nonrespondents, 38 (38.8%) and 48 (47.5%) were male, and the median age was 79.5 y (range, 2-105 y) and 78 y (range, 1-102 y), and 87 (88.8%) and 82 (81.2%) were at risk of flooding and/or landslides.

Perception of Flooding and Landslides Risk

Among the respondents, there were 87 (88.8%) respondents whose houses were at risk of flooding and/or landslides, and of those respondents, 47 (54.0%) were not fully aware of the flooding and landslides risks based on hazard maps (Table 2). The percentage of those who thought evacuation was important was 86.2% (75 respondents) in the group with risk of flooding and/or landslides and 63.6% (7 respondents) in the group without flooding and landslides risks (Figure 1). Of the 87 respondents at risk of flooding and/or landslides, 32.2% (28 respondents) were unaware of the risk yet considered evacuation necessary.

Factors Associated With Lack of Flooding and Landslides Risk Perception

Those who were at risk of flooding and/or landslides but failed to accurately assess their risk had the following in common: not placing importance on evacuation (odds ratio [OR]: 11.9; 95% confidence interval [CI]: 1.4-9.6), not using medical equipment (OR: 3.1; 95% CI: 1.1-9.0), and living on the first floor (OR: 4.71; 95% CI: 1.2-18.0) (Table 3). For not using medical equipment, 11 did not answer, and their data were excluded from the analysis.

Discussion

This survey revealed that 169 (84.9%) home-care patients were at risk of flooding and/or landslides. Of the respondents who lived there, 47 (54.0%) did not have an accurate risk awareness of flooding and landslides. Factors for this included not placing importance on evacuation, not using medical equipment, and living on the first floor.

Of 199 home-care patients, 169 (84.9%) were at risk of flooding and/or landslides, and among them, 141 (70.9%) were at risk of flooding. The Japanese Ministry of Land, Infrastructure, Transport and Tourism of Japan reports that the number of people at risk of flooding as of 2015 was 37.03 million across Japan, which was 29.1% of the total population.¹⁷

Table 1. Characteristics of home health-care patients in Fukui Prefecture, Japan, who responded to the survey and those who did not (2020–2021)

	Respondents (%) (N=98)	Non-respondents (%) (N=101)	Total (%) (N=199)
Sex			
Male	38 (38.8)	48 (47.5)	86 (43.2)
Female	60 (61.2)	53 (52.5)	113 (56.8)
Age			
100-	4 (4.1)	1 (1.0)	5 (2.5)
90-99	16 (16.3)	15 (14.9)	31 (15.6)
80-89	29 (29.6)	31 (30.7)	60 (30.2)
70-79	19 (19.4)	11 (10.9)	30 (15.1)
60-69	11 (11.2)	6 (5.9)	17 (8.5)
50-59	2 (2.0)	3 (3.0)	5 (2.5)
40-49	0 (0.0)	3 (3.0)	3 (1.5)
30-39	7 (7.1)	5 (5.0)	12 (6.0)
20-29	0 (0.0)	3 (3.0)	3 (1.5)
10-19	3 (3.1)	3 (3.0)	6 (3.0)
0-9	7 (7.1)	20 (19.8)	27 (13.6)
Level of the need for care or support			
Not certified	35 (35.7)	52 (51.5)	87 (43.7)
Support level 1	1 (1.0)	1 (1.0)	2 (1.0)
Support level 2	2 (2.0)	3 (3.0)	5 (2.5)
Care level 1	8 (8.2)	12 (11.9)	20 (10.1)
Care level 2	14 (14.3)	6 (5.9)	20 (10.1)
Care level 3	10 (10.2)	8 (7.9)	18 (9.0)
Care level 4	18 (18.4)	11 (10.9)	29 (14.6)
Care level 5	10 (10.2)	8 (7.9)	18 (9.0)
Primary disease			
Diseases of the nervous system	26 (26.5)	34 (33.7)	60 (30.2)
Mental, behavioral or neurodevelopmental disorders	29 (29.6)	28 (27.7)	57 (28.6)
Developmental anomalies	3 (3.1)	11 (10.9)	14 (7.0)
Injury, poisoning, or certain other consequences of external causes	3 (3.1)	10 (9.9)	13 (6.5)
Neoplasms	10 (10.2)	1 (1.0)	11 (5.5)
Diseases of the musculoskeletal system or connective tissue	7 (7.1)	4 (4.0)	11 (5.5)
Diseases of the circulatory system	6 (6.1)	4 (4.0)	10 (5.0)
Endocrine, nutritional or metabolic diseases	6 (6.1)	2 (2.0)	8 (4.0)
Others	8 (8.2)	7 (6.9)	15 (7.5)
Address			
Fukui City	86 (87.8)	79 (78.2)	165 (82.9)
Sakai City	4 (4.1)	7 (6.9)	11 (5.5)
Yoshida Gun	4 (4.1)	6 (5.9)	10 (5.0)
Sabae City	3 (3.1)	3 (3.0)	6 (3.0)
Ohno City	0 (0.0)	3 (3.0)	3 (1.5)
Katsuyama City	0 (0.0)	2 (2.0)	2 (1.0)
Echizen City	1 (1.0)	0 (0.0)	1 (0.5)
Awara City	0 (0.0)	1 (1.0)	1 (0.5)

(Continued)

Table 1. (Continued)

	Respondents (%) (N=98)	Non-respondents (%) (N=101)	Total (%) (N=199)
Disaster risk (flooding and/or landslides)			
Exists	87 (88.8)	82 (81.2)	169 (84.9)
Not exists	11 (11.2)	19 (18.8)	30 (15.1)
Depth of flooding (m)			
0	23 (23.5)	35 (34.7)	58 (29.1)
0~0.3	0 (0.0)	0 (0.0)	0 (0.0)
0.3~0.5	0 (0.0)	3 (3.0)	3 (1.5)
0.5~1.0	0 (0.0)	1 (1.0)	1 (0.5)
1.0~3.0	31 (31.6)	30 (29.7)	61 (30.7)
3.0~5.0	42 (42.9)	29 (28.7)	71 (35.7)
5.0~10.0	2 (2.0)	3 (3.0)	5 (2.5)
10.0~20.0	0 (0.0)	0 (0.0)	0 (0.0)
20.0~	0 (0.0)	0 (0.0)	0 (0.0)
Landslide disaster information			
None	84 (85.7)	83 (82.2)	167 (83.9)
Steep slope failure	0 (0.0)	1 (1.0)	1 (0.5)
Mudslide	4 (4.1)	4 (4.0)	8 (4.0)
Landslide	0 (0.0)	0 (0.0)	0 (0.0)
Streams in danger of mudslides	3 (3.1)	2 (2.0)	5 (2.5)
Steep slope failure hazard area	5 (5.1)	11 (10.9)	16 (8.0)
Landslide danger zone	2 (2.0)	2 (2.0)	4 (2.0)
Avalanche	12 (12.2)	12 (11.9)	24 (12.1)
Special alert area			
None	90 (91.8)	89 (88.1)	179 (90.4)
Warning area	6 (6.1)	9 (8.9)	15 (7.6)
Special warning area	0 (0.0)	2 (2.0)	2 (1.0)
Warning area (steep slope failure)	1 (1.0)	1 (1.0)	2 (1.0)
Warning area (mudslide)	2 (2.0)	1 (1.0)	3 (1.5)
Warning area (landslide)	0 (0.0)	1 (1.0)	1 (0.5)
Structure of the house			
Single house (1 story)	12 (12.2)		
Single house (2 stories)	62 (63.3)		
Single house (3 or more stories)	7 (7.1)		
Apartment (2 or 3 stories and wooden or lightweight steel frame)	5 (5.1)		
Apartment (3 or more stories and steel or reinforced concrete frame)	8 (8.2)		
Others	4 (4.1)		
Frame of the house			
Wooden frame □	71 (72.4)		
Heavy or lightweight steel frame	12 (12.2)		
Reinforced concrete frame	9 (9.2)		
Steel reinforced concrete frame	2 (2.0)		
Others	1 (1.0)		

(Continued)

Table 1. (Continued)

	Respondents (%) (N=98)	Non-respondents (%) (N=101)	Total (%) (N=199)
Unknown	4 (4.1)		
Not available	12 (12.2)		
Use of medical devices			
Ventilator	11 (11.2)		
Dialysis	0 (0.0)		
Intravenous drip	6 (6.1)		
Nutritional support thorough gastric or intestinal bypass	18 (18.4)		
Not used.	57 (58.2)		
Not available	20 (20.4)		

However, this study revealed that the percentage of home-care patients in Fukui under risk of flooding and landslides was higher than the Japanese average. Therefore, this suggests that the home-care patients included in this study may be living in geographically disadvantaged areas in terms of flooding and landslides risk. Fukui Prefecture has a population at risk of flooding reaching 410,000 (52.7%) and a risk of flooding area of 490,000 (62.5%),¹⁷ which is higher than the Japanese average, because of the ground being lower than the major rivers and sloping gradually.¹⁶ Previous studies reported geographical vulnerability among home-care patients both in economically developing countries and developed countries.^{18–21} The average income of Fukui residents was 4,450,000 yen, which puts them in the top 17% of the world's population,^{22,23} indicating that even though they were economically well off, they lived in an area vulnerable to flooding and landslides. Vulnerability to flooding and landslides of home-care patients may also need to be considered, in addition to the vulnerability to access to health care.

Of the 169 patients (84.9% of the total) who were at risk of flooding and/or landslides, 99 (58.6%) had been certified as needing nursing care. A previous study on Typhoon Hagibis indicates the vulnerability of those in need of supports for flooding, as 33.3% of all deaths were those who could have evacuated but failed to do so because of their own health problems.⁵ Furthermore, the mean age of those at risk of flooding and landslides were 62.8 and 81.9 y old, respectively. A previous study has pointed out vulnerabilities such as the fact that flooding victims are often concentrated in older people, and that even when they survive, older people are more likely to develop PTSD and general psychiatric symptoms than younger people.²⁴ In addition, patients with suppressed immune systems such as cancer may be at increased risk of infections caused by flooding.⁶ Considering the fact that the

frequency of flooding is expected to increase in the future, it is desirable to provide more support to home-care patients in times of flooding and landslides so that they can minimize the damage.²⁵

There were not many people who had an accurate assessment of the risks of flooding and landslides at their houses. Among 98 respondents, there were 87 (88.8%) whose houses were at risk of flooding and/or landslides, and among whom 47 (54.0%) had inadequate risk awareness of flooding and landslides. A previous study has highlighted the importance of being aware of the risks of disaster, such as the fact that simple preparations by family members and community members can increase the probability of surviving the first 72 h of a disaster before the arrival of government supports,²⁶ and that residents who were aware of hazard maps were able to start evacuating 1 h earlier than those who were not and had a 10% higher evacuation rate.^{27,28} It is true that the risk awareness of flooding and landslides among home-care patients in Fukui is higher than the Japanese national average of 38.6% for awareness of hazard maps,²⁹ probably because flooding and landslides occur more frequently in Fukui than in the rest of Japan, and most people are at risk of flooding.¹⁷ However, considering that their survival rates may increase with a little more flood preparedness (eg, 72 h of durable supplies such as an attic axe, freshwater, to allow survival until rescue arrives; improvement of the first floor structure to resist water damage such as converting the first floor from wood to cement or wooden floors to cement or tile; ability to run equipment with a generator with safety devices to prevent carbon monoxide poisoning), the awareness rate of risk of flooding and landslides may not be sufficient to minimize the damage to home-care patients during flooding and landslides.

Of the 87 respondents at risk of flooding and/or landslides, 32.2% (28 respondents) were unaware of the risk yet considered evacuation necessary, suggesting a lack of awareness regarding a hazard map. These findings warrants caution, given that an evacuation in flooding could be dangerous. Indeed, from the perspective of our previous study, in which we evaluated the fatalities associated with Typhoon Hagibis in 2019, some of the residents involved might have died outdoors in the process of evacuating from their homes.⁵ Of note, evacuation risk is higher among residents with impaired daily living activities, such as home-care patients. Therefore, home-care patients and their families need to be aware of the risks of floods and landslides in advance, and they should determine whether to evacuate while weighing the risk of staying at home.

We found that the factors resulting in lack of risk awareness of flooding and landslides include not placing importance on evacuation, not using medical equipment, and living on the first floor. In a previous study, it was mentioned that men over 55 y old (33%) and married people with children (27%) were the most well prepared for evacuation, and that the awareness rate of hazard maps improved as their age increased.³⁰ Furthermore, there was a

Table 2. Results regarding the accuracy of survey respondents' risk assessment of flooding and landslides, divided by the presence or absence of flooding and landslides risk

	Group at risk of flooding and/or landslides (%) (N=87)	Group at risk of flooding (%) (N=75)	Group at risk of landslide (%) (N=14)	Group at no risk of disaster (%) (N=11)	Total (%) (N=98)
Risk assessment of flooding and/or landslides					
Correct	40 (46.0)	34 (45.3)	6 (42.9)	6 (54.5)	46 (46.9)
Not correct	47 (54.0)	41 (54.7)	8 (57.1)	5 (45.5)	52 (53.1)

Table 3. Results of logistic analysis of factors that contributed to inadequate comprehension of risk of flooding and landslides

	OR univariate (95% confidence interval)	P-Value	OR multivariate (95% confidence interval)	P-Value
Sex				
Male	1			
Female	1.58 (0.66~3.79)	0.309		
Age				
0~75	1			
75~	1.59 (0.67~3.77)	0.298		
Primary disease				
Nervous	1			
Mental	2.93 (0.91~9.44)	0.073		
Neoplasm	4.55 (0.77~26.84)	0.094		
Others	0.92 (0.30~2.78)	0.879		
Presence of flooding risk				
Yes	1			
No	0.89 (0.24~2.81)	0.764		
Flooding depth (m)				
0~3	1			
3~	0.87 (0.37~2.02)	0.740		
Presence of landslide risk				
Yes	1			
No	0.86 (0.27~2.73)	0.798		
Presence of special alert				
Yes	1			
No	1.19 (0.28~5.12)	0.81		
Building count				
2 or more	1		1	
1	4.72 (1.24~18.00)	0.023	8.89 (1.66~47.68)	0.011
Home structure				
Wood	1			
Other	2.02 (0.77~5.27)	0.152		
Presence of medical device use				
Yes	1		1	
No	3.12 (1.09~8.89)	0.034	3.58 (1.01~12.71)	0.049
Placing importance on evacuation				
Yes	1		1	
No	11.92 (1.46~96.99)	0.021	11.87 (1.23~114.28)	0.032
Presence of difficulties in evacuating alone				
Yes	1			
No	0.41 (0.36~4.71)	0.475		
Bedroom floor				
2 or more	1			
1	1.56 (0.54~4.48)	0.410		
Presence of evacuation concern				
Yes	1			
No	0.48 (0.11~2.14)	0.333		
Respondents				
Patient	1			
Others	0.48 (0.18~1.29)	0.148		
Number of disaster risk				
1	1			
2	1 (empty)			




Figure 1. Perception of flooding and landslides among home-care patients.

difference of 25% in the awareness rate of hazard maps depending on whether they knew their neighbors or not.²⁹ Judging from the results of this study and previous studies, it is necessary to extend support to such people to make them more aware of risk of flooding and landslides.

We found that there are many people who need support during evacuation. However, it is necessary to survey what risks they face and what needs they have when they evacuate to consider flooding and landslides countermeasures for home-care patients. Specifically, it is desirable to research what problems exist during preparation, during evacuation, and after evacuation, respectively. It also needs to be examined whether risk awareness of flooding and landslides increases survival rates. It is desirable to investigate how risk awareness of flooding and landslides affects evacuation behavior among patients receiving home care.

Although we found some factors associated with inadequate risk awareness of flooding and landslides, there may be factors that were not detected due to the small sample size. Therefore, it is still unclear what interventions are effective in improving their risk assessment.

Limitations

There are several limitations in this study. First, the response rate for the questionnaire was 49.3%. Because of the small sample size, generalizability was low. Moreover, we do not know if these patients are representative of the population, and there may be undetected factors. In fact, information on the reasons for not responding to the questionnaire could not be obtained. These factors made it difficult to make accurate risk assessment to those who were at risk but did not respond to the survey.

Second, when comparing Fukui with other regions, it is necessary to consider the factors that are unique to Fukui Prefecture. Most of the respondents were residents of Fukui, so many of them lived in flooding areas where the ground is lower than the major rivers and sloping gradually.³¹ These environmental factors may have an influence on the awareness of evacuation. Therefore, its applicability to other communities, especially outside of Japan, is limited.

Third, we failed to consider the seasonal influence on the awareness of flooding and landslides. In Japan, flooding generally occurs during the rainy season in June and July and the typhoon season in August and September, so the awareness of flooding may be higher during these periods. The response period for this questionnaire was from December 1, 2020, to April 30, 2021, a period when flooding is relatively rare, while potentially allowing for more persons not to be evacuated during seasonal flooding, so seasonal factors could not be fully considered.

Fourth, this study was conducted at a single clinic only. Comparisons between clinics were not possible, thus making it difficult to generalize the results.

Fifth, bias may not be sufficiently reduced in the collection of survey responses. When handing out the survey, it was made clear that the responses would not affect future services. In addition, an opt-out format was used to disclose information about the study and to guarantee the opportunity to refuse. However, it might not be sufficient to reduce bias.

Sixth, there exist situations where family members and Orange Home-Care Clinic staff completed the survey. This made the assessment of the perception of flooding and landslides risk among the home-care patients challenging.

Conclusions

There was limited risk awareness of flooding and/or landslides among the home-care patients. This study reported the importance of discussing the vulnerability to flooding and landslides of home-care patients. We identified the risks flooding and landslides and awareness of it among home-care patients living in Fukui. We found that, in addition to 84.9% (169 of 199) of participants at risk of flooding and/or landslides, 58.6% (99 of 169) needed assistance during evacuation. Moreover, more than half of those exposed to risk of flooding and/or landslides did not accurately identify their risk. This result is important in discussing evacuation strategies for home-care patients. Further research in this area is needed in the future.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/dmp.2022.210>.

Data availability statement. Data available on request.

Acknowledgments. The authors express their sincere gratitude to all the clinic staff members, patients, and caregivers involved in this study. We also thank Ms. Tomoyo Nishimura, secretary of Medical Governance Research Institute, and Dr. Morihito Takita for their help with the various administrative procedures.

Author contributions. M.Y. analyzed the data. M.Y., M.K., Y.K., and A.O. wrote the manuscript. All authors conceptualized and designed the study and revised the manuscript.

Conflicts of interest. A.O. received a personal fee from MNES Inc, outside the submitted work. Y.N. received a personal fee from MRT Inc, outside the submitted work.

Ethical standards. The Institutional Review Board of Medical Governance Research Institute granted ethics approval of this study (MG2020-06), adhering to the Ministry of Health Labor and Welfare and The Ministry of Education guidelines, Culture, Sports, Science, and Technology in Japan. In addition, the conduct of this study was announced in writing, and participants were given an opt-out that guaranteed them the opportunity to refuse.

Reference

1. **Kenneth Research.** Home Healthcare Market - Global Revenue, Trends, Growth, Share, Size and Forecast. Accessed October 9, 2021. <https://prtimes.jp/main/html/rd/p/000001427.000059861.html>
2. **Alderman K, Turner LR, Tong S.** Floods and human health: a systematic review. *Environ Int.* 2012;47:37–47. doi: [10.1016/j.envint.2012.06.003](https://doi.org/10.1016/j.envint.2012.06.003)
3. **Fernandez LS, Byard D, Lin CC, et al.** Frail elderly as disaster victims: emergency management strategies. *Prehosp Disaster Med.* 2002;17(2):67–74. doi: [10.1017/s1049023x00000200](https://doi.org/10.1017/s1049023x00000200)
4. **Claver ML, Wyte-Lake T, Dobalian A.** Disaster preparedness in home-based primary care: policy and training. *Prehosp Disaster Med.* 2015;30(4):337–343. doi: [10.1017/s1049023x15004847](https://doi.org/10.1017/s1049023x15004847)
5. **Yoshida I, Ozaki A, Morita T, et al.** Characteristics of flood fatalities in Japan's Typhoon Hagibis in 2019: secondary analysis of public data and media reports. *Disaster Med Public Health Prep.* 2021;1-5. doi: [10.1017/dmp.2021.163](https://doi.org/10.1017/dmp.2021.163)
6. **Ozaki A, Kanemoto Y, Wada M, et al.** A call for individualized evacuation strategies for floods: a case report of secondary surgical site infection in a post-surgery breast cancer patient in Fukushima, Japan, following Typhoon Hagibis in 2019. *Clin Case Rep.* 2021;9(3):1212–1214. doi: [10.1002/ccr3.3727](https://doi.org/10.1002/ccr3.3727)
7. **Ministry of Health, Labour and Welfare.** Overview of patient surveys 2017. Accessed October 9, 2021. <https://www.mhlw.go.jp/toukei/saikin/hw/kanja/17/index.html>
8. **Hara H, Okamura S, Kondo T, et al.** Challenges and recommendations for home medical care in times of disaster: experiences from Typhoon No. 21 in 2008. Amagasaki Medical Association 2018. Accessed October 9, 2021. <https://www.amagasaki.hyogo.med.or.jp/news/1585/>
9. **Hata K, Hata M.** [Balance in Disaster Preparedness between Public, Self-, and Mutual Help for Home-Cared Individuals]. *Gan To Kagaku Ryoho.* 2018;45(Suppl 1):65–68.
10. **Wakui T, Agree EM, Saito T, et al.** Disaster preparedness among older Japanese adults with long-term care needs and their family caregivers. *Disaster Med Public Health Prep.* 2017;11(1):31–38. doi: [10.1017/dmp.2016.53](https://doi.org/10.1017/dmp.2016.53)
11. **Miyatake H, Kosaka M, Arita S, et al.** Videoconferencing for home care delivery in Japan: observational study. *J Med Internet Res.* 2021;23(9):e23539. doi: [10.2196/23539](https://doi.org/10.2196/23539)
12. **Kosaka M, Miyatake H, Kotera Y, et al.** The survival time of end-of-life home care patients in Fukui prefecture, Japan: a retrospective observational study. *Medicine (Baltimore).* 2021;100(38):e27225. doi: [10.1097/md.00000000000027225](https://doi.org/10.1097/md.00000000000027225)
13. **Kosaka M, Miyatake H, Arita S, et al.** Emergency transfers of home care patients in Fukui Prefecture, Japan: a retrospective observational study. *Medicine (Baltimore).* 2020;99(29):e21245. doi: [10.1097/md.00000000000021245](https://doi.org/10.1097/md.00000000000021245)
14. **Brannen DE, Branum M, Pawani S, et al.** Medical allocations to persons with special needs during a bioterrorism event. *Online J Public Health Inform.* 2016;8(3):e200. doi: [10.5210/ojphi.v8i3.6977](https://doi.org/10.5210/ojphi.v8i3.6977)
15. **Ministry of Land, Infrastructure, Transport and Tourism of Japan.** Overlapping hazard maps. 2020. Accessed December 1, 2020. <https://disaportal.gsi.go.jp/maps/?ll=36.003024,139.413972&z=15&base=pale&vs=c1j010u0>
16. **Yahoo Japan.** Yahoo!map. 2020. Accessed December 1, 2020. <https://map.yahoo.co.jp/>
17. **Ministry of Land, Infrastructure, Transport and Tourism of Japan.** Population living in disaster risk areas by prefecture; Accessed October 9, 2021. <https://www.mlit.go.jp/kokudoseisaku/content/001373119.pdf>
18. **McAuley WJ, Spector WD, Van Nostrand J, et al.** The influence of rural location on utilization of formal home care: the role of Medicaid. *Gerontologist.* 2004;44(5):655–664. doi: [10.1093/geront/44.5.655](https://doi.org/10.1093/geront/44.5.655)
19. **Lind JD, Fickel J, Cotner BA, et al.** Implementing Geographic Information Systems (GIS) into VHA home based primary care. *Geriatr Nurs.* 2020;41(3):282–289. doi: [10.1016/j.gerinurse.2019.10.010](https://doi.org/10.1016/j.gerinurse.2019.10.010)
20. **Rees-Roberts M, Williams P, Hashem F, et al.** Hospice at Home services in England: a national survey. *BMJ supportive & palliative care.* Dec 2021;11(4):454–460. doi: [10.1136/bmjspcare-2019-001818](https://doi.org/10.1136/bmjspcare-2019-001818)
21. **Forbes DA, Edge DS.** Canadian home care policy and practice in rural and remote settings: challenges and solutions. *J Agromedicine.* 2009;14(2):119–124. doi: [10.1080/10599240902724135](https://doi.org/10.1080/10599240902724135)
22. **STYLE Aai.** Average annual income and salary in Fukui Prefecture. March 24, 2022. Accessed March 24, 2022. <https://nensyu-style.com/130/>
23. **Database TWI.** World inequality database. March 24, 2022. Accessed March 24, 2022. <https://wid.world/income-comparator10.1080/10599240902724135>
24. **Jia Z, Tian W, Liu W, et al.** Are the elderly more vulnerable to psychological impact of natural disaster? A population-based survey of adult survivors of the 2008 Sichuan earthquake. *BMC Public Health.* 2010;10:172. doi: [10.1186/1471-2458-10-172](https://doi.org/10.1186/1471-2458-10-172)
25. **Hirabayashi Y, Mahendran R, Koirala S, et al.** Global flood risk under climate change. *Nat Clim Chang.* 2013;3(9):816–821. doi: [10.1038/nclimate1911](https://doi.org/10.1038/nclimate1911)
26. **Mahdaviyazad H, Abdolahifar GR.** Assessing hospital disaster preparedness in Shiraz, Iran 2011: teaching versus private hospitals. *Am J Disaster Med.* 2013;8(1):65–73. doi: [10.5055/ajdm.2013.0112](https://doi.org/10.5055/ajdm.2013.0112)
27. **Stewart M, Grahmann B, Fillmore A, et al.** Rural community disaster preparedness and risk perception in Trujillo, Peru. *Prehosp Disaster Med.* 2017;32(4):387–392. doi: [10.1017/s1049023x17006380](https://doi.org/10.1017/s1049023x17006380)
28. **Katada T, Kodama M, Saeki H.** Study on residents' recognition of a flood hazard map and promotion measures of its recognition. *J Proc Hydraul Eng.* 2004;48:433–438. doi: [10.2208/prohe.48.433](https://doi.org/10.2208/prohe.48.433)
29. **Mobile Society Research Institute.** Less than 40% are aware of hazard maps. Awareness of hazard maps tends to be low among those who do not have a relationship with their neighbors. Accessed October 5, 2021. <https://www.moba-ken.jp/project/disaster/disaster20210602.html>
30. **Kapucu N.** Culture of preparedness: household disaster preparedness: household disaster preparedness. *Disaster Prev Manage.* 2008;17(4):526–535. doi: [10.1108/09653560810901773](https://doi.org/10.1108/09653560810901773)
31. **Yamamoto H.** Characteristics of flood disaster by the Fukui heavy rainfall on July 18, 2004 at the Fukui city area on the left bank of the Asuwa River, Fukui Prefecture. *J JPN Soc Nat Disaster Sci.* 2007;26(1):41–53.