

## Evaluation of different methods to handle misreporting in obesity research: evidence from the Canadian national nutrition survey

Mahsa Jessri<sup>1</sup>, Wendy Y. Lou<sup>2</sup> and Mary R. L'Abbé<sup>1\*</sup>

<sup>1</sup>Department of Nutritional Sciences, Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada, M5S 3E2

<sup>2</sup>Dalla Lana School of Public Health, Biostatistics Division, University of Toronto, Toronto, Ontario, Canada, M5S 3M7

(Submitted 13 March 2015 – Final revision received 11 September 2015 – Accepted 28 September 2015 – First published online 2 November 2015)

### Abstract

The association of dietary exposures with health outcomes may be attenuated or reversed as a result of energy intake (EI) misreporting. This study evaluated several methods for dealing with implausible recalls when analysing the association between dietary factors and obesity. We examined data from 16 187 Canadians aged  $\geq 12$  years in the nationally representative Canadian Community Health Survey 2.2. Under- and over-reporting were defined as the ratio of EI:estimated energy requirement  $< 0.7$  and  $> 1.42$ , respectively. Multinomial logistic regression-generalised logit model was conducted to test the utility of different methods for handling misreporting, including (a) adjusting for variables related to misreporting, (b) excluding misreported recalls, (c) adjusting for reporting groups (under-, plausible and over-reporters), (d) adjusting for propensity score and (e) stratifying the analyses by reporting groups. In the basic model, EI showed a negative association with overweight (OR 0.988; 95 % CI 0.979, 0.998) and obesity (OR 0.989; 95 % CI 0.977, 0.999). Similarly, the association between total energy density and overweight (OR 0.670; 95 % CI 0.487, 0.923) and obesity (OR 0.709; 95 % CI 0.495, 1.016) was inverse. Among all methods of handling misreporting, adjusting for the reporting status revealed the most satisfactory results, where a positive association between EI and overweight (OR 1.037; 95 % CI 1.019, 1.055) and obesity (OR 1.109; 95 % CI 1.082, 1.137) was observed ( $P < 0.0001$ ), as well as direct positive associations between energy density and percentage energy from solid fats and added sugars with obesity ( $P < 0.05$ ). The results of this study can help advance knowledge about the relationship between dietary variables and obesity and demonstrate to researchers and nutrition policy makers the importance of adjusting for recall plausibility in obesity research, which is highly relevant in light of global obesity epidemic.

**Key words:** Implausible dietary recalls: Energy misreporting: Obesity: Energy intake

Nutritional studies often rely on self-reported dietary intakes because of the feasibility of this approach especially in large-scale national surveys<sup>(1)</sup>. Inevitably, self-reported dietary intakes involve misreporting (i.e. under- and over-reporting) and implausible intakes<sup>(2)</sup>. The prevalence of under-reporting varies between 10 and 50 % in different studies depending on the cut-off point used for identifying misreporters<sup>(3–5)</sup>. Misreporting of energy and nutrients can be both variable and substantial; hence, it poses a challenge for epidemiologists trying to find a clear relationship between dietary intakes and health outcomes<sup>(6)</sup>. Some studies have confirmed that there is a tendency towards omission of food items that are socially undesirable (i.e. high in fat, added sugars and alcohol), also referred to as selective misreporting<sup>(6–8)</sup>. In addition, energy intake (EI) reporting is influenced by subjects' characteristics<sup>(1,7,9,10)</sup>, for example, the magnitude of under-reporting increases with increasing BMI, which may falsely lead into the conclusion that overweight and obese individuals consume

less energy compared with their normal-weight counterparts (differential misreporting)<sup>(3,11)</sup>. As a result, misreporting is a particular problem for studies investigating the association of diet with obesity as it may render the relationship ambiguous or attenuated, diminishing the usefulness of nutrition data in informing public health policy (bias towards the null)<sup>(12)</sup>.

The misreporting phenomenon is still largely overlooked in obesity research. Several procedures have been proposed for identifying implausible dietary recalls<sup>(13,14)</sup>; even though methods of handling physiologically implausible recalls are less well-studied. Thus far, only few studies (none on adolescents) have investigated methods of handling implausible recalls<sup>(1,8,12,15)</sup>, and none have compared all available methods among different age groups, especially in a large-scale national survey. The purpose of this study is therefore to systematically compare the utility of seven different statistical approaches for handling inaccurate reports of dietary intakes among a nationally representative sample of Canadian adolescents

**Abbreviations:** CCHS, Canadian Community Health Survey; EER, estimated energy requirement; EI, energy intake; PAL, physical activity level; SoFAS, solid fats and added sugars.

\* **Corresponding author:** M. R. L'Abbé, fax +1 416 971-2366, email mary.labbe@utoronto.ca

(12–17 years) and adults ( $\geq 18$  years), when examining the relationship between dietary intakes and obesity. The following methods were used for correcting the misreporting: (1) adjusting for variables related to misreporting, (2) excluding misreported recalls, (3) adjusting for the reporting groups (under-reporter, plausible reporter and over-reporter), (4) adjusting for the propensity score<sup>(15,16)</sup>, (5) adjusting for both reporting groups and propensity score, (6) stratifying the analyses by reporting groups and (7) stratifying the analyses by reporting groups and adjusting for the propensity score<sup>(15,16)</sup>. Propensity score is a statistical technique aimed at reducing bias by equating groups based on variables associated with misreporting<sup>(15,16)</sup>. Adjustment for propensity score in nutrition surveys has only been used among children (proxy reports) and was found to be a useful tool for counteracting attenuation of risk estimates caused by misreporting<sup>(15)</sup>.

Additionally, we examined the validity of participants' self-estimated intake amount (less than, the same or more than the usual amount) collected as part of the 24-h recall procedure, through comparison with the calculated cut-off points for identifying misreporters. Recommendations on how best to counteract attenuation of risk estimates caused by misreporting bias in obesity research are also given.

## Methods

### Study population

Data for this study were collected under the authority of the Statistics Act of Canada, and all data analyses were conducted at the Statistics Canada's Research Data Center. The Canadian Community Health Survey (CCHS) cycle 2.2 (2004–2005) provides the most complete nutritional data on Canadian dietary intakes and is the only available national nutrition data in  $>30$  years<sup>(17)</sup>. CCHS 2.2 is a complex multistage nationally representative survey including cross-sectional nutritional and health data from 35 107 Canadians of all ages, representing  $>98\%$  of the Canadian population from ten provinces<sup>(18)</sup>. Details on the CCHS 2.2 study design, sample and procedures have been published previously<sup>(17)</sup>. For the purpose of this study, we excluded all pregnant ( $n$  175) and lactating ( $n$  92) women, those under 12 years of age ( $n$  8335) and participants with invalid self-reported dietary recalls (as defined by Statistics Canada) ( $n$  39). Data from all respondents with complete data on physical activity and measured weight and height were included, resulting in a final sample of 16 187 subjects. Only participants  $\geq 12$  years were included in this study as only this group had self-reported dietary recalls (as opposed to proxy reports for children). In order to evaluate the association of misreporting with lifestyle and socio-economic characteristics, missing values for these variables were additionally removed, leaving a total of 15 722 individuals for regression analyses. None of the socio-demographic or lifestyle characteristics of individuals included in the final analyses was significantly different from those of participants who were excluded because of missing variables (data not shown).

### Data collection

Detailed dietary intake data were collected through the 24-h recall method using a five-step modified version of the US Department of Agriculture (USDA) Automated Multiple Pass Method (AMPM)<sup>(17,19,20)</sup>. Respondents were asked to recall all foods and beverages consumed in the previous 24 h (midnight to midnight), and energy and nutrient compositions of reported foods were derived from the Health Canada's Canadian Nutrient File<sup>(21)</sup>. As part of the 24-h recall procedure, participants also responded to a question whether they ate less than the usual amount, the same as usual or more than the usual amount during the recall day<sup>(17)</sup>. We additionally used these data to test whether the self-reported usual intake (subjective measure of misreporting) was valid as compared with the cut-off points calculated to identify misreporters (objective measure of misreporting).

Trained interviewers measured height and weight according to standard protocols, and BMI in adults was used as a measure of body fatness using the standard cut-off points for overweight ( $\geq 25$ – $29.99$  kg/m<sup>2</sup>) and obesity ( $\geq 30$  kg/m<sup>2</sup>). For adolescents, categories of Cole *et al.*'s<sup>(22)</sup> were used. Data on socio-demographic characteristics, lifestyle behaviours and health history were collected using interviewer-administered questionnaires<sup>(17)</sup>. Anthropometric measurements and data collection interviews were carried out in person and at participants' homes<sup>(17)</sup>.

EI (kJ/d (kcal/d)) (in 100s of kcal), fibre density (g/4184 kJ (g/1000 kcal)), percentage of EI from solid fats and added sugars (SoFAS), percentage of EI from fruits and vegetables, and dietary energy density (kJ/g (kcal/g)) were used as exposure measures in this study as these have been repeatedly associated with overweight and obesity risk<sup>(23–25)</sup>. Fibre density was derived by calculating grams of NSP fibre intake (g) consumed per 4184 kJ (1000 kcal) EI. Fruits and vegetables were defined on the basis of the WHO Global Strategy on Diet, Physical Activity and Health, and included all fruits and vegetables reported by participants excluding potatoes, nuts and juices<sup>(26)</sup>. SoFAS were defined by the USDA Dietary Guidelines for Americans as high-energy low nutrient-dense food items that need to be limited<sup>(24)</sup>. Dietary energy density was calculated using two definitions: (i) dividing the total energy from foods/beverages (kJ (kcal)) by total weight of foods/beverages (g) or (ii) as above, using foods alone (excluding all drinks)<sup>(27–29)</sup>.

All analyses were performed in terms of EI and not the absolute amounts to reduce extraneous variability and to control for confounding<sup>(30,31)</sup>. The effect of EI was evaluated as a 100-unit offset from the mean, whereas for all other dietary variables a 1-unit change was applied. Descriptive analyses were stratified by sex and age categories, as defined in the Institute of Medicine (IOM) *Dietary Reference Intakes*<sup>(32)</sup>.

### Identification of implausible reporters

Each respondent was categorised as either under-reporter, plausible reporter or over-reporter on the basis of the comparison of their total estimated energy requirement (EER) with their reported EI. The EER was calculated using the IOM



factorial equations that were developed from meta-analyses of studies using double-labelled water as the criterion measure of EER<sup>(33)</sup>. These equations use participants' age, sex, BMI, weight, height and physical activity level (PAL) (sedentary, low active, moderately active, highly active) to estimate their EER<sup>(33)</sup>. As CCHS 2.2 only includes energy expenditure in terms of metabolic equivalents (MET) (kJ/kg per d (kcal/kg per d)), the IOM method was used to convert MET (intensity of an activity compared with the RMR) to the PAL (ratio of total energy expenditure:basal energy expenditure), which was then used in equations to predict EER<sup>(33)</sup>.

Among several methods developed for detecting implausible recalls, the method of McCrory *et al.* is currently the most detailed procedure by which EI is directly compared with EER using cut-off points for their agreements based on error propagation calculations<sup>(8,14,34)</sup>. This is important as other commonly used procedures to identify misreporters (e.g. Goldberg *et al.*<sup>(13)</sup>) are prone to several potential errors, especially in assigning appropriate PAL to individuals<sup>(14)</sup>. In this study, we applied McCrory *et al.*'s intervals for four different levels of physical activity, to data from adolescents and adults, using the level of physical activity each participant reported<sup>(8,34,35)</sup>. As the EI distribution was skewed, we constructed the CI in the log scale and exponentiated the cut-off points, in line with previous studies<sup>(8,35)</sup>. On the basis of our data set, individuals whose EI was <70 % of their EER were classified as under-reporters, and those whose EI was >142 % of their EER were classified as over-reporters ( $\pm 1$  sd). Equations used for this calculations have been published elsewhere<sup>(8,14,34,35)</sup>. We additionally classified individuals on the basis of the  $\pm 2$  sd cut-off points using the 50 and 198 % as the cut-off points for the EI:EER ratio.

### Statistical analyses

All statistical analyses were performed using SAS software (version 9.3; SAS Institute Inc.). To account for the complex multistage sampling frame of the CCHS 2.2, variance estimation was performed using the bootstrap balanced repeated replication technique<sup>(36,37)</sup> and the sample survey weights calculated by Statistics Canada. To maintain a nationally representative sample, a specific weight calculated by Statistics Canada was used in all analyses, which was based on respondent classes with similar socio-demographic characteristics. A two-tailed *P* value <0.05 was used to define statistical significance. Group comparison with Tukey's *post hoc* adjustment was used to evaluate the characteristics of participants classified as under-reporters, plausible reporters and over-reporters (PROC SURVEYREG).

**Calculation of the propensity score.** Step-wise elimination in the logistic regression was applied to identify lifestyle and socio-demographic factors significantly associated with under-reporting among adolescents and adults. We started fitting a model containing all determinants of under-reporting mentioned in previous studies, and the backward selection procedure was applied to screen out non-relevant factors.

The following variables were significant in the final model for adolescents: age, sex, physical activity, alcohol intake in the past 12 months, highest household education, self-reported health, smoking status, province of residence and income. For adults, variables that remained significant in the final model included the following: age, sex, physical activity, having a chronic disease, province of residence, highest household education, self-reported health and smoking status. BMI was not used in the construction of the propensity score as it was the main outcome in the present research<sup>(15)</sup>. The conditional probability of being classified as an under-reporter given the above-mentioned variables was calculated for adolescents and adults using two separate multiple logistic regression models, as follows<sup>(15)</sup>:

Propensity score = estimated probability (under-reporter | covariates).

**Statistical models for handling misreporting.** To compare the utility of different procedures for handling misreporting, the association of overweight/obesity was assessed in relation to a number of key food items identified by the WHO as the main determinants of obesity<sup>(23)</sup>. Multinomial logistic regression-generalised logit model (PROC SURVEYLOGISTIC) was conducted using a classification variable indicating overweight and obesity as outcomes of interest and six dietary variables as exposure measures: EI (kJ/d (kcal/d)), fibre density (g/4184 kJ (g/1000 kcal)), percentage of energy from SoFAS, percentage of energy from fruits and vegetables, total energy density (kJ/g (kcal/g)) and food-only energy density (kJ/g (kcal/g)). The following eight models were then examined and compared. The first model (basic model) was only adjusted for individuals' age and sex (model I). The second model was the same as the basic model but also adjusted for all confounding variables used in calculation of the propensity score (model II). Model III, however, was identical to the basic model with the recalls identified as under-reporter or over-reporter using the method of McCrory *et al.*<sup>(14)</sup> being removed (model III). Other models were the same as the basic model but additionally adjusted for the reporting group (model IV), propensity score (model V) and for both the reporting group and propensity score at the same time (model VI). Further analyses were conducted stratifying the analyses by the reporting group (model VII), and stratifying the analyses by the reporting group and adjusting for the propensity score simultaneously (model VIII).

## Results

### Part A: prevalence and determinants of misreporting

On the basis of the  $\pm 1$  sd cut-off point, 40.47 % of Canadian adolescents and 42.3 % of adults were categorised as misreporters, whereas the corresponding percentages using the  $\pm 2$  sd cut-off point were 12.18 and 13.6 %, respectively (online Supplementary Fig. S1). Throughout this study the more stringent cut-off point ( $\pm 1$  sd) was used for screening out implausible recalls.

**Table 1.** Weighted mean ratio of energy intake (EI):estimated energy requirement (EER) and disparity values (EI – EER) for Canadian adolescents (12–17 years) and adults (≥18 years) by BMI categories (*n* 16 187)\* (Mean values with their standard errors)

Age/sex groups	BMI categories	Weighted mean EI:EER	SE	Weighted mean EI – EER (kJ)†	SE	Weighted mean EI – EER (kcal)†	SE
<b>Adolescents</b>							
M	Normal weight	1.17‡§	0.02	1518‡§	243	363‡§	58
	Overweight	0.94§¶	0.03	–820§¶	414	–196§¶	99
	Obese	0.75 ¶	0.03	–3715 ¶  **	460	–888 ¶  **	110
F	Normal weight	1.20‡§	0.02	1431‡§	146	342‡§	35
	Overweight	0.86§¶	0.03	–1216§¶	243	–291§¶	58
	Obese	0.75 ¶	0.04	–2769 ¶  **	435	–662 ¶  **	104
<b>Adults</b>							
M	Normal weight	1.06‡§    ††	0.03	632‡§    ††	301	151‡§    ††	72
	Overweight	0.90§¶	0.02	–1151§¶	280	–275§¶	65
	Obese	0.81 ¶	0.02	–2409 ¶  ** ††	205	–576 ¶  ** ††	49
F	Normal weight	0.98‡§    ††	0.02	–247‡§    ††	188	–59‡§    ††	45
	Overweight	0.89§¶	0.02	–971§¶	159	–232§¶	38
	Obese	0.82 ¶	0.02	–1728 ¶  ** ††	150	–413 ¶  ** ††	36

F, females; M, males.

\* Estimates are weighted means and bootstrapped variances (balanced repeated replication technique); EI was from the Canadian Community Health Survey 2.2 24-h dietary recalls and the EER was from the Institute of Medicine<sup>(33)</sup> equations; for adolescents 12–17 years of age, Cole *et al.*'s<sup>(22)</sup> categories were used to define obesity.

† Negative values indicate total kJ/d (kcal/d) of under-reporting.

‡ Significantly different between adolescents and adults in normal-weight individuals in each sex (*P* < 0.05).

§ Significantly different between normal weight and overweight in each age and sex group (*P* < 0.001).

|| Significantly different between normal weight and obese in each age and sex group (*P* < 0.001).

¶ Significantly different between overweight and obese in each age and sex group (*P* < 0.05).

\*\* Significantly different between adolescents and adults in obese individuals in each sex (*P* < 0.05).

†† Significantly different between males and females in normal-weight adults (*P* < 0.01).

†† Significantly different between males and females in obese adults (*P* < 0.01).

Generally, the weighted mean ratio of EI:EER was significantly lower in overweight and obese individuals compared with their normal-weight counterparts (*P* < 0.0001) (Table 1). In addition, the ratio of EI:EER decreased by age among both sexes and was lower for females compared with males; however, after approximately 50 years of age, men consistently showed lower total EI:EER values (*P* < 0.024) (online Supplementary Fig. S2). In Table 1, disparity values between the reported EI and the recommended EER were also calculated in order to estimate the amount of EI being misreported among different age, sex and BMI categories. Disparity values were calculated by subtracting the IOM EER<sup>(33)</sup> from the EI reported in the CCHS 2.2. Negative disparity values represent the magnitude of energy under-reporting, whereas positive values show over-reporting. Among normal-weight males, disparity values were positive and significantly higher in adolescents compared with adults (*P* = 0.013). Disparity values among overweight and obese individuals were consistently negative, with the highest value being –3715 (SE 460) kJ (–888 (SE 110) kcal) among obese males aged 12–17 years, followed by –2769 (SE 435) kJ (–662 (SE 104) kcal) among obese females of the same age group (an under-reporting of approximately 25% of EER).

Table 2 presents the descriptive analyses of several covariates stratified by reporting group (under-reporters, plausible reporters and over-reporters) among adults (≥18 years). Under-reporters were more likely to be older (*P* = 0.0013) and to have higher BMI (*P* < 0.0001), diabetes (51.13 *v.* 30.33%; *P* < 0.0001), hypertension (36.64 *v.* 30.40%; *P* = 0.0003), heart disease (36.08 *v.* 31.11%; *P* = 0.035) and at least one chronic disease (34.53 *v.* 28.92%; *P* = 0.0031). In addition, the percentage of under-reporters was higher among residents of Prairie

provinces (Manitoba and Saskatchewan) and Ontario, those with secondary education or less and daily smokers (*P* < 0.006). Similar results were observed among adolescents (12–17 years) (online Supplementary Table S1).

The weighted mean values of dietary determinants of obesity by reporting status are reported for different age and sex categories to examine evidence of potential 'selective misreporting' (Table 3). Adult under-reporters reported substantially lower mean EI (5999 (SE 75) kJ/d (1434 (SE 18) kcal/d) in males and 4497 (SE 58) kJ/d (1075 (SE 14) kcal/d) in females) compared with the plausible (10 924 (SE 83) kJ/d (2611 (SE 20) kcal/d) in males and 8229 (SE 83) kJ/d (1967 (SE 20) kcal/d) in females) and over-reporters 18 756 (SE 297) kJ/d (4483 (SE 71) kcal/d) in males and 13 669 (SE 322) kJ/d (3267 (SE 77) kcal/d) in females (*P*<sub>trend</sub> < 0.0001). Similarly, percentage of energy from SoFAS, total energy density and food-only energy density were significantly higher in over-reporter males and females compared with under- and plausible reporters (*P*<sub>trend</sub> < 0.0001). In contrast, weighted mean fibre density and percentage of energy from fruits and vegetables were higher in under-reporters compared with plausible and over-reporters (*P*<sub>trend</sub> < 0.0064). Similar selective misreporting of dietary variables was also observed among adolescents, although the magnitude was not as large as in adults, probably because of the lower rate of misreporting in younger individuals.

*Part B: comparison of different methods to handle misreporting*

Table 4 shows the OR and 95% CI obtained from six different regression models for the association between overweight and



**Table 2.** Descriptive weighted analysis of covariates (row percentages) stratified by the reporting group (differential misreporting) among Canadian adults ( $\geq 18$  years) ( $n$  11 748)\* (Mean values with their standard errors)

Characteristics	Under-reporters†		Plausible reporters‡		Over-reporters§		P
	Weighted mean/%	SE	Weighted mean/%	SE	Weighted mean/%	SE	
Sex (%)							
Males	31.10	1.52	57.09	1.31	11.81	1.25	0.31
Females	31.61	1.46	58.47	1.32	9.91	0.95	
Age (years)	46.84	0.49	46.08	0.41	43.30	0.76	0.0013
BMI (kg/m <sup>2</sup> )	28.56	0.22	26.92	0.16	25.13	0.29	<0.0001
Self-reported diabetes (%)							
Yes	51.13	3.73	44.95	3.51	3.91	1.23	<0.0001
No	30.33	1.22	58.44	0.96	11.23	0.87	
Self-reported hypertension (%)							
Yes	36.64	1.76	56.28	1.82	7.08	1.12	0.0003
No	30.40	1.38	58.05	1.09	11.56	0.92	
Self-reported heart disease (%)							
Yes	36.08	3.18	57.25	3.36	6.67	1.31	0.035
No	31.11	1.27	57.80	0.98	11.08	0.88	
Has at least one chronic condition (%)							
Yes	34.53	1.66	55.96	1.31	9.51	1.03	0.0031
No	28.92	1.39	59.17	1.30	11.91	1.05	
Physical activity (%)							
Inactive	29.70	2.47	57.26	2.32	13.04	1.68	0.066
Moderately active	31.17	1.17	58.52	1.02	10.31	0.90	
High/very highly active	39.66	4.13	49.43	3.87	10.91	2.58	
Province of residence (%)							
NFLD, PEI, NS, NB	32.46	1.83	59.61	1.84	7.93	1.00	<0.0001
QC	24.03	1.72	60.97	2.04	15.00	1.65	
ON	35.24	1.79	56.11	1.74	8.65	1.17	
MB, SK	36.47	2.04	54.43	2.02	9.10	1.07	
AB	34.25	2.79	54.87	2.72	10.89	1.54	
BC	28.11	2.35	59.49	2.53	12.40	1.59	
Marital status (%)							
Never married	31.56	1.45	57.45	1.23	10.99	1.01	0.22
Married	34.47	2.02	56.26	2.03	9.27	1.03	
Widowed	28.81	1.67	59.67	1.71	11.53	1.45	
Highest household education (%)							
<Secondary education	34.76	1.94	54.76	1.94	10.47	1.49	0.006
Secondary education	38.39	3.50	53.68	3.19	7.93	1.26	
Some post-secondary education	32.70	2.91	58.64	2.96	8.66	1.47	
Post-secondary education	29.66	1.40	58.75	1.13	11.59	1.05	
Income adequacy (%)							
Lowest	34.12	2.83	57.30	3.43	8.58	1.95	0.22
Lower middle	34.17	2.43	56.82	2.21	9.00	1.28	
Upper middle	30.24	1.57	57.34	1.49	12.42	1.38	
Highest	30.54	1.89	59.33	1.73	10.12	1.26	
N/S	29.60	3.27	56.26	3.29	14.14	2.67	
Drank alcohol in past 12 months (%)							
Yes	30.47	1.39	58.35	1.03	11.18	0.94	0.093
No	35.29	2.03	55.22	2.10	9.49	1.35	
Self-perceived level of stress (%)							
Not at all	28.27	1.24	60.73	1.51	11.00	1.39	0.145
A bit stressful	33.35	2.03	56.25	1.77	10.40	1.05	
Quite a bit (extreme)	32.50	1.84	56.05	1.97	11.45	1.28	
Immigration status (%)							
Canadian born	31.41	1.43	57.46	1.18	11.12	0.79	0.782
Immigrant	31.17	2.00	58.76	2.33	10.07	1.80	
Smoking status (%)							
Daily	33.74	2.05	54.85	1.97	11.42	1.39	0.0053
Occasional	23.73	3.15	58.83	4.45	17.44	4.71	
Former	33.13	1.63	59.05	1.63	7.82	0.88	
Never smoked	29.91	1.61	58.23	1.40	11.86	1.28	
Self-perceived health status (%)							
Poor/fair	34.78	2.39	55.74	2.55	9.49	1.31	0.068
Good	33.04	1.63	58.25	1.59	8.71	1.19	
Very good	30.98	1.71	57.34	1.54	11.68	1.29	
Excellent	27.76	2.41	58.89	2.31	13.35	1.68	
Aboriginal of North America (%)							
Yes	30.06	3.51	58.11	4.09	11.83	3.32	0.928
No	31.37	1.26	57.77	0.97	10.86	0.85	

NFLD, Newfoundland; PEI, Prince Edward Island; NS, Nova Scotia; NB, New Brunswick; QC, Quebec; ON, Ontario; MB, Manitoba; SK, Saskatchewan; AB, Alberta; BC, British Columbia.

\* Estimates are weighted means and bootstrapped variances (balanced repeated replication technique).

† Under-reporters: individuals whose energy intake (EI) was <70% of their estimated energy requirement (EER).

‡ Plausible reporters: individuals whose EI was between 70 and 142% of their EER.

§ Over-reporters: individuals whose EI was >142% of their EER.

**Table 3.** Descriptive weighted analysis of dietary determinants of obesity as set by the WHO stratified by the reporting group (selective misreporting) among Canadian adults (≥18 years) and adolescents (12–17 years) (*n* 16 187)\* (Mean values with their standard errors)

Dietary variables	Under-reporters†		Plausible reporters‡		Over-reporters§		<i>P</i> <sub>trend</sub>
	Weighted mean	SE	Weighted mean	SE	Weighted mean	SE	
<b>Adult males</b>							
EI (kJ)	5999	75	10 924	83	18 756	297	<0.0001
EI (kcal)	1434	18	2611	20	4483	71	<0.0001
Fibre density (g/4184 kJ (1000 kcal))	9.18	0.40	7.83	0.13	7.24	0.29	0.0008
% Energy from SoFAS	25.00	0.80	30.39	0.63	32.26	1.45	<0.0001
% Energy from fruits and vegetables¶	4.74	0.52	3.92	0.23	2.59	0.21	<0.0001
Total energy density (kJ/g)**	2.51	0.04	3.30	0.04	4.05	0.08	<0.0001
Total energy density (kcal/g)**	0.60	0.01	0.79	0.01	0.96	0.02	<0.0001
Food-only energy density (kJ/g)††	5.77	0.13	6.40	0.08	7.03	0.17	<0.0001
Food-only energy density (kcal/g)††	1.38	0.03	1.53	0.02	1.68	0.04	<0.0001
<b>Adult females</b>							
EI (kJ)	4497	58	8230	83	13 669	322	<0.0001
EI (kcal)	1075	14	1967	20	3267	77	<0.0001
Fibre density (g/4184 kJ (1000 kcal))	10.44	0.30	8.82	0.14	7.67	0.22	<0.0001
% Energy from SoFAS	22.45	0.61	26.53	0.63	30.67	1.59	<0.0001
% Energy from fruits and vegetables	6.99	0.54	5.34	0.18	4.84	0.46	0.0064
Total energy density (kJ/g)	2.00	0.04	2.89	0.04	3.72	0.08	<0.0001
Total energy density (kcal/g)	0.48	0.01	0.69	0.01	0.89	0.02	<0.0001
Food-only energy density (kJ/g)	5.23	0.13	6.02	0.08	6.36	0.17	<0.0001
Food-only energy density (kcal/g)	1.25	0.03	1.44	0.02	1.52	0.04	<0.0001
<b>Adolescent males</b>							
EI (kJ)	6477	146	11 409	130	18 853	448	<0.0001
EI (kcal)	1548	35	2727	31	4506	107	<0.0001
Fibre density (g/4184 kJ (1000 kcal))	6.96	0.23	6.42	0.16	6.31	0.23	0.091
% Energy from SoFAS	30.45	1.37	31.30	0.76	36.11	1.47	0.0082
% Energy from fruits and vegetables	2.96	0.38	2.60	0.23	2.90	0.38	0.626
Total energy density (kJ/g)	3.43	0.17	4.01	0.08	4.60	0.08	<0.0001
Total energy density (kcal/g)	0.82	0.04	0.96	0.02	1.10	0.02	<0.0001
Food-only energy density (kJ/g)	7.69	0.25	7.69	0.003	8.07	0.25	0.356
Food-only energy density (kcal/g)	1.84	0.06	1.84	0.03	1.93	0.06	0.356
<b>Adolescent females</b>							
EI (kJ)	4644	96	8175	79	13 196	217	<0.0001
EI (kcal)	1110	23	1954	19	3154	52	<0.0001
Fibre density (g/4184 kJ (1000 kcal))	7.55	0.32	7.29	0.14	6.36	0.22	0.0007
% Energy from SoFAS	30.79	1.45	30.62	0.93	33.49	1.81	0.3809
% Energy from fruits and vegetables	3.64	0.59	3.77	0.21	2.79	0.29	0.0523
Total energy density (kJ/g)	2.80	0.12	3.64	0.04	4.51	0.12	<0.0001
Total energy density (kcal/g)	0.67	0.03	0.87	0.01	1.08	0.03	<0.0001
Food-only energy density (kJ/g)	2.80	0.12	6.94	0.12	7.86	0.25	0.0004
Food-only energy density (kcal/g)	0.67	0.03	1.66	0.03	1.88	0.06	0.0004

EI, energy intake; SoFAS, solid fats and added sugars.

\* Estimates are weighted means and bootstrapped variances (balanced repeated replication technique).

† Under-reporters: individuals whose EI was <70% of their estimated energy requirement (EER).

‡ Plausible reporters: individuals whose EI was between 70 and 142% of their EER.

§ Over-reporters: individuals whose EI was >142% of their EER.

|| SoFAS were defined by the US Department of Agriculture & US Department of Health and Human Services<sup>(24)</sup> as high-energy low nutrient-dense food items that need to be limited.

¶ Fruits and vegetables were defined based on the World Health Organization<sup>(26)</sup> excluding potatoes, nuts and juices.

\*\* Total energy density was calculated dividing the total EI from foods and beverages (kJ (kcal)) by total food and beverages weight (g)<sup>(27–29)</sup>.

†† Food-only energy density was calculated dividing the total EI from foods (kJ (kcal)) by total weight of foods (g)<sup>(28,29)</sup>.

obesity as outcomes and several dietary variables as exposures in adults. In the basic model (model I) adjusted for age and sex, a significant negative association was seen between EI and overweight (OR 0.988; 95% CI 0.979, 0.998; *P* = 0.0135) and obesity (OR 0.989; 95% CI 0.977, 0.999; *P* = 0.0553). A similar inverse association was observed between total energy density and overweight (OR 0.670; 95% CI 0.487, 0.923; *P* = 0.0142) and obesity (not significant). More specifically, only the association between fibre density and food-based energy density with obesity was significant and in the expected direction in model I

(*P* < 0.0019). Adjusting for covariates (model II) revealed very similar OR for all dietary exposures so that the direction and significance of none of the variables changed. The strongest relationship between overweight, obesity and dietary exposures was seen after excluding misreporters from the analyses (model III). In this model, significantly positive associations between EI and overweight (OR 1.045; 95% CI 1.021, 1.070) and obesity (OR 1.139; 95% CI 1.108, 1.171) were observed (*P* < 0.0001), as well as direct positive associations between percentage of energy from SoFAS, total energy density and food-only



**Table 4.** Association between overweight and obesity risk with dietary determinants of obesity as set by the WHO among Canadian adults ( $\geq 18$  years)\* (Odds ratios and 95 % confidence intervals)

Dietary variables	Basic model (n 11 748) (model I)†		Basic model adjusted for covariates (n 11 748) (model III)‡		Excluding misreporters (n 6725) (model III)§		Adjusting for the reporting group (n 11 748) (model IV)¶		Adjusting for propensity score (n 11 748) (model V)¶¶		Adjusting for the reporting group and propensity score (n 11 748) (model VI)**	
	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI	OR	95 % CI
EI (1 unit = 418 kJ (100 kcal)), overweight	0.988	0.979, 0.998	0.989	0.979, 0.999	1.045	1.021, 1.070	1.037	1.019, 1.055	0.989	0.980, 0.999	1.037	1.019, 1.056
EI (1 unit = 418 kJ (100 kcal)), obesity	0.989	0.977, 0.999	0.995	0.983, 1.007	1.139	1.108, 1.171	1.109	1.082, 1.137	0.993	0.981, 1.005	1.110	1.083, 1.139
Fibre density (g/4184 kJ (1000 kcal)), overweight	1.000	0.980, 1.021	1.002	0.981, 1.023	0.980	0.952, 1.009	0.992	0.972, 1.012	1.001	0.981, 1.021	0.993	0.973, 1.013
Fibre density (g/4184 kJ (1000 kcal)), obesity	0.963	0.942, 0.985	0.973	0.951, 0.995	0.933	0.898, 0.969	0.946	0.925, 0.968	0.964	0.943, 0.986	0.948	0.927, 0.970
%E from SoFAS, overweight	1.001	0.996, 1.007	1.001	0.995, 1.007	1.011	1.003, 1.019	1.003	0.998, 1.009	1.001	0.995, 1.007	1.003	0.997, 1.009
%E from SoFAS, obesity	1.003	0.997, 1.008	1.000	0.994, 1.006	1.012	1.004, 1.020	1.007	1.001, 1.012	1.002	0.996, 1.007	1.006	1.000, 1.011
%E from fruits and vegetables, overweight	1.002	0.986, 1.018	1.003	0.986, 1.020	0.982	0.961, 1.004	0.998	0.981, 1.014	1.003	0.987, 1.019	0.998	0.982, 1.015
%E from fruits and vegetables, obesity	0.984	0.968, 1.001	0.990	0.974, 1.007	0.966	0.938, 0.995	0.977	0.961, 0.994	0.987	0.971, 1.004	0.980	0.964, 0.999
Total energy density (kcal/g), overweight	0.670	0.487, 0.923	0.657	0.471, 0.917	1.039	0.645, 1.674	1.013	0.700, 1.464	0.685	0.496, 0.947	1.013	0.699, 1.467
Total energy density (kcal/g), obesity	0.709	0.495, 1.016	0.701	0.480, 1.023	2.453	1.385, 4.344	1.758	1.131, 2.731	0.774	0.537, 1.115	1.773	1.136, 2.769
Food-based energy density (kcal/g), overweight	1.073	0.930, 1.237	1.059	0.917, 1.222	1.314	1.064, 1.622	1.151	0.993, 1.334	1.063	0.922, 1.225	1.141	0.985, 1.321
Food-based energy density (kcal/g), obesity	1.273	1.093, 1.482	1.193	1.016, 1.399	1.727	1.360, 2.195	1.444	1.231, 1.695	1.238	1.059, 1.448	1.403	1.192, 1.650

SoFAS, solid fat and added sugar; EI, energy intake; %E, percentage of EI.

\* Estimates are weighted and variances are bootstrapped using the balance repeated replication technique. For the propensity score, 0.01 unit offset from mean was chosen due to its small scale and for the EI a 100-unit offset from mean was considered. All other continuous variables were assessed based on 1-unit offset from the mean. Under-reporters: individuals whose EI was <70 % of their estimated energy requirement (EER); plausible reporters: individuals whose EI was between 70 and 142 % of their EER; over-reporters: individuals whose EI was >142 % of their EER.

† Model I: weighted multinomial logistic regression adjusted for age and sex.

‡ Model II: model I additionally adjusted for physical activity, having a chronic disease, province of residence, highest household education, self-reported health and smoking status.

§ Model III: basic model but excluding under-reporters and over-reporters.

¶ Model IV: basic model adjusted for the reporting groups (under-reporters, plausible reporters, over-reporters).

¶¶ Model V: basic model adjusted for propensity score.

\*\* Model VI: basic model adjusted for both propensity score and the reporting group.

energy density with obesity risk ( $P < 0.0028$ ). Furthermore, the negative association between fibre density and percentage of energy from fruits and vegetables with density was changed to be strong and significant ( $P < 0.0205$ ). Including all respondents and adjusting for the reporting group (model IV) revealed similar results that were slightly less pronounced compared with the model excluding misreporters (model III). After adjusting for the propensity score (model V), all associations changed to be similar to the model II (adjusted for the covariates) and be no longer in the expected direction. Finally, adjustment for both the propensity score and the adjusting group (model VI) did not improve results compared with adjusting for the reporting group alone. Additional inclusion of dietary variables into the propensity score calculation did not improve the results (data not shown). The same results were confirmed among adolescents where excluding misreporters (model III) yielded the strongest association between most dietary exposures and overweight and obesity (online Supplementary Table S2).

In adults, when the basic model was stratified by the reporting group, only EI was significantly associated with obesity in all three groups (under-reporters, plausible reporters and over-reporters) (model VII) (Table 5). Additional adjustment for the propensity score did not improve statistical models (model VIII), except for a slight improvement in EI associations with obesity. Generally, the association of most dietary variables with overweight and obesity was significant and in the expected direction among plausible reporters, even though the strongest association between most dietary variables and overweight and obesity was observed among under- or over-reporters (Table 5 and online Supplementary Table S3). Graphical representations of the relationship between EI and BMI among under-reporters, plausible reporters and over-reporters are presented in Fig. 1(a)–(d).

### Part C: agreement of subjective and objective measures of recall validity

To test the validity of participants' self-reported usual intakes, we additionally compared the self-reported usual intake amounts with the  $\pm 1$  sd cut-off point for the agreement between EI and EER. As presented in online Supplementary Fig. S3(a) and (b), 58.95 (SE 1.04)% of adults and 60.06 (SE 1.55)% of adolescents who said they consumed 'their usual amount' were actually plausible reporters and 29.99 (SE 1.34)% and 20.23 (SE 1.22)% of these individuals under-reported their intakes. Among those who reported consuming 'less than the usual' amount of food, only 42.58 (SE 2.13)% of adults and 21.72 (SE 2.05)% of adolescents were under-reporters and 49.75 (SE 2.10)% and 58.62 (SE 3.17)% reported accurately. In addition, of those who reported that they consumed 'more than the usual' amount only 16.98 (SE 2.87)% of adults and 31.11 (SE 4.22)% of adolescents actually over-reported their intakes.

### Discussion

To our knowledge, this is the first study using a large-scale national survey data on adolescents and adults (self-reports)

to compare seven different statistical approaches to counteract attenuation of risk estimates caused by misreporting. Consistent with previous studies on differential misreporting by weight and disease status<sup>(12,15,38,39)</sup>, under-reporters were more likely to be obese and have higher rates of chronic diseases compared with the plausible and over-reporters. In addition, our results showed strong evidence of selective misreporting in line with others<sup>(4,40,41)</sup>, where under-reporters reported significantly higher intakes of healthy foods – such as fibre, fruits and vegetables – and lower intakes of energy and energy-dense foods. Given the high prevalence of such systematic differential and selective misreporting, statistical models that neglected misreporting of EI rendered the association of nearly all dietary exposures with obesity as insignificant or even reversed, even though the variables studied have been strongly suggested by the WHO as major determinants of overweight and obesity<sup>(23)</sup>. In addition, the nature of the relationship between dietary variables and obesity was different among different reporting groups (under-reporters, plausible reporters and over-reporters). Exclusion of misreporters, adjusting for the reporting groups and stratification resulted in risk estimates that were more consistent with the established hypotheses regarding the role of dietary variables in obesity<sup>(23,42,43)</sup>. Particularly, adjusting for the reporting group yielded more consistent results, even when compared with those from plausible reporters in stratified analyses, and it provided the maximum sample size while maintaining biological plausibility.

In line with a previous study<sup>(14)</sup>, findings of this research showed a significant disagreement between objective and subjective measures of intake validity, which suggests that, although individual's self-defined 'usual amount' may be within the normal range of day-to-day intake variation, this does not necessarily translate into the 'habitual' amount needed to maintain the current body weight<sup>(14)</sup>. In addition, this inconsistency confirms that individual's self-assessment of intake amounts cannot be used for identification of inaccurate recalls in nutritional surveys<sup>(14)</sup>.

Thus far, only a few studies have investigated methods of dealing with implausible recalls<sup>(1,8,12,15)</sup>. Huang *et al.*<sup>(8)</sup> in 2005 evaluated this issue and concluded that lack of exclusion of misreporters from the analyses results in non-significant, weak and misleading diet–obesity relationships. However, these authors did not consider the loss of statistical power that occurs as a result of excluding such large number of participants from the analyses and the fact that results would no longer be generalisable to the entire population, because misreporters have unique characteristics that are not shared by the plausible reporters (i.e. differential misreporting)<sup>(8,39,44)</sup>, as also clearly demonstrated in our study. In addition, extreme observations and outliers usually contain valuable information about the outcome of interest, and their exclusion may introduce an unknown bias<sup>(45)</sup>. Even though excluding misreporters may strengthen the diet–disease relationships, as was also seen in the present study, it is not an appropriate methodology and may lead to a selection bias<sup>(12,15,45,46)</sup>. Generally, results from the study by Huang *et al.*<sup>(8)</sup> should be interpreted with caution as all individuals were assumed to be low active for EER calculations because of the lack of data on PAL, which could potentially lead







**Table 5.** Association between overweight and obesity with dietary determinants of obesity as set by the WHO in different models stratified by the reporting group among Canadian adults ( $\geq 18$  years)\* (Odds ratios and 95% confidence intervals)


Dietary variables	Stratification (model VII)†						Stratification and adjustment for propensity score (model VIII)‡					
	Under-reporter (n 3847)		Plausible reporter (n 6725)		Over-reporter (n 1176)		Under-reporter (n 3847)		Plausible reporter (n 6725)		Over-reporter (n 1176)	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
EI (1 unit = 418 kJ (100 kcal)), overweight	1.044	0.990, 1.101	1.045	1.021, 1.070	1.046	1.006, 1.088	1.047	0.993, 1.105	1.046	1.021, 1.071	1.046	1.005, 1.088
EI (1 unit = 418 kJ (100 kcal)), obesity	1.154	1.094, 1.217	1.139	1.108, 1.171	1.083	1.023, 1.147	1.156	1.097, 1.219	1.141	1.109, 1.174	1.085	1.027, 1.147
Fibre density (g/4184 kJ (1000 kcal)) overweight	0.997	0.953, 1.044	0.980	0.952, 1.009	1.070	0.980, 1.168	1.000	0.956, 1.046	0.980	0.952, 1.009	1.070	0.979, 1.168
Fibre density (g/4184 kJ (1000 kcal)) obesity	0.959	0.919, 1.001	0.933	0.898, 0.969	0.941	0.796, 1.114	0.963	0.925, 1.004	0.933	0.899, 0.969	0.951	0.817, 1.108
%E from SoFAS, overweight	0.991	0.982, 1.001	1.011	1.003, 1.019	0.992	0.977, 1.007	0.991	0.981, 1.000	1.011	1.003, 1.019	0.992	0.977, 1.008
%E from SoFAS, obesity	0.994	0.985, 1.004	1.012	1.004, 1.020	1.022	1.002, 1.042	0.993	0.984, 1.002	1.011	1.003, 1.019	1.020	1.000, 1.040
%E from fruits and vegetables, overweight	1.014	0.987, 1.043	0.982	0.961, 1.004	0.999	0.944, 1.056	1.016	0.989, 1.044	0.983	0.962, 1.004	0.997	0.941, 1.055
%E from fruits and vegetables, obesity	0.994	0.972, 1.017	0.966	0.938, 0.995	0.834	0.736, 0.944	0.997	0.976, 1.020	0.969	0.941, 0.998	0.842	0.745, 0.952
Total energy density (kcal/g), overweight	0.574	0.242, 1.359	1.039	0.645, 1.674	2.189	0.831, 5.765	0.580	0.245, 1.371	1.039	0.643, 1.679	2.197	0.825, 5.851
Total energy density (kcal/g), obesity	0.689	0.332, 1.430	2.453	1.385, 4.344	3.576	0.959, 13.34	0.701	0.340, 1.444	2.461	1.378, 4.394	3.460	0.949, 12.613
Food-based energy density (kcal/g), overweight	0.926	0.706, 1.213	1.314	1.064, 1.622	1.073	0.664, 1.734	0.908	0.696, 1.185	1.307	1.059, 1.614	1.093	0.674, 1.773
Food-based energy density (kcal/g), obesity	1.062	0.829, 1.361	1.727	1.360, 2.195	2.206	1.423, 3.418	1.032	0.809, 1.317	1.679	1.314, 2.146	2.023	1.300, 3.148

EI, energy intake; %E, percentage of EI; SoFAS, solid fats and added sugars.

\* Estimates are weighted and variances are bootstrapped using the balance repeated replication technique. Under-reporters: individuals whose EI was <70% of their estimated energy requirement (EER); plausible reporters: individuals whose EI was between 70 and 142% of their EER; over-reporters: individuals whose EI was >142% of their EER.

† Basic model (adjusted for age and sex) stratified by under-reporting, plausible reporting and over-reporting.

‡ Basic model (adjusted for age and sex) additionally adjusted for the propensity score and stratified by under-reporting, plausible reporting and over-reporting.



**Fig. 1.** The relationship between energy intake (EI) and BMI among under-reporters (---, UR), plausible reporters (—, PR) and over-reporters (....., OR) Canadian adults ( $\geq 18$  years) and adolescents (12–17 years). (a) Adult males; (b) adult females; (c) adolescent males; (d) adolescent females. Estimates are weighted and variances are bootstrapped using the Balance Repeated Replication technique.  $\pm 1$  SD cut-off point for plausible reporting:  $0.7 \leq \text{EI}/\text{estimated energy requirement} \leq 1.42$ . To convert kcal to kJ, multiply by 4.184.

to misclassification of additional subjects to the under-reporting group.

Another study on alternative methods of dealing with inaccurate recalls was based on the National Health and Nutrition Examination Survey (NHANES) 1999–2002 data and concluded that stratification by the intake level is more representative of population nutrient intakes compared with data elimination or exclusions<sup>(1)</sup>. These authors observed a significant association between EI and BMI only among plausible reporters and not the total sample<sup>(1)</sup>. This also supports our findings for the total group where no significant association was observed (model I). Nevertheless, the association of EI with BMI in our study was significant for nearly all reporting groups (under-reporters, plausible reporters and over-reporters), which is in line with a previous study<sup>(15)</sup>. Generally, the limitations of stratification (models VII and VIII in our study) are similar to those of data exclusion, as it results in reduced sample size and loss of statistical power, especially among the over-reporter group (smaller  $n$ )<sup>(15)</sup>. An important limitation of the study by Nielsen & Adair<sup>(1)</sup> is the use of a modified Goldberg *et al.*<sup>(13)</sup> method for identifying misreporters, which assumes a certain habitual PAL for individuals without accounting for the error in assigning this PAL. In our study, however, EI was directly compared with EER using cut-off points for their agreements based on error propagation calculations<sup>(8,34)</sup>. This is important as previous studies have noted a very low precision for assigning PAL to individuals, which may also explain the lack of sensitivity of the Goldberg *et al.*<sup>(13)</sup> cut-off point for identifying inaccurate dietary reports<sup>(47)</sup>. Another limitation of the Goldberg *et al.*<sup>(13)</sup> method is that only extremely inaccurate recalls ( $\pm 2$  SD) are identified, even though misreporting can occur to varying degrees.

In 2011, Mendez *et al.*<sup>(12)</sup> concluded that adjusting for the reporting status through inclusion of a dummy variable for reporting group resulted in stronger associations between diet and obesity and yielded results similar to when misreporters were excluded from the analyses. Our findings corroborate these conclusions; adjustment for the reporting group maintained the statistical power and shifted the association of dietary exposures with overweight and obesity to the expected direction among Canadian adolescents and adults. Although the study by Mendez *et al.*<sup>(12)</sup> was the first to systematically compare the effect of ‘adjusting for the reporting group’ with ‘excluding misreporters’, it suffers from the same limitation as other previous studies in the field, which is assumption of a habitual PAL for all participants without consideration of error in assigning this PAL.

The most recent study that explored different methods of handling misreporting was conducted on children aged 2–9 years (proxy reports) and was the first to calculate and apply a propensity score for handling inaccurate recalls<sup>(15)</sup>. These authors concluded that mutual adjustment for the reporting group and a propensity score is a useful tool for obtaining unbiased risk estimates in obesity research on children<sup>(15)</sup>. However, their findings may have been influenced by the proxy-reported nature of diet recalls and lack of consideration of children’s PAL in calculation of EER, for identifying misreporters and for developing the propensity score. Our study is the first on adolescents and adults to develop and apply the propensity score as a means of counteracting misreporting bias. We found that among adolescents and adults adjusting for the propensity score had no benefit over adjusting for the reporting group for

improving the association between dietary exposures and obesity. This discrepancy may reflect higher differential and selective misreporting among adolescents and adults compared with children, which may not be simply accounted for by inclusion of a propensity score or other calibration methods, which assume a linear (non-differential) measurement error with a constant variance<sup>(48–50)</sup>.

Future studies could test the applicability of constructing calibration scores based on biomarker data in large-scale national surveys where dietary measures are also available for the same subjects. Although exclusion of misreporters strengthened the diet–obesity relationships in this study, it is not an appropriate strategy because of the introduction of an unknown bias by exclusion of about 40% of the population (misreporters) who are systematically different from the plausible reporters (different lifestyle and higher obesity and chronic disease risk). In the absence of biomarker measurements in the Canadian national nutrition survey, our results suggest that simple adjustment for the reporting group is superior to other statistical techniques for handling the misreporting bias, retaining adequate power among adolescents and adults.

### Strengths and limitations

This is the largest known study to compare seven different statistical approaches to address the misreporting bias among adolescents and adults in a nationally representative sample, and it provides important knowledge on the critical role of handling misreporting in obesity research. Developing specific cut-off limits for defining misreporting on the basis of participants' PAL and the algorithm-based method of McCrory *et al.*<sup>(14)</sup> was one of the strengths of this research, compared with studies<sup>(38)</sup> that mistakenly apply the first cut-off points used by Goldberg *et al.*<sup>(13)</sup> in 1991 to identify misreporting<sup>(51)</sup>. This is problematic as cut-off points should be derived on the basis of characteristics of the population being studied to avoid subject misclassification. Inclusion of various covariates, use of a large nationally representative sample and measured anthropometry are some of the other strengths of this study. In addition, the likelihood of misreporting due to missing items or eating occasions was minimised in this research as dietary data were collected using the USDA AMPM; therefore, some of our results may not be applicable to surveys with less comprehensive methods of dietary data collection.

One limitation of this study is day-to-day variation (random non-differential error) associated with dietary recalls, which may have weakened the associations between dietary intakes and obesity. In addition, causal inference is limited owing to the cross-sectional nature of this research<sup>(1)</sup>.

### Conclusions and implications

The present study clearly demonstrated widespread prevalence of selective and differential misreporting across all age and sex groups in the Canadian national nutrition survey, which can undermine the validity of existing national dietary assessments,

diet–disease relationships and public health policies that are developed based on these data, unless appropriate statistical methods are used to deal with such misreporting. Unlike some groups that concluded that national surveys have extremely limited ability for estimating EI and explaining the obesity epidemic<sup>(38)</sup>, we suggest that rigorous methods to control for the misreporting bias are needed and should be applied to any such analysis.

In this study, 'adjusting for the reporting group' maintained the statistical power and shifted the association of dietary exposures with obesity to the expected direction. These results can help advance knowledge about the relationship between dietary variables and obesity and demonstrate to obesity researchers and nutrition policy makers the importance of adjusting for recall plausibility in obesity research. Future studies that assess the sensitivity and specificity of different statistical techniques for correcting the misreporting bias against reference biomarkers of dietary intakes will further advance our abilities to handle misreporting in epidemiological and national cross-sectional studies.

### Acknowledgements

The authors are grateful to staff at Statistics Canada's Research Data Centre, especially Carmina Ng, for their technical support and Didier Garriguet at Statistics Canada for his professional comments.

This research was supported by funds to the Canadian Research Data Centre Network from the Social Science and Humanities Research Council, the Canadian Institute for Health Research (CIHR), the Canadian Foundation for Innovation and Statistics Canada.

M. J. is supported by the Canadian Institute of Health Research (CIHR) Vanier Canada Graduate Scholarship, and the CIHR/Cancer Care Ontario (CCO) Population Intervention for Chronic Disease Prevention (PICDP): a Pan-Canadian Fellowship. M. L. is the Earle W. McHenry professor and is supported by the chair-endowed unrestricted research funds, University of Toronto. Funders had no role in the design, analysis or writing of this article.

Contributions of each author are described as follows: M. J. conceptualised and designed the study; M. J. conducted the statistical analyses and prepared the data tables; W. Y. L, M. J. and M. R. L provided scientific input into the statistical analyses and manuscript; M. J. led drafting of the manuscript; and all authors approved the final version to be published. All of the authors contributed intellectually to the final manuscript.

None of authors declared personal or financial conflicts of interest.

### Supplementary material

For supplementary material/s referred to in this article, please visit <http://dx.doi.org/doi:10.1017/S0007114515004237>

## References

- Nielsen SJ & Adair L (2007) An alternative to dietary data exclusions. *J Am Diet Assoc* **107**, 792–799.
- Black AE, Goldberg GR, Jebb SA, *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 2. Evaluating the results of published surveys. *Eur J Clin Nutr* **45**, 583–599.
- Schoeller DA (1995) Limitations in the assessment of dietary energy intake by self-report. *Metabolism* **44**, 18–22.
- Subar AF, Kipnis V, Troiano RP, *et al.* (2003) Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study. *Am J Epidemiol* **158**, 1–13.
- Schoeller DA, Bandini LG & Dietz WH (1990) Inaccuracies in self-reported intake identified by comparison with the doubly labelled water method. *Can J Physiol Pharmacol* **68**, 941–949.
- Freedman LS, Schatzkin A, Midthune D, *et al.* (2011) Dealing with dietary measurement error in nutritional cohort studies. *J Natl Cancer Inst* **103**, 1086–1092.
- Lioret S, Touvier M, Balin M, *et al.* (2011) Characteristics of energy under-reporting in children and adolescents. *Br J Nutr* **105**, 1671–1680.
- Huang TT, Roberts SB, Howarth NC, *et al.* (2005) Effect of screening out implausible energy intake reports on relationships between diet and BMI. *Obes Res* **13**, 1205–1217.
- Murakami K, Miyake Y, Sasaki S, *et al.* (2012) Characteristics of under- and over-reporters of energy intake among Japanese children and adolescents: the Ryukyus Child Health Study. *Nutrition* **28**, 532–538.
- Black AE & Cole TJ (2001) Biased over- or under-reporting is characteristic of individuals whether over time or by different assessment methods. *J Am Diet Assoc* **101**, 70–80.
- Lafay L, Basdevant A, Charles MA, *et al.* (1997) Determinants and nature of dietary underreporting in a free-living population: the Fleurbaix Laventie Ville Sante (FLVS) Study. *Int J Obes Relat Metab Disord* **21**, 567–573.
- Mendez MA, Popkin BM, Buckland G, *et al.* (2011) Alternative methods of accounting for underreporting and overreporting when measuring dietary intake-obesity relations. *Am J Epidemiol* **173**, 448–458.
- Goldberg GR, Black AE, Jebb SA, *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr* **45**, 569–581.
- McCrorry MA, McCrorry MA, Hajduk CL, *et al.* (2002) Procedures for screening out inaccurate reports of dietary energy intake. *Public Health Nutr* **5**, 873–882.
- Bornhorst C, Huybrechts I, Hebestreit A, *et al.* (2013) Diet-obesity associations in children: approaches to counteract attenuation caused by misreporting. *Public Health Nutr* **16**, 256–266.
- Rosenbaum PR & Rubin DB (1983) The central role of the propensity score in observational studies for causal effects. *Biometrika* **70**, 41–55.
- Health Canada (2006) Canadian Community Health Survey cycle 2.2 nutrition (2004). [http://www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs\\_guide\\_esc-eng.php](http://www.hc-sc.gc.ca/fn-an/surveill/nutrition/commun/cchs_guide_esc-eng.php) (accessed June 2015).
- Béland Y, Dale V, Dufour J, *et al.* (2005) The Canadian Community Health Survey: building on the success from the past. *Proceedings of the American Statistical Association Joint Statistical Meeting, Section on Survey Research Methods*, August 2005, pp. 2738–2746. Minneapolis, MA: American Statistical Association.
- US Department of Agriculture & Agricultural Research Service (2009) USDA automated multiple-pass method. <http://www.ars.usda.gov/Services/docs.htm?docid=7710> (accessed June 2015).
- Moshfegh AJ, Borud L, Perloff B, *et al.* (1999) Improved method for the 24-hour dietary recall for use in national surveys. *FASEB J* **13**, A603 (Abstract).
- Health Canada (2001) The Canadian Nutrient File. Nutrition Research Division, editor. [9]. Ref Type: Data File.
- Cole TJ, Bellizzi MC, Flegal KM, *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240–1243.
- World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases*. WHO Technical Report Series, no. 916. Geneva: WHO.
- US Department of Agriculture & US Department of Health and Human Services (2010) *Dietary Guidelines for Americans*, 7th ed. Washington, DC: US Government Printing Office.
- Howarth NC, Saltzman E & Roberts SB (2001) Dietary fiber and weight regulation. *Nutr Rev* **59**, 129–139.
- World Health Organization (2003) Global strategy on diet, physical activity and health promoting fruit and vegetable consumption around the world. <http://www.who.int/dietphysicalactivity/fruit/en/> (accessed June 2015).
- Ledikwe JH, Blanck HM, Kettel Khan L, *et al.* (2006) Dietary energy density is associated with energy intake and weight status in US adults. *Am J Clin Nutr* **83**, 1362–1368.
- Perez-Escamilla R, Obbagy JE, Altman JM, *et al.* (2012) Dietary energy density and body weight in adults and children: a systematic review. *J Acad Nutr Diet* **112**, 671–684.
- Johnson L, Mander AP, Jones LR, *et al.* (2008) A prospective analysis of dietary energy density at age 5 and 7 years and fatness at 9 years among UK children. *Int J Obes (Lond)* **32**, 586–593.
- Livingstone MB, Robson PJ, Black AE, *et al.* (2003) An evaluation of the sensitivity and specificity of energy expenditure measured by heart rate and the Goldberg cut-off for energy intake: basal metabolic rate for identifying mis-reporting of energy intake by adults and children: a retrospective analysis. *Eur J Clin Nutr* **57**, 455–463.
- Willett W & Stampfer MJ (1986) Total energy intake: implications for epidemiologic analyses. *Am J Epidemiol* **124**, 17–27.
- Institute of Medicine (2000) *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academies Press.
- Institute of Medicine (2005) *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington, DC: National Academies Press.
- ENVIRON International Corporation (2006) What America drinks. [http://www.lanimoo.com/assets/pdf/What\\_America\\_Drinks\\_Report.pdf](http://www.lanimoo.com/assets/pdf/What_America_Drinks_Report.pdf) (accessed June 2015).
- Garriguet D (2008) Impact of identifying plausible respondents on the under-reporting of energy intake in the Canadian Community Health Survey. *Health Rep* **19**, 47–55.
- Rao JNK, Wu CFJ & Yue K (1992) Some recent work on resampling methods for complex surveys. *Surv Methodol* **18**, 209–217.
- Yeo D, Mantel H & Liu TP (1999) Bootstrap variance estimation for the national population health survey. *Proceedings of the Annual Meeting of the American Statistical Association: Survey Research Methods Section*. Baltimore, MA: American Statistical Association.
- Archer E, Hand GA & Blair SN (2013) Validity of US nutritional surveillance: National Health and Nutrition Examination Survey caloric energy intake data, 1971–2010. *PLOS ONE* **8**, e76632.





39. Broyles ME, Harris R & Taren DL (2008) Diabetics under report energy intake in NHANES III greater than non-diabetics. *Open Nutr J* **2**, 54–62.
40. Heitmann BL, Lissner L & Osler M (2000) Do we eat less fat, or just report so? *Int J Obes Relat Metab Disord* **24**, 435–442.
41. Voss S, Kroke A, Klipstein-Grobusch K, *et al.* (1998) Is macronutrient composition of dietary intake data affected by underreporting? Results from the EPIC-Potsdam Study. European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* **52**, 119–126.
42. Tohill BC, Seymour J, Serdula M, *et al.* (2004) What epidemiologic studies tell us about the relationship between fruit and vegetable consumption and body weight. *Nutr Rev* **62**, 365–374.
43. Rolls BJ, Drewnowski A & Ledikwe JH (2005) Changing the energy density of the diet as a strategy for weight management. *J Am Diet Assoc* **105**, S98–S103.
44. Bornhorst C, Huybrechts I, Ahrens W, *et al.* (2013) Prevalence and determinants of misreporting among European children in proxy-reported 24 h dietary recalls. *Br J Nutr* **109**, 1257–1265.
45. Livingstone MB & Black AE (2003) Markers of the validity of reported energy intake. *J Nutr* **133**, 895S–920S.
46. Poslusna K, Ruprich J, de Vries JH, *et al.* (2009) Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *Br J Nutr* **101**, S73–S85.
47. Black AE (2000) The sensitivity and specificity of the Goldberg cut-off for EI:BMR for identifying diet reports of poor validity. *Eur J Clin Nutr* **54**, 395–404.
48. Spiegelman D, McDermott A & Rosner B (1997) Regression calibration method for correcting measurement-error bias in nutritional epidemiology. *Am J Clin Nutr* **65**, 1179S–1186S.
49. Freedman LS, Midthune D, Carroll RJ, *et al.* (2011) Using regression calibration equations that combine self-reported intake and biomarker measures to obtain unbiased estimates and more powerful tests of dietary associations. *Am J Epidemiol* **174**, 1238–1245.
50. Kaaks R, Riboli E & van Staveren W (1995) Calibration of dietary intake measurements in prospective cohort studies. *Am J Epidemiol* **142**, 548–556.
51. Black AE (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes Relat Metab Disord* **24**, 1119–1130.