

Methods: We searched MEDLINE databases using the key terms “refugees” and “global mental health” without language or date restriction. Articles were considered for inclusion in the review if they comprised a population of refugees. Three studies were identified.

Results: Our review showed a great heterogeneity in the prevalence of mental disorders that affect migrants showing an overall prevalence of 20% of these pathologies among them. War-related factors are more associated with Post Traumatic Stress Disorder and post-migration-related factors (acculturation, economic uncertainty and ethnic discrimination) are more associated with mood, anxiety and substance use disorders.

Conclusions: Existing evidence suggests that mental disorders tend to be highly prevalent in refugees many years after resettlement. The increased risk is not only caused by the past adversities in the country of origin but also by the post-migration-related factors. Thus, there is a need for more consistent and rigorous research from a methodological point of view on the mental health of refugees, allowing to find measures to protect and promote their mental health.

Disclosure: No significant relationships.

Keywords: Refugees; global mental health

EPV0572

Dysregulation of the hypothalamic-pituitary-adrenal system in Tuvan alcoholics is associated with a high level of stress in comparison with ethnic Russian patients

T. Shushpanova^{1*}, N. Bokhan^{2,3}, A. Mandel³ and T. Novozheeva¹

¹Mental Health Research Institute of Tomsk National Investigation Medical Center of Russian Academy of Sciences, Clinical Psychoneuroimmunology And Neurobiology, Tomsk, Russian Federation; ²Siberian State Medical University, Department Of Psychiatry, Narcology And Psychotherapy, Tomsk, Russian Federation and ³Mental Health Research Institute Tomsk National Research Medical Center Russia Academy of Science, Department Of Addictive States, Tomsk, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1395

Introduction: Alcoholization is considered as prolonged auto-aggression, low frustration tolerance, filling of the existential vacuum. In all cases, the use of a psychoactive substance that alters the state of consciousness is regarded as a way of escape from real life. **Objectives:** The study of clinically significant markers in alcoholism associated with the constitutional characteristics of craving for alcohol in people of different ethnicity is importance for the formation of new approaches to prevention and treatment.

Methods: 68 Russian alcoholics and 67 Tuvans alcoholics only men and 20 healthy male were monitored. Clinical assessment of the condition of patients was carried out with the traditional clinical description. Enzyme-linked immunosorbent assay kits were used to determine serum hormone levels in patients and volunteers.

Results: Deeper shifts in the increase in ACTH and Cortisol levels were found in alcoholic Tuvians compared with Russian patients, which is associated with a high risk of alcohol dependence and a highly progressive course of the disease. The index of the ratio Cortisol/ACTH (IR) in the blood of alcoholic patients of the Russian (IR - 10,36) and Tuvan (IR - 10,62) nationalities does not differ, but significantly (1.5 times) differs from the indicator in healthy individuals (IR - 15,12).

Conclusions: The background level of dysregulation of the hypothalamic-pituitary-adrenal axis in patients of Tuvan nationality is significantly more pronounced, however, the index of Cortisol/ACTH ratios (IR) in each ethnic group of patients is constant in this disease, which is characterized by a high level of stress.

Disclosure: No significant relationships.

EPV0573

Gender differences in the subjective evaluation of factors determining human health in Russians

R. Shilko*, L. Shaigerova, O. Almazova, A. Dolgikh and M. Rabeson Lomonosov Moscow State University, Psychology, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1396

Introduction: Research into sociocultural mediation of human health engages the role of gender differences in the subjective evaluation of factors that determine health status.

Objectives: The focus of the research was the respondents' opinion about the importance of various factors for human health and subjective well-being.

Methods: 210 men and 403 women aged 14 to 76 years (M = 26.9; SD = 13.7) from six regions of the Russian Federation participated in the study. Participants were asked to rank six factors: genetics, healthy lifestyle, good ecology, regular medical examination, absence of stress (ability to cope with them), financial well-being in terms of their impact on human health (1 is the most important, 6 is the least important).

Results: Both men and women consider “healthy lifestyle” to be the most important factor for human health, while financial well-being - most unimportant. Using the t-test for two independent samples, it was found that: women consider “absence of stress (the ability to cope with it)” significantly more important for health than men (t = -2.569; p = 0.010), while men consider “financial well-being” to be significantly more important than women (t = 2.807; p = 0.005).

Conclusions: It was revealed that men and women equally indicate the most and least important factors determining health. At the same time, subjective assessments of the importance of such factors as absence of stress and financial well-being for health have significant differences. The reported study was funded by the RFBR, project number 17-29-02506.

Disclosure: No significant relationships.

Keywords: gender; factors determining human health; subjective evaluation; sociocultural determination

EPV0574

Hysteria: history of a conceptual and clinical pathomorphosis

A. Sanz Giancola* and C. Alvarez Garcia

Hospital Universitario Príncipe de Asturias, Psychiatry, Alcalá de Henares, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1397

Introduction: Transhistorical psychiatry defends that a psychic alteration can be interpreted as a cultural, historical and personal construction, subject to incessant variations.

Objectives: A journey through the history of the disorder and the successive pathomorphoses it has undergone could provide us with a better understanding of it and explain the reason for the epidemiological trend towards a decrease in its diagnosis; and bring us closer to a universal definition of the phenomenon.

Methods: Bibliographic review

Results: The word hysteria and all its subsequent meanings, not only contain a particular conception of the pathology, but also reflect its different forms of presentation in specific periods of time. Hysteria is presented as a voluble material that can take on any form: from the wandering womb theory of classical Greece to the demonic possessions of the Middle Ages; from the neurological degeneration of Charcot (1825-1893) to the conversion and dissociation of Freud (1856-1939). With the entry of the 20th century, its dramatic clinic has been progressively overshadowed by somatoform disorders and emerging functional somatic syndromes. Today, it is practically unrecognisable and very difficult to diagnose, to the point of having disappeared as a term from the official classifications of our time.

Conclusions: Hysteria is an entity that has not always been the same, neither in its conception nor in its manifestations. Depending on the socio-cultural context in which it is framed, it will be interpreted and expressed in different ways.

Disclosure: No significant relationships.

Keywords: hysteria; history; pathomorphosis; transhistorical psychiatry

EPV0576

Couvade Syndrome: Origin, Characterization, and Frequency

J. Sá Couto*, M. Pão Trigo, B. Da Luz, J. Rodrigues and T. Ventura Gil

Centro Hospitalar Universitário do Algarve, Departamento De Psiquiatria E Saúde Mental, Unidade De Faro, Faro, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1398

Introduction: The word *couvade* originated from the French verb *couver*, meaning to hatch, nest, or brood. Custom of Couvade or Couvade Syndrome (CS) is a poorly understood phenomenon observed since ancient times, in which the expectant father experiences somatic and psychological symptoms of pregnancy.

Objectives: Defining what is CS. Identifying possible origin. Hypothesizing causes. Identifying CS frequency.

Methods: PubMed database search, with "Couvade syndrome" keyword expression. Seven articles were selected among the best matches. Reference lists of articles were reviewed to identify additional articles.

Results: Currently, there are several views on this phenomenon, including religious, cultural, medical, psychoanalytic, and psychological. CS is used in Psychiatry to describe somatic symptoms resembling pregnancy and/or childbirth in expecting fathers, such as weight gain, diarrhea or constipation, toothache, and headache. Lipkin and Lamb (1982) studied 300 couples from New York: they diagnosed Couvade Syndrome in 22,5% of fathers. Nevertheless, Brennan et al. (2007) found different incidence rates of CS diagnose

in different areas of the world: 20% in Sweden; 25–97% in United States; 61% in Thailand; 68% in China; 35% in Russia.

Conclusions: Whether CS constitutes a disease entity, or it should be considered a ritual or custom remains a matter of debate. Different rates of CS around the globe may indicate that culture plays an important role. It may be a way for fathers-to-be to cope with changes imposed by pregnancy in the mother and in the couple. Overall, it is a fascinating intersection between the physiological and psychological realms.

Disclosure: No significant relationships.

Keywords: Couvade; Somatic; Cultural; Pregnancy

EPV0577

A qualitative study on the symptoms and psychological characteristics of young Hwa-Byung patients

S.-A. Park*, J. Kim, S. Lee and S.-W. Choi

Duksung Women's University, Clinical Psychology Lab, Dobong-gu, Seoul, Republic of Korea

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1399

Introduction: Hwa-Byung is characterized by feelings of anger or resent about unreasonable social violence and trauma. Although Prevalence of Hwa-Byung is high in middle-aged, recently Hwa-Byung is occurred in young people. And it is suggested that young Hwa-Byung patients will show different patterns from middle-aged.

Objectives: The purpose of present study was to qualitatively analyze young Hwa-Byung patients' experiences of symptoms and psychological characteristics related to antecedent events of Hwa-Byung.

Methods: 10 women aged 20s were Interviewed using semi-structured questionnaires to in-depth study on their experiences related to Hwa-Byung. The interview data were analyzed using phenomenological approach in order to understand the essence of experiences. In particular, it was analyzed through five steps according to Giorgi (1985). First, by repeatedly reading the material, recurring themes were identified. Second, the meaning units were divided to capture important parts of participants' statements. Third, similar meaning units were grouped together. Fourth, the determined meaning units were described in psychological term. Finally, research data were integrated and presented according to the described meaning units. Reliability between coders was higher than the minimum reliability coefficient.

Results: In symptoms, two themes and seven sub-themes were emerged. In psychological characteristics, three components and five sub-themes were derived. In particular, the antecedent events of young Hwa-Byung patients were related to vertical social relationships, suggesting that Hwa-Byung need to be understood under social context which make them angry.

Conclusions: Present study revealed the social context of Hwa-Byung by discussing the differences between young and middle-aged patients, and furthermore, differences between Hwa-Byung and depression, PTSD patients.

Disclosure: No significant relationships.

Keywords: Hwa-Byung; Qualitative study; cultural-related syndrome; Anger