

---

## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance, but only in exceptional circumstances will systematic or taxonomic studies be acceptable.

Manuscripts, which must be in English or French (with an English summary) should be accompanied by a letter signed by *all* the authors and should be addressed to:

The Editors, Journal of Helminthology,  
395A Hatfield Road,  
St Albans, Herts AL4 0XU,  
England.

Three copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief Introduction; Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Research Notes should also be preceded by a brief abstract. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration will be printed is 12.0×20.0 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale (preferably a scale bar) should normally be given on the figure. Photocopies of illustrations should be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 12.0×20.0 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

ASHFORD, R. W. & BARNISH, G. (1989) *Strongyloides fuelleborni* and similar parasites in animals and man. In: *Strongyloidiasis: a major roundworm infection of man* (editor, D. I. Grove) pp. 271–286. Taylor & Francis: London.

GARSDIE, P. & BEHNKE, J. M. (1989) Primary infection of hamsters with *Ancylostoma ceylanicum*. *Parasitology*, **98**, 283–289.

# Contents

|   | Pages   |
|---|---------|
| Development of the alimentary tract during morphogenesis of the metacercariae of <i>Levinseniella brachysoma</i> . K. V. GALAKTIONOV and I. I. MALKOVA  | 87-94   |
| Studies on the life-cycle of <i>Tremiorchis ranarum</i> . K. V. RAJENDRAN and K. P. JANARDANAN  | 95-101  |
| Imported <i>Opisthorchis viverrini</i> and parasite infections from Thai labourers in Taiwan. H. W. PENG, H. L. CHAO and P. C. FAN  | 102-106 |
| A first instance of human philophthalmosis in Israel. Y. LANG, Y. WEISS, H. GARZOZI, D. GOLD and J. LENGY   | 107-111 |
| Distribution of some elements in hydatid cysts of <i>Echinococcus granulosus</i> from buffalo ( <i>Bubalus bubalis</i> ). N. CHOWDHURY and R. SINGH   | 112-114 |
| The prevalence and intensity of infection with helminth parasites in <i>Mus spretus</i> from the Setubal Peninsula of Portugal. J. M. BEHNKE, C. BARNARD, J. L. HURST, P. K. MCGREGOR, F. GILBERT and J. W. LEWIS | 115-122 |
| Onchocerciasis in British cattle: a study of the transmission of <i>Onchocerca</i> sp. in North Wales. P. J. MCCALL and A. J. TREES   | 123-135 |
| Antagonistic effect of predacious <i>Arthrobotrys</i> fungi on infective <i>Haemonchus placei</i> larvae. J. V. ARAÚJO, M. A. SANTOS, S. FERRAZ and A. S. MAIA  | 136-138 |
| Histopathology of the small intestinal mucosa in <i>Nematodirus spathiger</i> infection in rabbits. H. HOSTE, S. MALLETT and G. FORT  | 139-144 |
| Changing chemosusceptibility in the second-stage larvae of <i>Toxocara canis</i> by long-term incubation. N. AKAO, Y. GOTO, K. KONDO and Y. TSUDA   | 145-150 |
| Density-dependent effects on establishment of <i>Necator americanus</i> and <i>Ancylostoma ceylanicum</i> . S. M. B. NORUZIAN-AMIRI and J. M. BEHNKE  | 151-157 |
| Infectivity, growth and distribution of preovigerous adults of <i>Echinostoma caproni</i> in ICR mice. P. M. MANGER and B. FRIED (Research Note)  | 158-160 |
| <i>Strongyloides ratti</i> : dissociation of immunological memory of the protection against tissue migrating larvae and intestinal adult worms in mice. T. ABE, K. YOSHIMURA and Y. NAWA (Research Note)          | 161-164 |
| Vaccination with hatched but non-activated, non-viable oncospheres of <i>Taenia taeniaeformis</i> in rats. A. ITO and A. HASHIMOTO (Research Note)  | 165-168 |