

## EXPLICIT MERTENS' THEOREMS FOR NUMBER FIELDS

ETHAN SIMPSON LEE 

(Received 9 March 2023; first published online 24 April 2023)

2020 *Mathematics subject classification*: primary 11Y35; secondary 11N32, 11R44.

*Keywords and phrases*: Mertens' theorem, number field, Dedekind zeta function.

Throughout, suppose that  $\mathbb{K}$  is a number field such that  $\mathbb{K} \neq \mathbb{Q}$  with ring of integers  $\mathcal{O}_{\mathbb{K}}$ , degree  $n_{\mathbb{K}}$  and discriminant  $\Delta_{\mathbb{K}}$ . Moreover, let  $\mathfrak{a} \subset \mathcal{O}_{\mathbb{K}}$  denote *integral* ideals,  $\mathfrak{p} \subset \mathcal{O}_{\mathbb{K}}$  denote *prime* ideals,  $N(\mathfrak{p})$  be the *norm* of  $\mathfrak{p}$  and  $\kappa_{\mathbb{K}}$  be the residue of the simple pole at  $s = 1$  of the Dedekind zeta-function  $\zeta_{\mathbb{K}}(s)$  associated to  $\mathbb{K}$ . In what follows, we summarise the contributions of the author's PhD thesis [8].

To study the distribution of prime ideals in a number field, it is desirable to study the asymptotic behaviour of certain counting functions. In particular, we need the *prime ideal theorem* and *Mertens' theorems for number fields*; these are natural generalisations of the famous prime number theorem and Mertens' theorems. Landau originally proved the former in [7] and Rosen originally proved the latter in [11]. In what follows, we introduce all five of the results that this thesis proves; a conditional version (assuming the generalised Riemann hypothesis) of each result is also established to complement these unconditional results.

**Explicit results.** Das established the latest explicit prime ideal theorem in [1] by building upon earlier work from Lagarias and Odlyzko [6]. Throughout, an *explicit* result completely describes the order of growth of the error term therein *and* the associated implied constants. Without assuming conditions, there are significant technicalities in their results; the result only holds for an impractical range and an exceptional (or Landau–Siegel) zero may be present. Moreover, one can use the explicit prime ideal theorem to obtain explicit Mertens' theorems for number fields, although this approach would embed the same technical obstructions into the outcome. On the other hand, we were able to prove explicit Mertens' theorems for number fields with *no* technical obstructions, by making the steps in [11] completely explicit; this was joint work with Garcia (see [2]) and the main result of the thesis.

---

Thesis submitted to the University of New South Wales in October 2022; degree approved on 17 January 2023; supervisor Timothy Trudgian.

© The Author(s), 2023. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

**THEOREM 1.** *If  $x \geq 2$  and  $\mathbb{K} \neq \mathbb{Q}$ , then there is a computable constant  $\Upsilon_{\mathbb{K}}$  depending on  $n_{\mathbb{K}}$  and  $\Delta_{\mathbb{K}}$  only, such that*

$$\sum_{N(\mathfrak{p}) \leq x} \frac{\log N(\mathfrak{p})}{N(\mathfrak{p})} = \log x + A_{\mathbb{K}}(x), \tag{M1}$$

$$\sum_{N(\mathfrak{p}) \leq x} \frac{1}{N(\mathfrak{p})} = \log \log x + M_{\mathbb{K}} + B_{\mathbb{K}}(x), \tag{M2}$$

$$\prod_{N(\mathfrak{p}) \leq x} \left(1 - \frac{1}{N(\mathfrak{p})}\right) = \frac{e^{-\gamma}}{\kappa_{\mathbb{K}} \log x} (1 + C_{\mathbb{K}}(x)), \tag{M3}$$

in which we have  $|A_{\mathbb{K}}(x)| \leq \Upsilon_{\mathbb{K}}$ ,  $|B_{\mathbb{K}}(x)| \leq 2\Upsilon_{\mathbb{K}}/\log x$ ,  $|C_{\mathbb{K}}(x)| \leq |E_{\mathbb{K}}(x)|e^{|E_{\mathbb{K}}(x)|}$ ,  $|E_{\mathbb{K}}(x)| \leq n_{\mathbb{K}}/2(x - 1) + |B_{\mathbb{K}}(x)|$  and

$$M_{\mathbb{K}} = \gamma + \log \kappa_{\mathbb{K}} + \sum_{\mathfrak{p}} \left[ \frac{1}{N(\mathfrak{p})} + \log \left(1 - \frac{1}{N(\mathfrak{p})}\right) \right]$$

satisfies  $-n_{\mathbb{K}} \leq M_{\mathbb{K}} - \gamma - \log \kappa_{\mathbb{K}} \leq 0$ .

Arguably the most important ingredient in our proof of Theorem 1 is the following explicit estimate for the ideal-counting function  $I_{\mathbb{K}}(x) = \#\{a : N(a) \leq x\}$ , which is the number fields generalisation of the floor function. This result was *by far* the most technical result in the thesis to prove, requiring an entire chapter, and it was adapted from the content of the author’s published paper [9].

**THEOREM 2.** *If  $x > 0$  and  $\mathbb{K} \neq \mathbb{Q}$ , then there is a computable constant  $\Lambda_{\mathbb{K}}(n_{\mathbb{K}})$  depending on  $n_{\mathbb{K}}$  only, such that*

$$|I_{\mathbb{K}}(x) - \kappa_{\mathbb{K}}x| < \Lambda_{\mathbb{K}}(n_{\mathbb{K}})|\Delta_{\mathbb{K}}|^{1/(n_{\mathbb{K}}+1)}(\log |\Delta_{\mathbb{K}}|)^{n_{\mathbb{K}}-1}x^{(n_{\mathbb{K}}-1)/(n_{\mathbb{K}}+1)}.$$

To prove Theorem 1, we applied Theorem 2, so the definition of  $\Upsilon_{\mathbb{K}}$  will also depend on  $\Lambda_{\mathbb{K}}(n_{\mathbb{K}})$ . The explicit description for the constant  $\Lambda_{\mathbb{K}}(n_{\mathbb{K}})$  in Theorem 2 that we obtain significantly refines the previous best, which was established by Sunley in her thesis [12, Theorem 3.3.5]. To see the margin of our improvement, refer to Table 1. Further, Theorem 2 is independently interesting, because it has potential applications in studying the zeros, size and value-distribution of  $L$ -functions defined over number fields.

**Applications.** By circumventing the technical issues that would be present in an explicit prime ideal theorem, Theorem 1 unlocks three new applications, which are presented below. Note that Corollary 3 is an explicit version of Bertrand’s postulate for number fields (which was originally established inexplicitly in [5, Section 3]), Corollary 4 gives explicit versions of a result Nagell originally proved in [10] and Corollary 5 was jointly established with Garcia, Suh and Yu in [3].

**COROLLARY 3.** *For  $x \geq 2$ , there exists at least one prime ideal  $\mathfrak{p}$  in  $\mathbb{K}$  such that  $N(\mathfrak{p}) \in [x, Ax]$  when  $\log A \geq 2\Upsilon_{\mathbb{K}}$ .*

TABLE 1. Comparison between values of  $\Lambda_{\mathbb{K}}(n_{\mathbb{K}})$  using the explicit definitions proved in [12, Theorem 3.3.5] and Theorem 2 for several choices of  $n_{\mathbb{K}}$ .

$n_{\mathbb{K}}$	$\Lambda_{\mathbb{K}}(n_{\mathbb{K}})$	
	[12, Theorem 3.3.5]	Theorem 2
2	$1.75425 \times 10^{30}$	$2.49133 \times 10^{10}$
3	$8.57799 \times 10^{44}$	$8.45088 \times 10^{11}$
4	$7.88887 \times 10^{59}$	$9.84482 \times 10^{13}$
5	$1.20023 \times 10^{75}$	$1.41763 \times 10^{16}$
10	$1.90904 \times 10^{153}$	$9.65555 \times 10^{26}$
15	$1.10367 \times 10^{234}$	$5.27930 \times 10^{38}$

**COROLLARY 4.** Let  $g \in \mathbb{Z}[X]$  be irreducible with degree  $d \geq 1$ , leading coefficient  $c$ , discriminant  $D_g$  and weighted discriminant  $\mathbf{D}_g = |c|^{(d-1)(d-2)}|D_g|$ . If  $x \geq \max\{2, \sqrt{\mathbf{D}_g}\}$ , then there are computable constants  $Q_g = O(1)$  and  $\tilde{Q}_g(x) = o(1)$  which depend on  $c$ ,  $d$  and  $D_g$  such that

$$\left| \sum_{p \leq x} \frac{\omega_g(p) \log p}{p} - \log x \right| \leq Q_g \quad \text{and} \quad \left| \sum_{p \leq x} \frac{\omega_g(p)}{p} - \log \log x \right| \leq \tilde{Q}_g(x),$$

where  $\omega_g(p)$  denotes the number of solutions to the congruence  $g(X) \equiv 0 \pmod{p}$ .

**COROLLARY 5.** Let  $f = f_1 f_2 \cdots f_k \in \mathbb{Z}[X]$  be a product of distinct, irreducible non-constant polynomials  $f_1, f_2, \dots, f_k \in \mathbb{Z}[X]$  such that  $f$  has degree  $d \geq 1$ , leading coefficient  $c$  and discriminant  $D_f$ . If  $x \geq \max\{2, |D_f|, \sqrt{\mathbf{D}_f}\}$ , then there exist computable constants  $\mathbf{A}_f = O(1)$  and  $\mathbf{B}_f(x) = o(1)$  which depend on  $c$ ,  $d$  and  $D_f$  such that

$$\left| \frac{1}{\log \log x} \sum_{p \leq x} \frac{\omega_f(p)}{p} - k \right| \leq \frac{\mathbf{A}_f + \mathbf{B}_f(x)}{\log \log x}.$$

The broad-strokes description of our proof of Corollary 4 is that we observed that each of the objective sums is approximately equal to one of the sums in (M1) or (M2), then applied Theorem 1. The significance of Corollary 4 is that  $\omega_g$  is an important multiplicative function that arises in sieve methods. In particular, Halberstam and Richert tell us how to use the sums in Corollary 4 to give upper bounds on the number of primes representable by a polynomial in [4]. Corollary 5 is another consequence of Corollary 4; this result tells us that there is a finite list of primes that certifies the number of irreducible factors of a polynomial  $f \in \mathbb{Z}[X]$ .

### References

- [1] S. Das, *An Explicit Version of Chebotarev's Density Theorem*, MSc Thesis, University of Lethbridge, 2020.
- [2] S. R. Garcia and E. S. Lee, 'Unconditional explicit Mertens' theorems for number fields and Dedekind zeta residue bounds', *Ramanujan J.* **57**(3) (2022), 1169–1191.

- [3] S. R. Garcia, E. S. Lee, J. Suh and J. Yu, 'An effective analytic formula for the number of distinct irreducible factors of a polynomial', *J. Aust. Math. Soc.* **113** (2021), 339–356.
- [4] H. Halberstam and H.-E. Richert, *Sieve Methods*, London Mathematical Society Monographs, 4 (Academic Press, London–New York, 1974).
- [5] T. A. Hulse and M. R. Murty, 'Bertrand's postulate for number fields', *Colloq. Math.* **147**(2) (2017), 165–180.
- [6] J. C. Lagarias and A. M. Odlyzko, 'Effective versions of the Chebotarev density theorem', in: *Algebraic Number Fields: L-Functions and Galois Properties, Proc. Sympos., Univ. Durham, Durham, 1975* (ed. A. Fröhlich) (Academic Press, London, 1977), 409–464.
- [7] E. Landau, 'Neuer Beweis des Primzahlsatzes und Beweis des Primidealsatzes', *Math. Ann.* **56**(4) (1903), 645–670.
- [8] E. S. Lee, *Explicit Mertens' Theorems for Number Fields*, PhD Thesis, University of New South Wales Canberra at the Australian Defence Force Academy, 2023.
- [9] E. S. Lee, 'On the number of integral ideals in a number field', *J. Math. Anal. Appl.* **517**(1) (2023), Article no. 126585, 25 pages.
- [10] T. Nagell, 'Généralisation d'un théorème de Tchebycheff', *J. Math. Pures Appl. (9)* **4** (1921), 343–356.
- [11] M. Rosen, 'A generalization of Mertens' theorem', *J. Ramanujan Math. Soc.* **14**(1) (1999), 1–19.
- [12] J. E. S. Sunley, *On the Class Numbers of Totally Imaginary Quadratic Extensions of Totally Real Fields*, PhD Thesis, University of Maryland, College Park, 1971.

ETHAN SIMPSON LEE, School of Mathematics,  
University of Bristol, Fry Building, Woodland Road, Bristol, BS8 1UG, UK  
e-mail: [ethan.lee@bristol.ac.uk](mailto:ethan.lee@bristol.ac.uk)