

Diabetes prevention after gestational diabetes: challenges and opportunities

Pauline Dunne¹, Deirbhile Sherry², Sharleen O'Reilly¹

¹School of Agriculture and Food Science, University College Dublin, Dublin, D04 V1W8, Ireland

²Department of Clinical Nutrition & Dietetics, Rotunda Hospital, Parnell Square E, Rotunda, Dublin 1, D01 P5W9

Corresponding author: Sharleen O'Reilly, Room SS2.16, Science Centre South Building, University College Dublin, Dublin, D04 V1W8, Ireland, Email: sharleen.oreilly@ucd.ie, Telephone: 01 7167138

Short title: Diabetes prevention after gestational diabetes

Keywords/phrases: Diabetes prevention; women's health; gestational diabetes



This is an Accepted Manuscript for Proceedings of the Nutrition Society. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its

DOI 10.1017/S0029665124007456

Proceedings of the Nutrition Society is published by Cambridge University Press on behalf of The Nutrition Society.

Abstract

Gestational diabetes (GDM) poses significant health concerns for women and their offspring, with implications that extend beyond pregnancy. While GDM often resolves postpartum, a diagnosis of GDM confers a greater risk of future type 2 diabetes (T2D) and other chronic illnesses. Furthermore, the intergenerational impact of GDM predisposes offspring to increased chronic disease risk. Despite the awareness of the short- and long-term consequences of GDM, translating this knowledge into prevention strategies remains challenging. Challenges arise from a lack of clarity among health professionals regarding roles and responsibilities in chronic disease prevention and women's lack of awareness of the magnitude of associated health risks. These challenges are compounded by changes in the circumstances of new mothers as they adjust to balance the demands of infant and family care with their own needs. Insights into behaviour change strategies, coupled with advances in technology and digital healthcare delivery options, have presented new opportunities for diabetes prevention among women with a history of GDM. Additionally, there is growing recognition of the benefits of adopting an implementation science approach to intervention delivery, which seeks to enhance the effectiveness and scalability of interventions. Effective prevention of T2D following GDM requires a comprehensive person-centred approach that leverages technology, targeted interventions, and implementation science methodologies to address the complex needs of this population. Through a multifaceted approach, it is possible to improve the long-term health outcomes of women with prior GDM.

Introduction

Gestational diabetes (GDM) is the most common pregnancy-related complication, affecting 10.9% of European pregnancies annually⁽¹⁾ and 12.4% of pregnancies in Ireland⁽²⁾. The reported prevalence rates of GDM vary (8.9% in northern Europe and 31.5% in eastern Europe)⁽¹⁾, and the International Diabetes Federation reports a global prevalence of GDM of 14%⁽³⁾. These variations in the prevalence rates of GDM are believed to be due to differences in access to and levels of maternal care, screening practices and diagnostic practices, i.e., timing of screening, type of test used, and thresholds used⁽⁴⁾.

GDM results in an increased risk of short- and long-term complications for women and their offspring. In the short term, there is an increased risk of adverse pregnancy outcomes^(5,6), and in the longer term, there is an increased risk of chronic disease, including cardiovascular disease and type 2 diabetes (T2D)^(7,8). Pregnancy affords a unique opportunity for intervention and has been described as a ‘teachable moment’⁽⁹⁾. However, this opportunity does not seem to extend into the postpartum period when challenges arise in participation in and implementation of diabetes prevention interventions^(10,11).

Evidence-based guidelines, such as those provided by the American Diabetes Association (ADA)⁽¹²⁾ and the National Institute for Clinical Excellence (NICE)⁽¹³⁾, support health professionals (HCPs) in providing standardised care for individuals with GDM and in postpartum diabetes prevention, yet a lack of consensus regarding which guidelines to adhere to persists^(14,15). Moreover, the challenges in diabetes prevention after GDM treatment highlight the need for innovative approaches. This review will explore how implementation science, with its focus on theory-driven strategies and robust evaluation processes, can enhance diabetes prevention activities⁽¹⁶⁾.

This review will first provide an overview of GDM. Second, it will outline current challenges in diabetes prevention after gestational diabetes and opportunities to address them. Finally, the review will explore an implementation science approach, and how it may benefit future diabetes prevention research.

Background

GDM is characterised by high blood glucose levels during pregnancy. Commonly, the condition develops in the third trimester (weeks 24 onwards), when a combination of increased demands for insulin and insulin resistance because of pregnancy hormones cause the woman's glucose levels to rise outside of acceptable target ranges⁽¹³⁾. Consistently higher glucose levels are problematic because of their ability to result in short- and long-term complications affecting both mothers and their offspring. In the short-term, higher glucose levels have the potential to result in accelerated growth of the foetus (macrosomia) and the potential for admission to the neonatal intensive care unit. Difficulties at birth are also a risk – birthing a larger infant can result in injury, such as shoulder dystocia, and increased requirements for caesarean section⁽⁵⁾. In the long-term, GDM confers an increased risk of chronic disease, including obesity, T2D and cardiovascular disease (CVD), for mothers and offspring. Women aged 30 years and older have a twofold increased risk of developing GDM, and women with obesity are more than six times more likely to develop GDM⁽¹⁷⁾. Children exposed to higher blood glucose in utero may also be at greater risk of overweight, obesity, and T2D⁽¹⁸⁾.

Commonly, GDM is identified through early or mid-pregnancy screening activity, typically using the International Association of the Diabetes and Pregnancy Study Groups criteria^(4,13). These criteria diagnose GDM if one or more values equal or exceed thresholds of fasting plasma glucose of 5.1mmol/l and/or a 2-hour plasma glucose level of 8.5mmol/l following a 75g oral glucose tolerance test (OGTT)⁽¹³⁾. A lack of consensus persists internationally as to who is screened during pregnancy for GDM. Variations persist between health systems in the use of universal versus risk factor-based screening, with cost implications and heterogeneity among populations among considerations. Selective risk factor-based screening is recommended in the United Kingdom (UK)⁽¹³⁾ and is in place across Ireland⁽¹⁹⁾. However, the recent model of care for diabetes in pregnancy in Ireland has moved to recommend universal screening for GDM while acknowledging the resulting resource implications⁽⁴⁾.

GDM management should include input from an experienced and suitably qualified multidisciplinary team^(4,20). Once diagnosis is confirmed, regular blood glucose monitoring (commonly four times daily for most individuals managing their GDM with diet and physical activity changes alone, increasing to eight-ten times daily when treated with insulin), medical

nutrition therapy (MNT), physical activity and weight management are recommended as first-line treatments to optimise glycaemic management for the duration of pregnancy^(13,21). In addition, 15-30% of women will require pharmacologic intervention, such as metformin or insulin, to achieve the target glycaemic range⁽¹²⁾. Physical activity interventions with exercise components that are in line with general population-based activity recommendations (resistance exercise, aerobic exercise at moderate intensity or a blend of both), have shown overall glycaemic management improvements and reduced need for insulin initiation and reduced insulin dose requirements during pregnancy with GDM^(12,22). Dietetic input of a minimum of three appointments of tailored MNT has been shown to reduce medication needs in GDM patients⁽²³⁻²⁵⁾ and has been proposed as a minimum standard during pregnancy with GDM⁽²³⁻²⁵⁾. However, many dietetic departments fall short of this level of service⁽²⁵⁻²⁷⁾. Despite the evidence supporting the central role that dietetics plays, there are no universal nutritional guidelines for the management of GDM. The Academy of Nutrition and Dietetics published MNT and GDM guidelines back in 2018⁽²¹⁾ but they have not been adopted universally. Certain preconception dietary patterns are associated with a lower risk of GDM; limited evidence from systematic review suggests high intakes of fruits, vegetables, wholegrains, nuts, legumes, and fish with lower intakes of red and processed meats are associated with reduced GDM⁽²⁸⁾. After a pregnancy with GDM, where strict dietary measures with emphasis on macronutrient distribution, meal timing and portion sizes inform treatment, the postpartum dietary messages can appear to be somewhat conflicting for women regarding their return to 'normal' eating⁽²⁹⁾.

Risk factors for GDM

GDM risk factors are well documented and include body mass index (BMI), pregnancy history and ethnicity⁽¹³⁾ (see Table 1). Weight is a modifiable GDM risk factor, as overweight and obesity contribute to insulin resistance⁽³⁰⁾. Greater levels of visceral adipose tissue in the first trimester are associated with hyperglycaemia at 24-28 weeks gestation⁽³¹⁾. Women with overweight and obesity are more likely to develop GDM, compared with women in a healthy BMI range, and those living with higher classes of obesity are at the highest risk^(32,33). Additional risk factors for T2D are well established and include family history, older age, sedentary behaviours, particular ethnic groups, and a history of GDM⁽³⁴⁾. Factors associated with a lower risk of developing T2D after GDM include gestational weight gain (GWG) within recommended guidelines, healthy eating, breastfeeding, and regular physical activity⁽³⁵⁾.

It is estimated that 30 - 50% of European women begin pregnancy living with overweight or obesity⁽³⁶⁾. While weight gain during pregnancy is normal and expected⁽³⁷⁾, meta-analysis indicates that less than one-third (30%) of pregnant women gained weight within the recommended GWG guidelines, and 47% of women gained weight exceeding the recommendations⁽³⁸⁾. The impact of weight gain differs between women with overweight or obesity and women with normal weights. There is a linear increase in the prevalence of GDM in women with obesity whose weight gain goes from low, adequate to high⁽³³⁾. Excessive GWG, generally defined as weight gain outside the American Institute of Medicine guidelines⁽³⁹⁾, has the potential to increase postpartum weight retention (PPWR), which is a contributing factor for long-term obesity development in women⁽⁴⁰⁾. PPWR predisposes women to higher weight status in subsequent pregnancies and is associated with GDM and T2D in the longer term⁽⁴¹⁾. A large retrospective cohort study demonstrated that significant PPWR after a first pregnancy was related to GDM in a consecutive pregnancy⁽⁴²⁾. Researchers have also reported that similar weight gain between two consecutive pregnancies increases the risk for GDM in the second pregnancy^(43,44). An increase in BMI between pregnancies increases the risk of GDM for all women; however, women with a healthy BMI are at a greater risk⁽⁴⁵⁾. An increase of >1 BMI unit from pre-pregnancy to 18 months postpartum is linked to hypertension risk⁽⁴⁶⁾, and PPWR is predictive of a higher BMI after 10 years⁽⁴⁷⁾ and of T2D⁽⁴⁸⁾. There are unique opportunities during pregnancy and postpartum follow-up, to identify risk and implement primary preventative management strategies for weight management⁽⁴⁸⁾. Weight loss is effective in reducing T2D risk. Haw et al.⁽⁴⁹⁾ reported that every kilogram of weight lost was associated with an additional 7% reduction in the risk of progression to T2D⁽⁴⁹⁾. However, HCPs report concerns that talking about weight could damage the HCP/patient relationship and often avoid the topic because they fear patients may feel they are being stigmatised^(50,51). Similarly, women living with obesity and GDM have reported feeling stigmatised throughout their pregnancies due to frequent HCP commentary on their weight and the need for diet and physical activity change⁽⁵²⁾.

Breastfeeding reduces T2D risk postpartum⁽⁵³⁾. Breastfeeding for more than three months has the potential to reduce the risk of future T2D by more than 40% and delay the onset of T2D for up to ten years⁽⁵⁴⁾. Atlantic DIP study data⁽⁵⁵⁾ highlighted that while ethnicity, family history of diabetes, elevated BMI at pregnancy and insulin treatment during pregnancy were all significant predictors of persistent glucose intolerance, lactation at the time of a postpartum OGTT (at 12 weeks postpartum) was significantly associated with reduced odds

of persistent dysglycaemia⁽⁵⁵⁾. A recent meta-analysis demonstrated that breastfeeding for any duration reduced diastolic blood pressure, serum triglycerides, blood glucose, and T2D risk in women with prior GDM⁽⁵⁶⁾. A further systematic review showed that exclusive breastfeeding for six to nine weeks significantly reduced T2D risk compared with formula feeding at the two-year follow-up⁽⁵⁷⁾. A meta-analysis revealed a 34% decrease in diabetes incidence, and breastfeeding for at least 12 weeks significantly reduced T2D risk at the 15-year follow-up^(58,59). In addition to improving glycemia and reducing cardiovascular disease and breast and ovarian cancer risk in mothers, the benefits of breastfeeding also extend to infants through a reduced risk of childhood illness and chronic disease⁽⁶⁰⁾. Antenatal breastfeeding education combined with maternal support after pregnancy in women with GDM is vital⁽⁶¹⁾. There are persistent gaps in breastfeeding promotion for women with GDM and dietitians report a lack of resources and a need for additional training to enhance skills knowledge and confidence^(26,62,63). Staff within GDM services have tended to focus on the short-term benefits of breastfeeding and its benefit for a baby's health, rather than longer-term benefits such as T2D prevention⁽⁶⁴⁾.

Postpartum follow-up in women with GDM

Up to 50% of women with GDM develop T2D within 10 years of delivery⁽¹⁸⁾. The likelihood of future T2D is highest within 5 years, and the prevalence/incidence of T2D varies depending on the diagnostic criteria applied⁽⁶⁵⁾. Such rates suggest that a systematic approach to long-term follow-up after delivery should be applied to mitigate the level of conversion to T2D and/or intervene at the earliest possible opportunity; however, such systematic approaches or monitoring programmes are not standard practices in routine clinical settings⁽¹⁸⁾. In diagnosing GDM, women with a high-risk cardiometabolic profile are identified⁽⁷⁾. Additionally, a GDM diagnosis has been shown to have a profound effect on individuals, resulting in fear, anxiety, and depression⁽⁶⁵⁾. The diagnosis is linked to both stigma^(66,67) and weight stigma, which are widely reported in these populations^(68,69).

Postpartum diabetes screening

Early postpartum identification of abnormal blood glucose levels is important at the population health level, as it allows for preventative approaches to be adopted that can ultimately impact T2D diagnosis rates. Clinical practice guidelines (CPGs) are intended to optimise patient care and are informed by systematic review of the evidence and expert consensus⁽⁷⁰⁾. Two internationally recognised CPGs of note are the ADA "Management of

Diabetes in Pregnancy: Standards of Care in Diabetes”⁽¹²⁾ and NICE guidance for “Diabetes in pregnancy: management from preconception to the postnatal period”⁽¹³⁾. International recommendations from the ADA⁽¹²⁾ recommend using an OGTT at 4-12 weeks postpartum after a GDM pregnancy, this is owing to its increased sensitivity at detecting glucose intolerance (for both prediabetes and T2D). It is suggested that women are tested every 1-3 years thereafter, if the initial 4–12-week postpartum results are normal. These further tests can be performed using any suitable glycaemic test, such as glycated haemoglobin (HbA1c), fasting plasma glucose or OGTT, using non-pregnant thresholds⁽¹²⁾. The NICE guidelines differ and advise not to routinely perform an OGTT but instead from 6-13 weeks postpartum to use fasting plasma glucose to exclude diabetes and, after 13 weeks use HbA1c if fasting plasma glucose is not possible. NICE guidance also states that the level of future diabetes risk advice should be provided in line with early postpartum test results, and an annual HbA1c test should be offered to women with GDM whose postpartum test is within accepted parameters, as the risk of future T2D persists beyond the initial postpartum tests⁽¹³⁾. The differing approaches to the recommendations outlined in these clinical practice guidelines (such as approaches to postpartum screening) may reflect the influence of local contextual factors (e.g., health system infrastructure and funding). Despite differences between their approaches to postpartum screening, the guidelines share a vital common goal in recommending regular postpartum monitoring to address the woman’s future chronic disease risk and prevent their progression to T2D.

Postpartum glucose screening attendance varies from 5-60%⁽⁷¹⁾, with the literature showing a year-on-year decrease after the initial year⁽⁷²⁾. This low uptake of postpartum glucose screening is of concern, as patients with T2D may go undiagnosed and present future health issues, and there is a missed opportunity to establish positive health behaviours to support the prevention of chronic disease. Postpartum screening rates are not captured at the regional or national level; therefore, estimating uptake outside of the local level is difficult. The successful state-funded Health Service Executive chronic disease management (CDM) programme in Ireland recently saw additional support added, or ‘enhancements of care’. One notable enhancement was the addition of women with a prior GDM history (from January 2023 onwards) to be routinely offered annual follow-up with their general practitioners (GPs) and practice nurses to address the risk of future chronic disease (T2D and CVD). These visits include preventative advice, relevant blood work and onwards referrals to intervention programmes or speciality services as appropriate. Previously, this care was offered in an ad

hoc fashion and incurred out of pocket expenses for the woman, as no state funding was provided. While the move is undoubtedly positive, challenges persist with implementation. GDM is generally managed during pregnancy by specialist maternity services, while postpartum follow-up occurs in primary care. Awareness of roles, and responsibilities for diabetes prevention activities and risk of associated chronic disease are low across both settings^(73–76). A danger exists for the CDM programme where the absence of an accompanying care pathway, a tailored education programme specifically for post-GDM women and the requisite primary care resources might place greater emphasis on monitoring without addressing the broader risk factors and preventing future chronic disease.

Further postpartum screening facilitators include implementing electronic reminders through text messaging, email, automated letters, and HCP-mediated communications; all of which can increase screening, monitoring, and preventative practices engagement^(77,78). Tools supporting HCP and patient conversations are helpful in determining a person's priorities. This approach is particularly relevant for establishing awareness and increasing the perceived importance of annual reviews. The woman's identity and priorities shift in the postpartum period with the needs/wants of their children/family becoming their priority and their own needs taking second or even last place. Clear messaging is needed on the woman's increased risk of developing T2D and the risk factors that she can modify are imperative to increase engagement in diabetes screening and prevention programmes. Conveying the impact of a GDM diagnosis on offspring is a further essential message. This insight may motivate women to adopt a family approach to health behaviour change, more so than if the behaviour change were to solely benefit the woman herself. FINDRISC is a well-established tool for assessing and communicating the risk of developing T2D, which includes many of the important determinants for women with previous GDM⁽⁷⁹⁾. In addition to being validated in the general population, it is also effective in determining those more likely to have diabetes/pre-diabetes postpartum⁽⁸⁰⁾. Competing demands, lack of childcare support, inconvenience of testing location (often located within secondary care alongside maternity services), and low self-perceived risk of developing T2D are consistently cited as reasons for a woman's lack of engagement in postpartum diabetes screening. The uptake of the OGTT is poor. It is constrained by the practicalities of the test requiring set appointment times, lasting up to 3 hours, causing a financial burden, the glucose test being an unpleasant procedure and challenges in finding locations to perform the test^(74,75,81).

HCPs cite health system challenges and a low awareness of the increased risk that GDM confers for T2D as barriers to optimum care^(75,76,82). Lithgow et al., 2021⁽⁷⁶⁾ identified hospital and primary care clinician views on their roles in GDM management. Hospital service providers believed that their primary focus was managing GDM, and future prevention was viewed as a lesser priority. As a result, communication regarding the woman's long-term risk was avoided because of the limited time available and a perception that it would overburden women. Postpartum follow-up and screening were seen as important, but they were perceived as being outside of the hospital service remit⁽⁷⁶⁾. Primary care service providers whose core role involves prevention, frequently lack the necessary data to support their work. For instance, the maternity setting may fail to provide them with the relevant information required to inform how best to support and guide the woman after GDM⁽⁸³⁾. GPs view postpartum glucose screening as a key prevention activity and that it is best placed within primary care, however there are several barriers that reduce engagement with testing⁽⁷⁶⁾. Lake et al.⁽⁸⁴⁾ highlighted in their recent systematic review that the barriers to postpartum glucose screening outnumber the facilitators by a ratio of three to one. The recommendations from their evidence synthesis include enhancing communication between HCPs and patients and putting screening forward as a “role modelling opportunity” to set the example of positive health seeking behaviours for their wider family⁽⁸⁴⁾. A subsequent qualitative study also by Lake et al.⁽⁸⁵⁾ used a theoretical approach utilising the behaviour change wheel (BCW)⁽⁸⁶⁾ and theoretical domains framework⁽⁸⁷⁾ to explore the determinants of uptake of T2D screening among women with a history of GDM. Modifiable barriers to and enablers of T2D screening were identified (Table 3). Barriers to T2D screening included lack of knowledge, low awareness of short- and long-term T2D risk, and multiple competing demands, while antenatal education, screening reminders, and social supports were perceived as enablers. The identified barriers and enablers were subsequently mapped to the BCW to highlight the relevant intervention functions and behaviour change techniques which should form the foundation of future intervention strategies⁽⁸⁵⁾.

Cost implications

The financial burden of GDM is substantial – it directly increases maternity care costs by about 34% and will have projected future T2D costs^(88–90). T2D significantly increases health services usage across the health system, in Ireland there is an average of 1.49 GP visits

and approximately one additional outpatient visit for a person with T2D each year⁽⁹¹⁾. Werbrouck et al., 2019⁽⁹²⁾ systematically reviewed the literature to explore the diabetes screening and prevention cost effectiveness in women with prior GDM. The evidence suggested that an OGTT every three years would lead to the lowest cost per case of T2D detected, and prevention of future T2D that may be cost effective or cost saving⁽⁹²⁾. The implementation of a structured diet and physical activity intervention during pregnancy has demonstrated healthcare cost savings through the reduction of GWG and associated GDM and T2D incidence compared with usual care⁽⁹³⁾. Effective interventions targeting the prevention of T2D can simultaneously reduce health burdens and healthcare costs so investment in future prevention programmes has the potential to considerably alleviate future strain on already limited health resources⁽⁹⁴⁾.

Postpartum diabetes prevention after GDM

UK nutrition guidelines for prevention and management of diabetes in the general population promote weight loss in high-risk groups, lifestyle intervention, dietary patterns such as the Mediterranean diet, Dietary Approaches to Stop Hypertension (DASH) diet, or plant-based diets, and culturally tailored multicomponent interventions for better long-term health outcomes⁽⁹⁵⁾. The options for lifestyle interventions include restricted energy intake, reduced total and saturated fat intake, increased fibre intake and increased physical activity⁽⁹⁵⁾. Postpartum dietary quality is a key factor influencing future chronic disease and as such, higher quality dietary patterns, such as the Mediterranean diet⁽⁹⁵⁾, rich in wholegrains, fibre, unsaturated fats, vegetables and fruit, while low in red and processed meats and sugar sweetened beverages, are associated with less long-term weight gain among women with prior GDM⁽⁹⁶⁾, reduced T2D risk⁽⁹⁷⁾, and are recommended⁽⁹⁵⁾. However, this population's dietary quality is generally suboptimal^(98,99) with time constraints, childcare responsibilities, and the cost of healthy eating all reported to be barriers to consuming a higher quality dietary pattern^(100,101).

Diet and physical activity interventions are effective in preventing or delaying T2D in high-risk individuals⁽¹⁰²⁾. Notably, the original diabetes prevention programme showed achieving and maintaining a 7% weight loss through a healthy diet and physical activity could reduce future T2D risk by half^(102,103). When diet and physical activity intervention was

compared with metformin, the number needed to prevent one case of T2D in 3 years was 7, whereas the number needed to prevent T2D for the same duration using the oral hypoglycaemic agent metformin was 14⁽¹⁰⁴⁾. Other interventions have addressed the future T2D risk through targeting health behaviour⁽¹⁰⁵⁾ and whole family-based approaches^(106,107). Reviews of diet- and physical activity-based interventions have demonstrated significant reductions in T2D incidence; one reported a 58% reduction⁽¹⁰⁸⁾, another demonstrated such interventions yielded a 24% lower T2D incidence when compared to routine care⁽¹⁰⁹⁾, and a third examining randomised controlled trials (RCTs) showed a 53% reduction in T2D incidence⁽¹¹⁰⁾. Effective intervention components include energy intake reduction (total calories) coupled with regular physical activity (150 minutes moderate intensity per week), and the programme lasting six weeks or longer⁽⁵³⁾. While many interventions demonstrate positive effects, long-term follow up data is lacking; a recent systematic review and meta-analysis highlighted that two-thirds of included studies reported one year or less of follow-up⁽¹¹¹⁾. Research on diet and physical activity interventions has almost exclusively been conducted in high-income countries; however, a recent systematic review and meta-analysis synthesised findings from RCTs over the past two decades in low- and middle-income countries. While extensive heterogeneity was evident across interventions, evidence suggests that diet and physical activity-based interventions are effective in delaying T2D and improving cardiometabolic health among at-risk populations in low- and middle-income countries⁽¹¹²⁾. Online communication platforms offer increasing opportunities for connection and access to information^(113,114) and women have expressed a preference for postpartum information to be delivered online⁽¹¹⁵⁾, valuing its flexibility⁽¹¹⁶⁾. Social support during the postpartum period can facilitate behavioural change^(116,117).

The postpartum period is a significant time with emotional and physical changes, presenting barriers to physical activity recommendations⁽¹¹⁸⁾. Diet and physical activity modification resulting in weight loss has been shown to be challenging in this population^(102,119,120). GDM requires women to implement significant antenatal diet and physical activity alterations to achieve normoglycaemia⁽¹²⁾. This radical adjustment and pressure to conform to more restrictive dietary and physical activity behaviours can have a significant opposite impact on postpartum behaviours⁽⁷¹⁾. To mitigate future T2D, cardiovascular disease and obesity risk, women are advised to achieve a healthy postpartum weight⁽¹²¹⁾. Peacock et al., reported that dietary interventions resulted in weight loss and altered dietary intake but achieved greater success in women who had not experienced

excessive GWG⁽¹⁰⁸⁾. Intervention timing is critical to effectiveness, with three potential time periods for interventions proposed—prenatal, early postpartum, and late postpartum⁽¹²²⁾. The use of postpartum interventions is a well-documented health promotion challenge⁽¹¹⁾, with barriers that include attending programmes not specifically tailored to the needs of postpartum mothers, unpredictability of the baby's routine and difficulty getting out of the house⁽¹²³⁾. Physical activity is a known T2D protector⁽¹²⁴⁾ and works for women with prior GDM⁽⁵³⁾. However, despite its benefits, physical activity barriers are commonly cited^(125,126). Family-based physical activity interventions, and various forms of social support (family, partner, or community), and organisational support (childcare, time) can influence physical activity levels^(75,125).

Diabetes prevention programmes use clinical practice guidelines to inform their structured education that will address the modifiable risk factors associated with its prevalence, they also use approaches that can be tailored to meet population needs, e.g., cultural considerations, language, and location⁽¹²⁷⁾. Many of the available clinical practice guidelines for GDM management include nutritional care^(13,20,128) but they have different levels of nutrition guidance provided^(129,130). Clinical practice guidelines should provide appropriate nutrition information for HCPs involved in GDM care and should cover the woman's pregnancy to postpartum journey with advice tailored to each stage. Dietetic practice requires much more detailed MNT covering standardised structured antenatal and postpartum diabetes prevention education. Delivering MNT requires comprehensive collaboration with the multidisciplinary care team, and multidisciplinary higher-level management support to develop the implementation plan alongside procuring adequate levels of associated staffing⁽¹³¹⁾. The challenge for GDM is that this process requires increased communication and collaboration across specialised maternity services and primary care settings where streamlined nutrition and dietetic pathways typically do not exist. GPs, practice nurses and other primary care HCPs (such as community dietitians and public health nurses) are likely to have frequent contact with new parents through routine clinical care, childhood immunisations, breastfeeding support, and other health promotion initiatives. Many women will also attend organised group activities for information sharing and peer support⁽¹³²⁾. These interactions all offer opportunities for brief interventions in promoting optimal nutrition, physical activity, and future health. Therefore, it is essential that HCP messages are aligned with best practices and consistent and relevant to the woman's life stage to support optimum behaviour change.

Implementation science and diabetes prevention

The significant gap between research evidence and its translation into real-world practice is often referred to as the "know-do" gap that hinders progress in healthcare⁽¹³³⁾. This challenge is particularly pronounced in diabetes prevention interventions designed to support women after GDM^(84,134,135). The field of implementation science is growing rapidly, offering an opportunity to more rapidly scale interventions into healthcare practices through evidence and theory-informed approaches to bridge this gap. Implementation science in nutrition and dietetics is an emerging field and offers a robust structure and foundation for design implementation, and evaluation of interventions^(136,137).

The complex intervention framework of the UK Medical Research Council and National Institute for Health Research highlights the need for a flexible approach that considers the specific needs and challenges of targeted setting^(133,138). The framework, which was updated in 2019 and 2021, provides a valuable roadmap for developing and evaluating complex interventions aimed at improving health outcomes^(133,138). However, intervention complexity itself can significantly impact implementation and effectiveness across diverse settings. Challenges arise from standardising intervention design and delivery, adapting to local contexts, considering the people involved (staff and patients), navigating the organisational environment, and developing appropriate outcome measures⁽¹³⁹⁾. A key update within the framework emphasises how interventions interact with their context and with system-level changes. This shift aims to identify the conditions necessary for achieving intended change mechanisms and ensure effectiveness in real-world settings⁽¹³³⁾. The framework advocates for the systematic development and evaluation of interventions based on evidence and theory. There are four iterative phases: 1) development/identification, which involves defining the intervention based on evidence and theory; 2) exploration of feasibility and acceptability, which assesses how realistic and acceptable the intervention is for target users and settings; 3) evaluation, which formally tests the intervention's effectiveness in achieving desired outcomes; and 4) implementation, which focuses on integrating the intervention into routine practice and ensuring its sustainability⁽¹³³⁾. Each component addresses core elements such as contextual considerations, program theory, stakeholder engagement, identifying key uncertainties, intervention refinement, and economic considerations. Troughton et al.⁽¹⁴⁰⁾, demonstrate a worked example of how the MRC framework supported the development feasibility and piloting phases of their group-based

lifestyle intervention to prevent T2D in people identified as 'at risk'⁽¹⁴⁰⁾. The MRC framework encourages early consideration of implementation throughout the development and evaluation process, increasing the intervention's potential for successful adoption across settings. This approach aligns with effectiveness-implementation hybrid designs⁽¹⁴¹⁾, where implementation and effectiveness are evaluated concurrently across three hybrid designs, each with a varying balance between implementation and effectiveness assessment.

Intervention complexity can be defined in various ways, including the number of interacting components, targeted groups or organisational levels, variability of outcomes, the degree of tailoring, and whether the intervention has a nonlinear causal pathway⁽¹⁴²⁾. To address real-world complexities, healthcare research increasingly utilises systems thinking and conceptual mapping approaches. These methods focus on the dynamic relationships between "people, processes, activities, settings and structures" within a system^(143,144). The Consolidated Framework for Implementation Research (CFIR) provides a comprehensive taxonomy of factors influencing intervention implementation, allowing researchers to tailor strategies to address specific barriers within different healthcare systems and settings⁽¹⁴⁵⁾. A 2016 qualitative interview study by Bernstein et al.⁽¹⁴⁶⁾, set out to identify barriers and facilitators to postpartum glucose testing and referral, including patient and provider perspectives. The authors used the CFIR to provide a systematic approach to their analysis and mapped the data to four domains: intervention attributes, the characteristics of individuals, the inner and outer contexts⁽¹⁴⁶⁾. More recently, implementation factors influencing the UK National Diabetes Prevention Program (National DPP) were explored using the CFIR inner and outer setting domain constructs⁽¹⁴⁷⁾. Inner setting domain constructs offer insights into factors influencing implementation from within the organisation, such as culture, preparedness for implementation, and networks and communications. Outer setting domain constructs explore external contexts which influence intervention implementation, such as population needs and resources, peer pressure, and external policies and incentives. The findings of this study indicated multiple factors internal to the organisation which can impact on intervention implementation success, in particular leadership engagement and implementation climate, offering valuable insights into facilitating successful implementation of the National DPP⁽¹⁴⁷⁾. The use of frameworks such as CFIR enables researchers to use a common language and approach to synthesise findings on factors influencing implementation.

Implementation theories offer a framework for developing implementation strategies and exploring factors influencing successful or unsuccessful intervention implementation⁽¹⁴⁸⁾. These theories have diverse aims, including guiding evidence-based practice application, exploring implementation influences, and evaluating intervention implementation itself⁽¹⁴⁸⁾. The exploration, preparation, implementation, sustainment (EPIS) framework⁽¹⁴⁹⁾ and normalisation process theory (NPT)⁽¹⁵⁰⁾ fall within the first two categories and help identify barriers and facilitators to implementation. The EPIS framework, developed from the public sector and allied health service implementation literature, outlines four phases: exploration (assessing health needs and identifying best practices), preparation (identifying potential contextual barriers and facilitators), implementation (adopting evidence-based practice), and sustainment (ongoing support structures and processes to ensure delivery achieves the intended impacts)^(149,151). The EPIS framework also describes common and unique factors within the outer system context, inner organisational context, and innovation itself, all of which influence implementation across different phases and is like the CFIR framework in this respect. Bridging factors, which acknowledge the interconnectedness of these contexts, are key components of the EPIS framework. Systematic reviews further highlight the flexibility and robustness for use of the EPIS framework across various income settings⁽¹⁵¹⁾. The “Bump2Baby and me” project⁽¹⁵²⁾ investigated the effectiveness of a mHealth coaching programme including nutrition and physical activity supports in pregnancy and postpartum for women with high risk of developing GDM. Researchers used the EPIS framework to frame the project, such as in evaluating needs to ensure intervention fit (exploration phase), and in planning for intervention readiness (preparation phase)⁽¹⁵²⁾.

On the other hand, the NPT provides an explanatory model for understanding the influence of implementation. It can be used alongside the EPIS to explore relevant contextual and innovation-related factors. Developed within diverse healthcare systems, the NPT focuses on the individual and collective behaviours observed in implementation processes⁽¹⁵³⁾. As an action theory, it describes the mechanisms of social action involved in implementing a new practice. NPT identifies four key constructs that are crucial for successful implementation: coherence, which is the degree to which stakeholders understand and make sense of the new intervention, ensuring buy-in and reducing resistance; cognitive participation, which assesses the level of buy-in and engagement from stakeholders involved in implementing the intervention, highlighting the importance of addressing potential concerns; collective action, which examines the resources and support structures available to

facilitate implementation; and refractory monitoring, which focuses on the ongoing appraisal and feedback mechanisms employed to assess the intervention's effectiveness and make necessary adjustments during implementation⁽¹⁵⁴⁾. Regular monitoring allows course correction and adaptation to ensure that the intervention remains effective within the real-world healthcare environment⁽¹⁵⁵⁾. NPT was applied to identify factors contributing to the successful implementation of a recent postpartum diabetes prevention quality improvement study based in general practice⁽¹⁵⁶⁾. The study found the NPT constructs to be more visible within actively participating practices, and these practices exhibited higher levels of change in diabetes screening, diabetes prevention planning and weight monitoring activities, when compared to moderately participating practices⁽¹⁵⁶⁾.

Theory plays a crucial role in intervention development, particularly in behaviour change. The BCW is a popular framework that ensures comprehensive intervention coverage, focusing on the core determinants of behaviour: capability, opportunity, and motivation⁽¹⁵⁷⁾. This foundation is further elaborated upon in the capability-opportunity-motivation behaviour model (COM-B), which provides a detailed breakdown of these determinants⁽¹⁵⁸⁾. Lake et al.⁽⁸⁵⁾ used the COM-B to map barriers and enablers to postpartum T2D screening to identify which intervention functions and behaviour change techniques (BCTs) could inform the development of an effective intervention to increase T2D screening post-GDM⁽⁸⁵⁾. The BCT taxonomy provides a standardised way to describe the specific BCT components within interventions using 93 distinct techniques⁽¹⁵⁹⁾. This taxonomy, derived from BCW development and physical activity/nutrition BCTs, facilitates clear communication, replication, and systematic evidence syntheses to identify the most effective BCTs for specific behaviours and contexts.

Evaluation plays a critical role in understanding implementation success, and several frameworks guide this process. However, the chosen frameworks should place a strong emphasis on implementation aspects. RE-AIM^(160,161) is a commonly used framework that provides a comprehensive approach to evaluating programme implementation, encompassing five key dimensions: reach (percentage of target population reached), effectiveness (program impact on intended outcomes), adoption (setting or organisation uptake), implementation (fidelity to program design), and maintenance (sustained delivery over time)⁽¹⁶¹⁾. This framework offers a valuable tool for researchers to assess the public health impact of interventions, considering not only intervention effectiveness but also real-world factors

influencing program success. Studies have shown its utility in evaluating diverse programs across various settings, including community health initiatives, chronic disease management interventions, and diabetes prevention interventions^(162,163). By applying the RE-AIM framework, researchers can gain a holistic understanding of program strengths and weaknesses, informing strategies to optimise public health impacts.

Evidence-based guidelines serve as a critical foundation for effective interventions. These guidelines synthesise robust scientific evidence into practical recommendations for healthcare professionals, providing a standardised approach to interventions and ensuring consistency and quality in patient care. However, simply disseminating guidelines is insufficient. Studies have identified several barriers to guideline use, including lack of awareness, lack of familiarity with recommendations, disagreement with guidance, and perceived lack of applicability in real-world practice^(12,134,164). Conversely, factors that enable guideline use include strong leadership support, clear and concise recommendations, and integration with existing workflows⁽⁸⁴⁾.

Quality improvement collaboratives (QICs) provide another powerful tool for driving implementation⁽¹⁶⁵⁾. QICs bring together healthcare professionals from different practices to share best practices, troubleshoot challenges, and collaboratively implement evidence-based interventions. A multilevel QIC approach including provider education, written supporting materials and a variety of reminder methods (electronic health provider as well as patient) increased postpartum glucose screening rates⁽¹⁶⁶⁾. O'Reilly et al.⁽¹⁶⁷⁾ demonstrated improved diabetes screening and BMI monitoring in women with previous gestational diabetes at postpartum follow-up through a quality improvement multicomponent intervention approach within primary care general practice where following the implementation of an intervention with plan–do–study–act feedback components to support ongoing engagement, rates of postpartum diabetes screening increased from 43% to 60%, annual diabetes screening rates increased from 26% to 61%, and a 20% increase in BMI monitoring was observed⁽¹⁶⁷⁾. By fostering a culture of learning and continuous improvement, QICs can significantly enhance the uptake and effectiveness of interventions within general practice settings, yielding cost savings to the health system when applied at scale⁽¹⁶⁸⁾.

Conclusion

There are several challenges and opportunities for diabetes prevention after pregnancy complicated by GDM. Ambiguity in healthcare roles and responsibilities, coupled with limited awareness of associated risks, complicates intervention efforts. However, advances in knowledge, intervention strategies and implementation science offer promise. Primary care is the ideal location for health promoting interventions, postpartum T2D risk assessment and ongoing follow-up for pregnant women with GDM. However, guidelines outlining the role of primary care staff (particularly general practitioners and practice nurses) lack consistency, which may negatively influence the provision of T2D risk management for women with GDM. Coordinated efforts, guided by clinical guidelines and public health policy, are crucial for mitigating T2D risk and promoting lifelong health after GDM.

Acknowledgements: We would like to thank the Maternity Dietitians Ireland group and the HSE Self-Management and Education Supports (SMES) team for their ongoing support.

Financial support: This work was completed as a component of doctoral research for PD and was funded by a Health Research Board (HRB) Collaborative Doctoral Award, 2019 (CDA-2019-001).

Conflicts of interest: None.

Authorship: Pauline Dunne: Conceptualisation, writing – original draft, review, and editing. Deirbhile Sherry: writing, review, and editing. Sharleen O’ Reilly: Conceptualisation, writing, review and editing, supervision. All authors have been involved in revising the manuscript and approved the final version of the manuscript submitted for publication. All authors agree to be accountable for all aspects of the work.

References

1. Paulo MS, Abdo NM, Bettencourt-Silva R, et al. Gestational Diabetes Mellitus in Europe: A Systematic Review and Meta-Analysis of Prevalence Studies. *Front Endocrinol (Lausanne)*. 2021 Dec 1;12. <https://www.frontiersin.org/articles/10.3389/fendo.2021.691033/full> (accessed June 2024)
2. Crowe C, Noctor E, Carmody L, et al. ATLANTIC DIP: The prevalence of pre-diabetes/type 2 diabetes in an Irish population with gestational diabetes mellitus 1-5 years post index pregnancy. *BMC Proc*. 2012 Jul 9;6(4):O35. doi: 10.1186/1753-6561-6-S4-O35
3. Wang H, Li N, Chivese T, et al. IDF Diabetes Atlas: Estimation of Global and Regional Gestational Diabetes Mellitus Prevalence for 2021 by International Association of Diabetes in Pregnancy Study Group's Criteria. *Diabetes Res Clin Pract*. 2022 Jan 1;183:109050. doi: 10.1016/j.diabres.2021.109050
4. HSE Diabetes in Pregnancy Model of Care Working Group. Diabetes in Pregnancy: A Model of Care for Ireland. 2024 Feb. p. 68. <https://www.hse.ie/eng/about/who/cspd/ncps/diabetes/moc/diabetes-in-pregnancy-a-model-of-care-for-ireland.pdf> (accessed June 2024)
5. Ye W, Luo C, Huang J, Li C, et al. Gestational diabetes mellitus and adverse pregnancy outcomes: systematic review and meta-analysis. *BMJ*. 2022 May 25;377:e067946. doi: 10.1136/bmj-2021-067946
6. Metzger B, Lowe L, Dyer A, et al. Hyperglycemia and Adverse Pregnancy Outcomes. *N Engl J Med*. 2008;358(19):1991–2002. doi: 10.1056/NEJMoa0707943
7. Kramer CK, Campbell S, Retnakaran R. Gestational diabetes and the risk of cardiovascular disease in women: a systematic review and meta-analysis. *Diabetologia*. 2019 Jun 1;62(6):905–14. doi: 10.1007/s00125-019-4840-2
8. Vounzoulaki E, Khunti K, Abner SC, et al. Progression to type 2 diabetes in women with a known history of gestational diabetes: systematic review and meta-analysis. *BMJ*. 2020 May 13;369:m1361. doi: 10.1136/bmj.m1361
9. Phelan S. Pregnancy: A “teachable moment” for weight control and obesity prevention. *Am J Obstet Gynecol*. 2010 Feb;202(2):135. doi: 10.1016/j.ajog.2009.06.008

10. Makama M, Awoke MA, Skouteris H, et al. Barriers and facilitators to a healthy lifestyle in postpartum women: A systematic review of qualitative and quantitative studies in postpartum women and healthcare providers. *Obes Rev.* 2021;22(4):e13167. doi: 10.1111/obr.13167
11. Dasgupta K, Terkildsen Maindal H, Kragelund Nielsen K, et al. Achieving penetration and participation in diabetes after pregnancy prevention interventions following gestational diabetes: A health promotion challenge. *Diabetes Res Clin Pract.* 2018 Nov 1;145:200–13. doi: 10.1016/j.diabres.2018.04.016
12. American Diabetes Association Professional Practice Committee. 15. Management of Diabetes in Pregnancy: Standards of Care in Diabetes—2024. *Diabetes Care.* 2024 Jan;47(Suppl. 1):S282–94. doi: 10.2337/dc24-S015
13. National Institute for Health and Care Excellence. Diabetes in pregnancy: management from preconception to the postnatal period. NICE; 2020. <https://www.nice.org.uk/guidance/ng3/chapter/Recommendations> (accessed June 2024)
14. O'Reilly S, Janus E, Dunbar J. Diabetes prevention in high-risk women: many guidelines do not make light work 'Creation and Innovation: Guidelines in the Digital Age'. In: Proceedings of the 11th Guidelines International Network Conference. Melbourne, Australia; 2014. p. 72.
15. Haran C, van Driel M, Mitchell BL, et al. Clinical guidelines for postpartum women and infants in primary care—a systematic review. *BMC Pregnancy Childbirth.* 2014 Jan 29;14(1):51. doi: 10.1186/1471-2393-14-51
16. Aziz Z, Absetz P, Oldroyd J, et al. A systematic review of real-world diabetes prevention programs: learnings from the last 15 years. *Implementation Sci.* 2015 Dec 15;10(1):172. doi: 10.1186/s13012-015-0354-6
17. Wang MC, Shah NS, Petit LC, et al. Gestational Diabetes and Overweight/Obesity: Analysis of Nulliparous Women in the U.S., 2011–2019. *Am J Prev Med.* 2021 Dec;61(6):863–71. doi: 10.1016/j.amepre.2021.05.036
18. Damm P, Houshmand-Oeregaard A, et al. Gestational diabetes mellitus and long-term consequences for mother and offspring: a view from Denmark. *Diabetologia.* 2016 Jul 1;59(7):1396–9. doi: 10.1007/s00125-016-3985-5

19. Smyth S, Dale C, Tully E, et al. In pursuit of consensus – a national review of gestational diabetes. *Ir Med J.* 2023 May 8;116(1):714. https://repository.rcsi.com/articles/journal_contribution/In_pursuit_of_consensus_a_national_review_of_gestational_diabetes/22341097/1 (accessed June 2024)
20. Health Service Executive. Guidelines for the management of pre-gestational and gestational diabetes mellitus from preconception to the postnatal period. Office of the Nursing & Midwifery Services, Director Quality & Clinical Care Directorate, Health Service Executive, Dr Steevens' Hospital, Dublin 8, Ireland: Health Service Executive; 2010 Jul. Available from: <https://www.hse.ie/eng/services/list/2/primarycare/east-coast-diabetes-service/management-of-type-2-diabetes/diabetes-and-pregnancy/guidelines-for-the-management-of-pre-gestational-and-gestational-diabetes-mellitus-from-pre-conception-to-the-postnatal-period.pdf> (accessed June 2024)
21. Duarte-Gardea MO, Gonzales-Pacheco DM, Reader DM, et al. Academy of Nutrition and Dietetics Gestational Diabetes Evidence-Based Nutrition Practice Guideline. *J Acad Nutr Diet.* 2018 Sep;118(9):1719–42. doi: 10.1016/j.jand.2018.03.014
22. Laredo-Aguilera JA, Gallardo-Bravo M, Rabanales-Sotos JA, et al. Physical Activity Programs during Pregnancy Are Effective for the Control of Gestational Diabetes Mellitus. *Int J Environ Res Public Health.* 2020 Sep;17(17):6151. doi: 10.3390%2Fijerph17176151
23. Wilkinson SA, McCray S, Beckmann M, et al. Evaluation of a process of implementation of a gestational diabetes nutrition model of care into practice. *Nutr Diet.* 2016;73(4):329–35. doi: 10.1111/1747-0080.12233
24. Reader D, Splett P, Gunderson EP. Impact of Gestational Diabetes Mellitus Nutrition Practice Guidelines Implemented by Registered Dietitians on Pregnancy Outcomes. *J Am Diet Assoc.* 2006 Sep 1;106(9):1426–33. doi: 10.1016/j.jada.2006.06.009
25. Wilkinson SA, McCray SJ, Kempe A, et al. Clinically relevant improvements achieved from a facilitated implementation of a gestational diabetes model of care. *Nutr Diet.* 2018;75(3):271–82. doi: 10.1111/1747-0080.12404
26. Dunne P, Carvalho M, Byrne M, et al. Maternity Dietitians' Perspectives on Delivering Gestational Diabetes Care, and Postpartum Follow-Up in Ireland: A Qualitative Study. *J Acad Nutr Diet.* 2023 Dec 28; doi: 10.1016/j.jand.2023.12.012

27. Lawrence RL, Wall CR, Bloomfield FH, et al. Dietetic management of gestational diabetes in New Zealand: A cross-sectional survey. *Nutr Diet*. 2017;74(1):95–104. doi: 10.1111/1747-0080.12311
28. Raghavan R, Dreibelbis C, Kingshipp BL, et al. Dietary patterns before and during pregnancy and maternal outcomes: a systematic review. *Am J Clin Nutr*. 2019 Mar 1;109:705S-728S. doi: 10.1093/ajcn/nqy216
29. Lawrence RL, Ward K, Wall CR, et al. New Zealand women's experiences of managing gestational diabetes through diet: a qualitative study. *BMC Pregnancy Childbirth*. 2021 Dec 10;21(1):819. doi: 10.1186/s12884-021-04297-0
30. Wondmkun YT. Obesity, Insulin Resistance, and Type 2 Diabetes: Associations and Therapeutic Implications. *Diabetes Metab Syndr Obes*. 2020 Oct 9;13:3611–6. doi: 10.2147%2FDMSO.S275898
31. Martin AM, Berger H, Nisenbaum R, et al. Abdominal Visceral Adiposity in the First Trimester Predicts Glucose Intolerance in Later Pregnancy. *Diabetes Care*. 2009 Apr 23;32(7):1308–10. doi: 10.2337/dc09-0290
32. Chu SY, Callaghan WM, Kim SY, et al. Maternal Obesity and Risk of Gestational Diabetes Mellitus. *Diabetes Care*. 2007 Aug 1;30(8):2070–6. doi: 10.2337/dc06-2559a
33. Santos S, Voerman E, Amiano P, et al. Impact of maternal body mass index and gestational weight gain on pregnancy complications: an individual participant data meta-analysis of European, North American and Australian cohorts. *BJOG*. 2019;126(8):984–95. doi: 10.1111/1471-0528.15661
34. Kyrou I, Tsigos C, Mavrogianni C, et al. Sociodemographic and lifestyle-related risk factors for identifying vulnerable groups for type 2 diabetes: a narrative review with emphasis on data from Europe. *BMC Endocr Disord*. 2020 Mar 12;20(1):134. doi: 10.1186/s12902-019-0463-3
35. Kennelly MA, McAuliffe FM. Prediction and prevention of Gestational Diabetes: an update of recent literature. *Eur J Obstet Gynecol Reprod Biol*. 2016 Jul 1;202:92–8. doi: 10.1016/j.ejogrb.2016.03.032
36. Euro-Peristat Project. European Perinatal Health Report. Core indicators of the health and care of pregnant women and babies in Europe in 2015. 2018 p. 180.

<http://www.europeristat.com/index.php/reports/european-perinatal-health-report-2015.html>

(accessed June 2024)

37. Institute of Obstetricians and Gynaecologists, Royal College of Physicians of Ireland, Directorate of Clinical Strategy and Programmes, Health Service Executive. Nutrition During Pregnancy Clinical Practice Guideline. 2019. <https://www.hse.ie/eng/about/who/acute-hospitals-division/woman-infants/clinical-guidelines/national-clinical-guidelines-nutrition-in-pregnancy-guideline.pdf> (accessed June 2024)
38. Goldstein RF, Abell SK, Ranasinha S, et al. Association of Gestational Weight Gain With Maternal and Infant Outcomes. *JAMA*. 2017 Jun 6;317(21):2207–25. doi: 10.1001%2Fjama.2017.3635
39. Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. Weight Gain During Pregnancy: Reexamining the Guidelines [Internet]. Rasmussen KM, Yaktine AL, editors. Washington (DC): National Academies Press (US); 2009. <http://www.ncbi.nlm.nih.gov/books/NBK32813/> (accessed June 2024)
40. Farpour-Lambert NJ, Ells LJ, Martinez de Tejada B, et al. Obesity and Weight Gain in Pregnancy and Postpartum: an Evidence Review of Lifestyle Interventions to Inform Maternal and Child Health Policies. *Front Endocrinol (Lausanne)*. 2018 Sep 26;9:546. doi: 10.3389%2Ffendo.2018.00546
41. Endres LK, Straub H, McKinney C, et al. Postpartum Weight Retention Risk Factors and Relationship to Obesity at One Year. *Obstet Gynecol*. 2015 Jan;125(1):144–52. doi: 10.1097%2FAOG.0000000000000565
42. Makama M, Skouteris H, Moran LJ, et al. Reducing Postpartum Weight Retention: A Review of the Implementation Challenges of Postpartum Lifestyle Interventions. *J Clin Med*. 2021 Apr 27;10(9):1891. doi: 10.3390/jcm10091891
43. Sorbye L, Cnattingius S, Skjaerven R, et al. Interpregnancy weight change and recurrence of gestational diabetes mellitus: a population-based cohort study. *BJOG*. 2020;127(13):1608–16. doi: 10.1111/1471-0528.16364

44. Liu J, Song G, Meng T, Zhao G, Guo S. Weight retention at six weeks postpartum and the risk of gestational diabetes mellitus in a second pregnancy. *BMC Pregnancy Childbirth*. 2019 Aug 1;19(1):272. doi: 10.1186/s12884-019-2423-3
45. Timmermans YEG, van de Kant KDG, Oosterman EO, et al. The impact of interpregnancy weight change on perinatal outcomes in women and their children: A systematic review and meta-analysis. *Obes Rev*. 2020;21(3):e12974. doi: 10.1111/obr.12974
46. Kirkegaard H, Bliddal M, Støvring H, et al. Maternal weight change from prepregnancy to 18 months postpartum and subsequent risk of hypertension and cardiovascular disease in Danish women: A cohort study. *PLOS Medicine*. 2021 Apr 2;18(4):e1003486. doi: 10.1371/journal.pmed.1003486
47. Rooney BL, Schauburger CW. Excess pregnancy weight gain and long-term obesity: one decade later. *Obstet Gynecol*. 2002 Aug 1;100(2):245–52. doi: 10.1016/S0029-7844(02)02125-7
48. McAuliffe FM, Killeen SL, Jacob CM, et al. Management of prepregnancy, pregnancy, and postpartum obesity from the FIGO Pregnancy and Non-Communicable Diseases Committee: A FIGO (International Federation of Gynecology and Obstetrics) guideline. *Int J Gynaecol Obstet*. 2020 Sep;151(Suppl 1):16–36. doi: 10.1002%2Fijgo.13334
49. Haw JS, Galaviz KI, Straus AN, et al. Long-term Sustainability of Diabetes Prevention Approaches: A Systematic Review and Meta-analysis of Randomized Clinical Trials. *JAMA Intern Med*. 2017 Dec 1;177(12):1808. doi: 10.1001/jamainternmed.2017.6040
50. Warr W, Aveyard P, Albury C, et al. A systematic review and thematic synthesis of qualitative studies exploring GPs' and nurses' perspectives on discussing weight with patients with overweight and obesity in primary care. *Obes Rev*. 2021 Apr;22(4):e13151. doi: 10.1111%2Fobr.13151
51. Glenister KM, Malatzky CA, Wright J. Barriers to effective conversations regarding overweight and obesity in regional Victoria. 2017; <https://www.racgp.org.au/afp/2017/october/barriers-to-effective-conversations-regarding-over> (accessed June 2024)
52. Jarvie R. Lived experiences of women with co-existing BMI \geq 30 and Gestational Diabetes Mellitus. *Midwifery*. 2017 Jun 1;49:79–86. doi: 10.1016/j.midw.2016.12.009

53. Hedeager Mømsen AM, Høftoft D, Ørtenblad L, et al. Diabetes prevention interventions for women after gestational diabetes mellitus: an overview of reviews. *Endocrinol Diabetes Metab*. 2021;4(3):e00230. doi: 10.1002/edm2.230
54. Gunderson EP, Lewis CE, Lin Y, et al. Lactation Duration and Progression to Diabetes in Women Across the Childbearing Years: The 30-Year CARDIA Study. *JAMA Intern Med*. 2018 Mar 1;178(3):328–37. doi: 10.1001/jamainternmed.2017.7978
55. O'Reilly MW, Avalos G, Denny MC, et al. Atlantic DIP: high prevalence of abnormal glucose tolerance post partum is reduced by breast-feeding in women with prior gestational diabetes mellitus. *Eur J Endocrinol*. 2011 Dec;165(6):953–9. doi: 10.1530/EJE-11-0663
56. Pathirana MM, Ali A, Lassi ZS, et al. Protective Influence of Breastfeeding on Cardiovascular Risk Factors in Women With Previous Gestational Diabetes Mellitus and Their Children: A Systematic Review and Meta-Analysis. *J Hum Lact*. 2022 Aug 1;38(3):501–12. doi: 10.1177/08903344211034779
57. Tanase-Nakao K, Arata N, Kawasaki M, et al. Potential protective effect of lactation against incidence of type 2 diabetes mellitus in women with previous gestational diabetes mellitus: A systematic review and meta-analysis. *Diabetes Metab Res Rev*. 2017;33(4):e2875. doi: 10.1002/dmrr.2875
58. Feng L, Xu Q, Hu Z, et al. Lactation and progression to type 2 diabetes in patients with gestational diabetes mellitus: A systematic review and meta-analysis of cohort studies. *J Diabetes Investig*. 2018 Nov;9(6):1360–9. doi: 10.1111/1522-4049.12838
59. Morton S, Kirkwood S, Thangaratnam S. Interventions to modify the progression to type 2 diabetes mellitus in women with gestational diabetes: a systematic review of literature. *Curr Opin in Obstet and Gyn*. 2014 Dec;26(6):476. doi: 10.1097/GCO.0000000000000127
60. Victora CG, Bahl R, Barros AJD, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet*. 2016 Jan 30;387(10017):475–90. doi: 10.1016/S0140-6736(15)01024-7
61. Jagiello KP, Azulay Chertok IR. Women's Experiences With Early Breastfeeding After Gestational Diabetes. *J Obstet Gynecol Neonatal Nurs*. 2015;44(4):500–9. doi: 10.1111/1552-6909.12658

62. Payne J, Radcliffe B, Blank E, et al. Breastfeeding: the neglected guideline for future Dietitian-Nutritionists? *Nutr Diet.* 2007;64(2):93–8. doi: 10.1111/j.1747-0080.2007.00094.x
63. Becker GE, Quinlan G, Ward F, et al. Dietitians supporting breastfeeding: a survey of education, skills, knowledge and attitudes. *Ir J Med Sci.* 2021 May 1;190(2):711–22. doi: 10.1007/s11845-020-02384-3
64. Lucas HR, Williams RC, Hollar LN, et al. Understanding Gestational Diabetes, Future Diabetes Risk, and Diabetes Prevention: A Qualitative Study of Patient, Provider, and Staff Perspectives. *Clin Diabetes.* 2022 Jan 1;40(1):39–50. doi: 10.2337/cd21-0016
65. Lawson EJ, Rajaram S. A transformed pregnancy: the psychosocial consequences of gestational diabetes. *Sociol Health Illn.* 1994;16(4):536–62. doi: 10.1111/1467-9566.ep11347644
66. Davidsen E, Maindal HT, Rod MH, et al. The stigma associated with gestational diabetes mellitus: A scoping review. *eClinicalMedicine.* 2022 Oct 1;52:101614. doi: 10.1016/j.eclinm.2022.101614
67. Davidsen E, Terkildsen Maindal H, Byrne M, et al. A qualitative investigation into the perceptions and experiences of the stigma attached to gestational diabetes mellitus among women in Denmark. *Diabetes Res Clin Pract.* 2023 Sep;203:110858. doi: 10.1016/j.diabres.2023.110858
68. Hill B, Incollingo Rodriguez AC. Weight Stigma across the Preconception, Pregnancy, and Postpartum Periods: A Narrative Review and Conceptual Model. *Semin Reprod Med.* 2020 Nov;38(6):414–22. doi: 10.1055/s-0041-1723775
69. Hailu H, Skouteris H, Incollingo Rodriguez AC, et al. Drivers and facilitators of weight stigma among preconception, pregnant, and postpartum women: A systematic review. *Obes Rev.* 2024 May;25(5):e13710. doi: 10.1111/obr.13710
70. Zhang M, Zhou Y, Zhong J, et al. Current guidelines on the management of gestational diabetes mellitus: a content analysis and appraisal. *BMC Pregnancy Childbirth.* 2019 Jun 13;19(1):200. doi: 10.1186/s12884-019-2343-2
71. Tierney M, O’Dea A, Danyliv A, et al. Factors influencing lifestyle behaviours during and after a gestational diabetes mellitus pregnancy. *Health Psychol Behav Med.* 2015 Jan 1;3(1):204–16. doi: 10.1080/21642850.2015.1073111

72. McGovern A, Butler L, Jones S, et al. Diabetes screening after gestational diabetes in England: a quantitative retrospective cohort study. *Br J Gen Pract.* 2014 Jan 1;64(618):e17–23. doi: 10.3399/bjgp14X676410
73. Van Ryswyk E, Middleton P, Hague W, et al. Clinician views and knowledge regarding healthcare provision in the postpartum period for women with recent gestational diabetes: A systematic review of qualitative/survey studies. *Diabetes Res Clin Pract.* 2014 Dec 1;106(3):401–11. doi: 10.1016/j.diabres.2014.09.001
74. Van Ryswyk E, Middleton P, Shute E, et al. Women’s views and knowledge regarding healthcare seeking for gestational diabetes in the postpartum period: A systematic review of qualitative/survey studies. *Diabetes Res Clin Pract.* 2015 Nov 1;110(2):109–22. doi: 10.1016/j.diabres.2015.09.010
75. Dennison RA, Fox RA, Ward RJ, et al. Women’s views on screening for Type 2 diabetes after gestational diabetes: a systematic review, qualitative synthesis and recommendations for increasing uptake. *Diabet Med.* 2020;37(1):29–43. doi: 10.1111/dme.14081
76. Lithgow GE, Rossi J, Griffin SJ, et al. Barriers to postpartum diabetes screening: a qualitative synthesis of clinicians’ views. *Br J Gen Pract.* 2021 May 5;71(707):e473–82. doi: 10.3399%2FBJGP.2020.0928
77. Carson MP, Frank MI, Keely E. Original research: Postpartum testing rates among women with a history of gestational diabetes—Systematic review. *Prim Care Diabetes.* 2013 Oct 1;7(3):177–86. doi: 10.1016/j.pcd.2013.04.007
78. Jeppesen C, Kristensen JK, Ovesen P, et al. The forgotten risk? A systematic review of the effect of reminder systems for postpartum screening for type 2 diabetes in women with previous gestational diabetes. *BMC Res Notes.* 2015 Aug 26;8. doi: 10.1186%2Fs13104-015-1334-2
79. Lindström J, Tuomilehto J. The Diabetes Risk Score: A practical tool to predict type 2 diabetes risk. *Diabetes Care.* 2003 Mar 1;26(3):725–31. doi: 10.2337/diacare.26.3.725
80. Crowe C, Noctor E, Carmody L, et al. Validation of a diabetes risk score in identifying patients at risk of progression to abnormal glucose tolerance post partum. *BMC Proc.* 2012 Jul 9;6(Suppl 4):O36. doi: 10.1186%2F1753-6561-6-S4-O36

81. Dennison RA, Griffin SJ, Usher-Smith JA, et al. ‘Post-GDM support would be really good for mothers’; A qualitative interview study exploring how to support a healthy diet and physical activity after gestational diabetes. *PLoS ONE*. 2022 Jan 21;17(1):e0262852–e0262852. doi: 10.1371/journal.pone.0262852
82. Dunne P, Culliney L, O’Mahony L, et al. Exploring health professionals’ knowledge, practices and attitudes regarding gestational diabetes: A cross-sectional Irish national survey. *Diabet Med*. 2024 May 31: e15373. doi: 10.1111/dme.15373
83. Timm A, Nielsen KK, Christensen U, et al. Healthcare Professionals’ Perspectives on the Cross-Sectoral Treatment Pathway for Women with Gestational Diabetes during and after Pregnancy—A Qualitative Study. *J Clin Med*. 2021 Jan;10(4):843. doi: 10.3390/jcm10040843
84. Lake AJ, Neven ACH, Williams A, et al. Barriers to and enablers of type 2 diabetes screening among women with prior gestational diabetes: A systematic review update and qualitative synthesis applying the Theoretical Domains Framework. *Diabet Med*. 2022;39(4):e14772. doi: 10.1111/dme.14772
85. Lake AJ, Williams A, Neven ACH, et al. Barriers to and enablers of type 2 diabetes screening among women with prior gestational diabetes: A qualitative study applying the Theoretical Domains Framework. *Front Clin Diabetes Healthc*. 2023 Feb 24;4. doi: 10.3389/fcdhc.2023.1086186
86. Michie S, Atkins L, West R. The behaviour change wheel: a guide to designing interventions. United Kingdom: Silverback Publishing; 2014.
87. Cane J, O’Connor D, Michie S. Validation of the theoretical domains framework for use in behaviour change and implementation research. *Implement Science*. 2012 Apr 24;7(1):37. doi: 10.1186/1748-5908-7-37
88. Gillespie P, Cullinan J, O’Neill C, et al. Modeling the Independent Effects of Gestational Diabetes Mellitus on Maternity Care and Costs. *Diabetes Care*. 2013 May;36(5):1111–6. doi: 10.1111/j.1464-5491.2011.03293.x
89. Danyliv A, Gillespie P, O’Neill C, et al. Short- and long-term effects of gestational diabetes mellitus on healthcare cost: a cross-sectional comparative study in the ATLANTIC DIP cohort. *Diabet Med*. 2015;32(4):467–76. doi: 10.1111/dme.12678

90. Gillespie P, O'Neill C, Avalos G, et al. The cost of universal screening for gestational diabetes mellitus in Ireland. *Diabet Med*. 2011;28(8):912–8. doi: 10.2337/dc12-0461
91. O'Neill KN, McHugh SM, Tracey ML, et al. Health service utilization and related costs attributable to diabetes. *Diabet Med*. 2018;35(12):1727–34. doi: 10.1111/dme.13806
92. Werbrouck A, Schmidt M, Putman K, et al. A systematic review on costs and cost-effectiveness of screening and prevention of type 2 diabetes in women with prior gestational diabetes: Exploring uncharted territory. *Diabetes Res Clin Pract*. 2019 Jan 1;147:138–48. doi: 10.1016/j.diabres.2018.11.012
93. Lloyd M, Morton J, Teede H, et al. Long-term cost-effectiveness of implementing a lifestyle intervention during pregnancy to reduce the incidence of gestational diabetes and type 2 diabetes. *Diabetologia*. 2023 Jul 1 ;66(7):1223–34. doi: 10.1007/s00125-023-05897-5
94. Zhou X, Siegel KR, Ng BP, et al. Cost-effectiveness of Diabetes Prevention Interventions Targeting High-risk Individuals and Whole Populations: A Systematic Review. *Diabetes Care*. 2020 Jun 11 43(7):1593–616. doi: 10.2337/dci20-0018
95. Dyson PA, Twenefour D, Breen C, et al. Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. *Diabet Med*. 2018 May;35(5):541–7. doi: 10.1111/dme.13603
96. Tobias DK, Zhang C, Chavarro J, et al. Healthful dietary patterns and long-term weight change among women with a history of gestational diabetes mellitus. *Int J Obes (Lond)*. 2016 Nov 1;40(11):1748–1748. doi: 10.1038/ijo.2016.156
97. D'Arcy E, Rayner J, Hodge A, et al. The Role of Diet in the Prevention of Diabetes among Women with Prior Gestational Diabetes: A Systematic Review of Intervention and Observational Studies. *J Acad Nutr Diet*. 2020 Jan;120(1):69-85.e7. doi: 10.1016/j.jand.2019.07.021
98. Morrison MK, Koh D, Lowe JM, et al. Postpartum diet quality in Australian women following a gestational diabetes pregnancy. *Eur J Clin Nutr*. 2012 Oct;66(10):1160–5. doi: 10.1038/ejcn.2012.84
99. O'Reilly S, Versace V, Mohebbi M, et al. The effect of a diabetes prevention program on dietary quality in women with previous gestational diabetes. *BMC Womens Health*. 2019 Jul 3;19(1):88. doi: 10.1186/s12905-019-0788-0

100. Nicklas JM, Zera CA, Seely EW, et al. Identifying postpartum intervention approaches to prevent type 2 diabetes in women with a history of gestational diabetes. *BMC Pregnancy Childbirth*. 2011 Mar 24;11(1):23. doi: 10.1186/1471-2393-11-23
101. Teh K, Quek IP, Tang WE. Postpartum dietary and physical activity-related beliefs and behaviors among women with recent gestational diabetes mellitus: a qualitative study from Singapore. *BMC Pregnancy Childbirth*. 2021 Dec;21(1):1–12. doi: 10.1186/s12884-021-04089-6
102. Ratner RE, Christophi CA, Metzger BE, et al. Prevention of Diabetes in Women with a History of Gestational Diabetes: Effects of Metformin and Lifestyle Interventions. *J Clin Endocrinol Metab*. 2008 Dec 1;93(12):4774–9. doi: 10.1210/jc.2008-0772
103. Aroda VR, Christophi CA, Edelstein SL et al. The Effect of Lifestyle Intervention and Metformin on Preventing or Delaying Diabetes Among Women With and Without Gestational Diabetes: The Diabetes Prevention Program Outcomes Study 10-Year Follow-Up. *J Clin Endocrinol Metab*. 2015 Apr;100(4):1646–53. doi: 10.1210/jc.2014-3761
104. American Diabetes Association. Prevention or Delay of Type 2 Diabetes: Position Statement. *Diabetes Care*. 2004 Jan 1;27(suppl 1):s47–s47. doi: 10.2337/diacare.27.2007.S47
105. Pedersen ALW, Terkildsen Maindal H, Juul L. How to prevent type 2 diabetes in women with previous gestational diabetes? A systematic review of behavioural interventions. *Prim Care Diabetes*. 2017 Oct 1;11(5):403–13. doi: 10.1016/j.pcd.2017.05.002
106. Maindal HT, Timm A, Dahl-Petersen IK, et al. Systematically developing a family-based health promotion intervention for women with prior gestational diabetes based on evidence, theory and co-production: the Face-it study. *BMC Public Health*. 2021 Sep 3;21:1616. doi: 10.1186/s12889-021-11655-2
107. Torenholt R, Schwennesen N, Willaing I. Lost in translation—the role of family in interventions among adults with diabetes: a systematic review. *Diabet Med*. 2014;31(1):15–23. doi: 10.1111/dme.12290
108. Peacock AS, Bogossian F, McIntyre HD, et al. A review of interventions to prevent Type 2 Diabetes after Gestational Diabetes. *Women Birth*. 2014 Dec 1;27(4):e7–15. doi: 10.1016/j.wombi.2014.09.002

109. Retnakaran M, Viana LV, Kramer CK. Lifestyle intervention for the prevention of type 2 diabetes in women with prior gestational diabetes: A systematic review and meta-analysis. *Diabetes Obes Metab.* 2023 May;25(5):1196–202. doi: 10.1111/dom.14966
110. Uusitupa M, Khan TA, Vigiliouk E, et al. Prevention of Type 2 Diabetes by Lifestyle Changes: A Systematic Review and Meta-Analysis. *Nutrients.* 2019 Nov 1;11(11):2611. doi: 10.3390%2Fnu11112611
111. Dennison RA, Oliver-Williams C, Qi HLJ, et al. The effectiveness of pharmacological and lifestyle interventions to reduce the risk of diabetes and hyperglycaemia following gestational diabetes: A systematic review and meta-analysis. *Diabet Med.* 2024 Jun;41(6):e15316. doi: 10.1111/dme.15316
112. Sagastume D, Siero I, Mertens E, et al. The effectiveness of lifestyle interventions on type 2 diabetes and gestational diabetes incidence and cardiometabolic outcomes: A systematic review and meta-analysis of evidence from low- and middle-income countries. *eClinicalMedicine.* 2022 Nov;53. doi: 10.1016%2Fj.eclinm.2022.101650
113. Suarez-Lledo V, Alvarez-Galvez J. Prevalence of Health Misinformation on Social Media: Systematic Review. *J Med Internet Res.* 2021 Jan 20;23(1):e17187. doi: 10.2196/17187
114. Cheng Z, Hao H, Tsofliou F, et al. Effects of online support and social media communities on gestational diabetes: A systematic review. *Int J Med Inform.* 2023 Dec;180:105263. doi: 10.1016/j.ijmedinf.2023.105263
115. Dunne P, O'Reilly S. Exploring the lived experience and views of women with gestational diabetes in Ireland: a cross-sectional national survey. *Proc Nut Soc.* 2023 Jan;82(OCE5):E296. doi: 10.1017/S0029665123003956
116. Lim S, Tan A, Madden S, et al. Health Professionals' and Postpartum Women's Perspectives on Digital Health Interventions for Lifestyle Management in the Postpartum Period: A Systematic Review of Qualitative Studies. *Front Endo (Lausanne).* 2019;10:767. doi: 10.3389/fendo.2019.00767
117. Negron R, Martin A, Almog M, et al. Social support during the postpartum period: Mothers' views on needs, expectations, and mobilization of support. *Matern Child Health J.* 2013 May;17(4):616–23. doi: 10.1007%2Fs10995-012-1037-4

118. Parsons J, Sparrow K, Ismail K, et al. A qualitative study exploring women's health behaviours after a pregnancy with gestational diabetes to inform the development of a diabetes prevention strategy. *Diabet Med*. 2019;36(2):203–13. doi: 10.1111/dme.13794
119. Gilinsky AS, Kirk AF, Hughes AR, et al. Lifestyle interventions for type 2 diabetes prevention in women with prior gestational diabetes: A systematic review and meta-analysis of behavioural, anthropometric and metabolic outcomes. *Prev Med Rep*. 2015 Jan 1;2:448–61. doi: 10.1016/j.pmedr.2015.05.009
120. Gray KL, McKellar L, O'Reilly SL, et al. Women's Barriers to Weight Loss, Perception of Future Diabetes Risk and Opinions of Diet Strategies Following Gestational Diabetes: An Online Survey. *Int J Environ Res Public Health*. 2020 Dec;17(24):9180. doi: 10.3390%2Fijerph17249180
121. Lim S, O'Reilly S, Behrens H, Skinner T, et al. Effective strategies for weight loss in post-partum women: a systematic review and meta-analysis. *Obes Rev*. 2015;16(11):972–87. doi: 10.1111/obr.12312
122. Jones EJ, Fraley HE, Mazzawi J. Appreciating Recent Motherhood and Culture: A Systematic Review of Multimodal Postpartum Lifestyle Interventions to Reduce Diabetes Risk in Women with Prior Gestational Diabetes. *Matern Child Health J*. 2017 Jan 1;21(1):45–57. doi: 10.1007/s10995-016-2092-z
123. Christie HE, Roach LA, Kennedy M, et al. 'Beyond the Bump': an online wellbeing and lifestyle pilot program during COVID-19 for first year postpartum mothers: a research article. *BMC Pregnancy Childbirth*. 2022 Jul 25;22:591. doi: 10.1186%2Fs12884-022-04913-7
124. Kriska AM, Rockette-Wagner B, Edelstein SL, et al. The Impact of Physical Activity on the Prevention of Type 2 Diabetes: Evidence and Lessons Learned From the Diabetes Prevention Program, a Long-Standing Clinical Trial Incorporating Subjective and Objective Activity Measures. *Diabetes Care*. 2021 Jan;44(1):43–9. doi: 10.2337%2Fdc20-1129
125. Ioannou E, Humphreys H, Homer C, et al. A systematic review and thematic synthesis of the barriers and facilitators to physical activity for women after gestational diabetes: a socio-ecological approach. *Br J Diabetes*. 2023 Jun 28;23(1):2–13. doi: 10.15277/bjd.2023.413

126. Ioannou E, Humphreys H, Homer C, et al. Beyond the individual: Socio-ecological factors impacting activity after gestational diabetes mellitus. *Diabet Med.* 2024;41(6):e15286. doi: 10.1111/dme.15286
127. Lagisetty PA, Priyadarshini S, Terrell S, et al. Culturally Targeted Strategies for Diabetes Prevention in Minority Populations: A Systematic Review and Framework. *Diabetes Educ.* 2017 Feb;43(1):54–77. doi: 10.1177/0145721716683811
128. Hod M, Kapur A, Sacks DA, et al. The International Federation of Gynecology and Obstetrics (FIGO) Initiative on gestational diabetes mellitus: A pragmatic guide for diagnosis, management, and care. *Int J Gynaecol Obstet.* 2015 Oct 1;131:S173–211. doi: 10.1016/S0020-7292(15)30033-3
129. Tsirou E, Grammatikopoulou MG, Theodoridis X, et al. Guidelines for Medical Nutrition Therapy in Gestational Diabetes Mellitus: Systematic Review and Critical Appraisal. *J Acad Nutr Diet.* 2019 Aug 1;119(8):1320–39. doi: 10.1016/j.jand.2019.04.002
130. Mustafa ST, Hofer OJ, Harding JE, et al. Dietary recommendations for women with gestational diabetes mellitus: a systematic review of clinical practice guidelines. *Nutr Rev.* 2021 Sep 1;79(9):988–1021. doi: 10.1093/nutrit/nuab005
131. Wilkinson SA, O'Brien M, McCray S, et al. Implementing a best-practice model of gestational diabetes mellitus care in dietetics: a qualitative study. *BMC Health Serv Res.* 2019 Feb 14;19(1):122. doi: 10.1186/s12913-019-3947-y
132. Ingstrup MS, Wozniak LA, Mathe N, et al. Women's experience with peer counselling and social support during a lifestyle intervention among women with a previous gestational diabetes pregnancy. *Health Psychol Behav Med.* 2019 Jan 1;7(1):147–59. doi: 10.1080/21642850.2019.1612750
133. Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ.* 2021 Sep 30;374:n2061. doi: 10.1136/bmj.n2061
134. Adam S, McIntyre HD, Tsoi KY, et al. Pregnancy as an opportunity to prevent type 2 diabetes mellitus: FIGO Best Practice Advice. *Int J Gynaecol Obstet.* 2023;160(S1):56–67. doi: 10.1002/ijgo.14537

135. Neven ACH, Lake AJ, Williams A, et al. Barriers to and enablers of postpartum health behaviours among women from diverse cultural backgrounds with prior gestational diabetes: A systematic review and qualitative synthesis applying the theoretical domains framework. *Diabet Med.* 2022;39(11):e14945. doi: 10.1111/dme.14945
136. Murofushi K, Badaracco C, County C, et al. Implement Sci in Evidence-based Nutrition Practice: Considerations for the Registered Dietitian Nutritionist. *J Acad Nutr Diet.* 2021 Jul 1;121(7):1392–400. doi: 10.1016/j.jand.2020.08.093
137. Young AM, Hickman I, Campbell K, et al. Implementation science for dietitians: The ‘what, why and how’ using multiple case studies. *Nutr Diet.* 2021 Jul;78(3):276–85. doi: 10.1111/1747-0080.12677
138. Craig P, Dieppe P, Macintyre S, et al. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ.* 2008 Sep 29;337:a1655. doi: 10.1136/bmj.a1655
139. Datta J, Petticrew M. Challenges to evaluating complex interventions: a content analysis of published papers. *BMC Public Health.* 2013 Dec;13(1):1–18. doi: 10.1186/1471-2458-13-568
140. Troughton J, Chatterjee S, Hill SE, et al. Development of a lifestyle intervention using the MRC framework for diabetes prevention in people with impaired glucose regulation. *J Public Health (Oxf).* 2016 Sep;38(3):493–501. doi: 10.1093/pubmed/fdv110
141. Curran GM, Bauer M, Mittman B, et al. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care.* 2012 Mar;50(3):217–26. doi: 10.1097/MLR.0b013e3182408812
142. Petticrew M. When are complex interventions ‘complex’? When are simple interventions ‘simple’? *Eur J Public Health.* 2011 Aug 1;21(4):397–8. doi: 10.1093/eurpub/ckr084
143. Baugh Littlejohns L, Near E, McKee G, et al. A scoping review of complex systems methods used in population physical activity research: do they align with attributes of a whole system approach? *Health Res Policy Syst.* 2023 Mar 2;21(1):18. doi: 10.1186/s12961-023-00961-3

144. Wutzke S, Morrice E, Benton M, et al. Systems approaches for chronic disease prevention: sound logic and empirical evidence, but is this view shared outside of academia? *Public Health Res Pract.* 2016 15;26(3). doi: 10.17061/phrp2631632
145. Damschroder LJ, Aron DC, Keith RE, et al. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci.* 2009 Dec;4(1):1–15. doi: 10.1186/1748-5908-4-50
146. Bernstein JA, McCloskey L, Gebel CM, et al. Lost opportunities to prevent early onset type 2 diabetes mellitus after a pregnancy complicated by gestational diabetes. *BMJ Open Diabetes Res Care.* 2016 Jun 1;4(1):e000250. doi: 10.1136/bmjdr-2016-000250
147. Madrigal L, Manders OC, Kegler M, et al. Inner and outer setting factors that influence the implementation of the National Diabetes Prevention Program (National DPP) using the Consolidated Framework for Implementation Research (CFIR): a qualitative study. *Implement Sci Commun.* 2022 Oct 1;3(1):104. doi: 10.1186/s43058-022-00350-x
148. Nilsen P. Making sense of implementation theories, models and frameworks. *Implement Science.* 2015 Apr 21;10(1):53. doi: 10.1186/s13012-015-0242-0
149. Aarons GA, Hurlburt M, Horwitz SM. Advancing a Conceptual Model of Evidence-Based Practice Implementation in Public Service Sectors. *Adm Policy Ment Health.* 2011 Jan;38(1):4–23. doi: 10.1007/s10488-010-0327-7
150. May CR, Cummings A, Girling M, et al. Using Normalization Process Theory in feasibility studies and process evaluations of complex healthcare interventions: a systematic review. *Implement Science.* 2018 Jun 7;13(1):80. doi: 10.1186/s13012-018-0758-1
151. Moullin JC, Dickson KS, Stadnick NA, et al. Systematic review of the Exploration, Preparation, Implementation, Sustainment (EPIS) framework. *Implement Science.* 2019 Jan 1;14(1):1–16. doi: 10.1186/s13012-018-0842-6
152. O'Reilly SL, Laws R, Maindal HT, et al. A Complex mHealth Coaching Intervention to Prevent Overweight, Obesity, and Diabetes in High-Risk Women in Antenatal Care: Protocol for a Hybrid Type 2 Effectiveness-Implementation Study. *JMIR Res Protoc.* 2023 Sep 18;12(1):e51431. doi: 10.2196/51431
153. May C. Towards a general theory of implementation. *Implement Science.* 2013 Feb 13;8(1):18. doi: 10.1186/1748-5908-8-18

154. May CR, Finch T, Ballini L, et al. Evaluating complex interventions and health technologies using normalization process theory: development of a simplified approach and web-enabled toolkit. *BMC Health Serv Res*. 2011 Sep 30;11(1):245. doi: 10.1186/1472-6963-11-245
155. Finch TL, Rapley T, Girling M, et al. Improving the normalization of complex interventions: measure development based on normalization process theory (NoMAD): study protocol. *Implement Science*. 2013 Apr 11;8(1):43. doi: 10.1186/1748-5908-8-43
156. O'Reilly SL, May CR, Ford D, et al, the MAGDA Study Group. Implementing Prim Care Diabetes prevention for women with previous gestational diabetes: a mixed-methods study. *Fam Pract*. 2022 Apr 12;cmac022. doi: 10.1093/fampra/cmac022
157. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implement Science*. 2011 Apr 23;6(1):42. doi: 10.1186/1748-5908-6-42
158. Michie S, Ashford S, Sniehotta FF, et al. A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating behaviours: the CALO-RE taxonomy. *Psychol Health*. 2011 Nov;26(11):1479–98. doi: 10.1080/08870446.2010.540664
159. Michie S, Richardson M, Johnston M, et al. The Behavior Change Technique Taxonomy (v1) of 93 Hierarchically Clustered Techniques: Building an International Consensus for the Reporting of Behavior Change Interventions. *Ann Behav Med*. 2013 Aug 1;46(1):81–95. doi: 10.1007/s12160-013-9486-6
160. Glasgow RE. Evaluation of theory-based interventions: the RE-AIM model. Health behavior and health education [K Glanz, FM Lewis and BK Rimer, editors]. San Francisco: Jossey-Bass.
161. Glasgow RE, Vogt TM, Boles SM. Evaluating the public health impact of health promotion interventions: the RE-AIM framework. *Am J Public Health*. 1999 Sep;89(9):1322–7. doi: 10.2105/AJPH.89.9.1322
162. Glasgow RE, Harden SM, Gaglio B, et al. RE-AIM Planning and Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. *Front Public Health*. 2019 Mar 29;7:64. doi: 10.3389/fpubh.2019.00064

163. Aziz Z, Mathews E, Absetz P, et al. A group-based lifestyle intervention for diabetes prevention in low- and middle-income country: implementation evaluation of the Kerala Diabetes Prevention Program. *Implement Sci.* 2018 Dec;13(1):97. doi: 10.1186/s13012-018-0791-0
164. Lim S, Chen M, Makama M, et al. Preventing Type 2 Diabetes in Women with Previous Gestational Diabetes: Reviewing the Implementation Gaps for Health Behavior Change Programs. *Semin Reprod Med.* 2020 Nov;38(6):377–83. doi: 10.1055/s-0040-1722315
165. Knight AW, Ford D, Audehm R, et al. The Australian Primary Care Collaboratives Program: improving diabetes care. *BMJ Qual Saf.* 2012 Nov;21(11):956–63. doi: 10.1136/bmjqs-2011-000460
166. Vesco KK, Dietz PM, Bulkley J, et al. A system-based intervention to improve postpartum diabetes screening among women with gestational diabetes. *Am J Obstet Gynecol.* 2012 Oct 1;207(4):283.e1-283.e6. doi: 10.1016/j.ajog.2012.08.017
167. O'Reilly SL, Dunbar JA, Best JD, et al. GooD4Mum: A general practice-based quality improvement collaborative for diabetes prevention in women with previous gestational diabetes. *Prim Care Diabetes.* 2019 Apr 1;13(2):134–41. doi: 10.1016/j.pcd.2018.10.006
168. de la Perrelle L, Radisic G, Cations M, et al. Costs and economic evaluations of Quality Improvement Collaboratives in healthcare: a systematic review. *BMC Health Serv Res.* 2020 Mar 2;20(1):155. doi: 10.1186/s12913-020-4981-5

Table 1. Risk factors for developing GDM. NICE, 2020

Gestational Diabetes risk factors

- BMI above 30 kg/m²
- Previous gestational diabetes
- Previous macrosomic baby weighing 4.5 kg or more
- Family history of diabetes (first-degree relative with diabetes)
- An ethnicity with a high prevalence of diabetes.

Table 2. Improving awareness and reducing future diabetes risk after gestational diabetes mellitus (GDM) *Adapted from Marschner et al., 2023*

Raise awareness of level of magnitude of health risk	Optimising management activities	Potential outcomes
<p>GDM is:</p> <ul style="list-style-type: none"> • 6 times more likely in women with obesity 	<p>Multidisciplinary input:</p> <ul style="list-style-type: none"> - Endocrinologist - Obstetrician - Diabetes in Pregnancy Midwifery/Nursing Staff - Dietitian - General practitioner - Practice nurse - Access to mental health care - Infant feeding support 	<p>Increased awareness of risk and its significance</p> <p>Monitoring of T2D risk factors</p>
<p>GDM confers:</p> <ul style="list-style-type: none"> • Seven to tenfold increased risk of future type 2 diabetes • Twofold increased risk of cardiovascular disease • Twice as likely to have GDM in subsequent pregnancy 	<p>Consistent messaging and information from healthcare professionals</p> <p>Structured care pathway after GDM to include risk assessment and follow up</p> <p>Diet and physical activity intervention tailored to women's needs after GDM, considering:</p> <ul style="list-style-type: none"> - Social support - Timing - Education - Ethnicity - Language - Learning styles - Mode of delivery. 	<p>Engagement with prevention programmes and active diet and physical activity</p> <p style="text-align: center;">↓</p> <p>Reduced risk of T2D and cardiovascular disease</p>

GDM: gestational diabetes, T2D: type 2 diabetes.

Table 3. TDF domains and COM-B components linked to perceived barriers and enablers of type 2 diabetes screening among women with prior gestational diabetes.*Adapted from Lake et al., 2023*

COM-B component	TDF domain	Barriers and Enablers
COM-B: Capability (Psychological)	Knowledge	Low awareness of magnitude of risk for T2D (barrier)
	Memory, attention and decision processes	Receiving reminders from hospital or GP (enabler)
COM-B: Opportunity (Physical, Social)	Environmental context and resources	Lack of time, competing demands (barrier) Prevention programme (enabler)
	Social influence	Lack of childcare (barrier) Family supports (enabler)
COM-B: Motivation (Automatic, Reflective)	Emotion	Concern for future health (barrier) Reassurance from regular screening (enabler)
	Beliefs about consequences	Consequences of screening (perceived as a barrier and an enabler)
	Social role and identity	Low prioritisation of own health (barrier) Role modelling health promoting behaviours for the family (enabler)
	Beliefs about capabilities	Confidence in ability to manage own health (enabler)

COM-B: Capability, Opportunity, Motivation-Behaviour model, TDF: Theoretical Domains Framework, T2D: type 2 diabetes, GP: general practitioner.