

ED annual census of 10,000. These EDs were evaluated to compare the effects of health care reform on different populations. Admissions data were collected from both hospitals during the period of March 2009 and September 2013. In order to assess influence of the ACA on admission rates, a multi-variable binary logistic regression model was computed.

Results:

Time Period	Response	Suburban Hospital	Urban Hospital
Pre-ACA	Admit	17 (0.16%)	21,067 (23.61%)
	Non-Admit	10,321 (99.84%)	68,145 (76.39%)
Post-ACA	Admit	39 (0.12%)	68,654 (22.41%)
	Non-Admit	32,536 (99.88%)	237,736 (77.59%)

Table 1. Summary of admission rates by hospital and period. For patients visiting the suburban hospital, those who visited after the ACA came into effect had a (nonsignificant) 27% (OR: 0.73; 95% CI, 0.42–1.32; P value = .275) reduced likelihood of being admitted as compared to before the ACA came into effect. Finally, there was a (non-significant) 28.4% (OR: 1.28; 95% CI, 0.71–2.23; P value = .391) increase in the odds ratio of admitting post-ACA for the urban hospital as compared to post-ACA for the suburban hospital.

Conclusion: There was not a significant change in admissions rates post-ACA implementation for either hospital. Total annual census at both sites decreased the year post-ACA. While there has been a change in ED utilization in the short-term, it may be too early to see the long-term impact of the ACA.

Prehosp Disaster Med 2017;32(Suppl. 1):s36–s37
doi:10.1017/S1049023X17001133

Hemin Microglia NLRP3 Protein N-methyl-D-aspartic Acid Receptor 1 Inflammasome

Lei Li

Research Institute Of Surgery, Third Military Medical University, Chongqing/China

Study/Objective: In this study, we aim to test the hypothesis that NLRP3 signaling pathway takes a vital position in ICH-induced secondary inflammatory damage, and detect the role of N-methyl-D-aspartic acid receptor 1 (NMDAR1) in this progress.

Background: Inflammation plays a critical role in secondary brain damage after intracerebral hemorrhage (ICH). However, the mechanisms of inflammatory injury following ICH are still unclear, particularly the involvement of NLRP3 inflammasome, which are crucial to sterile inflammatory responses.

Methods: ICH was induced in mice by microinjection of hemin into the striatum. The protein levels of NMDAR1, NMDAR1 phosphorylation, NLRP3 and IL-1b were measured by Western blot. The binding of NMDAR1 to NLRP3 was detected by immunoprecipitation.

Results: The expression of NMDAR1, NMDAR1 phosphorylation, NLRP3 and IL-1b were rapidly increased after ICH. Hemin treatment enhanced NMDAR1 expression and NMDAR1 phosphorylation, as well in cultured microglial cells treated by hemin. Hemin up-regulated NLRP3 and IL-1b level, which was reversed by MK801 (NMDAR antagonist) in vitro. Hemin also promoted the binding of NMDAR1 to NLRP3.

Conclusion: Our findings suggest that NMDAR1 plays a pivotal role in hemin-induced NLRP3-mediated inflammatory damage through synergistic activation.

Prehosp Disaster Med 2017;32(Suppl. 1):s37
doi:10.1017/S1049023X17001145

Improving Efficiencies with Real Time Locating Systems in Emergency Departments

Gary M. Mann

Emergency Medicine Program, Rouge Valley Health System-Ajax and Pickering Health Centre, Thornhill/ON/Canada

Study/Objective: To demonstrate improved efficiencies in Emergency Department patient flow and performance metrics using Real-Time Locating Systems.

Background: Efficiency is key in managing ED overcrowding. Utilizing a Real-Time Locating System (RTLS), our ED effectively manages ever-increasing volumes and acuity of emergency patients, including Mass-Casualty Event (MCE) victims.

Methods: RTLS systems are fully automated; they continually monitor locations of patients, staff, and assets, giving instantaneous visual status, measuring key metrics, and reporting key performance indicators. In addition to displaying patients' locations, RTLS measures and displays milestones based on time, location, acuity, and interaction; these are automated via customizable, intuitive software and low maintenance hardware. RTLS allows to identify treatment, space status, and availability, management of waiting room times, wait time to the physician's initial assessment (PIA), and overall length of stay (OLOS). Key assets are easily located when needed. Enhanced RTLS safety features include ability to track staff contacts with Infectious patients, as well as button-press alerts when staff need help urgently. The system displays data in different formats, including overhead floor-plan views displaying treatment space status, patient name, staff locations, and patient assignments. List views give information concerning complaint, acuity, disposition, location, and length of stay. Color flags provide visual warnings when approaching mandated targets for PIA and OLOS. This system is particularly useful in MCE's, where victims can rapidly be tagged and assigned triage levels.

Results: Despite the annual volume of ED patient growth of 35% since introduction of RTLS (in addition to other efficiencies), PIA time dropped by 85%. Patient and staff satisfaction improved dramatically. RTLS has been shown to be a useful modality in mass-casualty events.

Conclusion: RTLS is a low-cost, highly effective solution to help manage ED overcrowding by increasing efficiency, safety, and satisfaction among ED patients and staff by continuous